

# The Enzyme List

## Class 3 — Hydrolases

Nomenclature Committee  
of the  
International Union of Biochemistry and Molecular Biology  
(NC-IUBMB)

L<sup>A</sup>T<sub>E</sub>X version prepared by Andrew McDonald,  
School of Biochemistry and Immunology, Trinity College Dublin, Ireland

Generated from the [ExplorEnz](#) database, March 2019

© 2019 IUBMB

### Contents

<b>EC 3.1 Acting on ester bonds</b>	<b>3</b>
EC 3.1.1 Carboxylic-ester hydrolases	3
EC 3.1.2 Thioester hydrolases	25
EC 3.1.3 Phosphoric-monoester hydrolases	31
EC 3.1.4 Phosphoric-diester hydrolases	54
EC 3.1.5 Triphosphoric-monoester hydrolases	63
EC 3.1.6 Sulfuric-ester hydrolases	64
EC 3.1.7 Diphosphoric-monoester hydrolases	68
EC 3.1.8 Phosphoric-triester hydrolases	70
EC 3.1.11 Exodeoxyribonucleases producing 5'-phosphomonoesters	71
EC 3.1.12 Exodeoxyribonucleases producing 3'-phosphomonoesters	73
EC 3.1.13 Exoribonucleases producing 5'-phosphomonoesters	73
EC 3.1.14 Exoribonucleases producing 3'-phosphomonoesters	74
EC 3.1.15 Exonucleases that are active with either ribo- or deoxyribonucleic acids and produce 5'-phosphomonoesters	75
EC 3.1.16 Exonucleases that are active with either ribo- or deoxyribonucleic acids and produce 3'-phosphomonoesters	75
EC 3.1.21 Endodeoxyribonucleases producing 5'-phosphomonoesters	75
EC 3.1.22 Endodeoxyribonucleases producing 3'-phosphomonoesters	78
EC 3.1.25 Site-specific endodeoxyribonucleases that are specific for altered bases	85
EC 3.1.26 Endoribonucleases producing 5'-phosphomonoesters	86
EC 3.1.27 Endoribonucleases producing 3'-phosphomonoesters	89
EC 3.1.30 Endoribonucleases that are active with either ribo- or deoxyribonucleic acids and produce 5'-phosphomonoesters	90
EC 3.1.31 Endoribonucleases that are active with either ribo- or deoxyribonucleic acids and produce 3'-phosphomonoesters	90
<b>EC 3.2 Glycosylases</b>	<b>91</b>
EC 3.2.1 Glycosidases, i.e. enzymes that hydrolyse <i>O</i> - and <i>S</i> -glycosyl compounds	91
EC 3.2.2 Hydrolysing <i>N</i> -glycosyl compounds	138
<b>EC 3.3 Acting on ether bonds</b>	<b>145</b>
EC 3.3.1 Thioether and trialkylsulfonium hydrolases	145
EC 3.3.2 Ether hydrolases	146

<b>EC 3.4 Acting on peptide bonds (peptidases)</b>	<b>150</b>
EC 3.4.11 Aminopeptidases	153
EC 3.4.13 Dipeptidases	159
EC 3.4.14 Dipeptidyl-peptidases and tripeptidyl-peptidases	162
EC 3.4.15 Peptidyl-dipeptidases	165
EC 3.4.16 Serine-type carboxypeptidases	166
EC 3.4.17 Metallocoarboxypeptidases	167
EC 3.4.18 Cysteine-type carboxypeptidases	172
EC 3.4.19 Omega peptidases	172
EC 3.4.21 Serine endopeptidases	176
EC 3.4.22 Cysteine endopeptidases	202
EC 3.4.23 Aspartic endopeptidases	218
EC 3.4.24 Metalloendopeptidases	228
EC 3.4.25 Threonine endopeptidases	248
EC 3.4.99 Endopeptidases of unknown catalytic mechanism (sub-subclass is currently empty)	248
<b>EC 3.5 Acting on carbon-nitrogen bonds, other than peptide bonds</b>	<b>251</b>
EC 3.5.1 In linear amides	251
EC 3.5.2 In cyclic amides	279
EC 3.5.3 In linear amidines	283
EC 3.5.4 In cyclic amidines	288
EC 3.5.5 In nitriles	298
EC 3.5.99 In other compounds	300
<b>EC 3.6 Acting on acid anhydrides</b>	<b>302</b>
EC 3.6.1 In phosphorus-containing anhydrides	302
EC 3.6.2 In sulfonyl-containing anhydrides	315
EC 3.6.3 Acting on acid anhydrides to catalyse transmembrane movement of substances	316
EC 3.6.4 Acting on acid anhydrides to facilitate cellular and subcellular movement	320
EC 3.6.5 Acting on GTP to facilitate cellular and subcellular movement	322
<b>EC 3.7 Acting on carbon-carbon bonds</b>	<b>323</b>
EC 3.7.1 In ketonic substances	323
<b>EC 3.8 Acting on halide bonds</b>	<b>328</b>
EC 3.8.1 In carbon-halide compounds	328
<b>EC 3.9 Acting on phosphorus-nitrogen bonds</b>	<b>331</b>
EC 3.9.1 Acting on phosphorus-nitrogen bonds (only sub-subclass identified to date)	331
<b>EC 3.10 Acting on sulfur-nitrogen bonds</b>	<b>332</b>
EC 3.10.1 Acting on sulfur-nitrogen bonds (only sub-subclass identified to date)	332
<b>EC 3.11 Acting on carbon-phosphorus bonds</b>	<b>332</b>
EC 3.11.1 Acting on carbon-phosphorus bonds (only sub-subclass identified to date)	332
<b>EC 3.12 Acting on sulfur-sulfur bonds</b>	<b>333</b>
EC 3.12.1 Acting on sulfur-sulfur bonds (only sub-subclass identified to date)	333
<b>EC 3.13 Acting on carbon-sulfur bonds</b>	<b>333</b>
EC 3.13.1 Acting on carbon-sulfur bonds (only sub-subclass identified to date)	333
<b>References</b>	<b>336</b>
<b>Index</b>	<b>502</b>

## EC 3.1 Acting on ester bonds

This subclass contains the esterase enzymes. The esterases are subdivided into: carboxylic-ester hydrolases (EC 3.1.1), thioester hydrolases (EC 3.1.2), phosphoric-monoester hydrolases, the phosphatases (EC 3.1.3), phosphoric-diester hydrolases (EC 3.1.4), triphosphoric-monoester hydrolases (EC 3.1.5), sulfuric-ester hydrolases, the sulfatases (EC 3.1.6), diphosphoric monoesterases (EC 3.1.7) and phosphoric-triester hydrolases (EC 3.1.8). The nucleases, previously included under EC 3.1.4, are now placed in a number of new sub-subclasses: the exonucleases (EC 3.1.11-16) and the endonucleases (EC 3.1.21-31).; EC 3.1.23 and EC 3.1.24

In a previous edition, site-specific endodeoxyribonucleases were set out individually in subclasses EC 3.1.23 and EC 3.1.24 (since deleted), with 113 separate entries. These are now included in three entries EC 3.1.21.3, EC 3.1.21.4 and EC 3.1.21.5. A complete listing of all of these enzymes has been produced by R.J. Roberts and is available at <http://rebase.neb.com/rebase/rebase.html>.

### EC 3.1.1 Carboxylic-ester hydrolases

#### EC 3.1.1.1

- Accepted name:** carboxylesterase  
**Reaction:** a carboxylic ester + H<sub>2</sub>O = an alcohol + a carboxylate  
**Other name(s):** ali-esterase; B-esterase; monobutyrase; cocaine esterase; procaine esterase; methylbutyrase; vitamin A esterase; butyryl esterase; carboxyesterase; carboxylate esterase; carboxylic esterase; methylbutyrate esterase; triacetin esterase; carboxyl ester hydrolase; butyrate esterase; methylbutyrase;  $\alpha$ -carboxylesterase; propionyl esterase; nonspecific carboxylesterase; esterase D; esterase B; esterase A; serine esterase; carboxylic acid esterase; cocaine esterase  
**Systematic name:** carboxylic-ester hydrolase  
**Comments:** Wide specificity. The enzymes from microsomes also catalyse the reactions of EC 3.1.1.2 (arylesterase), EC 3.1.1.5 (lysophospholipase), EC 3.1.1.6 (acetyesterase), EC 3.1.1.23 (acylglycerol lipase), EC 3.1.1.28 (acylcarnitine hydrolase), EC 3.1.2.2 (palmitoyl-CoA hydrolase), EC 3.5.1.4 (amidase) and EC 3.5.1.13 (aryl-acylamidase). Also hydrolyses vitamin A esters.  
**References:** [97, 141, 207, 325, 1177, 1790, 1889, 2479]

[EC 3.1.1.1 created 1961]

#### EC 3.1.1.2

- Accepted name:** arylesterase  
**Reaction:** a phenyl acetate + H<sub>2</sub>O = a phenol + acetate  
**Other name(s):** A-esterase; paraoxonase; aromatic esterase  
**Systematic name:** aryl-ester hydrolase  
**Comments:** Acts on many phenolic esters. The reactions of EC 3.1.8.1 aryldialkylphosphatase, were previously attributed to this enzyme. It is likely that the three forms of human paraoxonase are lactonases rather than aromatic esterases [1434, 645]. The natural substrates of the paraoxonases are lactones [1434, 645], with ( $\pm$ )-5-hydroxy-6*E*,8*Z*,11*Z*,4*Z*-eicostetraenoic-acid 1,5-lactone being the best substrate [645].  
**References:** [30, 102, 267, 1445, 1768, 1, 1434, 645]

[EC 3.1.1.2 created 1961, modified 1989]

#### EC 3.1.1.3

- Accepted name:** triacylglycerol lipase  
**Reaction:** triacylglycerol + H<sub>2</sub>O = diacylglycerol + a carboxylate

**Other name(s):** lipase (ambiguous); butyrinase; tributyrinase; Tween hydrolase; steapsin; triacetinase; tributyrin esterase; Tweenase; amno *N*-AP; Takedo 1969-4-9; Meito MY 30; Tweenesterase; GA 56; capalase L; triglyceride hydrolase; triolein hydrolase; tween-hydrolyzing esterase; amano CE; cacordase; triglyceridase; triacylglycerol ester hydrolase; amano P; amano AP; PPL; glycerol-ester hydrolase; GEH; meito Sangyo OF lipase; hepatic lipase; lipazin; post-heparin plasma protamine-resistant lipase; salt-resistant post-heparin lipase; heparin releasable hepatic lipase; amano CES; amano B; tributyrase; triglyceride lipase; liver lipase; hepatic monoacylglycerol acyltransferase

**Systematic name:** triacylglycerol acylhydrolase

**Comments:** The pancreatic enzyme acts only on an ester-water interface; the outer ester links are preferentially hydrolysed.

**References:** [1511, 1760, 2528, 2669, 2670]

[EC 3.1.1.3 created 1961]

#### EC 3.1.1.4

**Accepted name:** phospholipase A<sub>2</sub>

**Reaction:** phosphatidylcholine + H<sub>2</sub>O = 1-acylglycerophosphocholine + a carboxylate

**Other name(s):** lecithinase A; phosphatidase; phosphatidolipase; phospholipase A

**Systematic name:** phosphatidylcholine 2-acylhydrolase

**Comments:** Also acts on phosphatidylethanolamine, choline plasmalogen and phosphatides, removing the fatty acid attached to the 2-position. Requires Ca<sup>2+</sup>.

**References:** [628, 801, 1042, 1957, 2497, 3032]

[EC 3.1.1.4 created 1961, modified 1976, modified 1983]

#### EC 3.1.1.5

**Accepted name:** lysophospholipase

**Reaction:** 2-lysophosphatidylcholine + H<sub>2</sub>O = glycerophosphocholine + a carboxylate

**Other name(s):** lecithinase B; lysolecithinase; phospholipase B; lysophosphatidase; lecitholipase; phosphatidase B; lysophosphatidylcholine hydrolase; lysophospholipase A1; lysophospholipase L2; lysophospholipase transacylase; neuropathy target esterase; NTE; NTE-LysoPLA; NTE-lysophospholipase

**Systematic name:** 2-lysophosphatidylcholine acylhydrolase

**References:** [5, 483, 551, 732, 2622, 3033, 3035, 3048, 2345, 1756, 3198]

[EC 3.1.1.5 created 1961, modified 1976, modified 1983]

#### EC 3.1.1.6

**Accepted name:** acetylerase

**Reaction:** an acetic ester + H<sub>2</sub>O = an alcohol + acetate

**Other name(s):** C-esterase (in animal tissues); acetic ester hydrolase; chloroesterase; *p*-nitrophenyl acetate esterase; *Citrus* acetylerase

**Systematic name:** acetic-ester acetylhydrolase

**References:** [30, 196, 1308]

[EC 3.1.1.6 created 1961]

#### EC 3.1.1.7

**Accepted name:** acetylcholinesterase

**Reaction:** acetylcholine + H<sub>2</sub>O = choline + acetate

**Other name(s):** true cholinesterase; choline esterase I; cholinesterase; acetylthiocholinesterase; acetylcholine hydrolase; acetyl.β-methylcholinesterase; AcCholE

**Systematic name:** acetylcholine acetylhydrolase

**Comments:** Acts on a variety of acetic esters; also catalyses transacetylations.

**References:** [98, 197, 453, 1653, 2021, 3344]

[EC 3.1.1.7 created 1961]

#### EC 3.1.1.8

**Accepted name:** cholinesterase  
**Reaction:** an acylcholine + H<sub>2</sub>O = choline + a carboxylate  
**Other name(s):** pseudocholinesterase; butyrylcholine esterase; non-specific cholinesterase; choline esterase II (un-specific); benzoylcholinesterase; choline esterase; butyrylcholinesterase; propionylcholinesterase; BtChoEase  
**Systematic name:** acylcholine acylhydrolase  
**Comments:** Acts on a variety of choline esters and a few other compounds.  
**References:** [98, 102, 1493, 2021, 2548, 2780]

[EC 3.1.1.8 created 1961]

[3.1.1.9 Deleted entry. benzoylcholinesterase; a side reaction of EC 3.1.1.8 cholinesterase]

[EC 3.1.1.9 created 1961, deleted 1972]

#### EC 3.1.1.10

**Accepted name:** tropinesterase  
**Reaction:** atropine + H<sub>2</sub>O = tropine + tropate  
**Other name(s):** tropine esterase; atropinase; atropine esterase  
**Systematic name:** atropine acylhydrolase  
**Comments:** Also acts on cocaine and other tropine esters.  
**References:** [930, 1956]

[EC 3.1.1.10 created 1961, deleted 1972, reinstated 1976]

#### EC 3.1.1.11

**Accepted name:** pectinesterase  
**Reaction:** pectin + *n* H<sub>2</sub>O = *n* methanol + pectate  
**Other name(s):** pectin demethoxylase; pectin methoxylase; pectin methylesterase; pectase; pectin methyl esterase; pectinoesterase  
**Systematic name:** pectin pectylhydrolase  
**References:** [593, 1704, 1918]

[EC 3.1.1.11 created 1961]

[3.1.1.12 Deleted entry. vitamin A esterase, now believed to be identical with EC 3.1.1.1 carboxylesterase]

[EC 3.1.1.12 created 1961, deleted 1972]

#### EC 3.1.1.13

**Accepted name:** sterol esterase  
**Reaction:** a steryl ester + H<sub>2</sub>O = a sterol + a fatty acid  
**Other name(s):** cholesterol esterase; cholesteryl ester synthase; triterpenol esterase; cholesteryl esterase; cholesteryl ester hydrolase; sterol ester hydrolase; cholesterol ester hydrolase; cholesterase; acylcholesterol lipase  
**Systematic name:** steryl-ester acylhydrolase  
**Comments:** A group of enzymes of broad specificity, acting on esters of sterols and long-chain fatty acids, that may also bring about the esterification of sterols. Activated by bile salts.  
**References:** [1216, 2185, 3028, 3127]

[EC 3.1.1.13 created 1961, modified 1990]

#### EC 3.1.1.14

**Accepted name:** chlorophyllase  
**Reaction:** chlorophyll + H<sub>2</sub>O = phytol + chlorophyllide  
**Other name(s):** CLH; Chlase  
**Systematic name:** chlorophyll chlorophyllidohydrolase  
**Comments:** Chlorophyllase has been found in higher plants, diatoms, and in the green algae *Chlorella* [2977]. This enzyme forms part of the chlorophyll degradation pathway and is thought to take part in de-greening processes such as fruit ripening, leaf senescence and flowering, as well as in the turnover and homeostasis of chlorophyll [2187]. This enzyme acts preferentially on chlorophyll *a* but will also accept chlorophyll *b* and pheophytins as substrates [1180]. Ethylene and methyl jasmonate, which are known to accelerate senescence in many species, can enhance the activity of the hormone-inducible form of this enzyme [1180].  
**References:** [1162, 1473, 2977, 2187, 1180]

[EC 3.1.1.14 created 1961, modified 2007]

#### EC 3.1.1.15

**Accepted name:** L-arabinonolactonase  
**Reaction:** L-arabinono-1,4-lactone + H<sub>2</sub>O = L-arabinonate  
**Systematic name:** L-arabinono-1,4-lactone lactonohydrolase  
**References:** [3153]

[EC 3.1.1.15 created 1961]

[3.1.1.16 Deleted entry. 4-carboxymethyl-4-hydroxyisocrotonolactonase. This reaction was due to a mixture of EC 5.3.3.4 (muconolactone Δ-isomerase) and EC 3.1.1.24 (3-oxoadipate enol-lactonase)]

[EC 3.1.1.16 created 1961, deleted 1972]

#### EC 3.1.1.17

**Accepted name:** gluconolactonase  
**Reaction:** D-glucono-1,5-lactone + H<sub>2</sub>O = D-gluconate  
**Other name(s):** lactonase; aldonolactonase; glucono-δ-lactonase; gulonolactonase  
**Systematic name:** D-glucono-1,5-lactone lactonohydrolase  
**Comments:** Acts on a wide range of hexose-1,5-lactones. The hydrolysis of L-gulono-1,5-lactone was previously listed separately.  
**References:** [298, 322, 2812]

[EC 3.1.1.17 created 1961 (EC 3.1.1.18 created 1961, incorporated 1982)]

[3.1.1.18 Deleted entry. aldonolactonase. Now included with EC 3.1.1.17 gluconolactonase]

[EC 3.1.1.18 created 1961, deleted 1982]

#### EC 3.1.1.19

**Accepted name:** uronolactonase  
**Reaction:** D-glucurono-6,2-lactone + H<sub>2</sub>O = D-glucuronate  
**Other name(s):** glucuronolactonase  
**Systematic name:** D-glucurono-6,2-lactone lactonohydrolase  
**References:** [3196]

[EC 3.1.1.19 created 1961]

#### EC 3.1.1.20

**Accepted name:** tannase  
**Reaction:** digallate + H<sub>2</sub>O = 2 gallate  
**Other name(s):** tannase S; tannin acetylhydrolase  
**Systematic name:** tannin acylhydrolase  
**Comments:** Also hydrolyses ester links in other tannins.  
**References:** [672]

[EC 3.1.1.20 created 1961]

[3.1.1.21 Deleted entry. retinyl-palmitate esterase. Now known to be catalysed by EC 3.1.1.1, carboxylesterase and EC 3.1.1.3, triacylglycerol lipase.]

[EC 3.1.1.21 created 1972, deleted 2011]

#### EC 3.1.1.22

**Accepted name:** hydroxybutyrate-dimer hydrolase  
**Reaction:** (R)-3-((R)-3-hydroxybutanoyloxy)butanoate + H<sub>2</sub>O = 2 (R)-3-hydroxybutanoate  
**Other name(s):** D-(-)-3-hydroxybutyrate-dimer hydrolase  
**Systematic name:** (R)-3-((R)-3-hydroxybutanoyloxy)butanoate hydroxybutanoylhydrolase  
**References:** [573]

[EC 3.1.1.22 created 1972]

#### EC 3.1.1.23

**Accepted name:** acylglycerol lipase  
**Reaction:** Hydrolyses glycerol monoesters of long-chain fatty acids  
**Other name(s):** monoacylglycerol lipase; monoacylglycerolipase; monoglyceride lipase; monoglyceride hydrolase; fatty acyl monoester lipase; monoacylglycerol hydrolase; monoglyceridyllipase; monoglyceridase  
**Systematic name:** glycerol-ester acylhydrolase  
**References:** [1887, 2311]

[EC 3.1.1.23 created 1972]

#### EC 3.1.1.24

**Accepted name:** 3-oxoadipate enol-lactonase  
**Reaction:** 3-oxoadipate enol-lactone + H<sub>2</sub>O = 3-oxoadipate  
**Other name(s):** carboxymethylbutenolide lactonase; β-ketoadipic enol-lactone hydrolase; 3-ketoadipate enol-lactonase; 3-oxoadipic enol-lactone hydrolase; β-ketoadipate enol-lactone hydrolase  
**Systematic name:** 4-carboxymethylbut-3-en-4-olide enol-lactonohydrolase  
**Comments:** The enzyme acts on the product of EC 4.1.1.44 4-carboxymuconolactone decarboxylase.  
**References:** [2202, 2203]

[EC 3.1.1.24 created 1961 as EC 3.1.1.16, part transferred 1972 to EC 3.1.1.24]

#### EC 3.1.1.25

**Accepted name:** 1,4-lactonase  
**Reaction:** a 1,4-lactone + H<sub>2</sub>O = a 4-hydroxyacid  
**Other name(s):** γ-lactonase  
**Systematic name:** 1,4-lactone hydroxyacylhydrolase  
**Comments:** The enzyme is specific for 1,4-lactones with 4-8 carbon atoms. It does not hydrolyse simple aliphatic esters, acetylcholine, sugar lactones or substituted aliphatic lactones, e.g. 3-hydroxy-4-butyrolactone; requires Ca<sup>2+</sup>.  
**References:** [775, 776]

[EC 3.1.1.25 created 1972, modified 1981]

#### EC 3.1.1.26

**Accepted name:** galactolipase  
**Reaction:**  $1,2\text{-diacyl-3-}\beta\text{-D-galactosyl-}sn\text{-glycerol} + 2 \text{H}_2\text{O} = 3\text{-}\beta\text{-D-galactosyl-}sn\text{-glycerol} + 2 \text{carboxylates}$   
**Other name(s):** galactolipid lipase; polygalactolipase; galactolipid acylhydrolase  
**Systematic name:** 1,2-diacyl-3- $\beta$ -D-galactosyl-*sn*-glycerol acylhydrolase  
**Comments:** Also acts on 2,3-di-*O*-acyl-1-*O*-(6-*O*- $\alpha$ -D-galactosyl- $\beta$ -D-galactosyl)-D-glycerol, and phosphatidyl-choline and other phospholipids.  
**References:** [1108, 1150]

[EC 3.1.1.26 created 1972]

#### EC 3.1.1.27

**Accepted name:** 4-pyridoxolactonase  
**Reaction:**  $4\text{-pyridoxolactone} + \text{H}_2\text{O} = 4\text{-pyridoxate}$   
**Systematic name:** 4-pyridoxolactone lactonohydrolase  
**References:** [327]

[EC 3.1.1.27 created 1972]

#### EC 3.1.1.28

**Accepted name:** acylcarnitine hydrolase  
**Reaction:**  $O\text{-acylcarnitine} + \text{H}_2\text{O} = \text{a fatty acid} + \text{L-carnitine}$   
**Other name(s):** high activity acylcarnitine hydrolase; HACH; carnitine ester hydrolase; palmitoylcarnitine hydrolase; palmitoyl-L-carnitine hydrolase; long-chain acyl-L-carnitine hydrolase; palmitoyl carnitine hydrolase  
**Systematic name:** *O*-acylcarnitine acylhydrolase  
**Comments:** Acts on higher fatty acid (C<sub>6</sub> to C<sub>18</sub>) esters of L-carnitine; highest activity is with *O*-decanoyl-L-carnitine.  
**References:** [1773, 1888]

[EC 3.1.1.28 created 1972]

#### EC 3.1.1.29

**Accepted name:** aminoacyl-tRNA hydrolase  
**Reaction:**  $N\text{-substituted aminoacyl-tRNA} + \text{H}_2\text{O} = N\text{-substituted amino acid} + \text{tRNA}$   
**Other name(s):** aminoacyl-transfer ribonucleate hydrolase; *N*-substituted aminoacyl transfer RNA hydrolase; peptidyl-tRNA hydrolase  
**Systematic name:** aminoacyl-tRNA aminoacylhydrolase  
**References:** [1342]

[EC 3.1.1.29 created 1972]

#### EC 3.1.1.30

**Accepted name:** D-arabinonolactonase  
**Reaction:**  $D\text{-arabinono-1,4-lactone} + \text{H}_2\text{O} = D\text{-arabinonate}$   
**Systematic name:** D-arabinono-1,4-lactone lactonohydrolase  
**References:** [2227]

[EC 3.1.1.30 created 1972]

#### EC 3.1.1.31



**Accepted name:** 6-phosphogluconolactonase  
**Reaction:** 6-phospho-D-glucono-1,5-lactone + H<sub>2</sub>O = 6-phospho-D-gluconate  
**Other name(s):** phosphogluconolactonase; 6-PGL  
**Systematic name:** 6-phospho-D-glucono-1,5-lactone lactonohydrolase  
**References:** [1405, 1906]

[EC 3.1.1.31 created 1972]

#### EC 3.1.1.32

**Accepted name:** phospholipase A<sub>1</sub>  
**Reaction:** phosphatidylcholine + H<sub>2</sub>O = 2-acylglycerophosphocholine + a carboxylate  
**Systematic name:** phosphatidylcholine 1-acylhydrolase  
**Comments:** This enzyme has a much broader specificity than EC 3.1.1.4 phospholipase A<sub>2</sub>. Requires Ca<sup>2+</sup>.  
**References:** [883, 2549, 3032, 3034]

[EC 3.1.1.32 created 1972, modified 1976]

#### EC 3.1.1.33

**Accepted name:** 6-acetylglucose deacetylase  
**Reaction:** 6-acetyl-D-glucose + H<sub>2</sub>O = D-glucose + acetate  
**Other name(s):** 6-O-acetylglucose deacetylase  
**Systematic name:** 6-acetyl-D-glucose acetylhydrolase  
**References:** [663]

[EC 3.1.1.33 created 1972]

#### EC 3.1.1.34

**Accepted name:** lipoprotein lipase  
**Reaction:** triacylglycerol + H<sub>2</sub>O = diacylglycerol + a carboxylate  
**Other name(s):** clearing factor lipase; diglyceride lipase; diacylglycerol lipase; postheparin esterase; diglyceride lipase; postheparin lipase; diacylglycerol hydrolase; lipemia-clearing factor  
**Systematic name:** triacylglycero-protein acylhydrolase  
**Comments:** Hydrolyses triacylglycerols in chylomicrons and low-density lipoproteins. Also hydrolyses diacylglycerol.  
**References:** [681, 765, 977, 1981, 2089]

[EC 3.1.1.34 created 1972, modified 1976]

#### EC 3.1.1.35

**Accepted name:** dihydrocoumarin hydrolase  
**Reaction:** dihydrocoumarin + H<sub>2</sub>O = melilotate  
**Systematic name:** dihydrocoumarin lactonohydrolase  
**Comments:** Also hydrolyses some other benzenoid 1,4-lactones.  
**References:** [1517]

[EC 3.1.1.35 created 1972]

#### EC 3.1.1.36

**Accepted name:** limonin-D-ring-lactonase  
**Reaction:** limonoate D-ring-lactone + H<sub>2</sub>O = limonoate  
**Other name(s):** limonin-D-ring-lactone hydrolase; limonin lactone hydrolase  
**Systematic name:** limonoate-D-ring-lactone lactonohydrolase

**Comments:** Limonoate is a triterpenoid.

**References:** [1776]

[EC 3.1.1.36 created 1972]

#### EC 3.1.1.37

**Accepted name:** steroid-lactonase

**Reaction:** testololactone + H<sub>2</sub>O = testolate

**Systematic name:** testololactone lactonohydrolase

**References:** [1164]

[EC 3.1.1.37 created 1972]

#### EC 3.1.1.38

**Accepted name:** triacetate-lactonase

**Reaction:** triacetate lactone + H<sub>2</sub>O = triacetate

**Other name(s):** triacetic lactone hydrolase; triacetic acid lactone hydrolase; TAL hydrolase; triacetate lactone hydrolase

**Systematic name:** triacetolactone lactonohydrolase

**References:** [1401]

[EC 3.1.1.38 created 1972]

#### EC 3.1.1.39

**Accepted name:** actinomycin lactonase

**Reaction:** actinomycin + H<sub>2</sub>O = actinomycinic monolactone

**Systematic name:** actinomycin lactonohydrolase

**References:** [1186]

[EC 3.1.1.39 created 1972]

#### EC 3.1.1.40

**Accepted name:** orsellinate-depside hydrolase

**Reaction:** orsellinate depside + H<sub>2</sub>O = 2 orsellinate

**Other name(s):** lecanorate hydrolase

**Systematic name:** orsellinate-depside hydrolase

**Comments:** The enzyme will only hydrolyse those substrates based on the 2,4-dihydroxy-6-methylbenzoate structure that also have a free hydroxy group *ortho* to the depside linkage.

**References:** [2588]

[EC 3.1.1.40 created 1976]

#### EC 3.1.1.41

**Accepted name:** cephalosporin-C deacetylase

**Reaction:** cephalosporin C + H<sub>2</sub>O = deacetylcephalosporin C + acetate

**Other name(s):** cephalosporin C acetyl-hydrolase; cephalosporin C acetylase; cephalosporin acylesterase; cephalosporin C acylesterase; cephalosporin C acetyl-esterase; cephalosporin C deacetylase

**Systematic name:** cephalosporin-C acetylhydrolase

**Comments:** Hydrolyses the acetyl ester bond on the 10-position of the antibiotic cephalosporin C.

**References:** [848]

[EC 3.1.1.41 created 1976]

#### EC 3.1.1.42

**Accepted name:** chlorogenate hydrolase  
**Reaction:** chlorogenate + H<sub>2</sub>O = caffeate + quinate  
**Other name(s):** chlorogenase; chlorogenic acid esterase  
**Systematic name:** chlorogenate hydrolase  
**Comments:** Also acts, more slowly, on isochlorogenate. No other substrates are known.  
**References:** [2577, 2578]

[EC 3.1.1.42 created 1981]

#### EC 3.1.1.43

**Accepted name:** α-amino-acid esterase  
**Reaction:** an α-amino acid ester + H<sub>2</sub>O = an α-amino acid + an alcohol  
**Other name(s):** α-amino acid ester hydrolase  
**Systematic name:** α-amino-acid-ester aminoacylhydrolase  
**Comments:** Also catalyses α-aminoacyl transfer to a number of amine nucleophiles.  
**References:** [1399, 1400, 2848]

[EC 3.1.1.43 created 1983]

#### EC 3.1.1.44

**Accepted name:** 4-methyloxaloacetate esterase  
**Reaction:** oxaloacetate 4-methyl ester + H<sub>2</sub>O = oxaloacetate + methanol  
**Systematic name:** oxaloacetate-4-methyl-ester oxaloacetohydrolase  
**References:** [639]

[EC 3.1.1.44 created 1983]

#### EC 3.1.1.45

**Accepted name:** carboxymethylenebutenolidase  
**Reaction:** 4-carboxymethylenebut-2-en-4-olide + H<sub>2</sub>O = 4-oxohex-2-enedioate  
**Other name(s):** maleylacetate enol-lactonase; dienelactone hydrolase; carboxymethylene butenolide hydrolase  
**Systematic name:** 4-carboxymethylenebut-2-en-4-olide lactonohydrolase  
**References:** [2572]

[EC 3.1.1.45 created 1983]

#### EC 3.1.1.46

**Accepted name:** deoxylimonate A-ring-lactonase  
**Reaction:** deoxylimonate + H<sub>2</sub>O = deoxylimononic acid D-ring-lactone  
**Systematic name:** deoxylimonate A-ring-lactonohydrolase  
**Comments:** The enzyme opens the A-ring-lactone of the triterpenoid deoxylimononic acid, leaving the D-ring-lactone intact.  
**References:** [1066]

[EC 3.1.1.46 created 1983]

#### EC 3.1.1.47

**Accepted name:** 1-alkyl-2-acetyl-glycerophosphocholine esterase  
**Reaction:** 1-alkyl-2-acetyl-*sn*-glycero-3-phosphocholine + H<sub>2</sub>O = 1-alkyl-*sn*-glycero-3-phosphocholine + acetate  
**Other name(s):** 1-alkyl-2-acetyl-*sn*-glycero-3-phosphocholine acetylhydrolase; alkylacetyl-GPC:acetylhydrolase

**Systematic name:** 1-alkyl-2-acetyl-*sn*-glycero-3-phosphocholine acetohydrolase

**References:** [240]

[EC 3.1.1.47 created 1984]

#### EC 3.1.1.48

**Accepted name:** fusarinine-C ornithinesterase

**Reaction:**  $N^5$ -acyl-L-ornithine ester + H<sub>2</sub>O =  $N^5$ -acyl-L-ornithine + an alcohol

**Other name(s):** ornithine esterase; 5-*N*-acyl-L-ornithine-ester hydrolase

**Systematic name:**  $N^5$ -acyl-L-ornithine-ester hydrolase

**Comments:** Hydrolyses the three ornithine ester bonds in fusarinine C. Also acts on  $N^5$ -dinitrophenyl-L-ornithine methyl ester.

**References:** [697]

[EC 3.1.1.48 created 1984]

#### EC 3.1.1.49

**Accepted name:** sinapine esterase

**Reaction:** sinapoylcholine + H<sub>2</sub>O = sinapate + choline

**Other name(s):** aromatic choline esterase

**Systematic name:** sinapoylcholine sinapohydrolase

**References:** [2121]

[EC 3.1.1.49 created 1984]

#### EC 3.1.1.50

**Accepted name:** wax-ester hydrolase

**Reaction:** a wax ester + H<sub>2</sub>O = a long-chain alcohol + a long-chain carboxylate

**Other name(s):** jojoba wax esterase; WEH

**Systematic name:** wax-ester acylhydrolase

**Comments:** Also acts on long-chain acylglycerol, but not diacyl- or triacylglycerols.

**References:** [1201, 1963]

[EC 3.1.1.50 created 1984]

#### EC 3.1.1.51

**Accepted name:** phorbol-diester hydrolase

**Reaction:** phorbol 12,13-dibutanoate + H<sub>2</sub>O = phorbol 13-butanoate + butanoate

**Other name(s):** diacylphorbate 12-hydrolase; diacylphorbate 12-hydrolase; phorbol-12,13-diester 12-ester hydrolase; PDEH

**Systematic name:** 12,13-diacylphorbate 12-acylhydrolase

**Comments:** Hydrolyses the 12-ester bond in a variety of 12,13-diacylphorbols (phorbol is a diterpenoid); this reaction inactivates the tumour promotor 12-*O*-tetradecanoylphorbol-13-acetate from croton oil.

**References:** [2647]

[EC 3.1.1.51 created 1984]

#### EC 3.1.1.52

**Accepted name:** phosphatidylinositol deacylase

**Reaction:** 1-phosphatidyl-*D*-*myo*-inositol + H<sub>2</sub>O = 1-acylglycerophosphoinositol + a carboxylate

**Other name(s):** phosphatidylinositol phospholipase A<sub>2</sub>; phospholipase A<sub>2</sub>

**Systematic name:** 1-phosphatidyl-*D*-*myo*-inositol 2-acylhydrolase

**References:** [970, 969]

[EC 3.1.1.52 created 1984]

#### EC 3.1.1.53

**Accepted name:** sialate *O*-acetyltransferase  
**Reaction:** *N*-acetyl-*O*-acetylneuraminate + H<sub>2</sub>O = *N*-acetylneuraminate + acetate  
**Other name(s):** *N*-acetylneuraminate acetyltransferase; sialate 9(4)-*O*-acetyltransferase; sialidase  
**Systematic name:** *N*-acyl-*O*-acetylneuraminate *O*-acetylhydrolase  
**Comments:** Acts on free and glycosidically bound *N*-acetyl- or *N*-glycoloyl-neuraminic acid; acts mainly on the 4-*O*- and 9-*O*-acetyl groups. Also acts on some other *O*-acetyl esters, both cyclic and acyclic compounds, which are not sialic acids.  
**References:** [877, 2648]

[EC 3.1.1.53 created 1984]

#### EC 3.1.1.54

**Accepted name:** acetoxybutynylbithiophene deacetylase  
**Reaction:** 5-(4-acetoxybut-1-ynyl)-2,2'-bithiophene + H<sub>2</sub>O = 5-(4-hydroxybut-1-ynyl)-2,2'-bithiophene + acetate  
**Other name(s):** acetoxybutynylbithiophene esterase; 5-(4-acetoxy-1-butynyl)-2,2'-bithiophene:acetate esterase  
**Systematic name:** 5-(4-acetoxybut-1-ynyl)-2,2'-bithiophene *O*-acetylhydrolase  
**Comments:** The enzyme is highly specific.  
**References:** [2807]

[EC 3.1.1.54 created 1986]

#### EC 3.1.1.55

**Accepted name:** acetylsalicylate deacetylase  
**Reaction:** acetylsalicylate + H<sub>2</sub>O = salicylate + acetate  
**Other name(s):** aspirin esterase; aspirin esterase; acetylsalicylic acid esterase; aspirin hydrolase  
**Systematic name:** acetylsalicylate *O*-acetylhydrolase  
**Comments:** Not identical with EC 3.1.1.1 (carboxylesterase), EC 3.1.1.2 (arylesterase), EC 3.1.1.7 (acetylcholinesterase) or EC 3.1.1.8 (cholinesterase). The activity of the liver cytosol enzyme is highest with acetyl esters of aryl alcohols, and thioesters are also hydrolysed; the microsomal enzyme also hydrolyses some other negatively charged esters, with highest activity on esters of salicylate with long-chain alcohols.  
**References:** [35, 1444, 3173]

[EC 3.1.1.55 created 1986, modified 1989]

#### EC 3.1.1.56

**Accepted name:** methylumbelliferyl-acetate deacetylase  
**Reaction:** 4-methylumbelliferyl acetate + H<sub>2</sub>O = 4-methylumbelliferone + acetate  
**Other name(s):** esterase D  
**Systematic name:** 4-methylumbelliferyl-acetate acylhydrolase  
**Comments:** Acts on short-chain acyl esters of 4-methylumbelliferone, but not on naphthyl, indoxyl or thiocholine esters.  
**References:** [1174]

[EC 3.1.1.56 created 1986]

#### EC 3.1.1.57

**Accepted name:** 2-pyrone-4,6-dicarboxylate lactonase

**Reaction:** 2-oxo-2*H*-pyran-4,6-dicarboxylate + H<sub>2</sub>O = (1*E*)-4-oxobut-1-ene-1,2,4-tricarboxylate  
**Other name(s):** 2-pyrone-4,6-dicarboxylate hydrolase; 2-pyrone-4,6-dicarboxylate lactonohydrolase  
**Systematic name:** 2-oxo-2*H*-pyran-4,6-dicarboxylate lactonohydrolase  
**Comments:** The product is most likely the keto-form of 4-oxalomesaconate (as shown in the reaction) [1422, 1827]. It can be converted to the enol-form, 4-hydroxybuta-1,3-diene-1,2,4-trioate, either spontaneously or by EC 5.3.2.8, 4-oxalomesaconate tautomerase [2103].  
**References:** [1422, 1827, 2103]

[EC 3.1.1.57 created 1986, modified 2010]

#### EC 3.1.1.58

**Accepted name:** *N*-acetylgalactosaminoglycan deacetylase  
**Reaction:** *N*-acetyl-D-galactosaminoglycan + H<sub>2</sub>O = D-galactosaminoglycan + acetate  
**Other name(s):** polysaccharide deacetylase (misleading); Vi-polysaccharide deacetylase; *N*-acetyl galactosaminoglycan deacetylase  
**Systematic name:** *N*-acetyl-D-galactosaminoglycan acetylhydrolase  
**References:** [1339]

[EC 3.1.1.58 created 1986]

#### EC 3.1.1.59

**Accepted name:** juvenile-hormone esterase  
**Reaction:** (1) juvenile hormone I + H<sub>2</sub>O = juvenile hormone I acid + methanol  
(2) juvenile hormone III + H<sub>2</sub>O = juvenile hormone III acid + methanol  
**Other name(s):** JH-esterase; juvenile hormone analog esterase; juvenile hormone carboxyesterase; methyl-(2*E*,6*E*)-(10*R*,11*S*)-10,11-epoxy-3,7,11-trimethyltrideca-2,6-dienoate acylhydrolase  
**Systematic name:** methyl-(2*E*,6*E*,10*R*)-10,11-epoxy-3,7,11-trimethyltrideca-2,6-dienoate acylhydrolase  
**Comments:** Demethylates the insect juvenile hormones JH1 and JH3, but does not hydrolyse the analogous ethyl or isopropyl esters.  
**References:** [558, 1932]

[EC 3.1.1.59 created 1989, modified 2015]

#### EC 3.1.1.60

**Accepted name:** bis(2-ethylhexyl)phthalate esterase  
**Reaction:** bis(2-ethylhexyl)phthalate + H<sub>2</sub>O = 2-ethylhexyl phthalate + 2-ethylhexan-1-ol  
**Other name(s):** DEHP esterase  
**Systematic name:** bis(2-ethylhexyl)phthalate acylhydrolase  
**Comments:** Also acts on 4-nitrophenyl esters, with optimum chain-length C<sub>6</sub> to C<sub>8</sub>.  
**References:** [1011]

[EC 3.1.1.60 created 1989]

#### EC 3.1.1.61

**Accepted name:** protein-glutamate methylesterase  
**Reaction:** protein L-glutamate *O*<sup>5</sup>-methyl ester + H<sub>2</sub>O = protein L-glutamate + methanol  
**Other name(s):** chemotaxis-specific methylesterase; methyl-accepting chemotaxis protein methyl-esterase; CheB methylesterase; methylesterase CheB; protein methyl-esterase; protein carboxyl methylesterase; PME; protein methylesterase; protein-L-glutamate-5-*O*-methyl-ester acylhydrolase  
**Systematic name:** protein-L-glutamate-*O*<sup>5</sup>-methyl-ester acylhydrolase  
**Comments:** Hydrolyses the products of EC 2.1.1.77 (protein-L-isoaspartate(D-aspartate) *O*-methyltransferase), EC 2.1.1.78 (isoorientin 3'-*O*-methyltransferase), EC 2.1.1.80 (protein-glutamate *O*-methyltransferase) and EC 2.1.1.100 (protein-*S*-isoprenylcysteine *O*-methyltransferase).  
**References:** [868, 1411]

[EC 3.1.1.61 created 1989, modified 2002]

[3.1.1.62 Deleted entry. *N*-acetyldiaminopimelate deacylase. Now listed as EC 3.5.1.47, *N*-acetyldiaminopimelate deacetylase]

[EC 3.1.1.62 created 1989, deleted 1992]

#### EC 3.1.1.63

**Accepted name:** 11-*cis*-retinyl-palmitate hydrolase  
**Reaction:** 11-*cis*-retinyl palmitate + H<sub>2</sub>O = 11-*cis*-retinol + palmitate  
**Other name(s):** 11-*cis*-retinol palmitate esterase; RPH  
**Systematic name:** 11-*cis*-retinyl-palmitate acylhydrolase  
**Comments:** Activated by bile salts.  
**References:** [238, 239]

[EC 3.1.1.63 created 1989]

#### EC 3.1.1.64

**Accepted name:** retinoid isomerohydrolase  
**Reaction:** an *all-trans*-retinyl ester + H<sub>2</sub>O = 11-*cis*-retinol + a fatty acid  
**Other name(s):** *all-trans*-retinyl-palmitate hydrolase (ambiguous); retinol isomerase (ambiguous); *all-trans*-retinol isomerase:hydrolase (ambiguous); *all-trans*-retinylester 11-*cis* isomerohydrolase; RPE65 (gene name)  
**Systematic name:** *all-trans*-retinyl ester acylhydrolase, 11-*cis* retinol forming  
**Comments:** This enzyme, which operates in the retinal pigment epithelium (RPE), catalyses the cleavage and isomerization of *all-trans*-retinyl fatty acid esters to 11-*cis*-retinol, a key step in the regeneration of the visual chromophore in the vertebrate visual cycle [1946]. Interaction of the enzyme with the membrane is critical for its enzymic activity [942].  
**References:** [238, 203, 288, 1946, 2086, 942]

[EC 3.1.1.64 created 1989 (EC 5.2.1.7 created 1989, incorporated 2011), modified 2011]

#### EC 3.1.1.65

**Accepted name:** L-rhamnono-1,4-lactonase  
**Reaction:** L-rhamnono-1,4-lactone + H<sub>2</sub>O = L-rhamnonate  
**Other name(s):** L-rhamno- $\gamma$ -lactonase; L-rhamnono- $\gamma$ -lactonase; L-rhamnonate dehydratase  
**Systematic name:** L-rhamnono-1,4-lactone lactonohydrolase  
**References:** [2426]

[EC 3.1.1.65 created 1989]

#### EC 3.1.1.66

**Accepted name:** 5-(3,4-diacetoxybut-1-ynyl)-2,2'-bithiophene deacetylase  
**Reaction:** 5-(3,4-diacetoxybut-1-ynyl)-2,2'-bithiophene + H<sub>2</sub>O = 5-(3-hydroxy-4-acetoxybut-1-ynyl)-2,2'-bithiophene + acetate  
**Other name(s):** diacetoxybutynylbithiophene acetate esterase; 3,4-diacetoxybutynylbithiophene:4-acetate esterase  
**Systematic name:** 5-(3,4-diacetoxybut-1-ynyl)-2,2'-bithiophene acetylhydrolase  
**Comments:** A highly specific enzyme from *Tagetes patula*.  
**References:** [2256]

[EC 3.1.1.66 created 1989]

#### EC 3.1.1.67

**Accepted name:** fatty-acyl-ethyl-ester synthase

**Reaction:** a long-chain-fatty-acyl ethyl ester + H<sub>2</sub>O = a long-chain-fatty acid + ethanol  
**Other name(s):** FAEES  
**Systematic name:** long-chain-fatty-acyl-ethyl-ester acylhydrolase  
**Comments:** The reaction, forms ethyl esters from fatty acids and ethanol in the absence of coenzyme A or ATP. Best substrates are unsaturated octadecanoic acids; palmitate, stearate and arachidonate also act, but more slowly.  
**References:** [1945]

[EC 3.1.1.67 created 1989]

#### EC 3.1.1.68

**Accepted name:** xylono-1,4-lactonase  
**Reaction:** D-xylono-1,4-lactone + H<sub>2</sub>O = D-xylonate  
**Other name(s):** xylono- $\gamma$ -lactonase; xylonolactonase  
**Systematic name:** D-xylono-1,4-lactone lactonohydrolase  
**References:** [324]

[EC 3.1.1.68 created 1990]

[3.1.1.69 *Transferred entry. N-acetylglucosaminylphosphatidylinositol deacetylase. Now EC 3.5.1.89, N-acetylglucosaminylphosphatidyl deacetylase. Previously classified erroneously as an enzyme that hydrolysed an ester and not an amide*]

[EC 3.1.1.69 created 1992, deleted 2002]

#### EC 3.1.1.70

**Accepted name:** cetraxate benzylesterase  
**Reaction:** cetraxate benzyl ester + H<sub>2</sub>O = cetraxate + benzyl alcohol  
**Systematic name:** cetraxate-benzyl-ester benzylhydrolase  
**Comments:** Acts on a number of benzyl esters of substituted phenyl propanoates, and on the benzyl esters of phenylalanine and tyrosine.  
**References:** [1571]

[EC 3.1.1.70 created 1992]

#### EC 3.1.1.71

**Accepted name:** acetylalkylglycerol acetylhydrolase  
**Reaction:** 2-acetyl-1-alkyl-*sn*-glycerol + H<sub>2</sub>O = 1-alkyl-*sn*-glycerol + acetate  
**Other name(s):** alkylacetylglycerol acetylhydrolase  
**Systematic name:** 2-acetyl-1-alkyl-*sn*-glycerol acetylhydrolase  
**Comments:** Hydrolysis of the acetyl group from the 1-alkyl-2-acetyl and 1-alkyl-3-acetyl substrates occurs at apparently identical rates. The enzyme from Erlich ascites cells is membrane-bound. It differs from lipoprotein lipase (EC 3.1.1.34) since 1,2-diacetyl-*sn*-glycerols are not substrates. It also differs from EC 3.1.1.47, 1-acetyl-2-alkyl-glycerophosphocholine esterase.  
**References:** [241]

[EC 3.1.1.71 created 1999]

#### EC 3.1.1.72

**Accepted name:** acetylxylan esterase  
**Reaction:** Deacetylation of xylans and xylo-oligosaccharides  
**Systematic name:** acetylxylan esterase  
**Comments:** Catalyses the hydrolysis of acetyl groups from polymeric xylan, acetylated xylose, acetylated glucose,  $\alpha$ -naphthyl acetate, *p*-nitrophenyl acetate but not from triacetyl-glycerol. Does not act on acetylated mannan or pectin.



**References:** [2802, 2319, 1811]

[EC 3.1.1.72 created 1999]

#### EC 3.1.1.73

**Accepted name:** feruloyl esterase  
**Reaction:** feruloyl-polysaccharide + H<sub>2</sub>O = ferulate + polysaccharide  
**Other name(s):** ferulic acid esterase, hydroxycinnamoyl esterase, hemicellulase accessory enzymes; FAE-III, cinnamoyl ester hydrolase, FAEA, cinnAE, FAE-I, FAE-II  
**Systematic name:** 4-hydroxy-3-methoxycinnamoyl-sugar hydrolase  
**Comments:** Catalyses the hydrolysis of the 4-hydroxy-3-methoxycinnamoyl (feruloyl) group from an esterified sugar, which is usually arabinose in "natural" substrates. *p*-Nitrophenol acetate and methyl ferulate are poorer substrates. All microbial ferulate esterases are secreted into the culture medium. They are sometimes called hemicellulase accessory enzymes, since they help xylanases and pectinases to break down plant cell wall hemicellulose.  
**References:** [745, 746, 1538, 597, 379]

[EC 3.1.1.73 created 2000]

#### EC 3.1.1.74

**Accepted name:** cutinase  
**Reaction:** cutin + H<sub>2</sub>O = cutin monomers  
**Systematic name:** cutin hydrolase  
**Comments:** Cutin, a polymeric structural component of plant cuticles, is a polymer of hydroxy fatty acids that are usually C<sub>16</sub> or C<sub>18</sub> and contain up to three hydroxy groups. The enzyme from several fungal sources also hydrolyses the *p*-nitrophenyl esters of hexadecanoic acid. It is however inactive towards several esters that are substrates for non-specific esterases.  
**References:** [876, 2331, 2330]

[EC 3.1.1.74 created 2000]

#### EC 3.1.1.75

**Accepted name:** poly(3-hydroxybutyrate) depolymerase  
**Reaction:** [(*R*)-3-hydroxybutanoate]<sub>*n*</sub> + H<sub>2</sub>O = [(*R*)-3-hydroxybutanoate]<sub>*n-x*</sub> + [(*R*)-3-hydroxybutanoate]<sub>*x*</sub>; *x* = 1–5  
**Other name(s):** PHB depolymerase; poly(3HB) depolymerase; poly[(*R*)-hydroxyalkanoic acid] depolymerase; poly(HA) depolymerase; poly(HA<sub>SCL</sub>) depolymerase; poly[(*R*)-3-hydroxybutyrate] hydrolase  
**Systematic name:** poly[(*R*)-3-hydroxybutanoate] hydrolase  
**Comments:** Reaction also occurs with esters of other short-chain-length (C<sub>1</sub>–C<sub>5</sub>) hydroxyalkanoic acids (HA). There are two types of polymers: native (intracellular) granules are amorphous and have an intact surface layer; denatured (extracellular) granules either have no surface layer or a damaged surface layer and are partially crystalline.  
**References:** [1316, 874]

[EC 3.1.1.75 created 2001]

#### EC 3.1.1.76

**Accepted name:** poly(3-hydroxyoctanoate) depolymerase  
**Reaction:** Hydrolyses the polyester polyoxycarbonyl[(*R*)-2-pentylethylene] to oligomers  
**Other name(s):** PHO depolymerase; poly(3HO) depolymerase; poly[(*R*)-hydroxyalkanoic acid] depolymerase; poly(HA) depolymerase; poly(HA<sub>MCL</sub>) depolymerase; poly[(*R*)-3-hydroxyoctanoate] hydrolase  
**Systematic name:** polyoxycarbonyl[(*R*)-2-pentylethylene] hydrolase

**Comments:** The main product after prolonged incubation is the dimer [2566]. Besides hydrolysing polymers of 3-hydroxyoctanoic acid, the enzyme also hydrolyses other polymers derived from medium-chain-length (C<sub>6</sub>-C<sub>12</sub>) hydroxyalkanoic acids and copolymers of mixtures of these. It also hydrolyses *p*-nitrophenyl esters of fatty acids. Polymers of short-chain-length hydroxyalkanoic acids such as poly[(*R*)-3-hydroxybutanoic acid] and poly[(*R*)-3-hydroxypentanoic acid] are not hydrolysed.

**References:** [1316, 874, 2566]

[EC 3.1.1.76 created 2001, modified 2005]

#### EC 3.1.1.77

**Accepted name:** acyloxyacyl hydrolase

**Reaction:** 3-(acyloxy)acyl group of bacterial toxin + H<sub>2</sub>O = 3-hydroxyacyl group of bacterial toxin + a fatty acid

**Comments:** The substrate is lipid A on the reducing end of the toxic lipopolysaccharide (LPS) of *Salmonella typhimurium* and related organisms. It consists of diglucosamine, β-D-GlcN-(1→6)-D-GlcN, attached by glycosylation on O-6 of its non-reducing residue, phosphorylated on O-4 of this residue and on O-1 of its potentially reducing residue. Both residues carry 3-(acyloxy)acyl groups on N-2 and O-3. The enzyme from human leucocytes detoxifies the lipid by hydrolysing the secondary acyl groups from O-3 of the 3-hydroxyacyl groups on the disaccharide (LPS). It also possesses a wide range of phospholipase and acyltransferase activities [e.g. EC 3.1.1.4 (phospholipase A<sub>2</sub>), EC 3.1.1.5 (lysophospholipase), EC 3.1.1.32 (phospholipase A<sub>1</sub>) and EC 3.1.1.52 (phosphatidylinositol deacylase)], hydrolysing diacylglycerol and phosphatidyl compounds, but not triacylglycerols. It has a preference for saturated C<sub>12</sub>-C<sub>16</sub> acyl groups.

**References:** [715, 1021, 2002]

[EC 3.1.1.77 created 2001]

#### EC 3.1.1.78

**Accepted name:** polyneuridine-aldehyde esterase

**Reaction:** polyneuridine aldehyde + H<sub>2</sub>O = 16-epivellosimine + CO<sub>2</sub> + methanol

**Other name(s):** polyneuridine aldehyde esterase; PNAE

**Systematic name:** polyneuridine aldehyde hydrolase (decarboxylating)

**Comments:** Following hydrolysis of this indole alkaloid ester the carboxylic acid decarboxylates spontaneously giving the sarpagan skeleton. The enzyme also acts on akuammidine aldehyde (the 16-epimer of polyneuridine aldehyde).

**References:** [2270, 2271, 629, 1845]

[EC 3.1.1.78 created 2002]

#### EC 3.1.1.79

**Accepted name:** hormone-sensitive lipase

**Reaction:** (1) diacylglycerol + H<sub>2</sub>O = monoacylglycerol + a carboxylate

(2) triacylglycerol + H<sub>2</sub>O = diacylglycerol + a carboxylate

(3) monoacylglycerol + H<sub>2</sub>O = glycerol + a carboxylate

**Other name(s):** HSL

**Systematic name:** diacylglycerol acylhydrolase

**Comments:** This enzyme is a serine hydrolase. Compared with other lipases, hormone-sensitive lipase has a uniquely broad substrate specificity. It hydrolyses all acylglycerols (triacylglycerol, diacylglycerol and monoacylglycerol) [2,3,4] as well as cholesteryl esters [804, 2207], steroid fatty acid esters [1623], retinyl esters [3148] and *p*-nitrophenyl esters [2207, 2981]. It exhibits a preference for the 1- or 3-ester bond of its acylglycerol substrate compared with the 2-ester bond [3282]. The enzyme shows little preference for the fatty acids in the triacylglycerol, although there is some increase in activity with decreasing chain length. The enzyme activity is increased in response to hormones that elevate intracellular levels of cAMP.

**References:** [1163, 804, 3056, 2207, 1623, 3148, 2981, 3282]

[EC 3.1.1.79 created 2004]

#### EC 3.1.1.80

- Accepted name:** acetylajmaline esterase  
**Reaction:** (1) 17-*O*-acetylajmaline + H<sub>2</sub>O = ajmaline + acetate  
(2) 17-*O*-acetylnorajmaline + H<sub>2</sub>O = norajmaline + acetate  
**Other name(s):** AAE; 2β(*R*)-17-*O*-acetylajmalan:acetylesterease; acetylajmalan esterase  
**Systematic name:** 17-*O*-acetylajmaline *O*-acetylhydrolase  
**Comments:** This plant enzyme is responsible for the last stages in the biosynthesis of the indole alkaloid ajmaline. The enzyme is highly specific for the substrates 17-*O*-acetylajmaline and 17-*O*-acetylnorajmaline as the structurally related acetylated alkaloids vinorine, vomilenine, 1,2-dihydrovomilenine and 1,2-dihydroraucaffricine cannot act as substrates [2480]. This is a novel member of the GDSL family of serine esterases/lipases.  
**References:** [2308, 2480]

[EC 3.1.1.80 created 2006]

#### EC 3.1.1.81

- Accepted name:** quorum-quenching *N*-acyl-homoserine lactonase  
**Reaction:** an *N*-acyl-L-homoserine lactone + H<sub>2</sub>O = an *N*-acyl-L-homoserine  
**Other name(s):** acyl homoserine degrading enzyme; acyl-homoserine lactone acylase; AHL lactonase; AHL-degrading enzyme; AHL-inactivating enzyme; AHLase; AhlD; AhlK; AiiA; AiiA lactonase; AiiA-like protein; AiiB; AiiC; AttM; delactonase; lactonase-like enzyme; *N*-acyl homoserine lactonase; *N*-acyl homoserine lactone hydrolase; *N*-acyl-homoserine lactone lactonase; *N*-acyl-L-homoserine lactone hydrolase; quorum-quenching lactonase; quorum-quenching *N*-acyl homoserine lactone hydrolase  
**Systematic name:** *N*-acyl-L-homoserine-lactone lactonohydrolase  
**Comments:** Acyl-homoserine lactones (AHLs) are produced by a number of bacterial species and are used by them to regulate the expression of virulence genes in a process known as quorum-sensing. Each bacterial cell has a basal level of AHL and, once the population density reaches a critical level, it triggers AHL-signalling which, in turn, initiates the expression of particular virulence genes [637]. Plants or animals capable of degrading AHLs would have a therapeutic advantage in avoiding bacterial infection as they could prevent AHL-signalling and the expression of virulence genes in quorum-sensing bacteria [637]. *N*-(3-Oxohexanoyl)-L-homoserine lactone, *N*-(3-oxododecanoyl)-L-homoserine lactone, *N*-butanoyl-L-homoserine lactone and *N*-(3-oxooctanoyl)-L-homoserine lactone can act as substrates [637].  
**References:** [2915, 636, 3116, 638, 637, 1630, 2236, 3008, 1453, 1718, 3258]

[EC 3.1.1.81 created 2007]

#### EC 3.1.1.82

- Accepted name:** pheophorbidase  
**Reaction:** pheophorbide *a* + H<sub>2</sub>O = pyropheophorbide *a* + methanol + CO<sub>2</sub> (overall reaction)  
(1a) pheophorbide *a* + H<sub>2</sub>O = *C*-13<sup>2</sup>-carboxypyropheophorbide *a* + methanol  
(1b) *C*-13<sup>2</sup>-carboxypyropheophorbide *a* = pyropheophorbide *a* + CO<sub>2</sub> (spontaneous)  
**Other name(s):** phedase; PPD  
**Systematic name:** pheophorbide-*a* hydrolase

**Comments:** This enzyme forms part of the chlorophyll degradation pathway, and is found in higher plants and in algae. In higher plants it participates in de-greening processes such as fruit ripening, leaf senescence, and flowering. The enzyme exists in two forms: type 1 is induced by senescence whereas type 2 is constitutively expressed [2817, 2815]. The enzyme is highly specific for pheophorbide as substrate (with a preference for pheophorbide *a* over pheophorbide *b*) as other chlorophyll derivatives such as protochlorophyllide *a*, pheophytin *a* and *c*, chlorophyll *a* and *b*, and chlorophyllide *a* cannot act as substrates [2815]. Another enzyme, called pheophorbide demethoxycarbonylase (PDC), produces pyropheophorbide *a* from pheophorbide *a* without forming an intermediate although the precise reaction is not yet known [2817].

**References:** [2817, 2815, 1180]

[EC 3.1.1.82 created 2007]

### EC 3.1.1.83

**Accepted name:** monoterpene  $\epsilon$ -lactone hydrolase  
**Reaction:** (1) isoprop(en)ylmethyloxepan-2-one + H<sub>2</sub>O = 6-hydroxyisoprop(en)ylmethylhexanoate (general reaction)  
(2) 4-isopropenyl-7-methyloxepan-2-one + H<sub>2</sub>O = 6-hydroxy-3-isopropenylheptanoate  
(3) 7-isopropyl-4-methyloxepan-2-one + H<sub>2</sub>O = 6-hydroxy-3,7-dimethyloctanoate  
**Other name(s):** MLH  
**Systematic name:** isoprop(en)ylmethyloxepan-2-one lactonohydrolase  
**Comments:** The enzyme catalyses the ring opening of  $\epsilon$ -lactones which are formed during degradation of dihydrocarveol by the Gram-positive bacterium *Rhodococcus erythropolis* DCL14. The enzyme also acts on ethyl caproate, indicating that it is an esterase with a preference for lactones (internal cyclic esters). The enzyme is not stereoselective.  
**References:** [3040]

[EC 3.1.1.83 created 2008]

### EC 3.1.1.84

**Accepted name:** cocaine esterase  
**Reaction:** cocaine + H<sub>2</sub>O = ecgonine methyl ester + benzoate  
**Other name(s):** CocE; hCE2; hCE-2; human carboxylesterase 2  
**Systematic name:** cocaine benzoylhydrolase  
**Comments:** *Rhodococcus* sp. strain MB1 and *Pseudomonas maltophilia* strain MB11L can utilize cocaine as sole source of carbon and energy [287, 294].  
**References:** [871, 287, 294, 1605, 2285]

[EC 3.1.1.84 created 2010]

### EC 3.1.1.85

**Accepted name:** pimeloyl-[acyl-carrier protein] methyl ester esterase  
**Reaction:** pimeloyl-[acyl-carrier protein] methyl ester + H<sub>2</sub>O = pimeloyl-[acyl-carrier protein] + methanol  
**Other name(s):** BioH  
**Systematic name:** pimeloyl-[acyl-carrier protein] methyl ester hydrolase  
**Comments:** Involved in biotin biosynthesis in Gram-negative bacteria. The enzyme exhibits carboxylesterase activity, particularly toward substrates with short acyl chains [2518, 1644]. Even though the enzyme can interact with coenzyme A thioesters [2933], the *in vivo* role of the enzyme is to hydrolyse the methyl ester of pimeloyl-[acyl carrier protein], terminating the part of the biotin biosynthesis pathway that is catalysed by the fatty acid elongation enzymes [1693].  
**References:** [2518, 1644, 2933, 1693]

[EC 3.1.1.85 created 2011]

#### EC 3.1.1.86

**Accepted name:** rhamnogalacturonan acylesterase  
**Reaction:** Hydrolytic cleavage of 2-*O*-acetyl- or 3-*O*-acetyl groups of  $\alpha$ -D-galacturonic acid in rhamnogalacturonan I.  
**Other name(s):** RGAE  
**Systematic name:** rhamnogalacturonan 2/3-*O*-acetyl- $\alpha$ -D-galacturonate *O*-acetylhydrolase  
**Comments:** The degradation of rhamnogalacturonan by rhamnogalacturonases depends on the removal of the acetyl esters from the substrate [1404].  
**References:** [1404, 1950]

[EC 3.1.1.86 created 2011]

#### EC 3.1.1.87

**Accepted name:** fumonisin B1 esterase  
**Reaction:** fumonisin B1 + 2 H<sub>2</sub>O = aminopentol + 2 propane-1,2,3-tricarboxylate  
**Other name(s):** *fumD* (gene name)  
**Systematic name:** fumonisin B1 acylhydrolase  
**Comments:** The enzyme is involved in degradation of fumonisin B1 [1103].  
**References:** [1103]

[EC 3.1.1.87 created 2011]

#### EC 3.1.1.88

**Accepted name:** pyrethroid hydrolase  
**Reaction:** *trans*-permethrin + H<sub>2</sub>O = (3-phenoxyphenyl)methanol + (1*S*,3*R*)-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate  
**Other name(s):** pyrethroid-hydrolyzing carboxylesterase; pyrethroid-hydrolysing esterase; pyrethroid-hydrolyzing esterase; pyrethroid-selective esterase; pyrethroid-cleaving enzyme; permethrinase; PytH; EstP  
**Systematic name:** pyrethroid-ester hydrolase  
**Comments:** The enzyme is involved in degradation of pyrethroid pesticides. The enzymes from *Sphingobium* sp., *Klebsiella* sp. and *Aspergillus niger* hydrolyse *cis*-permethrin at approximately equal rate to *trans*-permethrin [3105, 3220, 1677]. The enzyme from mouse hydrolyses *trans*-permethrin at a rate about 22-fold higher than *cis*-permethrin [2778].  
**References:** [3105, 3220, 1677, 2778, 1794, 1001]

[EC 3.1.1.88 created 2011]

#### EC 3.1.1.89

**Accepted name:** protein phosphatase methyltransferase-1  
**Reaction:** [phosphatase 2A protein]-leucine methyl ester + H<sub>2</sub>O = [phosphatase 2A protein]-leucine + methanol  
**Other name(s):** PME-1; PPME1  
**Systematic name:** [phosphatase 2A protein]-leucine ester acylhydrolase  
**Comments:** A key regulator of protein phosphatase 2A. The methyl ester is formed by EC 2.1.1.233 (leucine carboxy methyltransferase-1). Occurs mainly in the nucleus.  
**References:** [2153, 3229]

[EC 3.1.1.89 created 2011]

#### EC 3.1.1.90

**Accepted name:** *all-trans*-retinyl ester 13-*cis* isomerohydrolase  
**Reaction:** an *all-trans*-retinyl ester + H<sub>2</sub>O = 13-*cis*-retinol + a fatty acid  
**Systematic name:** *all-trans*-retinyl ester acylhydrolase, 13-*cis* retinol forming

**Comments:** All-*trans*-retinyl esters, which are a storage form of vitamin A, are generated by the activity of EC 2.3.1.135, phosphatidylcholine—retinol *O*-acyltransferase (LRAT). They can be hydrolysed to 11-*cis*-retinol by EC 3.1.1.64, retinoid isomerohydrolase (RPE65), or to 13-*cis*-retinol by this enzyme.

**References:** [2849]

[EC 3.1.1.90 created 2011]

#### EC 3.1.1.91

**Accepted name:** 2-oxo-3-(5-oxofuran-2-ylidene)propanoate lactonase

**Reaction:** 2-oxo-3-(5-oxofuran-2-ylidene)propanoate + H<sub>2</sub>O = maleylpyruvate

**Other name(s):** *naaC* (gene name)

**Systematic name:** 2-oxo-3-(5-oxofuran-2-ylidene)propanoate lactonohydrolase

**Comments:** This enzyme, characterized from the soil bacterium *Bradyrhizobium* sp. JS329, is involved in the pathway of 5-nitroanthranilate degradation.

**References:** [2340]

[EC 3.1.1.91 created 2012]

#### EC 3.1.1.92

**Accepted name:** 4-sulfomuconolactone hydrolase

**Reaction:** 4-sulfomuconolactone + H<sub>2</sub>O = maleylacetate + sulfite

**Systematic name:** 4-sulfomuconolactone sulfohydrolase

**Comments:** The enzyme was isolated from the bacteria *Hydrogenophaga intermedia* and *Agrobacterium radiobacter* S2. It catalyses a step in the degradation of 4-sulfocatechol.

**References:** [1024]

[EC 3.1.1.92 created 2012]

#### EC 3.1.1.93

**Accepted name:** mycophenolic acid acyl-glucuronide esterase

**Reaction:** mycophenolic acid *O*-acyl-glucuronide + H<sub>2</sub>O = mycophenolate + D-glucuronate

**Other name(s):** mycophenolic acid acyl-glucuronide deglucuronidase; AcMPAG deglucuronidase

**Systematic name:** mycophenolic acid *O*-acyl-glucuronide-ester hydrolase

**Comments:** This liver enzyme deglucuronidates mycophenolic acid *O*-acyl-glucuronide, a metabolite of the immunosuppressant drug mycophenolate that is thought to be immunotoxic.

**References:** [1282]

[EC 3.1.1.93 created 2012]

#### EC 3.1.1.94

**Accepted name:** versiconal hemiacetal acetate esterase

**Reaction:** (1) versiconal hemiacetal acetate + H<sub>2</sub>O = versiconal + acetate

(2) versiconol acetate + H<sub>2</sub>O = versiconol + acetate

**Other name(s):** VHA esterase

**Systematic name:** versiconal-hemiacetal-acetate *O*-acetylhydrolase

**Comments:** Isolated from the mold *Aspergillus parasiticus*. Involved in a metabolic grid that leads to aflatoxin biosynthesis.

**References:** [1577, 396]

[EC 3.1.1.94 created 2013]

#### EC 3.1.1.95

**Accepted name:** aclacinomycin methylesterase  
**Reaction:** aclacinomycin T + H<sub>2</sub>O = 15-demethylaclacinomycin T + methanol  
**Other name(s):** RdmC; aclacinomycin methyl esterase  
**Systematic name:** aclacinomycin T acylhydrolase  
**Comments:** The enzyme is involved in the modification of the aklavinone skeleton in the biosynthesis of anthracyclines in *Streptomyces* species.  
**References:** [3123, 1309]

[EC 3.1.1.95 created 2013]

#### EC 3.1.1.96

**Accepted name:** D-aminoacyl-tRNA deacylase  
**Reaction:** a D-aminoacyl-tRNA + H<sub>2</sub>O = a D-amino acid + tRNA  
**Other name(s):** Dtd2; D-Tyr-tRNA(Tyr) deacylase; D-Tyr-tRNA<sup>Tyr</sup> deacylase; D-tyrosyl-tRNA<sup>Tyr</sup> aminoacylhydrolase; *dtdA* (gene name)  
**Systematic name:** D-aminoacyl-tRNA aminoacylhydrolase  
**Comments:** The enzyme from *Escherichia coli* can cleave D-tyrosyl-tRNA<sup>Tyr</sup>, D-aspartyl-tRNA<sup>Asp</sup> and D-tryptophanyl-tRNA<sup>Trp</sup> [2733]. Whereas the enzyme from the archaeon *Pyrococcus abyssi* is a zinc protein, the enzyme from *Escherichia coli* does not carry any zinc [760].  
**References:** [2733, 760, 759, 3225]

[EC 3.1.1.96 created 2014]

#### EC 3.1.1.97

**Accepted name:** methylated diphthine methylhydrolase  
**Reaction:** diphthine methyl ester-[translation elongation factor 2] + H<sub>2</sub>O = diphthine-[translation elongation factor 2] + methanol  
**Other name(s):** Dph7; diphthine methylesterase (incorrect)  
**Systematic name:** diphthine methyl ester acylhydrolase  
**Comments:** The protein is only present in eukaryotes.  
**References:** [1698]

[EC 3.1.1.97 created 2014, modified 2015]

#### EC 3.1.1.98

**Accepted name:** [Wnt protein] *O*-palmitoleoyl-L-serine hydrolase  
**Reaction:** [Wnt]-*O*-(9Z)-hexadec-9-enoyl-L-serine + H<sub>2</sub>O = [Wnt]-L-serine + (9Z)-hexadec-9-enoate  
**Other name(s):** Notum  
**Systematic name:** [Wnt]-*O*-(9Z)-hexadec-9-enoyl-L-serine acylhydrolase  
**Comments:** The enzyme removes the palmitoleate modification that is introduced to specific L-serine residues in Wnt proteins by EC 2.3.1.250, [Wnt protein]-*O*-palmitoleoyl transferase.  
**References:** [1359]

[EC 3.1.1.98 created 2015]

#### EC 3.1.1.99

**Accepted name:** 6-deoxy-6-sulfogluconolactonase  
**Reaction:** 6-deoxy-6-sulfo-D-glucono-1,5-lactone + H<sub>2</sub>O = 6-deoxy-6-sulfo-D-gluconate  
**Other name(s):** SGL lactonase  
**Systematic name:** 6-deoxy-6-sulfo-D-glucono-1,5-lactone lactonohydrolase  
**Comments:** The enzyme, characterized from the bacterium *Pseudomonas putida* SQ1, participates in a sulfoquinovose degradation pathway.  
**References:** [753]



[EC 3.1.1.99 created 2016]

#### EC 3.1.1.100

- Accepted name:** chlorophyllide *a* hydrolase  
**Reaction:** chlorophyllide *a* + H<sub>2</sub>O = 8-ethyl-12-methyl-3-vinyl-bacteriochlorophyllide *d* + methanol + CO<sub>2</sub>  
**Other name(s):** *bciC* (gene name)  
**Systematic name:** chlorophyllide-*a* hydrolase  
**Comments:** This enzyme, found in green sulfur bacteria (*Chlorobiaceae*) and green filamentous bacteria (*Chloroflexaceae*), catalyses the first committed step in the biosynthesis of bacteriochlorophylls *c*, *d* and *e*, the removal of the C-13<sup>2</sup>-methylcarboxyl group from chlorophyllide *a*. The reaction is very similar to the conversion of pheophorbide *a* to pyropheophorbide *a* during chlorophyll *a* degradation, which is catalysed by EC 3.1.1.82, pheophorbidease.  
**References:** [1727]

[EC 3.1.1.100 created 2016]

#### EC 3.1.1.101

- Accepted name:** poly(ethylene terephthalate) hydrolase  
**Reaction:** (ethylene terephthalate)<sub>*n*</sub> + H<sub>2</sub>O = (ethylene terephthalate)<sub>*n*-1</sub> + 4-[(2-hydroxyethoxy)carbonyl]benzoate  
**Other name(s):** PETase; PET hydrolase  
**Systematic name:** poly(ethylene terephthalate) hydrolase  
**Comments:** The enzyme, isolated from the bacterium *Ideonella sakaiensis*, also produces small amounts of terephthalate (*cf.* EC 3.1.1.102, mono(ethylene terephthalate) hydrolase). The reaction takes place on PET-film placed in solution.  
**References:** [3301]

[EC 3.1.1.101 created 2016]

#### EC 3.1.1.102

- Accepted name:** mono(ethylene terephthalate) hydrolase  
**Reaction:** 4-[(2-hydroxyethoxy)carbonyl]benzoate + H<sub>2</sub>O = terephthalate + ethylene glycol  
**Other name(s):** MHET hydrolase; MHETase  
**Systematic name:** 4-[(2-hydroxyethoxy)carbonyl]benzoate acylhydrolase  
**Comments:** The enzyme, isolated from the bacterium *Ideonella sakaiensis*, has no activity with poly(ethylene terephthalate) PET (*cf.* EC 3.1.1.101, poly(ethylene terephthalate) hydrolase).  
**References:** [3301]

[EC 3.1.1.102 created 2016]

#### EC 3.1.1.103

- Accepted name:** teichoic acid D-alanine hydrolase  
**Reaction:** [(4-D-Ala)-(2-GlcNAc)-Rib-ol-*P*]<sub>*n*</sub>-[Gro-*P*]<sub>*m*</sub>-ManNAc-GlcNAc-*PP*-peptidoglycan + *n* H<sub>2</sub>O = [(2-GlcNAc)-Rib-ol-*P*]<sub>*n*</sub>-[Gro-*P*]<sub>*m*</sub>-ManNAc-GlcNAc-*PP*-peptidoglycan + *n* D-alanine  
**Other name(s):** *fmtA* (gene name)  
**Systematic name:** teichoic acid D-alanylhydrolase  
**Comments:** The enzyme, characterized from the bacterium *Staphylococcus aureus*, removes D-alanine groups from the teichoic acid produced by this organism, thus modulating the electrical charge of the bacterial surface. The activity greatly increases methicillin resistance in MRSA strains.  
**References:** [1501, 2335, 2358]

[EC 3.1.1.103 created 2018]



#### EC 3.1.1.104

**Accepted name:** 5-phospho-D-xylono-1,4-lactonase  
**Reaction:** (1) D-xylono-1,4-lactone 5-phosphate + H<sub>2</sub>O = 5-phospho-D-xylonate  
(2) L-arabino-1,4-lactone 5-phosphate + H<sub>2</sub>O = 5-phospho-L-arabininate  
**Systematic name:** 5-phospho-D-xylono-1,4-lactone hydrolase  
**Comments:** The enzyme, characterized from *Mycoplasma* spp., contains a binuclear metal center with two zinc cations. The enzyme is specific for the phosphorylated forms, and is unable to hydrolyse non-phosphorylated 1,4-lactones.  
**References:** [1510]

[EC 3.1.1.104 created 2018]

### EC 3.1.2 Thioester hydrolases

#### EC 3.1.2.1

**Accepted name:** acetyl-CoA hydrolase  
**Reaction:** acetyl-CoA + H<sub>2</sub>O = CoA + acetate  
**Other name(s):** acetyl-CoA deacylase; acetyl-CoA acylase; acetyl coenzyme A hydrolase; acetyl coenzyme A deacylase; acetyl coenzyme A acylase; acetyl-CoA thiol esterase  
**Systematic name:** acetyl-CoA hydrolase  
**References:** [894]

[EC 3.1.2.1 created 1961]

#### EC 3.1.2.2

**Accepted name:** palmitoyl-CoA hydrolase  
**Reaction:** palmitoyl-CoA + H<sub>2</sub>O = CoA + palmitate  
**Other name(s):** long-chain fatty-acyl-CoA hydrolase; palmitoyl coenzyme A hydrolase; palmitoyl thioesterase; palmitoyl coenzyme A hydrolase; palmitoyl-CoA deacylase; palmitoyl thioesterase; palmitoyl-CoA deacylase; fatty acyl thioesterase I; palmitoyl thioesterase I  
**Systematic name:** palmitoyl-CoA hydrolase  
**Comments:** Also hydrolyses CoA thioesters of other long-chain fatty acids.  
**References:** [146, 193, 1939, 2745, 3238]

[EC 3.1.2.2 created 1961]

#### EC 3.1.2.3

**Accepted name:** succinyl-CoA hydrolase  
**Reaction:** succinyl-CoA + H<sub>2</sub>O = CoA + succinate  
**Other name(s):** succinyl-CoA acylase; succinyl coenzyme A hydrolase; succinyl coenzyme A deacylase  
**Systematic name:** succinyl-CoA hydrolase  
**References:** [894]

[EC 3.1.2.3 created 1961]

#### EC 3.1.2.4

**Accepted name:** 3-hydroxyisobutyryl-CoA hydrolase  
**Reaction:** 3-hydroxy-2-methylpropanoyl-CoA + H<sub>2</sub>O = CoA + 3-hydroxy-2-methylpropanoate  
**Other name(s):** 3-hydroxy-isobutyryl CoA hydrolase; HIB CoA deacylase  
**Systematic name:** 3-hydroxy-2-methylpropanoyl-CoA hydrolase  
**Comments:** Also hydrolyses 3-hydroxypropanoyl-CoA.  
**References:** [2408]

[EC 3.1.2.4 created 1961]

#### EC 3.1.2.5

**Accepted name:** hydroxymethylglutaryl-CoA hydrolase  
**Reaction:** (S)-3-hydroxy-3-methylglutaryl-CoA + H<sub>2</sub>O = CoA + 3-hydroxy-3-methylglutarate  
**Other name(s):** β-hydroxy-β-methylglutaryl coenzyme A hydrolase; β-hydroxy-β-methylglutaryl coenzyme A deacylase; hydroxymethylglutaryl coenzyme A hydrolase; hydroxymethylglutaryl coenzyme A deacylase; 3-hydroxy-3-methylglutaryl-CoA hydrolase  
**Systematic name:** (S)-3-hydroxy-3-methylglutaryl-CoA hydrolase  
**References:** [570]

[EC 3.1.2.5 created 1961]

#### EC 3.1.2.6

**Accepted name:** hydroxyacylglutathione hydrolase  
**Reaction:** S-(2-hydroxyacyl)glutathione + H<sub>2</sub>O = glutathione + a 2-hydroxy carboxylate  
**Other name(s):** glyoxalase II; S-2-hydroxylacylglutathione hydrolase; hydroxyacylglutathione hydrolase; acetoacetylglutathione hydrolase  
**Systematic name:** S-(2-hydroxyacyl)glutathione hydrolase  
**Comments:** Also hydrolyses S-acetoacetylglutathione, but more slowly.  
**References:** [2352, 3014, 3015]

[EC 3.1.2.6 created 1961 (EC 3.1.2.8 created 1961, incorporated 1978)]

#### EC 3.1.2.7

**Accepted name:** glutathione thioesterase  
**Reaction:** S-acylglutathione + H<sub>2</sub>O = glutathione + a carboxylate  
**Other name(s):** citryl-glutathione thioesterhydrolase  
**Systematic name:** S-acylglutathione hydrolase  
**References:** [1438]

[EC 3.1.2.7 created 1961]

[3.1.2.8 Deleted entry. *S-acetoacetylglutathione hydrolase*. Now included with EC 3.1.2.6 hydroxyacylglutathione hydrolase]

[EC 3.1.2.8 created 1961, deleted 1978]

[3.1.2.9 Deleted entry. *S-acetoacetylhydrolipoate hydrolase*]

[EC 3.1.2.9 created 1961, deleted 1964]

#### EC 3.1.2.10

**Accepted name:** formyl-CoA hydrolase  
**Reaction:** formyl-CoA + H<sub>2</sub>O = CoA + formate  
**Other name(s):** formyl coenzyme A hydrolase  
**Systematic name:** formyl-CoA hydrolase  
**References:** [2691]

[EC 3.1.2.10 created 1965]

#### EC 3.1.2.11

**Accepted name:** acetoacetyl-CoA hydrolase  
**Reaction:** acetoacetyl-CoA + H<sub>2</sub>O = CoA + acetoacetate  
**Other name(s):** acetoacetyl coenzyme A hydrolase; acetoacetyl CoA deacylase; acetoacetyl coenzyme A deacylase

**Systematic name:** acetoacetyl-CoA hydrolase

**References:** [64, 655]

[EC 3.1.2.11 created 1972]

#### EC 3.1.2.12

**Accepted name:** S-formylglutathione hydrolase

**Reaction:** S-formylglutathione + H<sub>2</sub>O = glutathione + formate

**Systematic name:** S-formylglutathione hydrolase

**Comments:** Also hydrolyses S-acetylglutathione, but more slowly.

**References:** [3014, 3017, 1056]

[EC 3.1.2.12 created 1978]

#### EC 3.1.2.13

**Accepted name:** S-succinylglutathione hydrolase

**Reaction:** S-succinylglutathione + H<sub>2</sub>O = glutathione + succinate

**Systematic name:** S-succinylglutathione hydrolase

**References:** [3014, 3016]

[EC 3.1.2.13 created 1978]

#### EC 3.1.2.14

**Accepted name:** oleoyl-[acyl-carrier-protein] hydrolase

**Reaction:** an oleoyl-[acyl-carrier protein] + H<sub>2</sub>O = an [acyl-carrier protein] + oleate

**Other name(s):** acyl-[acyl-carrier-protein] hydrolase; acyl-ACP-hydrolase; acyl-acyl carrier protein hydrolase; oleoyl-ACP thioesterase; oleoyl-acyl carrier protein thioesterase; oleoyl-[acyl-carrier-protein] hydrolase

**Systematic name:** oleoyl-[acyl-carrier-protein] hydrolase

**Comments:** Acts on acyl-carrier-protein thioesters of fatty acids from C<sub>12</sub> to C<sub>18</sub>, but the derivative of oleic acid is hydrolysed much more rapidly than any other compound tested.

**References:** [2162, 2639]

[EC 3.1.2.14 created 1984]

[3.1.2.15 Deleted entry. This activity is covered by EC 3.4.19.12, ubiquitinyl hydrolase 1]

[EC 3.1.2.15 created 1986, deleted 2014]

#### EC 3.1.2.16

**Accepted name:** citrate-lyase deacetylase

**Reaction:** acetyl-[citrate (*pro*-3*S*)-lyase] + H<sub>2</sub>O = holo-[citrate (*pro*-3*S*)-lyase] + acetate

**Other name(s):** [citrate-(*pro*-3*S*)-lyase] thioesterase; acetyl-S-(acyl-carrier protein) enzyme thioester hydrolase; citrate lyase deacetylase; [citrate-(*pro*-3*S*)-lyase](acetyl-form) hydrolase

**Systematic name:** acetyl-[citrate-(*pro*-3*S*)-lyase] hydrolase

**Comments:** In the proteobacterium *Rubrivivax gelatinosus*, this enzyme modulates the activity of EC 4.1.3.6, citrate (*pro*-3*S*)-lyase, by converting it from its active acetyl form into its inactive thiol form by removal of its acetyl groups [915]. The activity of citrate-lyase deacetylase is itself inhibited by L-glutamate [915].

**References:** [914, 915]

[EC 3.1.2.16 created 1989]

#### EC 3.1.2.17

**Accepted name:** (S)-methylmalonyl-CoA hydrolase  
**Reaction:** (S)-methylmalonyl-CoA + H<sub>2</sub>O = methylmalonate + CoA  
**Other name(s):** D-methylmalonyl-coenzyme A hydrolase  
**Systematic name:** (S)-methylmalonyl-CoA hydrolase  
**References:** [1524]

[EC 3.1.2.17 created 1989]

#### EC 3.1.2.18

**Accepted name:** ADP-dependent short-chain-acyl-CoA hydrolase  
**Reaction:** acyl-CoA + H<sub>2</sub>O = CoA + a carboxylate  
**Other name(s):** short-chain acyl coenzyme A hydrolase; propionyl coenzyme A hydrolase; propionyl-CoA hydrolase; propionyl-CoA thioesterase; short-chain acyl-CoA hydrolase; short-chain acyl-CoA thioesterase  
**Systematic name:** ADP-dependent-short-chain-acyl-CoA hydrolase  
**Comments:** Requires ADP; inhibited by NADH. Maximum activity is shown with propanoyl-CoA.  
**References:** [32, 33]

[EC 3.1.2.18 created 1992]

#### EC 3.1.2.19

**Accepted name:** ADP-dependent medium-chain-acyl-CoA hydrolase  
**Reaction:** acyl-CoA + H<sub>2</sub>O = CoA + a carboxylate  
**Other name(s):** medium-chain acyl coenzyme A hydrolase; medium-chain acyl-CoA hydrolase; medium-chain acyl-thioester hydrolase; medium-chain hydrolase; myristoyl-CoA thioesterase  
**Systematic name:** ADP-dependent-medium-chain-acyl-CoA hydrolase  
**Comments:** Requires ADP; inhibited by NADH. Maximum activity is shown with nonanoyl-CoA.  
**References:** [32]

[EC 3.1.2.19 created 1992]

#### EC 3.1.2.20

**Accepted name:** acyl-CoA hydrolase  
**Reaction:** acyl-CoA + H<sub>2</sub>O = CoA + a carboxylate  
**Other name(s):** acyl coenzyme A thioesterase; acyl-CoA thioesterase; acyl coenzyme A hydrolase; thioesterase B; thioesterase II; acyl-CoA thioesterase  
**Systematic name:** acyl-CoA hydrolase  
**Comments:** Broad specificity for medium- to long-chain acyl-CoA. Insensitive to NAD<sup>+</sup> (*cf.* EC 3.1.2.19 ADP-dependent medium-chain-acyl-CoA hydrolase)  
**References:** [33]

[EC 3.1.2.20 created 1992]

#### EC 3.1.2.21

**Accepted name:** dodecanoyl-[acyl-carrier-protein] hydrolase  
**Reaction:** a dodecanoyl-[acyl-carrier protein] + H<sub>2</sub>O = an [acyl-carrier protein] + dodecanoate  
**Other name(s):** lauryl-acyl-carrier-protein hydrolase; dodecanoyl-acyl-carrier-protein hydrolase; dodecyl-acyl-carrier protein hydrolase; dodecanoyl-[acyl-carrier protein] hydrolase; dodecanoyl-[acyl-carrier-protein] hydrolase  
**Systematic name:** dodecanoyl-[acyl-carrier protein] hydrolase  
**Comments:** Acts on the acyl-carrier-protein thioester of C<sub>12</sub> and, with a much lower activity, C<sub>14</sub> fatty acids. The derivative of oleic acid is hydrolysed very slowly (*cf.* EC 3.1.2.14, oleoyl-[acyl-carrier-protein] hydrolase).  
**References:** [2305, 547]

[EC 3.1.2.21 created 1999]

#### EC 3.1.2.22

**Accepted name:** palmitoyl[protein] hydrolase  
**Reaction:** palmitoyl[protein] + H<sub>2</sub>O = palmitate + protein  
**Other name(s):** palmitoyl-protein thioesterase; palmitoyl-(protein) hydrolase  
**Systematic name:** palmitoyl[protein] hydrolase  
**Comments:** Specific for long-chain thioesters of fatty acids. Hydrolyses fatty acids from *S*-acylated cysteine residues in proteins, palmitoyl cysteine and palmitoyl-CoA.  
**References:** [358, 2584, 3061]

[EC 3.1.2.22 created 1999]

#### EC 3.1.2.23

**Accepted name:** 4-hydroxybenzoyl-CoA thioesterase  
**Reaction:** 4-hydroxybenzoyl-CoA + H<sub>2</sub>O = 4-hydroxybenzoate + CoA  
**Systematic name:** 4-hydroxybenzoyl-CoA hydrolase  
**Comments:** This enzyme is part of the bacterial 2,4-dichlorobenzoate degradation pathway.  
**References:** [394, 666]

[EC 3.1.2.23 created 1999]

[3.1.2.24 *Transferred entry. 2-(2-hydroxyphenyl)benzenesulfinate hydrolase. Now EC 3.13.1.3, 2'-hydroxybiphenyl-2-sulfinate desulfinase. The enzyme was incorrectly classified as a thioester hydrolase when the bond broken is a C-S bond, which is not an ester*]

[EC 3.1.2.24 created 2000, deleted 2005]

#### EC 3.1.2.25

**Accepted name:** phenylacetyl-CoA hydrolase  
**Reaction:** phenylglyoxyl-CoA + H<sub>2</sub>O = phenylglyoxylate + CoA  
**Systematic name:** phenylglyoxyl-CoA hydrolase  
**Comments:** This is the second step in the conversion of phenylacetyl-CoA to phenylglyoxylate, the first step being carried out by EC 1.17.5.1, phenylacetyl-CoA dehydrogenase.  
**References:** [2415, 2576]

[EC 3.1.2.25 created 2004]

[3.1.2.26 *Transferred entry. bile-acid-CoA hydrolase. Now EC 2.8.3.25, bile acid CoA transferase*]

[EC 3.1.2.26 created 2005, deleted 2016]

#### EC 3.1.2.27

**Accepted name:** choloyl-CoA hydrolase  
**Reaction:** choloyl-CoA + H<sub>2</sub>O = cholate + CoA  
**Other name(s):** PTE-2 (ambiguous); choloyl-coenzyme A thioesterase; chenodeoxycholoyl-coenzyme A thioesterase; peroxisomal acyl-CoA thioesterase 2  
**Systematic name:** choloyl-CoA hydrolase  
**Comments:** Also acts on chenodeoxycholoyl-CoA and to a lesser extent on short- and medium- to long-chain acyl-CoAs, and other substrates, including trihydroxycoprostanoyl-CoA, hydroxymethylglutaryl-CoA and branched chain acyl-CoAs, all of which are present in peroxisomes. The enzyme is strongly inhibited by CoA and may be involved in controlling CoA levels in the peroxisome [1208].  
**References:** [1208, 2710, 2481]

[EC 3.1.2.27 created 2005]

#### EC 3.1.2.28

**Accepted name:** 1,4-dihydroxy-2-naphthoyl-CoA hydrolase  
**Reaction:** 1,4-dihydroxy-2-naphthoyl-CoA + H<sub>2</sub>O = 1,4-dihydroxy-2-naphthoate + CoA  
**Other name(s):** *menI* (gene name); *ydiL* (gene name)  
**Systematic name:** 1,4-dihydroxy-2-naphthoyl-CoA hydrolase  
**Comments:** This enzyme participates in the synthesis of menaquinones [413], phyloquinone [3180], as well as several plant pigments [1999, 684]. The enzyme from the cyanobacterium *Synechocystis* sp. PCC 6803 does not accept benzoyl-CoA or phenylacetyl-CoA as substrates [3180].  
**References:** [1999, 684, 3180, 413]

[EC 3.1.2.28 created 2010]

#### EC 3.1.2.29

**Accepted name:** fluoroacetyl-CoA thioesterase  
**Reaction:** fluoroacetyl-CoA + H<sub>2</sub>O = fluoroacetate + CoA  
**Systematic name:** fluoroacetyl-CoA hydrolase  
**Comments:** Fluoroacetate is extremely toxic. It reacts with CoA to form fluoroacetyl-CoA, which substitutes for acetyl CoA and reacts with EC 2.3.3.1 (citrate synthase) to produce fluorocitrate, a metabolite of which binds very tightly to EC 4.2.1.3 (aconitase) and halts the TCA cycle. This enzyme hydrolyses fluoroacetyl-CoA before it can react with citrate synthase, and thus confers fluoroacetate resistance on the organisms that produce it. It has been described in the poisonous plant *Dichapetalum cymosum* and the bacterium *Streptomyces cattleya*, both of which are fluoroacetate producers.  
**References:** [1895, 1202, 600]

[EC 3.1.2.29 created 2011]

#### EC 3.1.2.30

**Accepted name:** (3S)-malyl-CoA thioesterase  
**Reaction:** (S)-malyl-CoA + H<sub>2</sub>O = (S)-malate + CoA  
**Other name(s):** *mcl2* (gene name)  
**Systematic name:** (S)-malyl-CoA hydrolase  
**Comments:** Stimulated by Mg<sup>2+</sup> or Mn<sup>2+</sup>. The enzyme has no activity with (2R,3S)-2-methylmalyl-CoA (*cf.* EC 4.1.3.24, malyl-CoA lyase) or other CoA esters.  
**References:** [710]

[EC 3.1.2.30 created 2014]

#### EC 3.1.2.31

**Accepted name:** dihydromonacolin L-[lovastatin nonaketide synthase] thioesterase  
**Reaction:** dihydromonacolin L-[lovastatin nonaketide synthase] + H<sub>2</sub>O = holo-[lovastatin nonaketide synthase] + dihydromonacolin L acid  
**Other name(s):** LovG  
**Systematic name:** dihydromonacolin L-[lovastatin nonaketide synthase] hydrolase  
**Comments:** Dihydromonacolin L acid is synthesized while bound to an acyl-carrier protein domain of the lovastatin nonaketide synthase (EC 2.3.1.161). Since that enzyme lacks a thioesterase domain, release of the dihydromonacolin L acid moiety from the polyketide synthase requires this dedicated enzyme.  
**References:** [3235]

[EC 3.1.2.31 created 2015]

#### EC 3.1.2.32

**Accepted name:** 2-aminobenzoylacetyl-CoA thioesterase  
**Reaction:** (2-aminobenzoyl)acetyl-CoA + H<sub>2</sub>O = (2-aminobenzoyl)acetate + CoA  
**Other name(s):** *pqsE* (gene name)  
**Systematic name:** (2-aminobenzoyl)acetyl-CoA hydrolase  
**Comments:** The enzyme, characterized from the bacterium *Pseudomonas aeruginosa*, participates in the production of the signal molecule 2-heptyl-4(1*H*)-quinolone (HHQ).  
**References:** [3310, 651]

[EC 3.1.2.32 created 2016]

### EC 3.1.3 Phosphoric-monoester hydrolases

#### EC 3.1.3.1

**Accepted name:** alkaline phosphatase  
**Reaction:** a phosphate monoester + H<sub>2</sub>O = an alcohol + phosphate  
**Other name(s):** alkaline phosphomonoesterase; phosphomonoesterase; glycerophosphatase; alkaline phosphohydrolase; alkaline phenyl phosphatase; orthophosphoric-monoester phosphohydrolase (alkaline optimum)  
**Systematic name:** phosphate-monoester phosphohydrolase (alkaline optimum)  
**Comments:** Wide specificity. Also catalyses transphosphorylations. The human placental enzyme is a zinc protein. Some enzymes hydrolyse diphosphate (*cf.* EC 3.6.1.1 inorganic diphosphatase)  
**References:** [706, 1054, 1785, 1984, 2752]

[EC 3.1.3.1 created 1961]

#### EC 3.1.3.2

**Accepted name:** acid phosphatase  
**Reaction:** a phosphate monoester + H<sub>2</sub>O = an alcohol + phosphate  
**Other name(s):** acid phosphomonoesterase; phosphomonoesterase; glycerophosphatase; acid monophosphatase; acid phosphohydrolase; acid phosphomonoester hydrolase; uteroferrin; acid nucleoside diphosphate phosphatase; orthophosphoric-monoester phosphohydrolase (acid optimum)  
**Systematic name:** phosphate-monoester phosphohydrolase (acid optimum)  
**Comments:** Wide specificity. Also catalyses transphosphorylations.  
**References:** [1345, 1561, 2976]

[EC 3.1.3.2 created 1961]

#### EC 3.1.3.3

**Accepted name:** phosphoserine phosphatase  
**Reaction:** *O*-phospho-L(or D)-serine + H<sub>2</sub>O = L(or D)-serine + phosphate  
**Systematic name:** *O*-phosphoserine phosphohydrolase  
**References:** [266, 344, 2070]

[EC 3.1.3.3 created 1961]

#### EC 3.1.3.4

**Accepted name:** phosphatidate phosphatase  
**Reaction:** a 1,2-diacylglycerol 3-phosphate + H<sub>2</sub>O = a 1,2-diacyl-*sn*-glycerol + phosphate  
**Other name(s):** phosphatic acid phosphatase; acid phosphatidyl phosphatase; phosphatic acid phosphohydrolase; PAP, Lipin  
**Systematic name:** diacylglycerol-3-phosphate phosphohydrolase

**Comments:** This enzyme catalyses the  $Mg^{2+}$ -dependent dephosphorylation of a 1,2-diacylglycerol-3-phosphate, yielding a 1,2-diacyl-*sn*-glycerol (DAG), the substrate for *de novo* lipid synthesis via the Kennedy pathway and for the synthesis of triacylglycerol. In lipid signalling, the enzyme generates a pool of DAG to be used for protein kinase C activation. The mammalian enzymes are known as lipins.

**References:** [2700, 373]

[EC 3.1.3.4 created 1961, modified 2010]

#### EC 3.1.3.5

**Accepted name:** 5'-nucleotidase

**Reaction:** a 5'-ribonucleotide + H<sub>2</sub>O = a ribonucleoside + phosphate

**Other name(s):** uridine 5'-nucleotidase; 5'-adenylic phosphatase; adenosine 5'-phosphatase; AMP phosphatase; adenosine monophosphatase; 5'-mononucleotidase; AMPase; UMPase; snake venom 5'-nucleotidase; thymidine monophosphate nucleotidase; 5'-AMPase; 5'-AMP nucleotidase; AMP phosphohydrolase; IMP 5'-nucleotidase

**Systematic name:** 5'-ribonucleotide phosphohydrolase

**Comments:** Wide specificity for 5'-nucleotides.

**References:** [999, 1121, 2603]

[EC 3.1.3.5 created 1961]

#### EC 3.1.3.6

**Accepted name:** 3'-nucleotidase

**Reaction:** a 3'-ribonucleotide + H<sub>2</sub>O = a ribonucleoside + phosphate

**Other name(s):** 3'-mononucleotidase; 3'-phosphatase; 3'-ribonucleotidase

**Systematic name:** 3'-ribonucleotide phosphohydrolase

**Comments:** Wide specificity for 3'-nucleotides.

**References:** [2650]

[EC 3.1.3.6 created 1961]

#### EC 3.1.3.7

**Accepted name:** 3'(2'),5'-bisphosphate nucleotidase

**Reaction:** adenosine 3',5'-bisphosphate + H<sub>2</sub>O = AMP + phosphate

**Other name(s):** phosphoadenylate 3'-nucleotidase; 3'-phosphoadenylylsulfate 3'-phosphatase; 3'(2'),5'-bisphosphonucleoside 3'(2')-phosphohydrolase

**Systematic name:** adenosine-3'(2'),5'-bisphosphate 3'(2')-phosphohydrolase

**Comments:** Also acts on 3'-phosphoadenylyl sulfate, and on the corresponding 2'-phosphates.

**References:** [318, 740, 2366, 2971]

[EC 3.1.3.7 created 1961]

#### EC 3.1.3.8

**Accepted name:** 3-phytase

**Reaction:** *myo*-inositol hexakisphosphate + H<sub>2</sub>O = 1D-*myo*-inositol 1,2,4,5,6-pentakisphosphate + phosphate

**Other name(s):** 1-phytase; phytase; phytate 1-phosphatase; phytate 6-phosphatase

**Systematic name:** *myo*-inositol-hexakisphosphate 3-phosphohydrolase

**References:** [491, 1328, 1255, 492]

[EC 3.1.3.8 created 1961, modified 1976, modified 2002]

#### EC 3.1.3.9



**Accepted name:** glucose-6-phosphatase  
**Reaction:** D-glucose 6-phosphate + H<sub>2</sub>O = D-glucose + phosphate  
**Other name(s):** glucose 6-phosphate phosphatase  
**Systematic name:** D-glucose-6-phosphate phosphohydrolase  
**Comments:** Wide distribution in animal tissues. Also catalyses potent transphosphorylations from carbamoyl phosphate, hexose phosphates, diphosphate, phospho*enol*pyruvate and nucleoside di- and triphosphates, to D-glucose, D-mannose, 3-methyl-D-glucose or 2-deoxy-D-glucose [*cf.* EC 2.7.1.62 (phosphoramidate—hexose phosphotransferase), EC 2.7.1.79 (diphosphate—glycerol phosphotransferase) and EC 3.9.1.1 (phosphoamidase)].  
**References:** [50, 466, 2110, 2111]

[EC 3.1.3.9 created 1961]

#### EC 3.1.3.10

**Accepted name:** glucose-1-phosphatase  
**Reaction:** α-D-glucose 1-phosphate + H<sub>2</sub>O = D-glucose + phosphate  
**Systematic name:** α-D-glucose-1-phosphate phosphohydrolase  
**Comments:** Also acts, more slowly, on D-galactose 1-phosphate.  
**References:** [747, 2999]

[EC 3.1.3.10 created 1961]

#### EC 3.1.3.11

**Accepted name:** fructose-bisphosphatase  
**Reaction:** D-fructose 1,6-bisphosphate + H<sub>2</sub>O = D-fructose 6-phosphate + phosphate  
**Other name(s):** hexose diphosphatase; FBPase; fructose 1,6-diphosphatase; fructose 1,6-diphosphate phosphatase; D-fructose 1,6-diphosphatase; fructose 1,6-bisphosphatase; fructose diphosphatase; fructose diphosphate phosphatase; fructose bisphosphate phosphatase; fructose 1,6-bisphosphate 1-phosphatase; fructose 1,6-bisphosphate phosphatase; hexose bisphosphatase; D-fructose-1,6-bisphosphate phosphatase  
**Systematic name:** D-fructose-1,6-bisphosphate 1-phosphohydrolase  
**Comments:** The animal enzyme also acts on sedoheptulose 1,7-bisphosphate.  
**References:** [690, 954, 1949, 2310]

[EC 3.1.3.11 created 1961, modified 1976]

#### EC 3.1.3.12

**Accepted name:** trehalose-phosphatase  
**Reaction:** α,α-trehalose 6-phosphate + H<sub>2</sub>O = α,α-trehalose + phosphate  
**Other name(s):** trehalose 6-phosphatase; trehalose 6-phosphate phosphatase; trehalose-6-phosphate phosphohydrolase  
**Systematic name:** α,α-trehalose-6-phosphate phosphohydrolase  
**References:** [348, 365]

[EC 3.1.3.12 created 1961]

[3.1.3.13 Deleted entry. bisphosphoglycerate phosphatase. Recent studies have shown that this is a partial activity of EC 5.4.2.11, phosphoglycerate mutase (2,3-diphosphoglycerate-dependent)]

[EC 3.1.3.13 created 1961, deleted 2016]

#### EC 3.1.3.14

**Accepted name:** methylphosphothioglycerate phosphatase  
**Reaction:** S-methyl-3-phospho-1-thio-D-glycerate + H<sub>2</sub>O = S-methyl-1-thio-D-glycerate + phosphate  
**Other name(s):** methylthiophosphoglycerate phosphatase  
**Systematic name:** S-methyl-3-phospho-1-thio-D-glycerate phosphohydrolase  
**References:** [229]

[EC 3.1.3.14 created 1961]

#### EC 3.1.3.15

**Accepted name:** histidinol-phosphatase  
**Reaction:** L-histidinol phosphate + H<sub>2</sub>O = L-histidinol + phosphate  
**Other name(s):** histidinol phosphate phosphatase; L-histidinol phosphate phosphatase; histidinolphosphate phosphatase; HPPase; histidinolphosphatase  
**Systematic name:** L-histidinol-phosphate phosphohydrolase  
**References:** [46]

[EC 3.1.3.15 created 1961]

#### EC 3.1.3.16

**Accepted name:** protein-serine/threonine phosphatase  
**Reaction:** [a protein]-serine/threonine phosphate + H<sub>2</sub>O = [a protein]-serine/threonine + phosphate  
**Other name(s):** phosphoprotein phosphatase (ambiguous); protein phosphatase-1; protein phosphatase-2A; protein phosphatase-2B; protein phosphatase-2C; protein D phosphatase; phosphospectrin phosphatase; casein phosphatase; *Aspergillus awamori* acid protein phosphatase; calcineurin; phosphatase 2A; phosphatase 2B; phosphatase II; phosphatase IB; phosphatase C-II; polycation modulated (PCM-) phosphatase; phosphopyruvate dehydrogenase phosphatase; phosphatase SP; branched-chain  $\alpha$ -keto acid dehydrogenase phosphatase; BCKDH phosphatase; 3-hydroxy 3-methylglutaryl coenzymeA reductase phosphatase; HMG-CoA reductase phosphatase; phosphatase H-II; phosphatase III; phosphatase I; protein phosphatase; phosphatase IV; phosphoprotein phosphohydrolase  
**Systematic name:** protein-serine/threonine-phosphate phosphohydrolase  
**Comments:** A group of enzymes removing the serine- or threonine-bound phosphate group from a wide range of phosphoproteins, including a number of enzymes that have been phosphorylated under the action of a kinase (*cf.* EC 3.1.3.48 protein-tyrosine-phosphatase). The spleen enzyme also acts on phenolic phosphates and phosphamides (*cf.* EC 3.9.1.1, phosphoamidase).  
**References:** [595, 1249, 2801, 2945]

[EC 3.1.3.16 created 1961, modified 1989, modified 2013]

#### EC 3.1.3.17

**Accepted name:** [phosphorylase] phosphatase  
**Reaction:** [phosphorylase *a*] + 4 H<sub>2</sub>O = 2 [phosphorylase *b*] + 4 phosphate  
**Other name(s):** PR-enzyme; phosphorylase *a* phosphatase; glycogen phosphorylase phosphatase; protein phosphatase C; type 1 protein phosphatase  
**Systematic name:** [phosphorylase *a*] phosphohydrolase  
**References:** [277, 966, 2363]

[EC 3.1.3.17 created 1961]

#### EC 3.1.3.18

**Accepted name:** phosphoglycolate phosphatase  
**Reaction:** 2-phosphoglycolate + H<sub>2</sub>O = glycolate + phosphate  
**Other name(s):** phosphoglycolate hydrolase; 2-phosphoglycolate phosphatase; P-glycolate phosphatase; phosphoglycollate phosphatase  
**Systematic name:** 2-phosphoglycolate phosphohydrolase  
**References:** [441]

[EC 3.1.3.18 created 1965]

#### EC 3.1.3.19

**Accepted name:** glycerol-2-phosphatase  
**Reaction:** glycerol 2-phosphate + H<sub>2</sub>O = glycerol + phosphate  
**Other name(s):** β-glycerophosphatase; β-glycerophosphate phosphatase; 2-glycerophosphatase  
**Systematic name:** glycerol-2-phosphate phosphohydrolase  
**References:** [2574, 2976]

[EC 3.1.3.19 created 1965]

#### EC 3.1.3.20

**Accepted name:** phosphoglycerate phosphatase  
**Reaction:** D-glycerate 2-phosphate + H<sub>2</sub>O = D-glycerate + phosphate  
**Other name(s):** D-2-phosphoglycerate phosphatase; glycerophosphate phosphatase  
**Systematic name:** D-glycerate-2-phosphate phosphohydrolase  
**References:** [735]

[EC 3.1.3.20 created 1972]

#### EC 3.1.3.21

**Accepted name:** glycerol-1-phosphatase  
**Reaction:** glycerol 1-phosphate + H<sub>2</sub>O = glycerol + phosphate  
**Other name(s):** α-glycerophosphatase; α-glycerol phosphatase; glycerol 3-phosphatase; glycerol-3-phosphate phosphatase; glycerol 3-phosphate phosphohydrolase  
**Systematic name:** glycerol-1-phosphate phosphohydrolase  
**Comments:** The *Dunaliella* enzyme acts more rapidly on *sn*-glycerol 1-phosphate than on the 3-phosphate. The enzyme from yeast also acts on propane-1,2-diol 1-phosphate, but not on a variety of other phosphate esters.  
**References:** [2806]

[EC 3.1.3.21 created 1972, modified 1986]

#### EC 3.1.3.22

**Accepted name:** mannitol-1-phosphatase  
**Reaction:** D-mannitol 1-phosphate + H<sub>2</sub>O = D-mannitol + phosphate  
**Other name(s):** mannitol-1-phosphate phosphatase  
**Systematic name:** D-mannitol-1-phosphate phosphohydrolase  
**References:** [2477, 3242]

[EC 3.1.3.22 created 1972]

#### EC 3.1.3.23

**Accepted name:** sugar-phosphatase  
**Reaction:** sugar phosphate + H<sub>2</sub>O = sugar + phosphate  
**Systematic name:** sugar-phosphate phosphohydrolase  
**Comments:** Has a wide specificity, acting on aldohexose 1-phosphates, ketohexose 1-phosphates, aldohexose 6-phosphates, ketohexose 6-phosphates, both phosphate ester bonds of fructose 1,6-bisphosphate, phosphoric esters of disaccharides, and on pentose and triose phosphates, but at a slower rate.  
**References:** [1637]

[EC 3.1.3.23 created 1972]

#### EC 3.1.3.24

**Accepted name:** sucrose-phosphate phosphatase

**Reaction:** sucrose 6<sup>F</sup>-phosphate + H<sub>2</sub>O = sucrose + phosphate  
**Other name(s):** sucrose 6-phosphate hydrolase; sucrose-phosphate hydrolase; sucrose-phosphate phosphohydrolase; sucrose-6-phosphatase; sucrose phosphatase; sucrose-6-phosphate phosphatase; SPP  
**Systematic name:** sucrose-6<sup>F</sup>-phosphate phosphohydrolase  
**Comments:** Requires Mg<sup>2+</sup> for maximal activity [1754]. This is the final step in the sucrose-biosynthesis pathway. The enzyme is highly specific for sucrose 6-phosphate, with fructose 6-phosphate unable to act as a substrate [1754]. Belongs in the haloacid dehydrogenase (HAD) superfamily. The F of sucrose 6<sup>F</sup>-phosphate is used to indicate that the fructose residue of sucrose carries the substituent.  
**References:** [1081, 1754, 1755, 767]

[EC 3.1.3.24 created 1972, modified 2008]

#### EC 3.1.3.25

**Accepted name:** inositol-phosphate phosphatase  
**Reaction:** *myo*-inositol phosphate + H<sub>2</sub>O = *myo*-inositol + phosphate  
**Other name(s):** *myo*-inositol-1(or 4)-monophosphatase; inositol 1-phosphatase; L-*myo*-inositol-1-phosphate phosphatase; *myo*-inositol 1-phosphatase; inositol phosphatase; inositol monophosphate phosphatase; inositol-1(or 4)-monophosphatase; *myo*-inositol-1(or 4)-phosphate phosphohydrolase; *myo*-inositol monophosphatase; *myo*-inositol-1-phosphatase  
**Systematic name:** *myo*-inositol-phosphate phosphohydrolase  
**Comments:** Acts on five of the six isomers of *myo*-inositol phosphate, all except *myo*-inositol 2-phosphate, but does not act on *myo*-inositol bearing more than one phosphate group. It also acts on adenosine 2'-phosphate (but not the 3'- or 5'- phosphates), *sn*-glycerol 3-phosphate and glycerol 2-phosphate. Two isoforms are known [3302].  
**References:** [687, 887, 1028, 3302, 3214, 11]

[EC 3.1.3.25 created 1972, modified 1990, modified 2002, modified 2004]

#### EC 3.1.3.26

**Accepted name:** 4-phytase  
**Reaction:** *myo*-inositol hexakisphosphate + H<sub>2</sub>O = 1D-*myo*-inositol 1,2,3,5,6-pentakisphosphate + phosphate  
**Other name(s):** 6-phytase (name based on 1L-numbering system and not 1D-numbering); phytase; phytate 6-phosphatase; *myo*-inositol-hexakisphosphate 6-phosphohydrolase (name based on 1L-numbering system and not 1D-numbering)  
**Systematic name:** *myo*-inositol-hexakisphosphate 4-phosphohydrolase  
**References:** [1328, 2938, 1688, 492]

[EC 3.1.3.26 created 1972, modified 1976, modified 2002]

#### EC 3.1.3.27

**Accepted name:** phosphatidylglycerophosphatase  
**Reaction:** phosphatidylglycerophosphate + H<sub>2</sub>O = phosphatidylglycerol + phosphate  
**Other name(s):** phosphatidylglycerol phosphate phosphatase; phosphatidylglycerol phosphatase; PGP phosphatase  
**Systematic name:** phosphatidylglycerophosphate phosphohydrolase  
**References:** [399]

[EC 3.1.3.27 created 1972]

#### EC 3.1.3.28

**Accepted name:** ADP-phosphoglycerate phosphatase  
**Reaction:** 3-(ADP)-2-phosphoglycerate + H<sub>2</sub>O = 3-(ADP)-glycerate + phosphate  
**Other name(s):** adenosine diphosphate phosphoglycerate phosphatase  
**Systematic name:** 3-(ADP)-2-phosphoglycerate phosphohydrolase

**Comments:** Also acts on 2,3-bisphosphoglycerate.

**References:** [3317]

[EC 3.1.3.28 created 1972]

#### EC 3.1.3.29

**Accepted name:** *N*-acylneuraminate-9-phosphatase

**Reaction:** *N*-acylneuraminate 9-phosphate + H<sub>2</sub>O = *N*-acylneuraminate + phosphate

**Other name(s):** acylneuraminate 9-phosphatase; *N*-acylneuraminic acid 9-phosphate phosphatase; *N*-acylneuraminic (sialic) acid 9-phosphatase

**Systematic name:** *N*-acylneuraminate-9-phosphate phosphohydrolase

**References:** [1344]

[EC 3.1.3.29 created 1972]

[3.1.3.30 Deleted entry. 3'-phosphoadenylylsulfate 3'-phosphatase. Now included with EC 3.1.3.31 nucleotidase]

[EC 3.1.3.30 created 1972, deleted 1992]

#### EC 3.1.3.31

**Accepted name:** nucleotidase

**Reaction:** a nucleotide + H<sub>2</sub>O = a nucleoside + phosphate

**Other name(s):** nucleotide phosphatase; nucleotide-specific phosphatase; NSP I; NSP II; deoxyribonucleoside-activated nucleotidase (DAN); deoxyinosine-activated nucleotidase (DIAN); acid nucleotidase

**Systematic name:** nucleotide phosphohydrolase

**Comments:** A wide specificity for 2', 3'- and 5'- nucleotides; also hydrolyses glycerol phosphate and 4-nitrophenyl phosphate.

**References:** [76]

[EC 3.1.3.31 created 1972 (EC 3.1.3.30 created 1972, incorporated 1992)]

#### EC 3.1.3.32

**Accepted name:** polynucleotide 3'-phosphatase

**Reaction:** a 3'-phosphopolynucleotide + H<sub>2</sub>O = a polynucleotide + phosphate

**Other name(s):** 2'(3')-polynucleotidase; DNA 3'-phosphatase; deoxyribonucleate 3'-phosphatase; 5'-polynucleotidekinase 3'-phosphatase

**Systematic name:** polynucleotide 3'-phosphohydrolase

**Comments:** Also hydrolyses nucleoside 2', 3'- and 5'-monophosphates, but only 2'- and 3'-phosphopolynucleotides.

**References:** [180]

[EC 3.1.3.32 created 1972]

#### EC 3.1.3.33

**Accepted name:** polynucleotide 5'-phosphatase

**Reaction:** a 5'-phosphopolynucleotide + H<sub>2</sub>O = a polynucleotide + phosphate

**Other name(s):** 5'-polynucleotidase

**Systematic name:** polynucleotide 5'-phosphohydrolase

**Comments:** Does not act on nucleoside monophosphates. Induced in *Escherichia coli* by T-even phages.

**References:** [180]

[EC 3.1.3.33 created 1972]

#### EC 3.1.3.34

**Accepted name:** deoxynucleotide 3'-phosphatase  
**Reaction:** a 2'-deoxyribonucleoside 3'-phosphate + H<sub>2</sub>O = a 2'-deoxyribonucleoside + phosphate  
**Other name(s):** 3'-deoxynucleotidase; 3'-deoxyribonucleotidase  
**Systematic name:** 2'-deoxyribonucleotide 3'-phosphohydrolase  
**Comments:** Also catalyses the selective removal of 3'-phosphate groups from DNA and oligodeoxyribonucleotides. Induced in *Escherichia coli* by T-even phages.  
**References:** [180]

[EC 3.1.3.34 created 1972]

#### EC 3.1.3.35

**Accepted name:** thymidylate 5'-phosphatase  
**Reaction:** thymidylate + H<sub>2</sub>O = thymidine + phosphate  
**Other name(s):** thymidylate 5'-nucleotidase; deoxythymidylate 5'-nucleotidase; thymidylate nucleotidase; deoxythymidylic 5'-nucleotidase; deoxythymidylate phosphohydrolase; dTMPase  
**Systematic name:** thymidylate 5'-phosphohydrolase  
**Comments:** Acts on 5-methyl-dCMP and on TMP, but more slowly than on dTMP.  
**References:** [63]

[EC 3.1.3.35 created 1972]

#### EC 3.1.3.36

**Accepted name:** phosphoinositide 5-phosphatase  
**Reaction:** 1-phosphatidyl-1D-*myo*-inositol 4,5-bisphosphate + H<sub>2</sub>O = 1-phosphatidyl-1D-*myo*-inositol 4-phosphate + phosphate  
**Other name(s):** type II inositol polyphosphate 5-phosphatase; triphosphoinositide phosphatase; IP<sub>3</sub> phosphatase; PtdIns(4,5)P<sub>2</sub> phosphatase; triphosphoinositide phosphomonoesterase; diphosphoinositide phosphatase; inositol 1,4,5-triphosphate 5-phosphomonoesterase; inositol triphosphate 5-phosphomonoesterase; phosphatidylinositol-bisphosphatase; phosphatidyl-*myo*-inositol-4,5-bisphosphate phosphatase; phosphatidylinositol 4,5-bisphosphate phosphatase; polyphosphoinositol lipid 5-phosphatase; phosphatidyl-inositol-bisphosphate phosphatase  
**Systematic name:** phosphatidyl-*myo*-inositol-4,5-bisphosphate 4-phosphohydrolase  
**Comments:** These enzymes can also remove the 5-phosphate from Ins(1,4,5)P<sub>3</sub> and/or Ins(1,3,4,5)P<sub>4</sub>. They are a diverse family of enzymes, with differing abilities to catalyse two or more of the four reactions listed. They are thought to use inositol lipids rather than inositol phosphates as substrates *in vivo*. All of them can use either or both of PtdIns(4,5)P<sub>2</sub> and PtdIns(3,4,5)P<sub>3</sub> as substrates; this is the main property that distinguishes them from EC 3.1.3.56, inositol-polyphosphate 5-phosphatase.  
**References:** [556, 2432, 3214]

[EC 3.1.3.36 created 1972, modified 2002]

#### EC 3.1.3.37

**Accepted name:** sedoheptulose-bisphosphatase  
**Reaction:** sedoheptulose 1,7-bisphosphate + H<sub>2</sub>O = sedoheptulose 7-phosphate + phosphate  
**Other name(s):** SBPase; sedoheptulose 1,7-diphosphate phosphatase; sedoheptulose 1,7-diphosphatase; sedoheptulose diphosphatase; sedoheptulose bisphosphatase; sedoheptulose 1,7-bisphosphatase  
**Systematic name:** sedoheptulose-1,7-bisphosphate 1-phosphohydrolase  
**References:** [2353, 2960]

[EC 3.1.3.37 created 1976]

#### EC 3.1.3.38

**Accepted name:** 3-phosphoglycerate phosphatase  
**Reaction:** D-glycerate 3-phosphate + H<sub>2</sub>O = D-glycerate + phosphate  
**Other name(s):** D-3-Phosphoglycerate phosphatase; 3-PGA phosphatase  
**Systematic name:** D-glycerate-3-phosphate phosphohydrolase  
**Comments:** Wide specificity, but 3-phosphoglycerate is the best substrate.  
**References:** [2369]

[EC 3.1.3.38 created 1976]

#### EC 3.1.3.39

**Accepted name:** streptomycin-6-phosphatase  
**Reaction:** streptomycin 6-phosphate + H<sub>2</sub>O = streptomycin + phosphate  
**Other name(s):** streptomycin 6-phosphate phosphatase; streptomycin 6-phosphate phosphohydrolase; streptomycin-6-*P* phosphohydrolase  
**Systematic name:** streptomycin-6-phosphate phosphohydrolase  
**Comments:** Also acts on dihydrostreptomycin 3' $\alpha$ ,6-bisphosphate and streptidine 6-phosphate.  
**References:** [3096, 3097]

[EC 3.1.3.39 created 1976]

#### EC 3.1.3.40

**Accepted name:** guanidinodeoxy-*scyllo*-inositol-4-phosphatase  
**Reaction:** 1-guanidino-1-deoxy-*scyllo*-inositol 4-phosphate + H<sub>2</sub>O = 1-guanidino-1-deoxy-*scyllo*-inositol + phosphate  
**Other name(s):** 1-guanidino-*scyllo*-inositol 4-phosphatase; 1-guanidino-1-deoxy-*scyllo*-inositol-4-*P* phosphohydrolase  
**Systematic name:** 1-guanidino-1-deoxy-*scyllo*-inositol-4-phosphate 4-phosphohydrolase  
**References:** [3097]

[EC 3.1.3.40 created 1976]

#### EC 3.1.3.41

**Accepted name:** 4-nitrophenylphosphatase  
**Reaction:** 4-nitrophenyl phosphate + H<sub>2</sub>O = 4-nitrophenol + phosphate  
**Other name(s):** nitrophenyl phosphatase; *p*-nitrophenylphosphatase; para-nitrophenyl phosphatase; K-pNPPase; NPPase; PNPPase; Ecto-*p*-nitrophenyl phosphatase; *p*-nitrophenylphosphate phosphohydrolase  
**Systematic name:** 4-nitrophenylphosphate phosphohydrolase  
**Comments:** A number of other substances, including phenyl phosphate, 4-nitrophenyl sulfate, acetyl phosphate and glycerol phosphate, are not substrates.  
**References:** [93, 94]

[EC 3.1.3.41 created 1976]

#### EC 3.1.3.42

**Accepted name:** [glycogen-synthase-D] phosphatase  
**Reaction:** [glycogen-synthase D] + H<sub>2</sub>O = [glycogen-synthase I] + phosphate  
**Other name(s):** uridine diphosphoglucose-glycogen glucosyltransferase phosphatase; UDP-glycogen glucosyltransferase phosphatase; UDPglucose-glycogen glucosyltransferase phosphatase; glycogen glucosyltransferase phosphatase; glycogen synthetase phosphatase; glycogen synthase phosphatase; glycogen synthase D phosphatase; Mg<sup>2+</sup> dependent glycogen synthase phosphatase; phosphatase type 2°C  
**Systematic name:** [UDP-glucose:glycogen 4- $\alpha$ -D-glucosyltransferase-D] phosphohydrolase  
**Comments:** The product is EC 2.4.1.11 glycogen(starch) synthase.  
**References:** [6]

[EC 3.1.3.42 created 1976]

#### EC 3.1.3.43

- Accepted name:** [pyruvate dehydrogenase (acetyl-transferring)]-phosphatase  
**Reaction:** [pyruvate dehydrogenase (acetyl-transferring)] phosphate + H<sub>2</sub>O = [pyruvate dehydrogenase (acetyl-transferring)] + phosphate  
**Other name(s):** pyruvate dehydrogenase phosphatase; phosphopyruvate dehydrogenase phosphatase; [pyruvate dehydrogenase (lipoamide)]-phosphatase; [pyruvate dehydrogenase (lipoamide)]-phosphate phosphohydrolase  
**Systematic name:** [pyruvate dehydrogenase (acetyl-transferring)]-phosphate phosphohydrolase  
**Comments:** A mitochondrial enzyme associated with EC 1.2.4.1 pyruvate dehydrogenase (acetyl-transferring), in the pyruvate dehydrogenase complex.  
**References:** [1708, 2395]

[EC 3.1.3.43 created 1978]

#### EC 3.1.3.44

- Accepted name:** [acetyl-CoA carboxylase]-phosphatase  
**Reaction:** [acetyl-CoA carboxylase] phosphate + H<sub>2</sub>O = [acetyl-CoA carboxylase] + phosphate  
**Systematic name:** [acetyl-CoA:carbon-dioxide ligase (ADP-forming)]-phosphate phosphohydrolase  
**Comments:** Simultaneously dephosphorylates and activates EC 6.4.1.2 acetyl-CoA carboxylase. Acts similarly on EC 1.1.1.88 (hydroxymethylglutaryl-CoA reductase), EC 2.4.1.1 (phosphorylase), EC 2.4.1.11 [glycogen(starch) synthase], and dephosphorylates phosphoprotamine and 4-nitrophenyl phosphate. Not identical to EC 3.1.3.17 ([phosphorylase] phosphatase) or EC 3.1.3.43 [pyruvate dehydrogenase (acetyl-transferring)]-phosphatase.  
**References:** [1527]

[EC 3.1.3.44 created 1983]

#### EC 3.1.3.45

- Accepted name:** 3-deoxy-*manno*-octulosonate-8-phosphatase  
**Reaction:** 3-deoxy-D-*manno*-octulosonate 8-phosphate + H<sub>2</sub>O = 3-deoxy-D-*manno*-octulosonate + phosphate  
**Systematic name:** 3-deoxy-D-*manno*-octulosonate-8-phosphate 8-phosphohydrolase  
**References:** [2385]

[EC 3.1.3.45 created 1983]

#### EC 3.1.3.46

- Accepted name:** fructose-2,6-bisphosphate 2-phosphatase  
**Reaction:** β-D-fructose 2,6-bisphosphate + H<sub>2</sub>O = D-fructose 6-phosphate + phosphate  
**Other name(s):** fructose-2,6-bisphosphatase; D-fructose-2,6-bisphosphate 2-phosphohydrolase  
**Systematic name:** β-D-fructose-2,6-bisphosphate 2-phosphohydrolase  
**Comments:** The enzyme copurifies with EC 2.7.1.105 6-phosphofructo-2-kinase. (*cf.* EC 3.1.3.54 fructose-2,6-bisphosphate 6-phosphatase).  
**References:** [2553]

[EC 3.1.3.46 created 1984]

#### EC 3.1.3.47

- Accepted name:** [hydroxymethylglutaryl-CoA reductase (NADPH)]-phosphatase  
**Reaction:** [hydroxymethylglutaryl-CoA reductase (NADPH)] phosphate + H<sub>2</sub>O = [hydroxymethylglutaryl-CoA reductase (NADPH)] + phosphate



**Other name(s):** reductase phosphatase  
**Systematic name:** [hydroxymethylglutaryl-CoA reductase (NADPH)]-phosphate phosphohydrolase  
**Comments:** Acts on the product of the reaction catalysed by EC 2.7.11.31 [hydroxymethylglutaryl-CoA reductase (NADPH)] kinase, simultaneously dephosphorylating and activating EC 1.1.1.34 hydroxymethylglutaryl-CoA reductase (NADPH).  
**References:** [917, 918]

[EC 3.1.3.47 created 1984]

#### EC 3.1.3.48

**Accepted name:** protein-tyrosine-phosphatase  
**Reaction:** [a protein]-tyrosine phosphate + H<sub>2</sub>O = [a protein]-tyrosine + phosphate  
**Other name(s):** phosphotyrosine phosphatase; phosphoprotein phosphatase (phosphotyrosine); phosphotyrosine histone phosphatase; protein phosphotyrosine phosphatase; tyrosylprotein phosphatase; phosphotyrosine protein phosphatase; phosphotyrosylprotein phosphatase; tyrosine *O*-phosphate phosphatase; PPT-phosphatase; PTPase; [phosphotyrosine]protein phosphatase; PTP-phosphatase  
**Systematic name:** protein-tyrosine-phosphate phosphohydrolase  
**Comments:** Dephosphorylates *O*-phosphotyrosine groups in phosphoproteins, such as the products of EC 2.7.10.2, non-specific protein-tyrosine kinase.  
**References:** [798, 870]

[EC 3.1.3.48 created 1984]

#### EC 3.1.3.49

**Accepted name:** [pyruvate kinase]-phosphatase  
**Reaction:** [pyruvate kinase] phosphate + H<sub>2</sub>O = [pyruvate kinase] + phosphate  
**Other name(s):** pyruvate kinase phosphatase  
**Systematic name:** [ATP:pyruvate 2-*O*-phosphotransferase]-phosphate phosphohydrolase  
**Comments:** Simultaneously dephosphorylates and activates EC 2.7.1.40 pyruvate kinase, that has been inactivated by protein kinase.  
**References:** [1321]

[EC 3.1.3.49 created 1984]

#### EC 3.1.3.50

**Accepted name:** sorbitol-6-phosphatase  
**Reaction:** sorbitol 6-phosphate + H<sub>2</sub>O = sorbitol + phosphate  
**Other name(s):** sorbitol-6-phosphate phosphatase  
**Systematic name:** sorbitol-6-phosphate phosphohydrolase  
**Comments:** Acts, very slowly, on hexose 6-phosphates.  
**References:** [964]

[EC 3.1.3.50 created 1984]

#### EC 3.1.3.51

**Accepted name:** dolichyl-phosphatase  
**Reaction:** dolichyl phosphate + H<sub>2</sub>O = dolichol + phosphate  
**Other name(s):** dolichol phosphate phosphatase; dolichol phosphatase; dolichol monophosphatase; dolichyl monophosphate phosphatase; dolichyl phosphate phosphatase; polyisoprenyl phosphate phosphatase; polyprenylphosphate phosphatase; Dol-*P* phosphatase  
**Systematic name:** dolichyl-phosphate phosphohydrolase  
**References:** [17, 2429, 3146]

[EC 3.1.3.51 created 1984]

#### EC 3.1.3.52

- Accepted name:** [3-methyl-2-oxobutanoate dehydrogenase (2-methylpropanoyl-transferring)]-phosphatase  
**Reaction:** [3-methyl-2-oxobutanoate dehydrogenase (2-methylpropanoyl-transferring)] phosphate + H<sub>2</sub>O = [3-methyl-2-oxobutanoate dehydrogenase (2-methylpropanoyl-transferring)] + phosphate  
**Other name(s):** branched-chain oxo-acid dehydrogenase phosphatase; branched-chain 2-keto acid dehydrogenase phosphatase; branched-chain  $\alpha$ -keto acid dehydrogenase phosphatase; BCKDH; [3-methyl-2-oxobutanoate dehydrogenase (lipoamide)]-phosphatase; [3-methyl-2-oxobutanoate dehydrogenase (lipoamide)]-phosphate phosphohydrolase  
**Systematic name:** [3-methyl-2-oxobutanoate dehydrogenase (2-methylpropanoyl-transferring)]-phosphate phosphohydrolase  
**Comments:** A mitochondrial enzyme associated with the 3-methyl-2-oxobutanoate dehydrogenase complex. Simultaneously dephosphorylates and activates EC 1.2.4.4 3-methyl-2-oxobutanoate dehydrogenase (2-methylpropanoyl-transferring), that has been inactivated by phosphorylation.  
**References:** [742, 2395]

[EC 3.1.3.52 created 1986]

#### EC 3.1.3.53

- Accepted name:** [myosin-light-chain] phosphatase  
**Reaction:** [myosin light-chain] phosphate + H<sub>2</sub>O = [myosin light-chain] + phosphate  
**Other name(s):** myosin light chain kinase phosphatase; myosin phosphatase; myosin phosphatase; protein phosphatase 2A; myosin-light-chain-phosphatase  
**Systematic name:** [myosin-light-chain]-phosphate phosphohydrolase  
**Comments:** The enzyme is composed of three subunits. The holoenzyme dephosphorylates myosin light chains and EC 2.7.11.18, myosin-light-chain kinase, but not myosin; the catalytic subunit acts on all three substrates.  
**References:** [2243]

[EC 3.1.3.53 created 1986]

#### EC 3.1.3.54

- Accepted name:** fructose-2,6-bisphosphate 6-phosphatase  
**Reaction:**  $\beta$ -D-fructose 2,6-bisphosphate + H<sub>2</sub>O =  $\beta$ -D-fructofuranose 2-phosphate + phosphate  
**Other name(s):** fructose 2,6-bisphosphate-6-phosphohydrolase; fructose-2,6-bisphosphate 6-phosphohydrolase; D-fructose-2,6-bisphosphate 6-phosphohydrolase  
**Systematic name:**  $\beta$ -D-fructose-2,6-bisphosphate 6-phosphohydrolase  
**Comments:** *cf.* EC 3.1.3.46 fructose-2,6-bisphosphate 2-phosphatase.  
**References:** [2332, 2333]

[EC 3.1.3.54 created 1989]

#### EC 3.1.3.55

- Accepted name:** caldesmon-phosphatase  
**Reaction:** caldesmon phosphate + H<sub>2</sub>O = caldesmon + phosphate  
**Other name(s):** SMP-I; smooth muscle caldesmon phosphatase  
**Systematic name:** caldesmon-phosphate phosphohydrolase  
**Comments:** Dephosphorylation activates the calmodulin- and actin-binding ability of the protein caldesmon.  
**References:** [2078]

[EC 3.1.3.55 created 1989]

### EC 3.1.3.56

- Accepted name:** inositol-polyphosphate 5-phosphatase
- Reaction:** (1) D-*myo*-inositol 1,4,5-trisphosphate + H<sub>2</sub>O = *myo*-inositol 1,4-bisphosphate + phosphate  
(2) 1D-*myo*-inositol 1,3,4,5-tetrakisphosphate + H<sub>2</sub>O = 1D-*myo*-inositol 1,3,4-trisphosphate + phosphate
- Other name(s):** type I inositol-polyphosphate phosphatase; inositol trisphosphate phosphomonoesterase; InsP<sub>3</sub>/Ins(1,3,4,5)P<sub>4</sub> 5-phosphatase; inosine triphosphatase; D-*myo*-inositol 1,4,5-trisphosphate 5-phosphatase; D-*myo*-inositol 1,4,5-trisphosphate 5-phosphatase; L-*myo*-inositol 1,4,5-trisphosphate-monoesterase; inositol phosphate 5-phosphomonoesterase; inositol-1,4,5-trisphosphate/1,3,4,5-tetrakisphosphate 5-phosphatase; Ins(1,4,5)P<sub>3</sub> 5-phosphatase; D-*myo*-inositol(1,4,5)/(1,3,4,5)-polyphosphate 5-phosphatase; inositol 1,4,5-trisphosphate phosphatase; inositol polyphosphate-5-phosphatase; *myo*-inositol-1,4,5-trisphosphate 5-phosphatase; inositol-1,4,5-trisphosphate 5-phosphatase
- Systematic name:** 1D-*myo*-inositol-1,4,5-trisphosphate 5-phosphohydrolase
- Comments:** One mammalian isoform is known. This enzyme is distinguished from the family of enzymes classified under EC 3.1.3.36, phosphoinositide 5-phosphatase, by its inability to dephosphorylate inositol lipids.
- References:** [643, 714, 3214, 3060]

[EC 3.1.3.56 created 1989, modified 2002]

### EC 3.1.3.57

- Accepted name:** inositol-1,4-bisphosphate 1-phosphatase
- Reaction:** 1D-*myo*-inositol 1,4-bisphosphate + H<sub>2</sub>O = 1D-*myo*-inositol 4-phosphate + phosphate
- Other name(s):** inositol-polyphosphate 1-phosphatase
- Systematic name:** 1D-*myo*-inositol-1,4-bisphosphate 1-phosphohydrolase
- Comments:** The enzyme acts on inositol 1,4-bisphosphate and inositol 1,3,4-trisphosphate (forming inositol 3,4-bisphosphate) with similar  $V_{max}$  values for both substrates, but with a five-times higher affinity for the bisphosphate. Does not act on inositol 1-phosphate, inositol 1,4,5-trisphosphate or inositol 1,3,4,5-tetrakisphosphate.
- References:** [204, 481, 1251]

[EC 3.1.3.57 created 1989, modified 2002]

### EC 3.1.3.58

- Accepted name:** sugar-terminal-phosphatase
- Reaction:** D-glucose 6-phosphate + H<sub>2</sub>O = D-glucose + phosphate
- Other name(s):** xylitol-5-phosphatase
- Systematic name:** sugar- $\omega$ -phosphate phosphohydrolase
- Comments:** Acts on sugars and polyols phosphorylated on the terminal carbon, with a preference for sugars with a D-*erythro*-configuration, e.g. good substrates are glucose 6-phosphate, mannose 6-phosphate, 6-phosphogluconate, erythrose 4-phosphate and xylitol 5-phosphate.
- References:** [1734]

[EC 3.1.3.58 created 1989]

### EC 3.1.3.59

- Accepted name:** alkylacetyl-glycerophosphatase
- Reaction:** 1-alkyl-2-acetyl-*sn*-glycero-3-phosphate + H<sub>2</sub>O = 1-alkyl-2-acetyl-*sn*-glycerol + phosphate
- Other name(s):** 1-alkyl-2-lyso-*sn*-glycero-3-P:acetyl-CoA acetyltransferase; alkylacetyl-glycerophosphate phosphatase
- Systematic name:** 1-alkyl-2-acetyl-*sn*-glycero-3-phosphate phosphohydrolase
- Comments:** Involved in the biosynthesis of thrombocyte activating factor in animal tissues.
- References:** [1632]

[EC 3.1.3.59 created 1989]

#### EC 3.1.3.60

**Accepted name:** phosphoenolpyruvate phosphatase  
**Reaction:** phosphoenolpyruvate + H<sub>2</sub>O = pyruvate + phosphate  
**Other name(s):** PEP phosphatase  
**Systematic name:** phosphoenolpyruvate phosphohydrolase  
**Comments:** Also acts, but more slowly, on a wide range of other monophosphates.  
**References:** [664, 1788, 1789]

[EC 3.1.3.60 created 1992]

[3.1.3.61 Deleted entry. inositol-1,4,5-trisphosphate 1-phosphatase, as its existence has not been established]

[EC 3.1.3.61 created 1992, deleted 2002]

#### EC 3.1.3.62

**Accepted name:** multiple inositol-polyphosphate phosphatase  
**Reaction:** *myo*-inositol hexakisphosphate + H<sub>2</sub>O = *myo*-inositol pentakisphosphate (mixed isomers) + phosphate  
**Other name(s):** inositol (1,3,4,5)-tetrakisphosphate 3-phosphatase; inositol 1,3,4,5-tetrakisphosphate 3-phosphomonoesterase; inositol 1,3,4,5-tetrakisphosphate-5-phosphomonoesterase; inositol tetrakisphosphate phosphomonoesterase; inositol-1,3,4,5-tetrakisphosphate 3-phosphatase; MIPP  
**Systematic name:** 1D-*myo*-inositol-hexakisphosphate 5-phosphohydrolase  
**Comments:** This enzyme exists in two isoforms. It also acts on Ins(1,3,4,5)P<sub>4</sub> to yield Ins(1,4,5)P<sub>3</sub>.  
**References:** [515, 501]

[EC 3.1.3.62 created 1992, modified 2002]

#### EC 3.1.3.63

**Accepted name:** 2-carboxy-D-arabinitol-1-phosphatase  
**Reaction:** 2-carboxy-D-arabinitol 1-phosphate + H<sub>2</sub>O = 2-carboxy-D-arabinitol + phosphate  
**Other name(s):** 2-carboxyarabinitol 1-phosphatase; 2-carboxy-D-arabinitol 1-phosphate phosphohydrolase  
**Systematic name:** 2-carboxy-D-arabinitol-1-phosphate 1-phosphohydrolase  
**References:** [2513]

[EC 3.1.3.63 created 1992]

#### EC 3.1.3.64

**Accepted name:** phosphatidylinositol-3-phosphatase  
**Reaction:** 1-phosphatidyl-1D-*myo*-inositol 3-phosphate + H<sub>2</sub>O = 1-phosphatidyl-1D-*myo*-inositol + phosphate  
**Other name(s):** inositol-1,3-bisphosphate 3-phosphatase; inositol 1,3-bisphosphate phosphatase; inositol-polyphosphate 3-phosphatase; D-*myo*-inositol-1,3-bisphosphate 3-phosphohydrolase; phosphatidyl-3-phosphate 3-phosphohydrolase  
**Systematic name:** 1-phosphatidyl-1D-*myo*-inositol-3-phosphate 3-phosphohydrolase  
**Comments:** This enzyme still works when the 2,3-bis(acyloxy)propyl group is removed, i.e., it hydrolyses Ins(1,3)P<sub>2</sub> to Ins-1-P.  
**References:** [1711, 355]

[EC 3.1.3.64 created 1992, [EC 3.1.3.65 created 1992, incorporated 2002], modified 2002]]

[3.1.3.65 Deleted entry. inositol-1,3-bisphosphate 3-phosphatase. Now included with EC 3.1.3.64, phosphatidylinositol-3-phosphatase]

[EC 3.1.3.65 created 1992, deleted 2002]

### EC 3.1.3.66

- Accepted name:** phosphatidylinositol-3,4-bisphosphate 4-phosphatase  
**Reaction:** 1-phosphatidyl-*myo*-inositol 3,4-bisphosphate + H<sub>2</sub>O = 1-phosphatidyl-1D-*myo*-inositol 3-phosphate + phosphate  
**Other name(s):** inositol-3,4-bisphosphate 4-phosphatase; D-*myo*-inositol-3,4-bisphosphate 4-phosphohydrolase; phosphoinositide 4-phosphatase; inositol polyphosphate 4-phosphatase; inositol polyphosphate 4-phosphatase type II  
**Systematic name:** 1-phosphatidyl-1D-*myo*-inositol-3,4-bisphosphate 4-phosphohydrolase  
**Comments:** Mg<sup>2+</sup>-independent. This enzyme still works when the 2,3-bis(acyloxy)propyl group is removed, i.e., it hydrolyses Ins(1,3,4)P<sub>3</sub> to Ins(1,3)P<sub>2</sub>. It also converts Ins(3,4)P<sub>2</sub> into Ins-3-*P*.  
**References:** [1194, 2114, 2113]

[EC 3.1.3.66 created 1992, modified 2002]

### EC 3.1.3.67

- Accepted name:** phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase  
**Reaction:** 1-phosphatidyl-1D-*myo*-inositol 3,4,5-trisphosphate + H<sub>2</sub>O = 1-phosphatidyl-1D-*myo*-inositol 4,5-bisphosphate + phosphate  
**Other name(s):** PTEN; MMAC1; phosphatidylinositol-3,4,5-trisphosphate 3-phosphohydrolase  
**Systematic name:** 1-phosphatidyl-1D-*myo*-inositol-3,4,5-trisphosphate 3-phosphohydrolase  
**Comments:** Requires Mg<sup>2+</sup>. Does not dephosphorylate inositol 4,5-bisphosphate. This enzyme still works when the 2,3-bis(acyloxy)propyl group is removed, i.e., it hydrolyses Ins(1,3,4,5)P<sub>4</sub> to Ins(1,4,5)P<sub>3</sub>  
**References:** [1351, 2824]

[EC 3.1.3.67 created 1999, modified 2002]

### EC 3.1.3.68

- Accepted name:** 2-deoxyglucose-6-phosphatase  
**Reaction:** 2-deoxy-D-glucose 6-phosphate + H<sub>2</sub>O = 2-deoxy-D-glucose + phosphate  
**Other name(s):** 2-deoxyglucose-6-phosphate phosphatase  
**Systematic name:** 2-deoxy-D-glucose-6-phosphate phosphohydrolase  
**Comments:** Also active towards fructose 1-phosphate  
**References:** [1332, 2371]

[EC 3.1.3.68 created 1999]

### EC 3.1.3.69

- Accepted name:** glucosylglycerol 3-phosphatase  
**Reaction:** 2-*O*-(α-D-glucosyl)-*sn*-glycerol-3-phosphate + H<sub>2</sub>O = 2-*O*-(α-D-glucopyranosyl)glycerol + phosphate  
**Other name(s):** salt tolerance protein A; StpA; 2-(β-D-glucosyl)-*sn*-glycerol-3-phosphate phosphohydrolase (incorrect)  
**Systematic name:** 2-*O*-(α-D-glucopyranosyl)-*sn*-glycerol-3-phosphate phosphohydrolase  
**Comments:** Acts with EC 2.4.1.213 (glucosylglycerol-phosphate synthase) to form glucosylglycerol, an osmolyte that endows cyanobacteria with resistance to salt.  
**References:** [1018, 1019, 1020]

[EC 3.1.3.69 created 2001, modified 2015]

### EC 3.1.3.70

- Accepted name:** mannosyl-3-phosphoglycerate phosphatase  
**Reaction:** 2-*O*-(α-D-mannosyl)-3-phosphoglycerate + H<sub>2</sub>O = 2-*O*-(α-D-mannosyl)-D-glycerate + phosphate  
**Systematic name:** 2-*O*-(α-D-mannosyl)-3-phosphoglycerate phosphohydrolase  
**Comments:** Requires Mg<sup>2+</sup>. The enzyme from *Pyrococcus horikoshii* is specific for α-D-mannosyl-3-phosphoglycerate and forms part of the pathway for the synthesis of mannosylglycerate.

**References:** [701]

[EC 3.1.3.70 created 2002]

#### EC 3.1.3.71

**Accepted name:** 2-phosphosulfolactate phosphatase  
**Reaction:** (2*R*)-2-phospho-3-sulfolactate + H<sub>2</sub>O = (2*R*)-3-sulfolactate + phosphate  
**Other name(s):** (2*R*)-phosphosulfolactate phosphohydrolase; ComB phosphatase  
**Systematic name:** (*R*)-2-phospho-3-sulfolactate phosphohydrolase  
**Comments:** Requires Mg<sup>2+</sup>. The enzyme from *Methanococcus jannaschii* acts on both stereoisomers of the substrate and also hydrolyses a number of phosphate monoesters of (*S*)-2-hydroxycarboxylic acids, including 2-phosphomalate, 2-phospholactate and 2-phosphoglycolate. This enzyme can also hydrolyse phosphate monoesters of (*R*)-2-hydroxycarboxylic acids such as (*S*)-2-phospho-3-sulfolactate and (*R*)-2-phosphomalate, which, presumably, bind to the enzyme in opposite orientations.  
**References:** [962]

[EC 3.1.3.71 created 2002]

#### EC 3.1.3.72

**Accepted name:** 5-phytase  
**Reaction:** *myo*-inositol hexakisphosphate + H<sub>2</sub>O = 1*L*-*myo*-inositol 1,2,3,4,6-pentakisphosphate + phosphate  
**Systematic name:** *myo*-inositol-hexakisphosphate 5-phosphohydrolase  
**Comments:** The enzyme attacks the product of the above reaction more slowly to yield Ins(1,2,3)P<sub>3</sub>.  
**References:** [163]

[EC 3.1.3.72 created 2002]

#### EC 3.1.3.73

**Accepted name:** adenosylcobalamin/ $\alpha$ -ribazole phosphatase  
**Reaction:** (1) adenosylcobalamin 5'-phosphate + H<sub>2</sub>O = coenzyme B<sub>12</sub> + phosphate  
(2)  $\alpha$ -ribazole 5'-phosphate + H<sub>2</sub>O =  $\alpha$ -ribazole + phosphate  
**Other name(s):** CobC; adenosylcobalamin phosphatase;  $\alpha$ -ribazole phosphatase  
**Systematic name:** adenosylcobalamin/ $\alpha$ -ribazole-5'-phosphate phosphohydrolase  
**Comments:** This enzyme catalyses the last step in the anaerobic (early cobalt insertion) pathway of adenosylcobalamin biosynthesis, characterized in *Salmonella enterica* [3318]. It also participates in a salvage pathway that recycles cobinamide into adenosylcobalamin [2213].  
**References:** [2213, 3131, 3318]

[EC 3.1.3.73 created 2004, modified 2011]

#### EC 3.1.3.74

**Accepted name:** pyridoxal phosphatase  
**Reaction:** pyridoxal 5'-phosphate + H<sub>2</sub>O = pyridoxal + phosphate  
**Other name(s):** vitamine B<sub>6</sub> (pyridoxine) phosphatase; PLP phosphatase; vitamin B<sub>6</sub>-phosphate phosphatase; PNP phosphatase  
**Systematic name:** pyridoxal-5'-phosphate phosphohydrolase  
**Comments:** Requires Mg<sup>2+</sup>. This enzyme is specific for phosphorylated vitamin B<sub>6</sub> compounds: it acts not only on pyridoxal phosphate (PLP), but also on pyridoxine phosphate (PNP), pyridoxamine phosphate (PMP), 4-pyridoxic acid phosphate and 4-deoxypyridoxine phosphate. This reaction can also be carried out by EC 3.1.3.1 (alkaline phosphatase) and EC 3.1.3.2 (acid phosphatase), but these enzymes have very broad substrate specificities.  
**References:** [794, 795, 1307]

[EC 3.1.3.74 created 2004]

### EC 3.1.3.75

- Accepted name:** phosphoethanolamine/phosphocholine phosphatase  
**Reaction:** (1) *O*-phosphoethanolamine + H<sub>2</sub>O = ethanolamine + phosphate  
(2) phosphocholine + H<sub>2</sub>O = choline + phosphate  
**Other name(s):** PHOSPHO1; 3X11A  
**Systematic name:** phosphoethanolamine phosphohydrolase  
**Comments:** Requires active site Mg<sup>2+</sup> but also works, to a lesser extent, with Co<sup>2+</sup> and Mn<sup>2+</sup>. The enzyme is highly specific for phosphoethanolamine and phosphocholine.  
**References:** [1189, 2771, 2438]

[EC 3.1.3.75 created 2004]

### EC 3.1.3.76

- Accepted name:** lipid-phosphate phosphatase  
**Reaction:** (9*S*,10*S*)-10-hydroxy-9-(phosphooxy)octadecanoate + H<sub>2</sub>O = (9*S*,10*S*)-9,10-dihydroxyoctadecanoate + phosphate  
**Other name(s):** hydroxy fatty acid phosphatase; dihydroxy fatty acid phosphatase; hydroxy lipid phosphatase; sEH (ambiguous); soluble epoxide hydrolase (ambiguous); (9*S*,10*S*)-10-hydroxy-9-(phosphonooxy)octadecanoate phosphohydrolase  
**Systematic name:** (9*S*,10*S*)-10-hydroxy-9-(phosphooxy)octadecanoate phosphohydrolase  
**Comments:** Requires Mg<sup>2+</sup> for maximal activity. The enzyme from mammals is a bifunctional enzyme: the N-terminal domain exhibits lipid-phosphate-phosphatase activity and the C-terminal domain has the activity of EC 3.3.2.10, soluble epoxide hydrolase (sEH) [2074]. The best substrates for this enzyme are 10-hydroxy-9-(phosphooxy)octadecanoates, with the *threo*- form being a better substrate than the *erythro*- form [2074]. The phosphatase activity is not found in plant sEH or in EC 3.3.2.9, microsomal epoxide hydrolase, from mammals [2074].  
**References:** [2074, 507, 1977, 2959, 2073, 2749, 952]

[EC 3.1.3.76 created 2006]

### EC 3.1.3.77

- Accepted name:** acireductone synthase  
**Reaction:** 5-(methylsulfanyl)-2,3-dioxopentyl phosphate + H<sub>2</sub>O = 1,2-dihydroxy-5-(methylsulfanyl)pent-1-en-3-one + phosphate (overall reaction)  
(1a) 5-(methylsulfanyl)-2,3-dioxopentyl phosphate = 2-hydroxy-5-(methylsulfanyl)-3-oxopent-1-enyl phosphate (probably spontaneous)  
(1b) 2-hydroxy-5-(methylsulfanyl)-3-oxopent-1-enyl phosphate + H<sub>2</sub>O = 1,2-dihydroxy-5-(methylsulfanyl)pent-1-en-3-one + phosphate  
**Other name(s):** E1; E-1 enolase-phosphatase; 5-(methylthio)-2,3-dioxopentyl-phosphate phosphohydrolase (isomerizing)  
**Systematic name:** 5-(methylsulfanyl)-2,3-dioxopentyl-phosphate phosphohydrolase (isomerizing)  
**Comments:** This bifunctional enzyme first enolizes the substrate to form the intermediate 2-hydroxy-5-(methylsulfanyl)-3-oxopent-1-enyl phosphate, which is then dephosphorylated to form the acireductone 1,2-dihydroxy-5-(methylsulfanyl)pent-1-en-3-one [3215]. The acireductone represents a branch point in the methionine-salvage pathway as it is used in the formation of formate, CO and 3-(methylsulfanyl)propanoate by EC 1.13.11.53 [acireductone dioxygenase (Ni<sup>2+</sup>-requiring)] and of formate and 4-(methylsulfanyl)-2-oxobutanoate either by a spontaneous reaction under aerobic conditions or by EC 1.13.11.54 acireductone dioxygenase [iron(II)-requiring] [2016, 3215].  
**References:** [2016, 3215, 3109]

[EC 3.1.3.77 created 2006]



### EC 3.1.3.78

- Accepted name:** phosphatidylinositol-4,5-bisphosphate 4-phosphatase  
**Reaction:** 1-phosphatidyl-1D-*myo*-inositol 4,5-bisphosphate + H<sub>2</sub>O = 1-phosphatidyl-1D-*myo*-inositol 5-phosphate + phosphate  
**Other name(s):** phosphatidylinositol-4,5-bisphosphate 4-phosphatase I; phosphatidylinositol-4,5-bisphosphate 4-phosphatase II; type I PtdIns-4,5-P<sub>2</sub> 4-Ptase; type II PtdIns-4,5-P<sub>2</sub> 4-Ptase; IpgD; PtdIns-4,5-P<sub>2</sub> 4-phosphatase type I; PtdIns-4,5-P<sub>2</sub> 4-phosphatase type II; type I phosphatidylinositol-4,5-bisphosphate 4-phosphatase; type 1 4-phosphatase  
**Systematic name:** 1-phosphatidyl-1D-*myo*-inositol-4,5-bisphosphate 4-phosphohydrolase  
**Comments:** Two pathways exist in mammalian cells to degrade 1-phosphatidyl-1D-*myo*-inositol 4,5-bisphosphate [PtdIns(4,5)P<sub>2</sub>] [3013]. One is catalysed by this enzyme and the other by EC 3.1.3.36, phosphoinositide 5-phosphatase, where the product is PtdIns4P. The enzyme from human is specific for PtdIns(4,5)P<sub>2</sub> as substrate, as it cannot use PtdIns(3,4,5)P<sub>3</sub>, PtdIns(3,4)P<sub>2</sub>, PtdIns(3,5)P<sub>2</sub>, PtdIns5P, PtdIns4P or PtdIns3P [3013]. In humans, the enzyme is localized to late endosomal/lysosomal membranes [3349]. It can control nuclear levels of PtdIns5P and thereby control p53-dependent apoptosis [3349].  
**References:** [2082, 3013, 3349, 1833]

[EC 3.1.3.78 created 2008]

### EC 3.1.3.79

- Accepted name:** mannosylfructose-phosphate phosphatase  
**Reaction:** β-D-fructofuranosyl-α-D-mannopyranoside 6<sup>F</sup>-phosphate + H<sub>2</sub>O = β-D-fructofuranosyl-α-D-mannopyranoside + phosphate  
**Other name(s):** mannosylfructose-6-phosphate phosphatase; MFPP  
**Systematic name:** β-D-fructofuranosyl-α-D-mannopyranoside-6F-phosphate phosphohydrolase  
**Comments:** This enzyme, from the soil proteobacterium and plant pathogen *Agrobacterium tumefaciens* strain C58, requires Mg<sup>2+</sup> for activity. Mannosylfructose is the major endogenous osmolyte produced by several α-proteobacteria in response to osmotic stress and is synthesized by the sequential action of EC 2.4.1.246 (mannosylfructose-phosphate synthase) followed by this enzyme. While mannosylfructose 6-phosphate is the physiological substrate, the enzyme can use sucrose 6-phosphate very efficiently. The F in mannosylfructose 6<sup>F</sup>-phosphate is used to indicate that the fructose residue of sucrose carries the substituent.  
**References:** [2948]

[EC 3.1.3.79 created 2009]

### EC 3.1.3.80

- Accepted name:** 2,3-bisphosphoglycerate 3-phosphatase  
**Reaction:** 2,3-bisphospho-D-glycerate + H<sub>2</sub>O = 2-phospho-D-glycerate + phosphate  
**Other name(s):** MIPP1; 2,3-BPG 3-phosphatase  
**Systematic name:** 2,3-bisphospho-D-glycerate 3-phosphohydrolase  
**Comments:** This reaction is a shortcut in the Rapoport-Luebering shunt. It bypasses the reactions of EC 5.4.2.11/EC 5.4.2.12 [phosphoglycerate mutases (2,3-diphosphoglycerate-dependent and independent)] and directly forms 2-phospho-D-glycerate by removing the 3-phospho-group of 2,3-diphospho-D-glycerate [436]. The MIPP1 protein also catalyses the reaction of EC 3.1.3.62 (multiple inositol polyphosphate phosphatase).  
**References:** [436]

[EC 3.1.3.80 created 2010]

### EC 3.1.3.81

- Accepted name:** diacylglycerol diphosphate phosphatase  
**Reaction:** 1,2-diacyl-*sn*-glycerol 3-diphosphate + H<sub>2</sub>O = 1,2-diacyl-*sn*-glycerol 3-phosphate + phosphate



**Other name(s):** DGPP phosphatase; DGPP phosphohydrolase; DPP1; DPPL1; DPPL2; PAP2; pyrophosphate phosphatase  
**Systematic name:** 1,2-diacyl-*sn*-glycerol 3-phosphate phosphohydrolase  
**Comments:** The bifunctional enzyme catalyses the dephosphorylation of diacylglycerol diphosphate to phosphatidate and the subsequent dephosphorylation of phosphatidate to diacylglycerol (*cf.* phosphatidate phosphatase (EC 3.1.3.4)). It regulates intracellular levels of diacylglycerol diphosphate and phosphatidate, phospholipid molecules believed to play a signalling role in stress response [1040]. The phosphatase activity of the bifunctional enzyme is Mg<sup>2+</sup>-independent and N-ethylmaleimide-insensitive and is distinct from the Mg<sup>2+</sup>-dependent and N-ethylmaleimide-sensitive enzyme EC 3.1.3.4 (phosphatidate phosphatase) [372]. The diacylglycerol pyrophosphate phosphatase activity in *Saccharomyces cerevisiae* is induced by zinc depletion, by inositol supplementation, and when cells enter the stationary phase [2206].  
**References:** [615, 614, 3223, 2206, 372, 1040]

[EC 3.1.3.81 created 2010]

#### EC 3.1.3.82

**Accepted name:** D-glycero-β-D-manno-heptose 1,7-bisphosphate 7-phosphatase  
**Reaction:** D-glycero-β-D-manno-heptose 1,7-bisphosphate + H<sub>2</sub>O = D-glycero-β-D-manno-heptose 1-phosphate + phosphate  
**Other name(s):** *gmhB* (gene name); *yaeD* (gene name)  
**Systematic name:** D-glycero-β-D-manno-heptose 1,7-bisphosphate 7-phosphohydrolase  
**Comments:** The enzyme is involved in biosynthesis of ADP-L-glycero-β-D-manno-heptose, which is utilized for assembly of the lipopolysaccharide inner core in Gram-negative bacteria. *In vitro* the catalytic efficiency with the β-anomer is 100-200-fold higher than with the α-anomer [3113].  
**References:** [1482, 3031, 3113]

[EC 3.1.3.82 created 2010]

#### EC 3.1.3.83

**Accepted name:** D-glycero-α-D-manno-heptose 1,7-bisphosphate 7-phosphatase  
**Reaction:** D-glycero-α-D-manno-heptose 1,7-bisphosphate + H<sub>2</sub>O = D-glycero-α-D-manno-heptose 1-phosphate + phosphate  
**Other name(s):** *gmhB* (gene name)  
**Systematic name:** D-glycero-α-D-manno-heptose 1,7-bisphosphate 7-phosphohydrolase  
**Comments:** The enzyme is involved in biosynthesis of GDP-D-glycero-α-D-manno-heptose, which is required for assembly of S-layer glycoprotein in some Gram-positive bacteria. The *in vitro* catalytic efficiency of the enzyme from *Bacteroides thetaiotaomicron* is 6-fold higher with the α-anomer than with the β-anomer [3113].  
**References:** [3113]

[EC 3.1.3.83 created 2010]

#### EC 3.1.3.84

**Accepted name:** ADP-ribose 1''-phosphate phosphatase  
**Reaction:** ADP-D-ribose 1''-phosphate + H<sub>2</sub>O = ADP-D-ribose + phosphate  
**Other name(s):** POA1; Appr1p phosphatase; Poa1p; ADP-ribose 1''-phosphate phosphohydrolase  
**Systematic name:** ADP-D-ribose 1''-phosphate phosphohydrolase  
**Comments:** The enzyme is highly specific for ADP-D-ribose 1''-phosphate. Involved together with EC 3.1.4.37, 2',3'-cyclic-nucleotide 3'-phosphodiesterase, in the breakdown of adenosine diphosphate ribose 1'',2''-cyclic phosphate (Appr<sub>i</sub>p), a by-product of tRNA splicing.  
**References:** [2649]

[EC 3.1.3.84 created 2011]

### EC 3.1.3.85

- Accepted name:** glucosyl-3-phosphoglycerate phosphatase  
**Reaction:** 2-*O*-( $\alpha$ -D-glucopyranosyl)-3-phospho-D-glycerate + H<sub>2</sub>O = 2-*O*-( $\alpha$ -D-glucopyranosyl)-D-glycerate + phosphate  
**Other name(s):** GpgP protein  
**Systematic name:**  $\alpha$ -D-glucosyl-3-phospho-D-glycerate phosphohydrolase  
**Comments:** The enzyme is involved in biosynthesis of 2-*O*-( $\alpha$ -D-glucopyranosyl)-D-glycerate via the two-step pathway in which EC 2.4.1.266 (glucosyl-3-phosphoglycerate synthase) catalyses the conversion of GDP-glucose and 3-phospho-D-glycerate into 2-*O*-( $\alpha$ -D-glucopyranosyl)-3-phospho-D-glycerate, which is then converted to 2-*O*-( $\alpha$ -D-glucopyranosyl)-D-glycerate by glucosyl-3-phosphoglycerate phosphatase. *In vivo* the enzyme catalyses the dephosphorylation of 2-*O*-( $\alpha$ -D-mannopyranosyl)-3-phospho-D-glycerate with lower efficiency [493, 494]. Divalent metal ions (Mg<sup>2+</sup>, Mn<sup>2+</sup> or Co<sup>2+</sup>) stimulate activity [493, 494].  
**References:** [493, 494, 1885]

[EC 3.1.3.85 created 2011]

### EC 3.1.3.86

- Accepted name:** phosphatidylinositol-3,4,5-trisphosphate 5-phosphatase  
**Reaction:** 1-phosphatidyl-1D-*myo*-inositol 3,4,5-trisphosphate + H<sub>2</sub>O = 1-phosphatidyl-1D-*myo*-inositol 3,4-bisphosphate + phosphate  
**Other name(s):** SHIP1; SHIP2; SHIP; p150Ship  
**Systematic name:** 1-phosphatidyl-1D-*myo*-inositol-3,4,5-trisphosphate 5-phosphohydrolase  
**Comments:** This enzyme hydrolyses 1-phosphatidyl-1D-*myo*-inositol 3,4,5-trisphosphate (PtdIns(3,4,5)P<sub>3</sub>) to produce PtdIns(3,4)P<sub>2</sub>, thereby negatively regulating the PI3K (phosphoinositide 3-kinase) pathways. The enzyme also shows activity toward (PtdIns(1,3,4,5)P<sub>4</sub>) [2258]. The enzyme is involved in several signal transduction pathways in the immune system leading to an adverse range of effects.  
**References:** [1709, 533, 925, 650, 2258]

[EC 3.1.3.86 created 2011]

### EC 3.1.3.87

- Accepted name:** 2-hydroxy-3-keto-5-methylthiopent-1-enyl phosphate phosphatase  
**Reaction:** 2-hydroxy-5-(methylsulfanyl)-3-oxopent-1-enyl phosphate + H<sub>2</sub>O = 1,2-dihydroxy-5-(methylsulfanyl)pent-1-en-3-one + phosphate  
**Other name(s):** HK-MTPenyl-1-*P* phosphatase; MtnX; YkrX; 2-hydroxy-5-(methylthio)-3-oxopent-1-enyl phosphate phosphohydrolase  
**Systematic name:** 2-hydroxy-5-(methylsulfanyl)-3-oxopent-1-enyl phosphate phosphohydrolase  
**Comments:** The enzyme participates in the methionine salvage pathway in *Bacillus subtilis* [86]. In some species a single bifunctional enzyme, EC 3.1.3.77, acireductone synthase, catalyses both this reaction and EC 5.3.2.5, 2,3-diketo-5-methylthiopent-1-enyl phosphate enolase [2016].  
**References:** [2016, 86]

[EC 3.1.3.87 created 2012]

### EC 3.1.3.88

- Accepted name:** 5''-phosphoribostamycin phosphatase  
**Reaction:** 5''-phosphoribostamycin + H<sub>2</sub>O = ribostamycin + phosphate  
**Other name(s):** *btP* (gene name); *neol* (gene name)  
**Systematic name:** 5''-phosphoribostamycin phosphohydrolase  
**Comments:** Involved in the biosynthetic pathways of several clinically important aminocyclitol antibiotics, including ribostamycin, neomycin and butirosin. No metal is required for activity.  
**References:** [1543]

[EC 3.1.3.88 created 2012]

#### EC 3.1.3.89

- Accepted name:** 5'-deoxynucleotidase  
**Reaction:** a 2'-deoxyribonucleoside 5'-monophosphate + H<sub>2</sub>O = a 2'-deoxyribonucleoside + phosphate  
**Other name(s):** *yfbR* (gene name)  
**Systematic name:** 2'-deoxyribonucleoside 5'-monophosphate phosphohydrolase  
**Comments:** The enzyme, characterized from the bacterium *Escherichia coli*, shows strict specificity towards deoxyribonucleoside 5'-monophosphates and does not dephosphorylate 5'-ribonucleotides or ribonucleoside 3'-monophosphates. A divalent metal cation is required for activity, with cobalt providing the highest activity.  
**References:** [2327, 3342]

[EC 3.1.3.89 created 2013]

#### EC 3.1.3.90

- Accepted name:** maltose 6'-phosphate phosphatase  
**Reaction:** maltose 6'-phosphate + H<sub>2</sub>O = maltose + phosphate  
**Other name(s):** maltose 6'-P phosphatase; *mapP* (gene name)  
**Systematic name:** maltose 6'-phosphate phosphohydrolase  
**Comments:** The enzyme from the bacterium *Enterococcus faecalis* also has activity with the sucrose isomer turanose 6'-phosphate (α-D-glucopyranosyl-(1→3)-D-fructose 6-phosphate).  
**References:** [1947]

[EC 3.1.3.90 created 2013]

#### EC 3.1.3.91

- Accepted name:** 7-methylguanosine nucleotidase  
**Reaction:** (1) *N*<sup>7</sup>-methyl-GMP + H<sub>2</sub>O = *N*<sup>7</sup>-methyl-guanosine + phosphate  
(2) CMP + H<sub>2</sub>O = cytidine + phosphate  
**Other name(s):** cytosolic nucleotidase III-like; cNIII-like; *N*<sup>7</sup>-methylguanylate 5'-phosphatase  
**Systematic name:** *N*<sup>7</sup>-methyl-GMP phosphohydrolase  
**Comments:** The enzyme also has low activity with *N*<sup>7</sup>-methyl-GDP, producing *N*<sup>7</sup>-methyl-GMP. Does not accept AMP or GMP, and has low activity with UMP.  
**References:** [334]

[EC 3.1.3.91 created 2013]

#### EC 3.1.3.92

- Accepted name:** kanosamine-6-phosphate phosphatase  
**Reaction:** kanosamine 6-phosphate + H<sub>2</sub>O = kanosamine + phosphate  
**Other name(s):** *ntdB* (gene name)  
**Systematic name:** kanosamine-6-phosphate phosphohydrolase  
**Comments:** The enzyme, found in the bacterium *Bacillus subtilis*, is involved in a kanosamine biosynthesis pathway.  
**References:** [3064]

[EC 3.1.3.92 created 2013]

#### EC 3.1.3.93

- Accepted name:** L-galactose 1-phosphate phosphatase  
**Reaction:** β-L-galactose 1-phosphate + H<sub>2</sub>O = L-galactose + phosphate

**Other name(s):** VTC4 (gene name) (ambiguous); IMPL2 (gene name) (ambiguous)  
**Systematic name:**  $\beta$ -L-galactose-1-phosphate phosphohydrolase  
**Comments:** The enzyme from plants also has the activity of EC 3.1.3.25, inositol-phosphate phosphatase. The enzymes have very low activity with D-galactose 1-phosphate (*cf.* EC 3.1.3.94, D-galactose 1-phosphate phosphatase).  
**References:** [1590, 2947, 2262]

[EC 3.1.3.93 created 2014]

#### EC 3.1.3.94

**Accepted name:** D-galactose 1-phosphate phosphatase  
**Reaction:**  $\alpha$ -D-galactose 1-phosphate + H<sub>2</sub>O = D-galactose + phosphate  
**Systematic name:**  $\alpha$ -D-galactose-1-phosphate phosphohydrolase  
**Comments:** The human enzyme also has the activity of EC 3.1.3.25, inositol-phosphate phosphatase. The enzyme has very low activity with L-galactose 1-phosphate (*cf.* EC 3.1.3.93, L-galactose 1-phosphate phosphatase).  
**References:** [2241]

[EC 3.1.3.94 created 2014]

#### EC 3.1.3.95

**Accepted name:** phosphatidylinositol-3,5-bisphosphate 3-phosphatase  
**Reaction:** 1-phosphatidyl-1D-*myo*-inositol 3,5-bisphosphate + H<sub>2</sub>O = 1-phosphatidyl-1D-*myo*-inositol 5-phosphate + phosphate  
**Other name(s):** MTMR; PtdIns-3,5-P<sub>2</sub> 3-Ptase  
**Systematic name:** 1-phosphatidyl-1D-*myo*-inositol-3,5-bisphosphate 3-phosphohydrolase  
**Comments:** The enzyme is found in both plants and animals. It also has the activity of EC 3.1.3.64 (phosphatidylinositol-3-phosphatase).  
**References:** [3093, 194, 619]

[EC 3.1.3.95 created 2014]

#### EC 3.1.3.96

**Accepted name:** pseudouridine 5'-phosphatase  
**Reaction:** pseudouridine 5'-phosphate + H<sub>2</sub>O = pseudouridine + phosphate  
**Other name(s):** pseudouridine 5'-monophosphatase; 5'-PsiMPase; HDHD1  
**Systematic name:** pseudouridine 5'-phosphohydrolase  
**Comments:** Requires Mg<sup>2+</sup> for activity.  
**References:** [2324]

[EC 3.1.3.96 created 2014]

#### EC 3.1.3.97

**Accepted name:** 3',5'-nucleoside bisphosphate phosphatase  
**Reaction:** nucleoside 3',5'-bisphosphate + H<sub>2</sub>O = nucleoside 5'-phosphate + phosphate  
**Systematic name:** nucleoside-3',5'-bisphosphate 3'-phosphohydrolase  
**Comments:** The enzyme, characterized from the bacterium *Chromobacterium violaceum*, has similar catalytic efficiencies with all the bases. The enzyme has similar activity with ribonucleoside and 2'-deoxyribonucleoside 3',5'-bisphosphates, but shows no activity with nucleoside 2',5'-bisphosphates (*cf.* EC 3.1.3.7, 3'(2'),5'-bisphosphate nucleotidase).  
**References:** [516]

[EC 3.1.3.97 created 2015]

[3.1.3.98 Transferred entry. geranyl diphosphate phosphohydrolase, transferred to EC 3.6.1.68, geranyl diphosphate phosphohydrolase]

[EC 3.1.3.98 created 2015, deleted 2016]

#### EC 3.1.3.99

**Accepted name:** IMP-specific 5'-nucleotidase  
**Reaction:** IMP + H<sub>2</sub>O = inosine + phosphate  
**Other name(s):** ISN1 (gene name)  
**Systematic name:** inosine 5'-phosphate phosphohydrolase  
**Comments:** The enzyme, isolated from the yeast *Saccharomyces cerevisiae*, is highly specific for inosine 5'-phosphate, and has no detectable activity with other purine and pyrimidine nucleotides. Requires divalent metals, such as Mg<sup>2+</sup>, Co<sup>2+</sup> or Mn<sup>2+</sup>.  
**References:** [1270, 1271]

[EC 3.1.3.99 created 2016]

#### EC 3.1.3.100

**Accepted name:** thiamine phosphate phosphatase  
**Reaction:** thiamine phosphate + H<sub>2</sub>O = thiamine + phosphate  
**Systematic name:** thiamine phosphate phosphohydrolase  
**Comments:** The enzyme participates in the thiamine biosynthesis pathway in eukaryotes and a few prokaryotes. These organisms lack EC 2.7.4.16, thiamine-phosphate kinase, and need to convert thiamine phosphate to thiamine diphosphate, the active form of the vitamin, by first removing the phosphate group, followed by diphosphorylation by EC 2.7.6.2, thiamine diphosphokinase.  
**References:** [2517, 1504, 2596, 1997, 1500, 1920]

[EC 3.1.3.100 created 2016]

#### EC 3.1.3.101

**Accepted name:** validoxylamine A 7'-phosphate phosphatase  
**Reaction:** validoxylamine A 7'-phosphate + H<sub>2</sub>O = validoxylamine A + phosphate  
**Other name(s):** *vldH* (gene name)  
**Systematic name:** validoxylamine-A 7'-phosphate phosphohydrolase  
**Comments:** The enzyme, characterized from the bacterium *Streptomyces hygroscopicus* subsp. *limoneus*, is involved in the biosynthesis of the antifungal agent validamycin A.  
**References:** [81]

[EC 3.1.3.101 created 2016]

#### EC 3.1.3.102

**Accepted name:** FMN hydrolase  
**Reaction:** FMN + H<sub>2</sub>O = riboflavin + phosphate  
**Other name(s):** FMN phosphatase; AtpcFHy1  
**Systematic name:** FMN phosphohydrolase  
**Comments:** Requires Mg<sup>2+</sup>. The enzyme, found in many isoforms purified from both bacteria and plants, is a member of the haloacid dehalogenase superfamily. Most of the isoforms have a wide substrate specificity [1579], but isoforms specific for FMN also exist [2381].  
**References:** [2516, 1579, 2381]

[EC 3.1.3.102 created 2016]

#### EC 3.1.3.103

**Accepted name:** 3-deoxy-D-*glycero*-D-*galacto*-nonulopyranosonate 9-phosphatase  
**Reaction:** 3-deoxy-D-*glycero*-D-*galacto*-non-2-ulopyranosonate 9-phosphate + H<sub>2</sub>O = 3-deoxy-D-*glycero*-D-*galacto*-non-2-ulopyranosonate + phosphate  
**Other name(s):** 3-deoxy-D-*glycero*-D-*galacto*-non-2-ulopyranosonate-9-phosphate phosphatase  
**Systematic name:** 3-deoxy-D-*glycero*-D-*galacto*-non-2-ulopyranosonate 9-phosphohydrolase  
**Comments:** The enzyme, characterized from the bacterium *Bacteroides thetaiotaomicron*, is part of the biosynthesis pathway of the sialic acid 3-deoxy-D-*glycero*-D-*galacto*-non-2-ulopyranosonate (Kdn). Kdn is abundant in extracellular glycoconjugates of lower vertebrates such as fish and amphibians, but is also found in the capsular polysaccharides of bacteria that belong to the *Bacteroides* genus.  
**References:** [3114, 1747]

[EC 3.1.3.103 created 2016]

#### EC 3.1.3.104

**Accepted name:** 5-amino-6-(5-phospho-D-ribitylamino)uracil phosphatase  
**Reaction:** 5-amino-6-(5-phospho-D-ribitylamino)uracil + H<sub>2</sub>O = 5-amino-6-(D-ribitylamino)uracil + phosphate  
**Other name(s):** 5-amino-6-ribitylamino-2,4(1*H*,3*H*)-pyrimidinedione 5'-phosphate phosphatase  
**Systematic name:** 5-amino-6-(5-phospho-D-ribitylamino)uracil phosphohydrolase  
**Comments:** Requires Mg<sup>2+</sup>. The enzyme, which is found in plants and bacteria, is part of a pathway for riboflavin biosynthesis. Most forms of the enzyme has a broad substrate specificity [1013, 2529].  
**References:** [1013, 1735, 2529]

[EC 3.1.3.104 created 2016]

#### EC 3.1.3.105

**Accepted name:** *N*-acetyl-D-muramate 6-phosphate phosphatase  
**Reaction:** *N*-acetyl-D-muramate 6-phosphate + H<sub>2</sub>O = *N*-acetyl-D-muramate + phosphate  
**Other name(s):** *mupP* (gene name)  
**Systematic name:** *N*-acetyl-D-muramate 6-phosphate phosphohydrolase  
**Comments:** The enzyme, characterized from *Pseudomonas* species, participates in a peptidoglycan salvage pathway.  
**References:** [265]

[EC 3.1.3.105 created 2017]

### EC 3.1.4 Phosphoric-diester hydrolases

#### EC 3.1.4.1

**Accepted name:** phosphodiesterase I  
**Reaction:** Hydrolytically removes 5'-nucleotides successively from the 3'-hydroxy termini of 3'-hydroxy-terminated oligonucleotides  
**Other name(s):** 5'-exonuclease; 5'-phosphodiesterase; 5'-nucleotide phosphodiesterase; oligonucleate 5'-nucleotidohydrolase; 5' nucleotide phosphodiesterase/alkaline phosphodiesterase I; 5'-NPDase; 5'-PDase; 5'-PDE; 5'NPDE; alkaline phosphodiesterase; nucleotide pyrophosphatase/phosphodiesterase I; orthophosphoric diester phosphohydrolase; PDE I; phosphodiesterase (ambiguous); exonuclease I  
**Systematic name:** oligonucleotide 5'-nucleotidohydrolase  
**Comments:** Hydrolyses both ribonucleotides and deoxyribonucleotides. Has low activity towards polynucleotides. A 3'-phosphate terminus on the substrate inhibits hydrolysis.  
**References:** [1435]

[EC 3.1.4.1 created 1961]

#### EC 3.1.4.2

**Accepted name:** glycerophosphocholine phosphodiesterase  
**Reaction:** *sn*-glycero-3-phosphocholine + H<sub>2</sub>O = choline + *sn*-glycerol 3-phosphate  
**Other name(s):** glycerophosphinicocholine diesterase; glycerylphosphorylcholinediesterase; *sn*-glycero-3-phosphorylcholine diesterase; glycerolphosphorylcholine phosphodiesterase; glycerophosphohydro-lase  
**Systematic name:** *sn*-glycero-3-phosphocholine glycerophosphohydrolase  
**Comments:** Also acts on *sn*-glycero-3-phosphoethanolamine.  
**References:** [550, 1084, 3144]

[EC 3.1.4.2 created 1961, modified 1976]

#### EC 3.1.4.3

**Accepted name:** phospholipase C  
**Reaction:** a phosphatidylcholine + H<sub>2</sub>O = 1,2-diacyl-*sn*-glycerol + phosphocholine  
**Other name(s):** lipophosphodiesterase I; lecithinase C; *Clostridium welchii* α-toxin; *Clostridium oedematiens* β- and γ-toxins; lipophosphodiesterase C; phosphatidase C; heat-labile hemolysin; α-toxin  
**Systematic name:** phosphatidylcholine cholinephosphohydrolase  
**Comments:** The bacterial enzyme, which is a zinc protein, also acts on sphingomyelin and phosphatidylinositol; that from seminal plasma does not act on phosphatidylinositol.  
**References:** [656, 1712, 2625, 2846]

[EC 3.1.4.3 created 1961]

#### EC 3.1.4.4

**Accepted name:** phospholipase D  
**Reaction:** a phosphatidylcholine + H<sub>2</sub>O = choline + a phosphatidate  
**Other name(s):** lipophosphodiesterase II; lecithinase D; choline phosphatase  
**Systematic name:** phosphatidylcholine phosphatidohydrolase  
**Comments:** Also acts on other phosphatidyl esters.  
**References:** [91, 686, 1043, 2946]

[EC 3.1.4.4 created 1961]

[3.1.4.5 Transferred entry. deoxyribonuclease. Now EC 3.1.21.1, deoxyribonuclease I]

[EC 3.1.4.5 created 1961, deleted 1978]

[3.1.4.6 Transferred entry. deoxyribonuclease II. Now EC 3.1.22.1, deoxyribonuclease II]

[EC 3.1.4.6 created 1961, deleted 1978]

[3.1.4.7 Transferred entry. micrococcal nuclease. Now EC 3.1.31.1, micrococcal nuclease]

[EC 3.1.4.7 created 1961, deleted 1978]

[3.1.4.8 Transferred entry. *Aspergillus oryzae* ribonuclease. Now EC 3.1.27.3, ribonuclease T<sub>1</sub>]

[EC 3.1.4.8 created 1961, transferred 1965 to EC 2.7.7.26, reinstated 1972, deleted 1978]

[3.1.4.9 Transferred entry. nucleate endonuclease. Now EC 3.1.30.2, *Serratia marcescens* nuclease]

[EC 3.1.4.9 created 1965, deleted 1978]

[3.1.4.10 Transferred entry. 1-phosphatidylinositol phosphodiesterase. Now EC 4.6.1.13, phosphatidylinositol diacylglycerol-lyase. As there is no hydrolysis of the inositol 1,2-cyclic phosphate formed, previous classification of the enzyme as a hydrolase was incorrect]

[EC 3.1.4.10 created 1972, modified 1976, deleted 2002]



#### EC 3.1.4.11

- Accepted name:** phosphoinositide phospholipase C  
**Reaction:** 1-phosphatidyl-1D-*myo*-inositol 4,5-bisphosphate + H<sub>2</sub>O = 1D-*myo*-inositol 1,4,5-trisphosphate + diacylglycerol  
**Other name(s):** triphosphoinositide phosphodiesterase; phosphoinositidase C; 1-phosphatidylinositol-4,5-bisphosphate phosphodiesterase; monophosphatidylinositol phosphodiesterase; phosphatidylinositol phospholipase C; PI-PLC; 1-phosphatidyl-D-*myo*-inositol-4,5-bisphosphate inositoltrisphosphohydrolyase  
**Systematic name:** 1-phosphatidyl-1D-*myo*-inositol-4,5-bisphosphate inositoltrisphosphohydrolase  
**Comments:** These enzymes form some of the cyclic phosphate Ins(cyclic1,2)P(4,5)P<sub>2</sub> as well as Ins(1,4,5)P<sub>3</sub>. They show activity towards phosphatidylinositol, i.e., the activity of EC 4.6.1.13, phosphatidylinositol diacylglycerol-lyase, in vitro at high [Ca<sup>2+</sup>]. Four β-isofoms regulated by G-proteins, two γ-forms regulated by tyrosine kinases, four δ-forms regulated at least in part by calcium and an ε-form, probably regulated by the oncogene ras, have been found.  
**References:** [642, 2917, 2414]

[EC 3.1.4.11 created 1972, modified 2002]

#### EC 3.1.4.12

- Accepted name:** sphingomyelin phosphodiesterase  
**Reaction:** a sphingomyelin + H<sub>2</sub>O = a ceramide + phosphocholine  
**Other name(s):** neutral sphingomyelinase  
**Systematic name:** sphingomyelin cholinephosphohydrolase  
**Comments:** Has very little activity on phosphatidylcholine.  
**References:** [149, 404, 1107, 1383]

[EC 3.1.4.12 created 1972]

#### EC 3.1.4.13

- Accepted name:** serine-ethanolaminephosphate phosphodiesterase  
**Reaction:** serine phosphoethanolamine + H<sub>2</sub>O = serine + ethanolamine phosphate  
**Other name(s):** serine ethanolamine phosphodiester phosphodiesterase; SEP diesterase  
**Systematic name:** serine-phosphoethanolamine ethanolaminephosphohydrolase  
**Comments:** Acts only on those phosphodiesters that have ethanolamine as a component part of the molecule.  
**References:** [1022]

[EC 3.1.4.13 created 1972, modified 1976]

#### EC 3.1.4.14

- Accepted name:** [acyl-carrier-protein] phosphodiesterase  
**Reaction:** holo-[acyl-carrier protein] + H<sub>2</sub>O = 4'-phosphopantetheine + apo-[acyl-carrier protein]  
**Other name(s):** ACP hydrolyase; ACP phosphodiesterase; AcpH; [acyl-carrier-protein] 4'-pantetheine-phosphohydrolase; holo-[acyl-carrier-protein] 4'-pantetheine-phosphohydrolase  
**Systematic name:** holo-[acyl-carrier protein] 4'-pantetheine-phosphohydrolase  
**Comments:** The enzyme cleaves acyl-[acyl-carrier-protein] species with acyl chains of 6-16 carbon atoms although it appears to demonstrate a preference for the unacylated acyl-carrier protein (ACP) and short-chain ACPs over the medium- and long-chain species [2913]. Deletion of the gene encoding this enzyme abolishes ACP prosthetic-group turnover in vivo [2913]. Activation of apo-ACP to form the holoenzyme is carried out by EC 2.7.8.7, holo-[acyl-carrier-protein] synthase.  
**References:** [2706, 3027, 2913]

[EC 3.1.4.14 created 1972, modified 2006]

[3.1.4.15 Transferred entry. *adenylyl-[glutamateammonia ligase] hydrolase*. As it has been shown that the enzyme catalyses



a transfer of the adenylyl group to phosphate, the enzyme has been transferred to EC 2.7.7.89, adenylyl-[glutamateammonia ligase] phosphorylase]

[EC 3.1.4.15 created 1972, deleted 2015]

#### EC 3.1.4.16

**Accepted name:** 2',3'-cyclic-nucleotide 2'-phosphodiesterase  
**Reaction:** nucleoside 2',3'-cyclic phosphate + H<sub>2</sub>O = nucleoside 3'-phosphate  
**Other name(s):** ribonucleoside 2',3'-cyclic phosphate diesterase; 2',3'-cyclic AMP phosphodiesterase; 2',3'-cyclic nucleotidase; cyclic 2',3'-nucleotide 2'-phosphodiesterase; cyclic 2',3'-nucleotide phosphodiesterase; 2',3'-cyclic nucleoside monophosphate phosphodiesterase; 2',3'-cyclic AMP 2'-phosphohydrolase; cyclic phosphodiesterase:3'-nucleotidase; 2',3'-cyclic nucleotide phosphohydrolase; 2':3'-cyclic phosphodiesterase; 2':3'-cyclic nucleotide phosphodiesterase:3'-nucleotidase  
**Systematic name:** nucleoside-2',3'-cyclic-phosphate 3'-nucleotidohydrolase  
**Comments:** Also hydrolyses 3'-nucleoside monophosphates and bis-4-nitrophenyl phosphate, but not 3'-deoxynucleotides. Similar reactions are carried out by EC 3.1.27.3 (ribonuclease T<sub>1</sub>) and EC 3.1.27.5 (pancreatic ribonuclease).  
**References:** [58, 59, 385, 2191, 3011]

[EC 3.1.4.16 created 1972, modified 1976]

#### EC 3.1.4.17

**Accepted name:** 3',5'-cyclic-nucleotide phosphodiesterase  
**Reaction:** nucleoside 3',5'-cyclic phosphate + H<sub>2</sub>O = nucleoside 5'-phosphate  
**Other name(s):** cyclic 3',5'-mononucleotide phosphodiesterase; PDE; cyclic 3',5'-nucleotide phosphodiesterase; cyclic 3',5'-phosphodiesterase; 3',5'-nucleotide phosphodiesterase; 3':5'-cyclic nucleotide 5'-nucleotidohydrolase; 3',5'-cyclonucleotide phosphodiesterase; cyclic nucleotide phosphodiesterase; 3', 5'-cyclic nucleoside monophosphate phosphodiesterase; 3': 5'-monophosphate phosphodiesterase (cyclic CMP); cytidine 3':5'-monophosphate phosphodiesterase (cyclic CMP); cyclic 3',5'-nucleotide monophosphate phosphodiesterase; nucleoside 3',5'-cyclic phosphate diesterase; nucleoside-3',5'-monophosphate phosphodiesterase  
**Systematic name:** 3',5'-cyclic-nucleotide 5'-nucleotidohydrolase  
**Comments:** Acts on 3',5'-cyclic AMP, 3',5'-cyclic dAMP, 3',5'-cyclic IMP, 3',5'-cyclic GMP and 3',5'-cyclic CMP.  
**References:** [773, 2029]

[EC 3.1.4.17 created 1972, modified 1976]

[3.1.4.18 Transferred entry. phosphodiesterase II. Now EC 3.1.16.1, spleen exonuclease]

[EC 3.1.4.18 created 1972, deleted 1978]

[3.1.4.19 Transferred entry. oligonucleotidase. Now EC 3.1.13.3, oligonucleotidase]

[EC 3.1.4.19 created 1972, deleted 1978]

[3.1.4.20 Transferred entry. exoribonuclease. Now EC 3.1.13.1, exoribonuclease II]

[EC 3.1.4.20 created 1972, deleted 1978]

[3.1.4.21 Transferred entry. single-stranded-nucleate endonuclease. Now EC 3.1.30.1, Aspergillus nuclease S<sub>1</sub>]

[EC 3.1.4.21 created 1972, deleted 1978]

[3.1.4.22 Transferred entry. ribonuclease I. Now EC 3.1.27.5, pancreatic ribonuclease]

[EC 3.1.4.22 created 1972, deleted 1978]

[3.1.4.23 Transferred entry. ribonuclease II. Now EC 3.1.27.1, ribonuclease T<sub>2</sub>]

[EC 3.1.4.23 created 1972, deleted 1978]

[3.1.4.24 Deleted entry. *endoribonuclease III*]

[EC 3.1.4.24 created 1972, deleted 1978]

[3.1.4.25 Transferred entry. *exodeoxyribonuclease I*. Now EC 3.1.11.1, *exodeoxyribonuclease I*]

[EC 3.1.4.25 created 1972, deleted 1978]

[3.1.4.26 Deleted entry. *exodeoxyribonuclease II*]

[EC 3.1.4.26 created 1972, deleted 1978]

[3.1.4.27 Transferred entry. *exodeoxyribonuclease III*. Now EC 3.1.11.2, *exodeoxyribonuclease III*]

[EC 3.1.4.27 created 1972, deleted 1978]

[3.1.4.28 Transferred entry. *exodeoxyribonuclease IV*. Now EC 3.1.11.3, *exodeoxyribonuclease (lambda-induced)*]

[EC 3.1.4.28 created 1972, deleted 1978]

[3.1.4.29 Deleted entry. *oligodeoxyribonucleate exonuclease*]

[EC 3.1.4.29 created 1972, deleted 1978]

[3.1.4.30 Transferred entry. *endodeoxyribonuclease*. Now EC 3.1.21.2, *deoxyribonuclease IV (phage-T<sub>4</sub>-induced)*]

[EC 3.1.4.30 created 1972, deleted 1978]

[3.1.4.31 Transferred entry. *DNA 5'-dinucleotidohydrolase*. Now EC 3.1.11.4, *exodeoxyribonuclease (phage SP<sub>3</sub>-induced)*]

[EC 3.1.4.31 created 1972, deleted 1978]

[3.1.4.32 Deleted entry. *endodeoxyribonuclease (ATP- and S-adenosylmethionine-dependent)*. See EC 3.1.21.3 *type I site-specific deoxyribonuclease* and EC 3.1.21.5 *type III site-specific deoxyribonuclease*]

[EC 3.1.4.32 created 1972, deleted 1978]

[3.1.4.33 Deleted entry. *endodeoxyribonuclease (ATP-hydrolysing)*. See EC 3.1.21.3 *type I site-specific deoxyribonuclease* and EC 3.1.21.5 *type III site-specific deoxyribonuclease*]

[EC 3.1.4.33 created 1972, deleted 1978]

[3.1.4.34 Deleted entry. *hybrid nuclease*. See subclasses EC 3.1.15, EC 3.1.16, EC 3.1.30 and EC 3.1.31]

[EC 3.1.4.34 created 1972, deleted 1978]

#### EC 3.1.4.35

**Accepted name:** 3',5'-cyclic-GMP phosphodiesterase  
**Reaction:** guanosine 3',5'-cyclic phosphate + H<sub>2</sub>O = GMP  
**Other name(s):** guanosine cyclic 3',5'-phosphate phosphodiesterase; cyclic GMP phosphodiesterase; cyclic 3',5'-GMP phosphodiesterase; cyclic guanosine 3',5'-monophosphate phosphodiesterase; cyclic guanosine 3',5'-phosphate phosphodiesterase; cGMP phosphodiesterase; cGMP-PDE  
**Systematic name:** 3',5'-cyclic-GMP 5'-nucleotidohydrolase  
**References:** [1815]

[EC 3.1.4.35 created 1976]

[3.1.4.36 Deleted entry. *1,2-cyclic-inositol-phosphate phosphodiesterase*. Now included with EC 3.1.4.43, *glycerophosphoinositol inositolphosphodiesterase*]

[EC 3.1.4.36 created 1976, deleted 2002]

#### EC 3.1.4.37

- Accepted name:** 2',3'-cyclic-nucleotide 3'-phosphodiesterase  
**Reaction:** nucleoside 2',3'-cyclic phosphate + H<sub>2</sub>O = nucleoside 2'-phosphate  
**Other name(s):** cyclic-CMP phosphodiesterase; 2',3'-cyclic AMP phosphodiesterase; cyclic 2',3'-nucleotide 3'-phosphodiesterase; cyclic 2',3'-nucleotide phosphodiesterase; 2',3'-cyclic nucleoside monophosphate phosphodiesterase; 2',3'-cyclic nucleotide 3'-phosphohydrolase; CNPase; 2',3'-cyclic nucleotide phosphohydrolase; 2':3'-cyclic nucleotide 3'-phosphodiesterase; 2':3'-CNMP-3'-ase  
**Systematic name:** nucleoside-2',3'-cyclic-phosphate 2'-nucleotidohydrolase  
**Comments:** The brain enzyme acts on 2',3'-cyclic AMP more rapidly than on the UMP or CMP derivatives. An enzyme from liver acts on 2',3'-cyclic CMP more rapidly than on the purine derivatives; it also hydrolyses the corresponding 3',5'-cyclic phosphates, but more slowly. This latter enzyme has been called cyclic-CMP phosphodiesterase.  
**References:** [654, 1105, 1106, 1568, 2098]

[EC 3.1.4.37 created 1976]

#### EC 3.1.4.38

- Accepted name:** glycerophosphocholine cholinephosphodiesterase  
**Reaction:** *sn*-glycero-3-phosphocholine + H<sub>2</sub>O = glycerol + phosphocholine  
**Other name(s):** L-3-glycerylphosphinicocholine cholinephosphohydrolase  
**Systematic name:** *sn*-glycero-3-phosphocholine cholinephosphohydrolase  
**Comments:** No activity on *sn*-3-glycerophosphoethanolamine.  
**References:** [7]

[EC 3.1.4.38 created 1976]

#### EC 3.1.4.39

- Accepted name:** alkylglycerophosphoethanolamine phosphodiesterase  
**Reaction:** 1-alkyl-*sn*-glycero-3-phosphoethanolamine + H<sub>2</sub>O = 1-alkyl-*sn*-glycerol 3-phosphate + ethanolamine  
**Other name(s):** lysophospholipase D  
**Systematic name:** 1-alkyl-*sn*-glycero-3-phosphoethanolamine ethanolaminehydrolase  
**Comments:** Also acts on acyl and choline analogues.  
**References:** [3226]

[EC 3.1.4.39 created 1976]

#### EC 3.1.4.40

- Accepted name:** CMP-*N*-acylneuraminate phosphodiesterase  
**Reaction:** CMP-*N*-acylneuraminate + H<sub>2</sub>O = CMP + *N*-acylneuraminate  
**Other name(s):** CMP-sialate hydrolase; CMP-sialic acid hydrolase; CMP-*N*-acylneuraminic acid hydrolase; cytidine monophosphosialic hydrolase; cytidine monophosphosialate hydrolase; cytidine monophosphate-*N*-acetylneuraminic acid hydrolase; CMP-*N*-acetylneuraminate hydrolase  
**Systematic name:** CMP-*N*-acylneuraminate *N*-acylneuraminohydrolase  
**References:** [1409]

[EC 3.1.4.40 created 1976]

#### EC 3.1.4.41

- Accepted name:** sphingomyelin phosphodiesterase D  
**Reaction:** sphingomyelin + H<sub>2</sub>O = ceramide phosphate + choline  
**Other name(s):** sphingomyelinase D  
**Systematic name:** sphingomyelin ceramide-phosphohydrolase  
**Comments:** Does not act on phosphatidylcholine, but hydrolyses 2-lysophosphatidylcholine to choline and 2-lysophosphatidate.

**References:** [375, 2731]

[EC 3.1.4.41 created 1978]

#### EC 3.1.4.42

**Accepted name:** glycerol-1,2-cyclic-phosphate 2-phosphodiesterase  
**Reaction:** glycerol 1,2-cyclic phosphate + H<sub>2</sub>O = glycerol 1-phosphate  
**Other name(s):** rac-glycerol 1:2-cyclic phosphate 2-phosphodiesterase  
**Systematic name:** rac-glycerol-1,2-cyclic-phosphate 2-glycerophosphohydrolase  
**Comments:** Acts on both stereoisomers of the substrate and also, more slowly, on 3',5'-cyclic AMP and on 2',3'-cyclic AMP.  
**References:** [459]

[EC 3.1.4.42 created 1984]

#### EC 3.1.4.43

**Accepted name:** glycerophosphoinositol inositolphosphodiesterase  
**Reaction:** 1-(*sn*-glycero-3-phospho)-1D-*myo*-inositol + H<sub>2</sub>O = glycerol + 1D-*myo*-inositol 1-phosphate  
**Other name(s):** 1,2-cyclic-inositol-phosphate phosphodiesterase; D-*myo*-inositol 1:2-cyclic phosphate 2-phosphohydrolase; D-inositol 1,2-cyclic phosphate 2-phosphohydrolase; D-*myo*-inositol 1,2-cyclic phosphate 2-phosphohydrolase; 1-D-*myo*-inositol-1,2-cyclic-phosphate 2-inositolphosphohydrolase; inositol-1,2-cyclic-phosphate 2-inositolphosphohydrolase  
**Systematic name:** 1-(*sn*-glycero-3-phospho)-1D-*myo*-inositol inositolphosphohydrolase  
**Comments:** This enzyme also hydrolyses Ins(cyclic1,2)*P* to Ins-1-*P*  
**References:** [554, 552, 553, 2463]

[EC 3.1.4.43 created 1984, (EC 3.1.4.36 created 1976, incorporated 2002), modified 2002]

#### EC 3.1.4.44

**Accepted name:** glycerophosphoinositol glycerophosphodiesterase  
**Reaction:** 1-(*sn*-glycero-3-phospho)-1D-*myo*-inositol + H<sub>2</sub>O = *myo*-inositol + *sn*-glycero 3-phosphate  
**Other name(s):** *sn*-glycero(3)phosphoinositol glycerophosphohydrolase; *sn*-glycero-3-phospho-1-inositol glycerophosphohydrolase  
**Systematic name:** 1-(*sn*-glycero-3-phospho)-1D-*myo*-inositol glycerophosphohydrolase  
**References:** [555]

[EC 3.1.4.44 created 1984, modified 2002]

#### EC 3.1.4.45

**Accepted name:** *N*-acetylglucosamine-1-phosphodiester  $\alpha$ -*N*-acetylglucosaminidase  
**Reaction:** glycoprotein *N*-acetyl-D-glucosaminyl-phospho-D-mannose + H<sub>2</sub>O = *N*-acetyl-D-glucosamine + glycoprotein phospho-D-mannose  
**Other name(s):**  $\alpha$ -*N*-acetylglucosaminyl phosphodiesterase; lysosomal  $\alpha$ -*N*-acetylglucosaminidase; phosphodiester glycosidase;  $\alpha$ -*N*-acetyl-D-glucosamine-1-phosphodiester *N*-acetylglucosaminidase; 2-acetamido-2-deoxy- $\alpha$ -D-glucose 1-phosphodiester acetamidodeoxyglucohydrolase  
**Systematic name:** glycoprotein-*N*-acetyl-D-glucosaminyl-phospho-D-mannose *N*-acetyl-D-glucosaminylphosphohydrolase  
**Comments:** Acts on a variety of compounds in which *N*-acetyl-D-glucosamine is  $\alpha$ -linked to a phosphate group, including the biosynthetic intermediates of the high mannose oligosaccharide components of some lysosomal enzymes and the products of EC 2.7.8.17 UDP-*N*-acetylglucosamine—lysosomal-enzyme *N*-acetylglucosaminephosphotransferase.  
**References:** [581, 3037, 3039, 3083]

[EC 3.1.4.45 created 1984]

#### EC 3.1.4.46

**Accepted name:** glycerophosphodiester phosphodiesterase  
**Reaction:** a glycerophosphodiester + H<sub>2</sub>O = an alcohol + *sn*-glycerol 3-phosphate  
**Other name(s):** gene hpd protein; glycerophosphoryl diester phosphodiesterase; IgD-binding protein D  
**Systematic name:** glycerophosphodiester glycerophosphohydrolase  
**Comments:** Broad specificity for glycerophosphodiesters; glycerophosphocholine, glycerophosphoethanolamine, glycerophosphoglycerol and bis(glycerophospho)-glycerol are hydrolysed.  
**References:** [1606]

[EC 3.1.4.46 created 1986]

[3.1.4.47 *Transferred entry. variant-surface-glycoprotein phospholipase C. Now EC 4.6.1.14, glycosylphosphatidylinositol diacylglycerol-lyase*]

[EC 3.1.4.47 created 1989, deleted 2002]

#### EC 3.1.4.48

**Accepted name:** dolichylphosphate-glucose phosphodiesterase  
**Reaction:** dolichyl β-D-glucosyl phosphate + H<sub>2</sub>O = dolichyl phosphate + D-glucose  
**Other name(s):** dolichol phosphoglucose phosphodiesterase; Dol-*P*-Glc phosphodiesterase  
**Systematic name:** dolichyl-β-D-glucosyl-phosphate dolichylphosphohydrolase  
**References:** [502]

[EC 3.1.4.48 created 1989]

#### EC 3.1.4.49

**Accepted name:** dolichylphosphate-mannose phosphodiesterase  
**Reaction:** dolichyl β-D-mannosyl phosphate + H<sub>2</sub>O = dolichyl phosphate + D-mannose  
**Other name(s):** mannosylphosphodolichol phosphodiesterase  
**Systematic name:** dolichyl-β-D-mannosyl-phosphate dolichylphosphohydrolase  
**References:** [2936]

[EC 3.1.4.49 created 1990]

#### EC 3.1.4.50

**Accepted name:** glycosylphosphatidylinositol phospholipase D  
**Reaction:** 6-(α-D-glucosaminyl)-1-phosphatidyl-1D-*myo*-inositol + H<sub>2</sub>O = 6-(α-D-glucosaminyl)-1D-*myo*-inositol + 3-*sn*-phosphatidate  
**Other name(s):** GPI-PLD; glycoprotein phospholipase D; phosphatidylinositol phospholipase D; phosphatidylinositol-specific phospholipase D  
**Systematic name:** glycoprotein-phosphatidylinositol phosphatidohydrolase  
**Comments:** This enzyme is also active when *O*-4 of the glucosamine is substituted by carrying the oligosaccharide that can link a protein to the structure. It therefore cleaves proteins from the lipid part of the glycosylphosphatidylinositol (GPI) anchors, but does so by hydrolysis, whereas glycosylphosphatidylinositol diacylglycerol-lyase (EC 4.6.1.14) does so by elimination. It acts on plasma membranes only after solubilization of the substrate with detergents or solvents, but it may act on intracellular membranes.  
**References:** [1742, 1791, 1665, 567]

[EC 3.1.4.50 created 1990, modified 2002]

#### EC 3.1.4.51

**Accepted name:** glucose-1-phospho-D-mannosylglycoprotein phosphodiesterase  
**Reaction:** 6-(D-glucose-1-phospho)-D-mannosylglycoprotein + H<sub>2</sub>O = α-D-glucose 1-phosphate + D-mannosylglycoprotein  
**Other name(s):** α-glucose-1-phosphate phosphodiesterase  
**Systematic name:** 6-(D-glucose-1-phospho)-D-mannosylglycoprotein glucose-1-phosphohydrolase  
**Comments:** The enzyme is specific for the product of EC 2.7.8.19 UDP-glucose—glycoprotein glucose phosphotransferase.  
**References:** [2748]

[EC 3.1.4.51 created 1992]

#### EC 3.1.4.52

**Accepted name:** cyclic-guanylate-specific phosphodiesterase  
**Reaction:** cyclic di-3',5'-guanylate + H<sub>2</sub>O = 5'-phosphoguanylyl(3'→5')guanosine  
**Other name(s):** cyclic bis(3'→5')diguanylate phosphodiesterase; c-di-GMP-specific phosphodiesterase; c-di-GMP phosphodiesterase; phosphodiesterase (misleading); phosphodiesterase A1; PDEA1; VieA  
**Systematic name:** cyclic bis(3'→5')diguanylate 3'-guanylylhydrolase  
**Comments:** Requires Mg<sup>2+</sup> or Mn<sup>2+</sup> for activity and is inhibited by Ca<sup>2+</sup> and Zn<sup>2+</sup>. Contains a heme unit. This enzyme linearizes cyclic di-3',5'-guanylate, the product of EC 2.7.7.65, diguanylate cyclase and an allosteric activator of EC 2.4.1.12, cellulose synthase (UDP-forming), rendering it inactive [392]. It is the balance between these two enzymes that determines the cellular level of c-di-GMP [392].  
**References:** [392, 442, 2570, 2866]

[EC 3.1.4.52 created 2008]

#### EC 3.1.4.53

**Accepted name:** 3',5'-cyclic-AMP phosphodiesterase  
**Reaction:** adenosine 3',5'-cyclic phosphate + H<sub>2</sub>O = AMP  
**Other name(s):** cAMP-specific phosphodiesterase; cAMP-specific PDE; PDE1; PDE2A; PDE2B; PDE4; PDE7; PDE8; PDEB1; PDEB2  
**Systematic name:** 3',5'-cyclic-AMP 5'-nucleotidohydrolase  
**Comments:** Requires Mg<sup>2+</sup> or Mn<sup>2+</sup> for activity [117]. This enzyme is specific for 3',5'-cAMP and does not hydrolyse other nucleoside 3',5'-cyclic phosphates such as cGMP (*cf.* EC 3.1.4.17, 3,5-cyclic-nucleotide phosphodiesterase and EC 3.1.4.35, 3,5-cyclic-GMP phosphodiesterase). It is involved in modulation of the levels of cAMP, which is a mediator in the processes of cell transformation and proliferation [2376].  
**References:** [39, 117, 2376, 1324, 1751, 1240]

[EC 3.1.4.53 created 2008, modified 2011]

#### EC 3.1.4.54

**Accepted name:** *N*-acetylphosphatidylethanolamine-hydrolysing phospholipase D  
**Reaction:** *N*-acylphosphatidylethanolamine + H<sub>2</sub>O = *N*-acylethanolamine + a 1,2-diacylglycerol 3-phosphate  
**Other name(s):** NAPE-PLD; anandamide-generating phospholipase D; *N*-acyl phosphatidylethanolamine phospholipase D; NAPE-hydrolyzing phospholipase D  
**Systematic name:** *N*-acetylphosphatidylethanolamine phosphatidohydrolase  
**Comments:** This enzyme is involved in the biosynthesis of anandamide. It does not hydrolyse phosphatidylcholine and phosphatidylethanolamine [2184]. No transphosphatidation [2184]. The enzyme contains Zn<sup>2+</sup> and is activated by Mg<sup>2+</sup> or Ca<sup>2+</sup> [3111].  
**References:** [2184, 3111]

[EC 3.1.4.54 created 2011]

#### EC 3.1.4.55

- Accepted name:** phosphoribosyl 1,2-cyclic phosphate phosphodiesterase  
**Reaction:** 5-phospho- $\alpha$ -D-ribose 1,2-cyclic phosphate + H<sub>2</sub>O =  $\alpha$ -D-ribose 1,5-bisphosphate  
**Other name(s):** *phnP* (gene name)  
**Systematic name:** 5-phospho- $\alpha$ -D-ribose 1,2-cyclic phosphate 2-phosphohydrolase ( $\alpha$ -D-ribose 1,5-bisphosphate-forming)  
**Comments:** Binds Mn<sup>2+</sup> and Zn<sup>2+</sup>. Isolated from the bacterium *Escherichia coli*, where it participates in the degradation of methylphosphonate.  
**References:** [2296, 1190, 1091]

[EC 3.1.4.55 created 2013]

#### EC 3.1.4.56

- Accepted name:** 7,8-dihydroneopterin 2',3'-cyclic phosphate phosphodiesterase  
**Reaction:** (1) 7,8-dihydroneopterin 2',3'-cyclic phosphate + H<sub>2</sub>O = 7,8-dihydroneopterin 3'-phosphate  
(2) 7,8-dihydroneopterin 2',3'-cyclic phosphate + H<sub>2</sub>O = 7,8-dihydroneopterin 2'-phosphate  
**Other name(s):** MptB  
**Systematic name:** 7,8-dihydroneopterin 2',3'-cyclic phosphate 2'/3'-phosphodiesterase  
**Comments:** Contains one zinc atom and one iron atom per subunit of the dodecameric enzyme. It hydrolyses 7,8-dihydroneopterin 2',3'-cyclic phosphate, a step in tetrahydromethanopterin biosynthesis. *In vitro* the enzyme forms 7,8-dihydroneopterin 2'-phosphate and 7,8-dihydroneopterin 3'-phosphate at a ratio of 4:1.  
**References:** [1832]

[EC 3.1.4.56 created 2013]

#### EC 3.1.4.57

- Accepted name:** phosphoribosyl 1,2-cyclic phosphate 1,2-diphosphodiesterase  
**Reaction:** (1) 5-phospho- $\alpha$ -D-ribose 1,2-cyclic phosphate + H<sub>2</sub>O = D-ribofuranose 2,5-bisphosphate  
(2) D-ribofuranose 2,5-bisphosphate + H<sub>2</sub>O = D-ribofuranose 5-phosphate + phosphate  
**Other name(s):** cyclic phosphate dihydrolase; *phnPP* (gene name)  
**Systematic name:** 5-phospho- $\alpha$ -D-ribose 1,2-cyclic phosphate 1,2-diphosphophosphohydrolase  
**Comments:** The enzyme, characterized from the bacterium *Eggerthella lenta*, is involved in degradation of methylphosphonate.  
**References:** [904]

[EC 3.1.4.57 created 2014]

#### EC 3.1.4.58

- Accepted name:** RNA 2',3'-cyclic 3'-phosphodiesterase  
**Reaction:** (ribonucleotide)<sub>n</sub>-2',3'-cyclic phosphate + H<sub>2</sub>O = (ribonucleotide)<sub>n</sub>-2'-phosphate  
**Other name(s):** *thpR* (gene name); *ligT* (gene name)  
**Systematic name:** (ribonucleotide)<sub>n</sub>-2',3'-cyclic phosphate 3'-nucleotidohydrolase  
**Comments:** The enzyme hydrolyses RNA 2',3'-cyclic phosphodiester to an RNA 2'-phosphomonoester. *In vitro* the enzyme can also ligate tRNA molecules with 2',3'-cyclic phosphate to tRNA with 5'-hydroxyl termini, forming a 2'-5' phosphodiester linkage. However, the ligase activity is unlikely to be relevant *in vivo*.  
**References:** [1376, 2407]

[EC 3.1.4.58 created 2017]

### EC 3.1.5 Triphosphoric-monoester hydrolases

#### EC 3.1.5.1

**Accepted name:** dGTPase  
**Reaction:**  $\text{dGTP} + \text{H}_2\text{O} = \text{deoxyguanosine} + \text{triphosphate}$   
**Other name(s):** deoxy-GTPase; deoxyguanosine 5-triphosphate triphosphohydrolase; deoxyguanosine triphosphatase; deoxyguanosine triphosphate triphosphohydrolase  
**Systematic name:** dGTP triphosphohydrolase  
**Comments:** Also acts on GTP.  
**References:** [1514]

[EC 3.1.5.1 created 1961]

### EC 3.1.6 Sulfuric-ester hydrolases

#### EC 3.1.6.1

**Accepted name:** arylsulfatase  
**Reaction:**  $\text{a phenol sulfate} + \text{H}_2\text{O} = \text{a phenol} + \text{sulfate}$   
**Other name(s):** sulfatase; nitrocatechol sulfatase; phenolsulfatase; phenylsulfatase; *p*-nitrophenyl sulfatase; arylsulfohydrolase; 4-methylumbelliferyl sulfatase; estrogen sulfatase  
**Systematic name:** aryl-sulfate sulfohydrolase  
**Comments:** A group of enzymes with rather similar specificities.  
**References:** [625, 2471, 2472, 3141]

[EC 3.1.6.1 created 1961, modified 2011]

#### EC 3.1.6.2

**Accepted name:** steryl-sulfatase  
**Reaction:**  $3\beta\text{-hydroxyandrost-5-en-17-one 3-sulfate} + \text{H}_2\text{O} = 3\beta\text{-hydroxyandrost-5-en-17-one} + \text{sulfate}$   
**Other name(s):** arylsulfatase; steroid sulfatase; sterol sulfatase; dehydroepiandrosterone sulfate sulfatase; arylsulfatase C; steroid 3-sulfatase; steroid sulfate sulfohydrolase; dehydroepiandrosterone sulfatase; pregnenolone sulfatase; phenolic steroid sulfatase; 3- $\beta$ -hydroxysteroid sulfate sulfatase  
**Systematic name:** steryl-sulfate sulfohydrolase  
**Comments:** Also acts on some related steryl sulfates.  
**References:** [2470, 2471, 2772]

[EC 3.1.6.2 created 1961]

#### EC 3.1.6.3

**Accepted name:** glycosulfatase  
**Reaction:**  $\text{D-glucose 6-sulfate} + \text{H}_2\text{O} = \text{D-glucose} + \text{sulfate}$   
**Other name(s):** glucosulfatase  
**Systematic name:** sugar-sulfate sulfohydrolase  
**Comments:** Also acts on other sulfates of monosaccharides and disaccharides and on adenosine 5'-sulfate.  
**References:** [624, 679, 2471]

[EC 3.1.6.3 created 1961]

#### EC 3.1.6.4

**Accepted name:** *N*-acetylgalactosamine-6-sulfatase  
**Reaction:** Hydrolysis of the 6-sulfate groups of the *N*-acetyl-D-galactosamine 6-sulfate units of chondroitin sulfate and of the D-galactose 6-sulfate units of keratan sulfate  
**Other name(s):** chondroitin sulfatase; chondroitinase; galactose-6-sulfate sulfatase; acetylgalactosamine 6-sulfatase; *N*-acetylgalactosamine-6-sulfate sulfatase; *N*-acetylgalactosamine 6-sulfatase



**Systematic name:** *N*-acetyl-D-galactosamine-6-sulfate 6-sulfohydrolase

**References:** [709, 931, 1687, 2726, 3316]

[EC 3.1.6.4 created 1961]

[3.1.6.5 Deleted entry. *sinigrin sulfohydrolase; myrosulfatase*]

[EC 3.1.6.5 created 1961, deleted 1964]

#### EC 3.1.6.6

**Accepted name:** choline-sulfatase

**Reaction:** choline sulfate + H<sub>2</sub>O = choline + sulfate

**Systematic name:** choline-sulfate sulfohydrolase

**References:** [2854]

[EC 3.1.6.6 created 1965]

#### EC 3.1.6.7

**Accepted name:** cellulose-polysulfatase

**Reaction:** Hydrolysis of the 2- and 3-sulfate groups of the polysulfates of cellulose and charonin

**Systematic name:** cellulose-sulfate sulfohydrolase

**References:** [2842]

[EC 3.1.6.7 created 1965]

#### EC 3.1.6.8

**Accepted name:** cerebroside-sulfatase

**Reaction:** a cerebroside 3-sulfate + H<sub>2</sub>O = a cerebroside + sulfate

**Other name(s):** arylsulfatase A; cerebroside sulfate sulfatase

**Systematic name:** cerebroside-3-sulfate 3-sulfohydrolase

**Comments:** Hydrolyses galactose-3-sulfate residues in a number of lipids. Also hydrolyses ascorbate 2-sulfate and many phenol sulfates.

**References:** [1877, 2472]

[EC 3.1.6.8 created 1972]

#### EC 3.1.6.9

**Accepted name:** chondro-4-sulfatase

**Reaction:** 4-deoxy-β-D-gluc-4-enuronosyl-(1→3)-*N*-acetyl-D-galactosamine 4-sulfate + H<sub>2</sub>O = 4-deoxy-β-D-gluc-4-enuronosyl-(1→3)-*N*-acetyl-D-galactosamine + sulfate

**Other name(s):** chondroitin-4-sulfatase; 4-deoxy-β-D-gluc-4-enuronosyl-(1,3)-*N*-acetyl-D-galactosamine-4-sulfate 4-sulfohydrolase

**Systematic name:** 4-deoxy-β-D-gluc-4-enuronosyl-(1→3)-*N*-acetyl-D-galactosamine-4-sulfate 4-sulfohydrolase

**Comments:** Also acts on the saturated analogue but not on higher oligosaccharides, nor any 6-sulfates.

**References:** [1104, 2472, 3244]

[EC 3.1.6.9 created 1972]

#### EC 3.1.6.10

**Accepted name:** chondro-6-sulfatase

**Reaction:** 4-deoxy-β-D-gluc-4-enuronosyl-(1→3)-*N*-acetyl-D-galactosamine 6-sulfate + H<sub>2</sub>O = 4-deoxy-β-D-gluc-4-enuronosyl-(1→3)-*N*-acetyl-D-galactosamine + sulfate

**Other name(s):** 4-deoxy-β-D-gluc-4-enuronosyl-(1,3)-*N*-acetyl-D-galactosamine-6-sulfate 6-sulfohydrolase

**Systematic name:** 4-deoxy- $\beta$ -D-gluc-4-enuronosyl-(1 $\rightarrow$ 3)-*N*-acetyl-D-galactosamine-6-sulfate 6-sulfohydrolase  
**Comments:** Also acts on the saturated analogue and *N*-acetyl-D-galactosamine 4,6-disulfate, but not higher oligosaccharides, nor any 4-sulfate  
**References:** [3244]

[EC 3.1.6.10 created 1972]

#### EC 3.1.6.11

**Accepted name:** disulfoglucosamine-6-sulfatase  
**Reaction:** 2-*N*,6-*O*-disulfo-D-glucosamine + H<sub>2</sub>O = 2-*N*-sulfo-D-glucosamine + sulfate  
**Other name(s):** *N*-sulfoglucosamine-6-sulfatase; 6,*N*-disulfoglucosamine 6-*O*-sulfohydrolase; *N*,6-*O*-disulfo-D-glucosamine 6-sulfohydrolase  
**Systematic name:** 2-*N*,6-*O*-disulfo-D-glucosamine 6-sulfohydrolase  
**Comments:** May be identical with EC 3.1.6.14 *N*-acetylglucosamine-6-sulfatase.  
**References:** [609]

[EC 3.1.6.11 created 1972, modified 1989]

#### EC 3.1.6.12

**Accepted name:** *N*-acetylgalactosamine-4-sulfatase  
**Reaction:** Hydrolysis of the 4-sulfate groups of the *N*-acetyl-D-galactosamine 4-sulfate units of chondroitin sulfate and dermatan sulfate  
**Other name(s):** chondroitinsulfatase; chondroitinase; arylsulfatase B; acetylgalactosamine 4-sulfatase; *N*-acetylgalactosamine 4-sulfate sulfohydrolase  
**Systematic name:** *N*-acetyl-D-galactosamine-4-sulfate 4-sulfohydrolase  
**Comments:** Acts also on *N*-acetylglucosamine 4-sulfate.  
**References:** [739, 959, 2979]

[EC 3.1.6.12 created 1984]

#### EC 3.1.6.13

**Accepted name:** iduronate-2-sulfatase  
**Reaction:** Hydrolysis of the 2-sulfate groups of the L-iduronate 2-sulfate units of dermatan sulfate, heparan sulfate and heparin  
**Other name(s):** chondroitinsulfatase; idurono-2-sulfatase; iduronide-2-sulfate sulfatase; L-iduronosulfatase; L-idurono sulfate sulfatase; iduronate sulfatase; sulfo-L-iduronate sulfatase; L-iduronate 2-sulfate sulfatase; sulfoiduronate sulfohydrolase; 2-sulfo-L-iduronate 2-sulfatase; iduronate-2-sulfate sulfatase; iduronate sulfate sulfatase  
**Systematic name:** L-iduronate-2-sulfate 2-sulfohydrolase  
**References:** [71, 113, 616, 3315]

[EC 3.1.6.13 created 1984]

#### EC 3.1.6.14

**Accepted name:** *N*-acetylglucosamine-6-sulfatase  
**Reaction:** Hydrolysis of the 6-sulfate groups of the *N*-acetyl-D-glucosamine 6-sulfate units of heparan sulfate and keratan sulfate  
**Other name(s):** chondroitinsulfatase; *O*,*N*-disulfate *O*-sulfohydrolase; acetylglucosamine 6-sulfatase; *N*-acetylglucosamine 6-sulfate sulfatase; acetylglucosamine 6-sulfatase; 2-acetamido-2-deoxy-D-glucose 6-sulfate sulfatase  
**Systematic name:** *N*-acetyl-D-glucosamine-6-sulfate 6-sulfohydrolase  
**Comments:** May be identical with EC 3.1.6.11 disulfoglucosamine-6-sulfatase.  
**References:** [169, 1534, 3156]

[EC 3.1.6.14 created 1984]

#### EC 3.1.6.15

**Accepted name:** *N*-sulfolglucosamine-3-sulfatase  
**Reaction:** Hydrolysis of the 3-sulfate groups of the *N*-sulfo-D-glucosamine 3-*O*-sulfate units of heparin  
**Other name(s):** chondroitinsulfatase  
**Systematic name:** *N*-sulfo-3-sulfolglucosamine 3-sulfohydrolase  
**Comments:** The enzyme from *Flavobacterium heparinum* also hydrolyses *N*-acetyl-D-glucosamine 3-*O*-sulfate; the mammalian enzyme acts only on the disulfated residue.  
**References:** [315, 1616]

[EC 3.1.6.15 created 1984, modified 1989]

#### EC 3.1.6.16

**Accepted name:** monomethyl-sulfatase  
**Reaction:** monomethyl sulfate + H<sub>2</sub>O = methanol + sulfate  
**Systematic name:** monomethyl-sulfate sulfohydrolase  
**Comments:** Highly specific; does not act on monoethyl sulfate, monoisopropyl sulfate or monododecyl sulfate.  
**References:** [903]

[EC 3.1.6.16 created 1989]

#### EC 3.1.6.17

**Accepted name:** D-lactate-2-sulfatase  
**Reaction:** (*R*)-2-*O*-sulfolactate + H<sub>2</sub>O = (*R*)-lactate + sulfate  
**Other name(s):** (*S*)-2-*O*-sulfolactate 2-sulfohydrolase (incorrect stereochemistry)  
**Systematic name:** (*R*)-2-*O*-sulfolactate 2-sulfohydrolase  
**Comments:** Highly specific.  
**References:** [504]

[EC 3.1.6.17 created 1989]

#### EC 3.1.6.18

**Accepted name:** glucuronate-2-sulfatase  
**Reaction:** Hydrolysis of the 2-sulfate groups of the 2-*O*-sulfo-D-glucuronate residues of chondroitin sulfate, heparin and heparitin sulfate  
**Other name(s):** glucurono-2-sulfatase  
**Systematic name:** polysaccharide-2-*O*-sulfo-D-glucuronate 2-sulfohydrolase  
**Comments:** Does not act on iduronate 2-sulfate residues (*cf.* EC 3.1.6.13 iduronate-2-sulfatase)  
**References:** [2617]

[EC 3.1.6.18 created 1989]

#### EC 3.1.6.19

**Accepted name:** (*R*)-specific secondary-alkylsulfatase  
**Reaction:** an (*R*)-secondary-alkyl sulfate + H<sub>2</sub>O = an (*S*)-secondary-alcohol + sulfate  
**Other name(s):** S3 secondary alkylsulphohydrolase; Pisa1; secondary alkylsulphohydrolase; (*R*)-specific *sec*-alkylsulfatase; *sec*-alkylsulfatase  
**Systematic name:** (*R*)-secondary-alkyl sulfate sulfohydrolase [(*S*)-secondary-alcohol forming]

**Comments:** The enzyme from *Rhodococcus ruber* is involved in the biodegradation of alkyl sulfate esters used as detergents and released into the environment. The preferred substrates are linear secondary-alkyl sulfate esters, particularly octan-2-yl, octan-3-yl, and octan-4-yl sulfates [2299]. The enzyme from *Pseudomonas* sp. DSM6611 utilizes a range of secondary-alkyl sulfate esters bearing aromatic, olefinic and acetylenic moieties. Perfect enantioselectivities are obtained for substrates bearing groups of different size adjacent to the sulfate moiety [2579]. The enzymic hydrolysis proceeds through inversion of the configuration at the stereogenic carbon atom [2299, 2579]. The enzyme contains a Zn<sup>2+</sup> ion [1481].

**References:** [2299, 3100, 1481, 2579]

[EC 3.1.6.19 created 2013]

#### EC 3.1.6.20

**Accepted name:** *S*-sulfosulfanyl-L-cysteine sulfohydrolase

**Reaction:** (1) [SoxY protein]-*S*-sulfosulfanyl-L-cysteine + H<sub>2</sub>O = [SoxY protein]-*S*-sulfanyl-L-cysteine + sulfate  
(2) [SoxY protein]-*S*-(2-sulfodisulfanyl)-L-cysteine + H<sub>2</sub>O = [SoxY protein]-*S*-disulfanyl-L-cysteine + sulfate

**Other name(s):** SoxB

**Systematic name:** [SoxY protein]-*S*-sulfosulfanyl-L-cysteine sulfohydrolase

**Comments:** Contains Mn<sup>2+</sup>. The enzyme is part of the Sox enzyme system, which participates in a bacterial thio-sulfate oxidation pathway that produces sulfate. It catalyses two reactions in the pathway. In both cases the enzyme hydrolyses a sulfonate moiety that is bound (either directly or via a sulfane) to a cysteine residue of a SoxY protein, releasing sulfate. The enzyme from *Paracoccus pantotrophus* contains a pyroglutamate (cycloglutamate) at its N-terminus.

**References:** [2343, 822, 2344, 707, 1118, 961]

[EC 3.1.6.20 created 2018]

### EC 3.1.7 Diphosphoric-monoester hydrolases

#### EC 3.1.7.1

**Accepted name:** prenyl-diphosphatase

**Reaction:** prenyl diphosphate + H<sub>2</sub>O = prenol + diphosphate

**Other name(s):** prenyl-pyrophosphatase; prenol pyrophosphatase; prenylphosphatase

**Systematic name:** prenyl-diphosphate diphosphohydrolase

**Comments:** Farnesyl diphosphate is the best substrate tested to date.

**References:** [2969]

[EC 3.1.7.1 created 1972]

#### EC 3.1.7.2

**Accepted name:** guanosine-3',5'-bis(diphosphate) 3'-diphosphatase

**Reaction:** guanosine 3',5'-bis(diphosphate) + H<sub>2</sub>O = GDP + diphosphate

**Other name(s):** guanosine-3',5'-bis(diphosphate) 3'-pyrophosphatase; PpGpp-3'-pyrophosphohydrolase; PpGpp phosphohydrolase

**Systematic name:** guanosine-3',5'-bis(diphosphate) 3'-diphosphohydrolase

**References:** [1102, 2422]

[EC 3.1.7.2 created 1980]

#### EC 3.1.7.3

**Accepted name:** monoterpenyl-diphosphatase

**Reaction:** a monoterpenyl diphosphate + H<sub>2</sub>O = a monoterpenol + diphosphate  
**Other name(s):** bornyl pyrophosphate hydrolase; monoterpenyl-pyrophosphatase  
**Systematic name:** monoterpenyl-diphosphate diphosphohydrolase  
**Comments:** A group of enzymes with varying specificity for the monoterpenol moiety. One has the highest activity on sterically hindered compounds such as (+)-bornyl diphosphate; another has highest activity on the diphosphates of primary allylic alcohols such as geraniol.  
**References:** [510]

[EC 3.1.7.3 created 1984]

[3.1.7.4 Deleted entry. Now recognized as two enzymes EC 4.2.1.133, copal-8-ol diphosphate synthase and EC 4.2.3.141 sclareol synthase]

[EC 3.1.7.4 created 2008, deleted 2013]

#### EC 3.1.7.5

**Accepted name:** geranylgeranyl diphosphate diphosphatase  
**Reaction:** geranylgeranyl diphosphate + H<sub>2</sub>O = geranylgeraniol + diphosphate  
**Other name(s):** geranylgeranyl diphosphate phosphatase  
**Systematic name:** geranyl-diphosphate diphosphohydrolase  
**Comments:** Involved in the biosynthesis of plaunotol. There are two isoenzymes with different ion requirements. Neither require Mg<sup>2+</sup> but in addition PII is inhibited by Zn<sup>2+</sup>, Mn<sup>2+</sup> and Co<sup>2+</sup>. It is not known which isoenzyme is involved in plaunotol biosynthesis.  
**References:** [2119]

[EC 3.1.7.5 created 2009]

#### EC 3.1.7.6

**Accepted name:** farnesyl diphosphatase  
**Reaction:** (2*E*,6*E*)-farnesyl diphosphate + H<sub>2</sub>O = (2*E*,6*E*)-farnesol + diphosphate  
**Other name(s):** FPP phosphatase  
**Systematic name:** (2*E*,6*E*)-farnesyl-diphosphate diphosphohydrolase  
**Comments:** The enzyme is involved in the biosynthesis of acyclic sesquiterpenoids [2717].  
**References:** [2717, 2969]

[EC 3.1.7.6 created 2010]

[3.1.7.7 Transferred entry. (-)-drimenol synthase. Now EC 4.2.3.194, (-)-drimenol synthase]

[EC 3.1.7.7 created 2011, deleted 2017]

#### EC 3.1.7.8

**Accepted name:** tuberculosinyl synthase  
**Reaction:** tuberculosinyl diphosphate + H<sub>2</sub>O = tuberculosinol + diphosphate  
**Other name(s):** Rv3378c  
**Systematic name:** tuberculosinyl diphosphate diphosphohydrolase (tuberculosinol forming)  
**Comments:** Only found in species of *Mycobacterium* that cause tuberculosis. In addition, it also gives isotuberculosinol in 1:1 mixture, cf. EC 3.1.7.9, isotuberculosinol synthase.  
**References:** [2049, 1184]

[EC 3.1.7.8 created 2011]

#### EC 3.1.7.9

**Accepted name:** isotuberculosinol synthase

**Reaction:** tuberculosinyl diphosphate + H<sub>2</sub>O = (13S)-isotuberculosinol + diphosphate  
**Other name(s):** Rv3378c  
**Systematic name:** tuberculosinyl diphosphate diphosphohydrolase (isotuberculosinol forming)  
**Comments:** Only found in species of *Mycobacterium* that cause tuberculosis. In addition, it also gives tuberculosinol in 1:1 mixture, cf. EC 3.1.7.8, tuberculosinol synthase. The isotuberculosinol form was a 3:1 mixture of the 13S and 13R forms, respectively.  
**References:** [2049, 1184]

[EC 3.1.7.9 created 2011]

#### EC 3.1.7.10

**Accepted name:** (13E)-labda-7,13-dien-15-ol synthase  
**Reaction:** geranylgeranyl diphosphate + H<sub>2</sub>O = (13E)-labda-7,13-dien-15-ol + diphosphate  
**Other name(s):** labda-7,13E-dien-15-ol synthase  
**Systematic name:** geranylgeranyl-diphosphate diphosphohydrolase [(13E)-labda-7,13-dien-15-ol-forming]  
**Comments:** The enzyme from the lycophyte *Selaginella moellendorffii* is bifunctional, initially forming (13E)-labda-7,13-dien-15-yl diphosphate, which is hydrolysed to the alcohol.  
**References:** [1770]

[EC 3.1.7.10 created 2012]

#### EC 3.1.7.11

**Accepted name:** geranyl diphosphate diphosphatase  
**Reaction:** geranyl diphosphate + H<sub>2</sub>O = geraniol + diphosphate  
**Other name(s):** geraniol synthase; geranyl pyrophosphate pyrophosphatase; GES; *CtGES*  
**Systematic name:** geranyl-diphosphate diphosphohydrolase  
**Comments:** Isolated from *Ocimum basilicum* (basil) and *Cinnamomum tenuipile* (camphor tree). Requires Mg<sup>2+</sup> or Mn<sup>2+</sup>. Geraniol is labelled when formed in the presence of [<sup>18</sup>O]H<sub>2</sub>O. Thus mechanism involves a geranyl cation [1225]. Neryl diphosphate is hydrolysed more slowly. May be the same as EC 3.1.7.3 monoterpenyl-diphosphatase.  
**References:** [1225, 3260]

[EC 3.1.7.11 created 2012]

#### EC 3.1.7.12

**Accepted name:** (+)-kolavelool synthase  
**Reaction:** (+)-kolavenyl diphosphate + H<sub>2</sub>O = (+)-kolavelool + diphosphate  
**Other name(s):** Haur\_2146  
**Systematic name:** kolavenyl-diphosphate diphosphohydrolase  
**Comments:** Isolated from the bacterium *Herpetosiphon aurantiacus*.  
**References:** [2050]

[EC 3.1.7.12 created 2017]

### EC 3.1.8 Phosphoric-triester hydrolases

#### EC 3.1.8.1

**Accepted name:** aryldialkylphosphatase  
**Reaction:** an aryl dialkyl phosphate + H<sub>2</sub>O = dialkyl phosphate + an aryl alcohol

**Other name(s):** organophosphate hydrolase; paraoxonase; A-esterase; aryltriphosphatase; organophosphate esterase; esterase B1; esterase E4; paraoxon esterase; pirimiphos-methyloxon esterase; OPA anhydrase; organophosphorus hydrolase; phosphotriesterase; paraoxon hydrolase; OPH; organophosphorus acid anhydrase

**Systematic name:** aryltriphosphate dialkylphosphohydrolase

**Comments:** Acts on organophosphorus compounds (such as paraoxon) including esters of phosphonic and phosphinic acids. Inhibited by chelating agents; requires divalent cations for activity. Previously regarded as identical with EC 3.1.1.2 arylesterase.

**References:** [30, 267, 1768, 1777, 1]

[EC 3.1.8.1 created 1989]

#### EC 3.1.8.2

**Accepted name:** diisopropyl-fluorophosphatase

**Reaction:** diisopropyl fluorophosphate + H<sub>2</sub>O = diisopropyl phosphate + fluoride

**Other name(s):** DFPase; tabunase; somanase; organophosphorus acid anhydrolase; organophosphate acid anhydrase; OPA anhydrase; diisopropylphosphofluoridase; dialkylfluorophosphatase; diisopropyl phosphorofluoridate hydrolase; isopropylphosphorofluoridase; diisopropylfluorophosphonate dehalogenase

**Systematic name:** diisopropyl-fluorophosphate fluorohydrolase

**Comments:** Acts on phosphorus anhydride bonds (such as phosphorus-halide and phosphorus-cyanide) in organophosphorus compounds (including 'nerve gases'). Inhibited by chelating agents; requires divalent cations. Related to EC 3.1.8.1 arylalkylphosphatase.

**References:** [99, 100, 101, 463, 1991, 1]

[EC 3.1.8.2 created 1961 as EC 3.8.2.1, transferred 1992 to EC 3.1.8.2]

### EC 3.1.11 Exodeoxyribonucleases producing 5'-phosphomonoesters

#### EC 3.1.11.1

**Accepted name:** exodeoxyribonuclease I

**Reaction:** Exonucleolytic cleavage in the 3'- to 5'-direction to yield nucleoside 5'-phosphates

**Other name(s):** *Escherichia coli* exonuclease I; *E. coli* exonuclease I; exonuclease I

**Comments:** Preference for single-stranded DNA. The *Escherichia coli* enzyme hydrolyses glucosylated DNA.

**References:** [236, 1415, 1640]

[EC 3.1.11.1 created 1972 as EC 3.1.4.25, transferred 1978 to EC 3.1.11.1]

#### EC 3.1.11.2

**Accepted name:** exodeoxyribonuclease III

**Reaction:** Exonucleolytic cleavage in the 3'- to 5'-direction to yield nucleoside 5'-phosphates

**Other name(s):** *Escherichia coli* exonuclease III; *E. coli* exonuclease III; endoribonuclease III

**Comments:** Preference for double-stranded DNA. Has endonucleolytic activity near apurinic sites on DNA.

**References:** [1700, 2419, 2420]

[EC 3.1.11.2 created 1972 as EC 3.1.4.27, transferred 1978 to EC 3.1.11.2]

#### EC 3.1.11.3

**Accepted name:** exodeoxyribonuclease (lambda-induced)

**Reaction:** Exonucleolytic cleavage in the 5'- to 3'-direction to yield nucleoside 5'-phosphates

**Other name(s):** lambda exonuclease; phage lambda-induced exonuclease; *Escherichia coli* exonuclease IV; *E. coli* exonuclease IV; exodeoxyribonuclease IV; exonuclease IV

**Comments:** Preference for double-stranded DNA. Does not attack single-strand breaks.

**References:** [1699, 1713]

[EC 3.1.11.3 created 1972 as EC 3.1.4.28, transferred 1978 to EC 3.1.11.3]

#### EC 3.1.11.4

**Accepted name:** exodeoxyribonuclease (phage SP<sub>3</sub>-induced)  
**Reaction:** Exonucleolytic cleavage in the 5'- to 3'-direction to yield nucleoside 5'-phosphates  
**Other name(s):** phage SP<sub>3</sub> DNase; DNA 5'-dinucleotidohydrolase; deoxyribonuclease 5'-dinucleotidase; deoxyribonucleic 5'-dinucleotidohydrolase; bacteriophage SP<sub>3</sub> deoxyribonuclease; deoxyribonuclease 5'-dinucleotidase  
**Comments:** Preference for single-stranded DNA.  
**References:** [2964]

[EC 3.1.11.4 created 1972 as EC 3.1.4.31, transferred 1978 to EC 3.1.11.4]

#### EC 3.1.11.5

**Accepted name:** exodeoxyribonuclease V  
**Reaction:** Exonucleolytic cleavage (in the presence of ATP) in either 5'- to 3'- or 3'- to 5'-direction to yield 5'-phosphooligonucleotides  
**Other name(s):** *Escherichia coli* exonuclease V; *E. coli* exonuclease V; gene recBC endoenzyme; RecBC deoxyribonuclease; gene recBC DNase; exonuclease V; gene recBCD enzymes  
**Comments:** Preference for double-stranded DNA. Possesses DNA-dependent ATPase activity. Acts endonucleolytically on single-stranded circular DNA.  
**References:** [685, 951, 2177, 3216]

[EC 3.1.11.5 created 1978]

#### EC 3.1.11.6

**Accepted name:** exodeoxyribonuclease VII  
**Reaction:** Exonucleolytic cleavage in either 5'- to 3'- or 3'- to 5'-direction to yield nucleoside 5'-phosphates  
**Other name(s):** *Escherichia coli* exonuclease VII; *E. coli* exonuclease VII; endodeoxyribonuclease VII; exonuclease VII  
**Comments:** Preference for single-stranded DNA.  
**References:** [402, 401]

[EC 3.1.11.6 created 1978]

#### EC 3.1.11.7

**Accepted name:** adenosine-5'-diphospho-5'-[DNA] diphosphatase  
**Reaction:** (1) adenosine-5'-diphospho-5'-[DNA] + H<sub>2</sub>O = AMP + phospho-5'-[DNA]  
(2) adenosine-5'-diphospho-5'-(ribonucleotide)-[DNA] + H<sub>2</sub>O = AMP + 5'-phospho-(ribonucleotide)-[DNA]  
**Other name(s):** aprataxin; 5'-App5'-DNA adenylate hydrolase; APTX (gene name); HNT3 (gene name)  
**Systematic name:** adenosine-5'-diphospho-5'-[DNA] hydrolase (adenosine 5'-phosphate-forming)  
**Comments:** Aprataxin is a DNA-binding protein involved in different types of DNA break repair. The enzyme acts (among other activities) on abortive DNA ligation intermediates that contain an adenylate covalently linked to the 5'-phosphate DNA terminus. It also acts when the adenylate is covalently linked to the 5'-phosphate of a ribonucleotide linked to a DNA strand, which is the result of abortive ligase activity on products of EC 3.1.26.4, ribonuclease H, an enzyme that cleaves RNA-DNA hybrids on the 5' side of the ribonucleotide found in the 5'-RNA-DNA-3' junction. Aprataxin binds the adenylate group to a histidine residue within the active site, followed by its hydrolysis from the nucleic acid and eventual release, leaving a 5'-phosphate terminus that can be efficiently rejoined. The enzyme also possesses the activities of EC 3.1.11.8, guanosine-5'-diphospho-5'-[DNA] diphosphatase, and EC 3.1.12.2, DNA-3'-diphospho-5'-guanosine diphosphatase.  
**References:** [22, 2996]



[EC 3.1.11.7 created 2017]

#### EC 3.1.11.8

- Accepted name:** guanosine-5'-diphospho-5'-[DNA] diphosphatase  
**Reaction:** guanosine-5'-diphospho-5'-[DNA] + H<sub>2</sub>O = phospho-5'-[DNA] + GMP  
**Other name(s):** aprataxin; pp5'G5'DNA diphosphatase; pp5'G5'-DNA guanylate hydrolase; APTX (gene name); HNT3 (gene name)  
**Systematic name:** guanosine-5'-diphospho-5'-[DNA] hydrolase (guanosine 5'-phosphate-forming)  
**Comments:** Aprataxin is a DNA-binding protein that catalyses (among other activities) the 5' decapping of Gpp-DNA (formed by homologs of RtcB3 from the bacterium *Myxococcus xanthus*). The enzyme binds the guanylate group to a histidine residue at its active site, forming a covalent enzyme-nucleotide phosphate intermediate, followed by the hydrolysis of the guanylate from the nucleic acid and eventual release. The enzyme forms a 5'-phospho terminus that can be efficiently joined by "classical" ligases. The enzyme also possesses the activity of EC 3.1.11.7, adenosine-5'-diphospho-5'-[DNA] diphosphatase and EC 3.1.12.2, DNA-3'-diphospho-5'-guanosine diphosphatase.  
**References:** [1849]

[EC 3.1.11.8 created 2017]

### EC 3.1.12 Exodeoxyribonucleases producing 3'-phosphomonoesters

#### EC 3.1.12.1

- Accepted name:** 5' to 3' exodeoxyribonuclease (nucleoside 3'-phosphate-forming)  
**Reaction:** exonucleolytic cleavage in the 5'- to 3'-direction to yield nucleoside 3'-phosphates  
**Other name(s):** Cas4; 5' to 3' single stranded DNA exonuclease  
**Comments:** Preference for single-stranded DNA. The enzyme from the archaeon *Sulfolobus solfataricus* contains a [4Fe-4S] cluster and requires a divalent metal cation, such as Mg<sup>2+</sup> or Mn<sup>2+</sup>, for activity.  
**References:** [3326, 1642]

[EC 3.1.12.1 created 2014]

#### EC 3.1.12.2

- Accepted name:** DNA-3'-diphospho-5'-guanosine diphosphatase  
**Reaction:** [DNA]-3'-diphospho-5'-guanosine + H<sub>2</sub>O = [DNA]-3'-phosphate + GMP  
**Other name(s):** aprataxin; DNA-3'pp5'G guanylate hydrolase; APTX (gene name); HNT3 (gene name)  
**Systematic name:** [DNA]-3'-diphospho-5'-guanosine hydrolase (guanosine 5'-phosphate-forming)  
**Comments:** Aprataxin is a DNA-binding protein that catalyses (among other activities) the 3' decapping of DNA-ppG (formed by EC 6.5.1.8, 3'-phosphate/5'-hydroxy nucleic acid ligase) [538]. The enzyme binds the guanylate group to a histidine residue at its active site, forming a covalent enzyme-nucleotide phosphate intermediate, followed by the hydrolysis of the guanylate from the nucleic acid and its eventual release. The enzyme also possesses the activity of EC 3.1.11.7, adenosine-5'-diphospho-5'-[DNA] diphosphatase, and EC 3.1.11.8, guanosine-5'-diphospho-5'-[DNA] diphosphatase.  
**References:** [538, 406]

[EC 3.1.12.2 created 2017]

### EC 3.1.13 Exoribonucleases producing 5'-phosphomonoesters

#### EC 3.1.13.1

- Accepted name:** exoribonuclease II  
**Reaction:** Exonucleolytic cleavage in the 3'- to 5'-direction to yield nucleoside 5'-phosphates

**Other name(s):** ribonuclease II; ribonuclease Q; BN ribonuclease; *Escherichia coli* exo-RNase II; RNase II; exoribonuclease (misleading); 5'-exoribonuclease (misleading)  
**Comments:** Preference for single-stranded RNA. The enzyme processes 3'-terminal extra-nucleotides of monomeric tRNA precursors, following the action of EC 3.1.26.5 ribonuclease P.  
**References:** [2117, 2573, 2635, 2742]

[EC 3.1.13.1 created 1972 as EC 3.1.4.20, transferred 1978 to EC 3.1.13.1]

#### EC 3.1.13.2

**Accepted name:** exoribonuclease H  
**Reaction:** 3'-end directed exonucleolytic cleavage of viral RNA-DNA hybrid  
**Comments:** This is a secondary reaction to the RNA 5'-end directed cleavage 13-19 nucleotides from the RNA end performed by EC 3.1.26.13 (retroviral ribonuclease H).  
**References:** [2555]

[EC 3.1.13.2 created 1978, modified 2010]

#### EC 3.1.13.3

**Accepted name:** oligonucleotidase  
**Reaction:** Exonucleolytic cleavage of oligonucleotides to yield nucleoside 5'-phosphates  
**Other name(s):** oligoribonuclease  
**Comments:** Also hydrolyses NAD<sup>+</sup> to NMN and AMP.  
**References:** [866]

[EC 3.1.13.3 created 1972 as EC 3.1.4.19, transferred 1978 to EC 3.1.13.3]

#### EC 3.1.13.4

**Accepted name:** poly(A)-specific ribonuclease  
**Reaction:** Exonucleolytic cleavage of poly(A) to 5'-AMP  
**Other name(s):** 3'-exoribonuclease; 2',3'-exoribonuclease  
**Comments:** Cleaves poly(A) in either the single- or double-stranded form.  
**References:** [2586]

[EC 3.1.13.4 created 1984]

#### EC 3.1.13.5

**Accepted name:** ribonuclease D  
**Reaction:** Exonucleolytic cleavage that removes extra residues from the 3'-terminus of tRNA to produce 5'-mononucleotides  
**Other name(s):** RNase D  
**Comments:** Requires divalent cations for activity (Mg<sup>2+</sup>, Mn<sup>2+</sup> or Co<sup>2+</sup>). Alteration of the 3'-terminal base has no effect on the rate of hydrolysis whereas modification of the 3'-terminal sugar has a major effect. tRNA terminating with a 3'-phosphate is completely inactive [512]. This enzyme can convert a tRNA precursor into a mature tRNA [513].  
**References:** [906, 513, 512, 3327]

[EC 3.1.13.5 created 2006]

### EC 3.1.14 Exoribonucleases producing 3'-phosphomonoesters

#### EC 3.1.14.1

**Accepted name:** yeast ribonuclease  
**Reaction:** Exonucleolytic cleavage to nucleoside 3'-phosphates  
**Comments:** Similar enzyme: RNase U<sub>4</sub>.  
**References:** [2212]

[EC 3.1.14.1 created 1978]

### EC 3.1.15 Exonucleases that are active with either ribo- or deoxyribonucleic acids and produce 5'-phosphomonoesters

#### EC 3.1.15.1

**Accepted name:** venom exonuclease  
**Reaction:** Exonucleolytic cleavage in the 3'- to 5'- direction to yield nucleoside 5'-phosphates  
**Other name(s):** venom phosphodiesterase  
**Comments:** Preference for single-stranded substrate.  
**References:** [1607]

[EC 3.1.15.1 created 1978]

### EC 3.1.16 Exonucleases that are active with either ribo- or deoxyribonucleic acids and produce 3'-phosphomonoesters

#### EC 3.1.16.1

**Accepted name:** spleen exonuclease  
**Reaction:** Exonucleolytic cleavage in the 5'- to 3'-direction to yield nucleoside 3'-phosphates  
**Other name(s):** 3'-exonuclease; spleen phosphodiesterase; 3'-nucleotide phosphodiesterase; phosphodiesterase II  
**Comments:** Preference for single-stranded substrate.  
**References:** [199]

[EC 3.1.16.1 created 1972 as EC 3.1.4.18, transferred 1978 to EC 3.1.16.1]

### EC 3.1.21 Endodeoxyribonucleases producing 5'-phosphomonoesters

#### EC 3.1.21.1

**Accepted name:** deoxyribonuclease I  
**Reaction:** Endonucleolytic cleavage to 5'-phosphodinucleotide and 5'-phosphooligonucleotide end-products  
**Other name(s):** pancreatic DNase; DNase; thymonuclease, dornase; dornava; dornavac; pancreatic deoxyribonuclease; pancreatic dornase; deoxyribonuclease (pancreatic); pancreatic DNase; DNAase; deoxyribonucleic phosphatase; DNase I; alkaline deoxyribonuclease; alkaline DNase; endodeoxyribonuclease I; DNA depolymerase; *Escherichia coli* endonuclease I; deoxyribonuclease A; DNA endonuclease; DNA nuclease  
**Comments:** Preference for double-stranded DNA.  
**References:** [557, 1555, 1608]

[EC 3.1.21.1 created 1961 as EC 3.1.4.5, transferred 1978 to EC 3.1.21.1, modified 1981]

#### EC 3.1.21.2

**Accepted name:** deoxyribonuclease IV  
**Reaction:** Endonucleolytic cleavage of ssDNA at apurinic/apyrimidinic sites to 5'-phosphooligonucleotide end-products

**Other name(s):** deoxyribonuclease IV (phage-T<sub>4</sub>-induced) (misleading); endodeoxyribonuclease IV (phage T<sub>4</sub>-induced) (misleading); *E. coli* endonuclease IV; endodeoxyribonuclease (misleading); redoxendonuclease; deoxriboendonuclease (misleading); endonuclease II; endonuclease IV; DNA-adenine-transferase; *nfo* (gene name)

**Comments:** The enzyme is an apurinic/apyrimidinic (AP) site endonuclease that primes DNA repair synthesis at AP sites. It specifically cleaves the DNA backbone at AP sites and also removes 3' DNA-blocking groups such as 3' phosphates, 3' phosphoglycolates, and 3'  $\alpha,\beta$ -unsaturated aldehydes that arise from oxidative base damage and the activity of combined glycosylase/lyase enzymes. It is also the only known repair enzyme that is able to cleave the DNA backbone 5' of the oxidative lesion  $\alpha$ -deoxyadenosine. The enzyme has a strong preference for single-stranded DNA.

**References:** [819, 820, 1016, 520, 1221, 1183]

[EC 3.1.21.2 created 1972 as EC 3.1.4.30, transferred 1978 to EC 3.1.21.2, modified 2014]

### EC 3.1.21.3

**Accepted name:** type I site-specific deoxyribonuclease

**Reaction:** Endonucleolytic cleavage of DNA to give random double-stranded fragments with terminal 5'-phosphates; ATP is simultaneously hydrolysed

**Other name(s):** type I restriction enzyme; deoxyribonuclease (ATP- and *S*-adenosyl-L-methionine-dependent); restriction-modification system; deoxyribonuclease (adenosine triphosphate-hydrolyzing); adenosine triphosphate-dependent deoxyribonuclease; ATP-dependent DNase; type 1 site-specific deoxyribonuclease

**Comments:** This is a large group of enzymes which, together with those now listed as EC 3.1.21.4 (type II site-specific deoxyribonuclease) and EC 3.1.21.5 (type III site-specific deoxyribonuclease), were previously listed separately in sub-subclasses EC 3.1.23 and EC 3.1.24. They have an absolute requirement for ATP (or dATP) and *S*-adenosyl-L-methionine. They recognize specific short DNA sequences and cleave at sites remote from the recognition sequence. They are multifunctional proteins that also catalyse the reactions of EC 2.1.1.72 [site-specific DNA-methyltransferase (adenine-specific)] and EC 2.1.1.37

**References:** [2437]

[EC 3.1.21.3 created 1984 from EC 3.1.23 and EC 3.1.24]

### EC 3.1.21.4

**Accepted name:** type II site-specific deoxyribonuclease

**Reaction:** Endonucleolytic cleavage of DNA to give specific double-stranded fragments with terminal 5'-phosphates

**Other name(s):** type II restriction enzyme

**Comments:** This is a large group of enzymes which, together with those now listed as EC 3.1.21.3 (type 1 site-specific deoxyribonuclease) and EC 3.1.21.5.

**References:** [2437]

[EC 3.1.21.4 created 1984 from EC 3.1.23 and EC 3.1.24]

### EC 3.1.21.5

**Accepted name:** type III site-specific deoxyribonuclease

**Reaction:** Endonucleolytic cleavage of DNA to give specific double-stranded fragments with terminal 5'-phosphates

**Other name(s):** type III restriction enzyme; restriction-modification system

**Comments:** This is a large group of enzymes which, together with those now listed as EC 3.1.21.3 (type I site-specific deoxyribonuclease) and EC 3.1.21.4 (type II site-specific deoxyribonuclease), were previously listed separately in sub-subclasses EC 3.1.23 and EC 3.1.24. They have an absolute requirement for ATP but do not hydrolyse it; *S*-adenosyl-L-methionine stimulates the reaction, but is not absolutely required. They recognize specific, short DNA sequences and cleave a short distance away from the recognition sequence. These enzymes exist as complexes with enzymes of similar specificity listed under EC 2.1.1.72 [site-specific DNA-methyltransferase (adenine-specific)] or EC 2.1.1.73

**References:** [2437]

[EC 3.1.21.5 created 1984 from EC 3.1.23 and EC 3.1.24]

#### EC 3.1.21.6

**Accepted name:** CC-preferring endodeoxyribonuclease  
**Reaction:** endonucleolytic cleavage to give 5'-phosphooligonucleotide end-products, with a preference for cleavage within the sequence CC  
**Other name(s):** *Streptomyces glaucescens* exocyttoplasmic dodeoxyribonuclease  
**Comments:** Prefers CC sites in double-stranded circular and linear DNA. Greater affinity for double-stranded than single-stranded DNA. Produces nicks, generating double-stranded fragments with 5'- and/or 3'-protruding single-stranded tails. Requires magnesium ions for activity. The endonuclease from *Chlorella*-like green algae infected with NYS-1 virus 4[3228] may be the same enzyme.  
**References:** [3228, 62]

[EC 3.1.21.6 created 1999]

#### EC 3.1.21.7

**Accepted name:** deoxyribonuclease V  
**Reaction:** Endonucleolytic cleavage at apurinic or apyrimidinic sites to products with a 5'-phosphate  
**Other name(s):** endodeoxyribonuclease V; DNase V; *Escherichia coli* endodeoxyribonuclease V  
**Comments:** Previously classified erroneously as EC 3.1.22.3.  
**References:** [882]

[EC 3.1.21.7 created 1978 as EC 3.1.22.3, transferred 2001 to EC 3.1.21.7]

#### EC 3.1.21.8

**Accepted name:** T<sub>4</sub> deoxyribonuclease II  
**Reaction:** Endonucleolytic nicking and cleavage of cytosine-containing double-stranded DNA.  
**Other name(s):** T<sub>4</sub> endonuclease II; EndoII (ambiguous); *denA* (gene name)  
**Comments:** Requires Mg<sup>2+</sup>. This phage T<sub>4</sub> enzyme is involved in degradation of host DNA. The enzyme primarily catalyses nicking of the bottom strand of double stranded DNA between the first and second base pair to the right of a top-strand CCGC motif. Double-stranded breaks are produced 5- to 10-fold less frequently [370]. It does not cleave the T4 native DNA, which contains 5-hydroxymethylcytosine instead of cytosine.  
**References:** [371, 369, 370, 53]

[EC 3.1.21.8 created 2014]

#### EC 3.1.21.9

**Accepted name:** T<sub>4</sub> deoxyribonuclease IV  
**Reaction:** Endonucleolytic cleavage of the 5' phosphodiester bond of deoxycytidine in single-stranded DNA.  
**Other name(s):** T<sub>4</sub> endonuclease IV; EndoIV (ambiguous); *denB* (gene name)  
**Comments:** This phage T<sub>4</sub> enzyme is involved in degradation of host DNA. The enzyme does not cleave double-stranded DNA or native T4 DNA, which contains 5-hydroxymethylcytosine instead of cytosine.  
**References:** [2489, 1705, 2488, 200, 1147, 2167]

[EC 3.1.21.9 created 2014]

## EC 3.1.22 Endodeoxyribonucleases producing 3'-phosphomonoesters

### EC 3.1.22.1

**Accepted name:** deoxyribonuclease II  
**Reaction:** Endonucleolytic cleavage to nucleoside 3'-phosphates and 3'-phosphooligonucleotide end-products  
**Other name(s):** DNase II; pancreatic DNase II; deoxyribonuclease 3'-nucleotidohydrolase; DNase II; pancreatic DNase II; acid deoxyribonuclease; acid DNase  
**Comments:** Preference for double-stranded DNA.  
**References:** [201]

[EC 3.1.22.1 created 1961 as EC 3.1.4.6, transferred 1978 to EC 3.1.22.1, modified 1981]

### EC 3.1.22.2

**Accepted name:** *Aspergillus* deoxyribonuclease K<sub>1</sub>  
**Reaction:** Endonucleolytic cleavage to nucleoside 3'-phosphates and 3'-phosphooligonucleotide end-products  
**Other name(s):** *Aspergillus* DNase K<sub>1</sub>  
**Comments:** Preference for single-stranded DNA.  
**References:** [1761, 2644]

[EC 3.1.22.2 created 1978, modified 1981]

[3.1.22.3 Transferred entry. deoxyribonuclease V. Now EC 3.1.21.7, deoxyribonuclease V]

[EC 3.1.22.3 created 1978, deleted 2001]

### EC 3.1.22.4

**Accepted name:** crossover junction endodeoxyribonuclease  
**Reaction:** Endonucleolytic cleavage at a junction such as a reciprocal single-stranded crossover between two homologous DNA duplexes (Holliday junction)  
**Other name(s):** Hje endonuclease; Holliday junction endonuclease CCE1; Holliday junction resolvase; Holliday junction-cleaving endonuclease; Holliday junction-resolving endonuclease; RusA Holliday junction resolvase; RusA endonuclease; RuvC endonuclease; SpCCE<sub>1</sub> Holliday junction resolvase; crossover junction endonuclease; cruciform-cutting endonuclease; endo X3; endonuclease RuvC; endonuclease VII; endonuclease X3; resolving enzyme CCE1  
**Comments:** The enzyme from *Saccharomyces cerevisiae* has no endonuclease or exonuclease activity on single-stranded or double-stranded DNA molecules that do not contain Holliday junctions.  
**References:** [2822, 2630, 2616, 787, 1907]

[EC 3.1.22.4 created 1989, modified 2003]

### EC 3.1.22.5

**Accepted name:** deoxyribonuclease X  
**Reaction:** Endonucleolytic cleavage of supercoiled plasma DNA to linear DNA duplexes  
**Other name(s):** *Escherichia coli* endodeoxyribonuclease; *Escherichia coli* endodeoxyribonuclease X  
**Comments:** Preference for supercoiled DNA; little activity on linear double-stranded DNA. Inhibited by single-stranded DNA, ATP and AMP.  
**References:** [907]

[EC 3.1.22.5 created 1992]

### **EC 3.1.23 Site-specific endodeoxyribonucleases: cleavage is sequence specific (deleted sub-subclass)**

- [3.1.23.1 *Transferred entry. endodeoxyribonuclease AluI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.1 created 1978, deleted 1984]
- [3.1.23.2 *Transferred entry. endodeoxyribonuclease AsuI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.2 created 1978, deleted 1984]
- [3.1.23.3 *Transferred entry. endodeoxyribonuclease AvaI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.3 created 1978, deleted 1984]
- [3.1.23.4 *Transferred entry. endodeoxyribonuclease AvaII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.4 created 1978, deleted 1984]
- [3.1.23.5 *Transferred entry. endodeoxyribonuclease Ball. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.5 created 1978, deleted 1984]
- [3.1.23.6 *Transferred entry. endodeoxyribonuclease BamHI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.6 created 1978, deleted 1984]
- [3.1.23.7 *Transferred entry. endodeoxyribonuclease BbvI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.7 created 1978, deleted 1984]
- [3.1.23.8 *Transferred entry. endodeoxyribonuclease BclI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.8 created 1978, deleted 1984]
- [3.1.23.9 *Transferred entry. endodeoxyribonuclease BglI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.9 created 1978, deleted 1984]
- [3.1.23.10 *Transferred entry. endodeoxyribonuclease BglII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.10 created 1978, deleted 1984]
- [3.1.23.11 *Transferred entry. endodeoxyribonuclease BpuI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.11 created 1978, deleted 1984]
- [3.1.23.12 *Transferred entry. endodeoxyribonuclease DpnI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.12 created 1978, modified 1982, deleted 1984]
- [3.1.23.13 *Transferred entry. endodeoxyribonuclease EcoRI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.13 created 1978, deleted 1984]
- [3.1.23.14 *Transferred entry. endodeoxyribonuclease EcoRII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.14 created 1978, deleted 1984]
- [3.1.23.15 *Transferred entry. endodeoxyribonuclease HaeI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.15 created 1978, deleted 1984]
- [3.1.23.16 *Transferred entry. endodeoxyribonuclease HaeII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.16 created 1978, deleted 1984]
- [3.1.23.17 *Transferred entry. endodeoxyribonuclease HaeIII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*

- [EC 3.1.23.17 created 1978, deleted 1984]
- [3.1.23.18 *Transferred entry. endodeoxyribonuclease HgaI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.18 created 1978, deleted 1984]
- [3.1.23.19 *Transferred entry. endodeoxyribonuclease HhaI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.19 created 1978, deleted 1984]
- [3.1.23.20 *Transferred entry. endodeoxyribonuclease HindII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.20 created 1978, deleted 1984]
- [3.1.23.21 *Transferred entry. endodeoxyribonuclease HindIII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.21 created 1978, deleted 1984]
- [3.1.23.22 *Transferred entry. endodeoxyribonuclease HinfI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.22 created 1978, deleted 1984]
- [3.1.23.23 *Transferred entry. endodeoxyribonuclease HpaI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.23 created 1978, deleted 1984]
- [3.1.23.24 *Transferred entry. endodeoxyribonuclease HpaII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.24 created 1978, deleted 1984]
- [3.1.23.25 *Transferred entry. endodeoxyribonuclease HphI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.25 created 1978, deleted 1984]
- [3.1.23.26 *Transferred entry. endodeoxyribonuclease KpnI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.26 created 1978, deleted 1984]
- [3.1.23.27 *Transferred entry. endodeoxyribonuclease MboI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.27 created 1978, deleted 1984]
- [3.1.23.28 *Transferred entry. endodeoxyribonuclease MboII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.28 created 1978, deleted 1984]
- [3.1.23.29 *Transferred entry. endodeoxyribonuclease MnlI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.29 created 1978, deleted 1984]
- [3.1.23.30 *Transferred entry. endodeoxyribonuclease PfaI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.30 created 1978, modified 1982, deleted 1984]
- [3.1.23.31 *Transferred entry. endodeoxyribonuclease PstI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.31 created 1978, deleted 1984]
- [3.1.23.32 *Transferred entry. endodeoxyribonuclease PvuI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.32 created 1978, modified 1982, deleted 1984]
- [3.1.23.33 *Transferred entry. endodeoxyribonuclease PvuII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.33 created 1978, deleted 1984]
- [3.1.23.34 *Transferred entry. endodeoxyribonuclease SacI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.34 created 1978, deleted 1984]



- [3.1.23.35] *Transferred entry. endodeoxyribonuclease SacII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.35 created 1978, deleted 1984]
- [3.1.23.36] *Transferred entry. endodeoxyribonuclease SacIII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.36 created 1978, deleted 1984]
- [3.1.23.37] *Transferred entry. endodeoxyribonuclease Sall. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.37 created 1978, deleted 1984]
- [3.1.23.38] *Transferred entry. endodeoxyribonuclease SgrI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.38 created 1978, deleted 1984]
- [3.1.23.39] *Transferred entry. endodeoxyribonuclease TaqI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.39 created 1978, deleted 1984]
- [3.1.23.40] *Transferred entry. endodeoxyribonuclease TaqII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.40 created 1978, deleted 1984]
- [3.1.23.41] *Transferred entry. endodeoxyribonuclease XbaI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.41 created 1978, deleted 1984]
- [3.1.23.42] *Transferred entry. endodeoxyribonuclease XhoI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.42 created 1978, deleted 1984]
- [3.1.23.43] *Transferred entry. endodeoxyribonuclease XhoII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.43 created 1978, modified 1982, deleted 1984]
- [3.1.23.44] *Transferred entry. endodeoxyribonuclease XmaI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.44 created 1978, deleted 1984]
- [3.1.23.45] *Transferred entry. endodeoxyribonuclease XniI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.45 created 1978, modified 1982, deleted 1984]
- [3.1.23.46] *Transferred entry. endodeoxyribonuclease AimI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.46 created 1982, deleted 1984]
- [3.1.23.47] *Transferred entry. endodeoxyribonuclease AccI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.47 created 1982, deleted 1984]
- [3.1.23.48] *Transferred entry. endodeoxyribonuclease AccII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.48 created 1982, deleted 1984]
- [3.1.23.49] *Transferred entry. endodeoxyribonuclease AtuAI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.49 created 1982, deleted 1984]
- [3.1.23.50] *Transferred entry. endodeoxyribonuclease AtuBVI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.50 created 1982, deleted 1984]
- [3.1.23.51] *Transferred entry. endodeoxyribonuclease AcaI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.51 created 1982, deleted 1984]
- [3.1.23.52] *Transferred entry. endodeoxyribonuclease AcyI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*

[EC 3.1.23.52 created 1982, deleted 1984]

[3.1.23.53 *Transferred entry. endodeoxyribonuclease Aosl. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*]

[EC 3.1.23.53 created 1982, deleted 1984]

[3.1.23.54 *Transferred entry. endodeoxyribonuclease AsuII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*]

[EC 3.1.23.54 created 1982, deleted 1984]

[3.1.23.55 *Transferred entry. endodeoxyribonuclease AvaIII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*]

[EC 3.1.23.55 created 1982, deleted 1984]

[3.1.23.56 *Transferred entry. endodeoxyribonuclease AvrII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*]

[EC 3.1.23.56 created 1982, deleted 1984]

[3.1.23.57 *Transferred entry. endodeoxyribonuclease BceI4579. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as endodeoxyribonuclease Bce4579I (see <http://rebase.neb.com/rebase/rebase.html>)*]

[EC 3.1.23.57 created 1982, deleted 1984]

[3.1.23.58 *Transferred entry. endodeoxyribonuclease BceI229. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as endodeoxyribonuclease BceI229I (see <http://rebase.neb.com/rebase/rebase.html>)*]

[EC 3.1.23.58 created 1982, deleted 1984]

[3.1.23.59 *Transferred entry. endodeoxyribonuclease Bme899. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as endodeoxyribonuclease Bme899I (see <http://rebase.neb.com/rebase/rebase.html>)*]

[EC 3.1.23.59 created 1982, deleted 1984]

[3.1.23.60 *Transferred entry. endodeoxyribonuclease Bme205. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as endodeoxyribonuclease Bme205I (see <http://rebase.neb.com/rebase/rebase.html>)*]

[EC 3.1.23.60 created 1982, deleted 1984]

[3.1.23.61 *Transferred entry. endodeoxyribonuclease BmeI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*]

[EC 3.1.23.61 created 1982, deleted 1984]

[3.1.23.62 *Transferred entry. endodeoxyribonuclease BspI286. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as endodeoxyribonuclease BspI286I (see <http://rebase.neb.com/rebase/rebase.html>)*]

[EC 3.1.23.62 created 1982, deleted 1984]

[3.1.23.63 *Transferred entry. endodeoxyribonuclease BstAI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*]

[EC 3.1.23.63 created 1982, deleted 1984]

[3.1.23.64 *Transferred entry. endodeoxyribonuclease BstEI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*]

[EC 3.1.23.64 created 1982, deleted 1984]

[3.1.23.65 *Transferred entry. endodeoxyribonuclease BstEIII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*]

[EC 3.1.23.65 created 1982, deleted 1984]

[3.1.23.66 *Transferred entry. endodeoxyribonuclease BstPI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*]

[EC 3.1.23.66 created 1982, deleted 1984]

[3.1.23.67 *Transferred entry. endodeoxyribonuclease BsuM. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as endodeoxyribonuclease BsuMI (see <http://rebase.neb.com/rebase/rebase.html>)*]

[EC 3.1.23.67 created 1982, deleted 1984]

[3.1.23.68 Transferred entry. *endodeoxyribonuclease Bsu6633*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. The name was misprinted in supplement 3 of the 1978 edition. Assumed to be the same as *endodeoxyribonuclease Bsu6633I* (see <http://rebase.neb.com/rebase/rebase.html>)]

[EC 3.1.23.68 created 1982, deleted 1984]

[3.1.23.69 Transferred entry. *endodeoxyribonuclease Bsu1145*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as *endodeoxyribonuclease Bsu1145I* (see <http://rebase.neb.com/rebase/rebase.html>)]

[EC 3.1.23.69 created 1982, deleted 1984]

[3.1.23.70 Transferred entry. *endodeoxyribonuclease Bsu1192*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as *endodeoxyribonuclease Bsu1192I* or see *Bsu1192II* (see <http://rebase.neb.com/rebase/rebase.html>)]

[EC 3.1.23.70 created 1982, deleted 1984]

[3.1.23.71 Transferred entry. *endodeoxyribonuclease Bsu1193*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as *endodeoxyribonuclease Bsu1193I* (see <http://rebase.neb.com/rebase/rebase.html>)]

[EC 3.1.23.71 created 1982, deleted 1984]

[3.1.23.72 Transferred entry. *endodeoxyribonuclease Bsu1231*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Not found in <http://rebase.neb.com/rebase/rebase.html>]

[EC 3.1.23.72 created 1982, deleted 1984]

[3.1.23.73 Transferred entry. *endodeoxyribonuclease Bsu1259*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as *endodeoxyribonuclease Bsu1259I* (see <http://rebase.neb.com/rebase/rebase.html>)]

[EC 3.1.23.73 created 1982, deleted 1984]

[3.1.23.74 Transferred entry. *endodeoxyribonuclease ClaI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.74 created 1982, deleted 1984]

[3.1.23.75 Transferred entry. *endodeoxyribonuclease CauII*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.75 created 1982, deleted 1984]

[3.1.23.76 Transferred entry. *endodeoxyribonuclease CviI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.76 created 1982, deleted 1984]

[3.1.23.77 Transferred entry. *endodeoxyribonuclease DdeI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.77 created 1982, deleted 1984]

[3.1.23.78 Transferred entry. *endodeoxyribonuclease EclI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.78 created 1982, deleted 1984]

[3.1.23.79 Transferred entry. *endodeoxyribonuclease EcaI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.79 created 1982, deleted 1984]

[3.1.23.80 Transferred entry. *endodeoxyribonuclease EcoRI'*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as *endodeoxyribonuclease EcoRI'* (see <http://rebase.neb.com/rebase/rebase.html>)]

[EC 3.1.23.80 created 1982, deleted 1984]

[3.1.23.81 Transferred entry. *endodeoxyribonuclease Fnu48I*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.81 created 1982, deleted 1984]

[3.1.23.82 Transferred entry. *endodeoxyribonuclease Fnu4H*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as *endodeoxyribonuclease Fnu4HI* (see <http://rebase.neb.com/rebase/rebase.html>)]

- [EC 3.1.23.82 created 1982, deleted 1984]
- [3.1.23.83 *Transferred entry. endodeoxyribonuclease HapI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.83 created 1982, deleted 1984]
- [3.1.23.84 *Transferred entry. endodeoxyribonuclease Hin1056II. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.84 created 1982, deleted 1984]
- [3.1.23.85 *Transferred entry. endodeoxyribonuclease HinfIII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.85 created 1982, deleted 1984]
- [3.1.23.86 *Transferred entry. endodeoxyribonuclease HgiAI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.86 created 1982, deleted 1984]
- [3.1.23.87 *Transferred entry. endodeoxyribonuclease HgiCI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.87 created 1982, deleted 1984]
- [3.1.23.88 *Transferred entry. endodeoxyribonuclease HgiDI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.88 created 1982, deleted 1984]
- [3.1.23.89 *Transferred entry. endodeoxyribonuclease HgiEII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.89 created 1982, deleted 1984]
- [3.1.23.90 *Transferred entry. endodeoxyribonuclease MstI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.90 created 1982, deleted 1984]
- [3.1.23.91 *Transferred entry. endodeoxyribonuclease MstII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.91 created 1982, deleted 1984]
- [3.1.23.92 *Transferred entry. endodeoxyribonuclease MglI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.92 created 1982, deleted 1984]
- [3.1.23.93 *Transferred entry. endodeoxyribonuclease MglIII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.93 created 1982, deleted 1984]
- [3.1.23.94 *Transferred entry. endodeoxyribonuclease MnoII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.94 created 1982, deleted 1984]
- [3.1.23.95 *Transferred entry. endodeoxyribonuclease MnnIII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.95 created 1982, deleted 1984]
- [3.1.23.96 *Transferred entry. endodeoxyribonuclease MviI. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.96 created 1982, deleted 1984]
- [3.1.23.97 *Transferred entry. endodeoxyribonuclease MviII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.97 created 1982, deleted 1984]
- [3.1.23.98 *Transferred entry. endodeoxyribonuclease OxaII. Now EC 3.1.21.4, type II site-specific deoxyribonuclease*  
[EC 3.1.23.98 created 1982, deleted 1984]
- [3.1.23.99 *Transferred entry. endodeoxyribonuclease PaeR7. Now EC 3.1.21.4, type II site-specific deoxyribonuclease. Assumed to be the same as endodeoxyribonuclease PaeR7I (see <http://rebase.neb.com/rebase/rebase.html>)*

[EC 3.1.23.99 created 1982, deleted 1984]

[3.1.23.100 Transferred entry. *endodeoxyribonuclease RspI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.100 created 1982, deleted 1984]

[3.1.23.101 Transferred entry. *endodeoxyribonuclease RsaI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.101 created 1982, deleted 1984]

[3.1.23.102 Transferred entry. *endodeoxyribonuclease SmaI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.102 created 1982, deleted 1984]

[3.1.23.103 Transferred entry. *endodeoxyribonuclease SspI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.103 created 1982, deleted 1984]

[3.1.23.104 Transferred entry. *endodeoxyribonuclease SnaI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.104 created 1982, deleted 1984]

[3.1.23.105 Transferred entry. *endodeoxyribonuclease SfaNI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.105 created 1982, deleted 1984]

[3.1.23.106 Transferred entry. *endodeoxyribonuclease SalI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.106 created 1982, deleted 1984]

[3.1.23.107 Transferred entry. *endodeoxyribonuclease SauI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.107 created 1982, deleted 1984]

[3.1.23.108 Transferred entry. *endodeoxyribonuclease SphI*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.108 created 1982, deleted 1984]

[3.1.23.109 Transferred entry. *endodeoxyribonuclease XmaIII*. Now EC 3.1.21.4, type II site-specific deoxyribonuclease]

[EC 3.1.23.109 created 1982, deleted 1984]

### **EC 3.1.24 Site specific endodeoxyribonucleases: cleavage is not sequence specific (deleted sub-subclass)**

[3.1.24.1 Transferred entry. *endodeoxyribonuclease EcoB*. Now EC 3.1.21.3, type I site-specific deoxyribonuclease. Assumed to be the same as *endodeoxyribonuclease EcoBI* (see <http://rebase.neb.com/rebase/rebase.html>)]

[EC 3.1.24.1 created 1978, modified 1982, deleted 1984]

[3.1.24.2 Transferred entry. *endodeoxyribonuclease EcoK*. Now EC 3.1.21.3, type I site-specific deoxyribonuclease. Assumed to be the same as *endodeoxyribonuclease EcoKI* (see <http://rebase.neb.com/rebase/rebase.html>)]

[EC 3.1.24.2 created 1978, modified 1982, deleted 1984]

[3.1.24.3 Transferred entry. *endodeoxyribonuclease EcoPI*. Now EC 3.1.21.5, type III site-specific deoxyribonuclease. The name is misprinted in supplement 3 of the 1978 edition]

[EC 3.1.24.3 created 1978, modified 1982, deleted 1984]

[3.1.24.4 Transferred entry. *endodeoxyribonuclease EcoP15*. Now EC 3.1.21.5, type III site-specific deoxyribonuclease. Assumed to be the same as *endodeoxyribonuclease EcoP15I* (see <http://rebase.neb.com/rebase/rebase.html>)]

[EC 3.1.24.4 created 1978, modified 1982, deleted 1984]

### **EC 3.1.25 Site-specific endodeoxyribonucleases that are specific for altered bases**

#### EC 3.1.25.1

**Accepted name:** deoxyribonuclease (pyrimidine dimer)  
**Reaction:** Endonucleolytic cleavage near pyrimidine dimers to products with 5'-phosphate  
**Other name(s):** endodeoxyribonuclease (pyrimidine dimer); endodeoxyribonuclease (pyrimidine dimer); bacteriophage T<sub>4</sub> endodeoxyribonuclease V; T4 endonuclease V  
**Comments:** Acts on a damaged strand, 5' from the damaged site.  
**References:** [279, 2416]

[EC 3.1.25.1 created 1978]

[3.1.25.2 *Transferred entry. endodeoxyribonuclease (apurinic or apyrimidinic). Now EC 4.2.99.18, DNA-(apurinic or apyrimidinic site) lyase*]

[EC 3.1.25.2 created 1978, deleted 1992]

### EC 3.1.26 Endoribonucleases producing 5'-phosphomonoesters

#### EC 3.1.26.1

**Accepted name:** *Physarum polycephalum* ribonuclease  
**Reaction:** Endonucleolytic cleavage to 5'-phosphomonoester  
**References:** [1145]

[EC 3.1.26.1 created 1978]

#### EC 3.1.26.2

**Accepted name:** ribonuclease  $\alpha$   
**Reaction:** Endonucleolytic cleavage to 5'-phosphomonoester  
**Other name(s):** 2'-O-methyl RNase  
**Comments:** Specific for O-methylated RNA.  
**References:** [2116]

[EC 3.1.26.2 created 1978]

#### EC 3.1.26.3

**Accepted name:** ribonuclease III  
**Reaction:** Endonucleolytic cleavage to a 5'-phosphomonoester  
**Other name(s):** RNase III; ribonuclease 3  
**Comments:** This is an endoribonuclease that cleaves double-stranded RNA molecules [994]. The cleavage can be either a single-stranded nick or double-stranded break in the RNA, depending in part upon the degree of base-pairing in the region of the cleavage site [497]. Specificity is conferred by negative determinants, i.e., the presence of certain Watson-Crick base-pairs at specific positions that strongly inhibit cleavage [3328]. RNase III is involved in both rRNA processing and mRNA processing and decay.  
**References:** [511, 2391, 2440, 994, 497, 3328]

[EC 3.1.26.3 created 1978, modified 2006]

#### EC 3.1.26.4

**Accepted name:** ribonuclease H  
**Reaction:** Endonucleolytic cleavage to 5'-phosphomonoester  
**Other name(s):** endoribonuclease H (calf thymus); RNase H; RNA\*DNA hybrid ribonucleotidohydrolase; hybrid ribonuclease; hybridase; hybridase (ribonuclease H); ribonuclease H; hybrid nuclease; calf thymus ribonuclease H  
**Comments:** Acts on RNA-DNA hybrids.

**References:** [1014, 2757]

[EC 3.1.26.4 created 1978, modified 2010]

#### EC 3.1.26.5

**Accepted name:** ribonuclease P

**Reaction:** Endonucleolytic cleavage of RNA, removing 5'-extranucleotides from tRNA precursor

**Other name(s):** RNase P

**Comments:** An RNA-containing enzyme, essential for tRNA processing; generates 5'-termini or mature tRNA molecules.

**References:** [215, 216, 2439]

[EC 3.1.26.5 created 1978, modified 1982]

#### EC 3.1.26.6

**Accepted name:** ribonuclease IV

**Reaction:** Endonucleolytic cleavage of poly(A) to fragments terminated by 3'-hydroxy and 5'-phosphate groups

**Other name(s):** endoribonuclease IV; poly(A)-specific ribonuclease

**Comments:** Forms oligonucleotides with an average chain length of 10.

**References:** [1995, 1996]

[EC 3.1.26.6 created 1984]

#### EC 3.1.26.7

**Accepted name:** ribonuclease P4

**Reaction:** Endonucleolytic cleavage of RNA, removing 3'-extranucleotides from tRNA precursor

**References:** [2610]

[EC 3.1.26.7 created 1984]

#### EC 3.1.26.8

**Accepted name:** ribonuclease M5

**Reaction:** Endonucleolytic cleavage of RNA, removing 21 and 42 nucleotides, respectively, from the 5'- and 3'-termini of a 5S-rRNA precursor

**Other name(s):** RNase M5; 5S ribosomal maturation nuclease; 5S ribosomal RNA maturation endonuclease

**Comments:** Converts the 5S-rRNA precursor from *Bacillus subtilis* into 5S-rRNA, with 5'-phosphate and 3'-hydroxy groups.

**References:** [2708]

[EC 3.1.26.8 created 1986]

#### EC 3.1.26.9

**Accepted name:** ribonuclease [poly-(U)-specific]

**Reaction:** Endonucleolytic cleavage of poly(U) to fragments terminated by 3'-hydroxy and 5'-phosphate groups

**Other name(s):** ribonuclease (uracil-specific); uracil-specific endoribonuclease; uracil-specific RNase

**Comments:** Forms oligonucleotides with chain lengths of 6 to 12.

**References:** [115]

[EC 3.1.26.9 created 1986]

#### EC 3.1.26.10

**Accepted name:** ribonuclease IX  
**Reaction:** Endonucleolytic cleavage of poly(U) or poly(C) to fragments terminated by 3'-hydroxy and 5'-phosphate groups  
**Other name(s):** poly(U)- and poly(C)-specific endoribonuclease  
**Comments:** Acts on poly(U) and poly(C), with a higher affinity for poly(C), but does not act on poly(A) or poly(G).  
**References:** [2652]

[EC 3.1.26.10 created 1992]

#### EC 3.1.26.11

**Accepted name:** tRNase Z  
**Reaction:** endonucleolytic cleavage of RNA, removing extra 3' nucleotides from tRNA precursor, generating 3' termini of tRNAs. A 3'-hydroxy group is left at the tRNA terminus and a 5'-phosphoryl group is left at the trailer molecule  
**Other name(s):** 3 tRNase; tRNA 3 endonuclease; RNase Z; 3' tRNase  
**Comments:** No cofactor requirements. An homologous enzyme to that found in *Arabidopsis thaliana* has been found in *Methanococcus janaschii*.  
**References:** [2565, 1853, 2564, 1558, 1979, 1921, 2851]

[EC 3.1.26.11 created 2002]

#### EC 3.1.26.12

**Accepted name:** ribonuclease E  
**Reaction:** Endonucleolytic cleavage of single-stranded RNA in A- and U-rich regions  
**Other name(s):** endoribonuclease E; RNase E; Rne protein  
**Comments:** RNase E is a bacterial ribonuclease that plays a role in the processing of ribosomal RNA (9S to 5S rRNA), the chemical degradation of bulk cellular RNA, the decay of specific regulatory, messenger and structural RNAs and the control of plasmid DNA replication [754]. The enzyme binds to monophosphorylated 5' ends of substrates but exhibits sequential cleavages in the 3' to 5' direction [754]. 2'-O-Methyl nucleotide substitutions at RNase E binding sites do not prevent binding but do prevent cleavage of non-modified target sequences 5' to that locus [754]. In *Escherichia coli*, the enzyme is found in the RNA degradosome. The C-terminal half of the protein contains binding sites for the three other major degradosomal components, the DEAD-box RNA helicase Rh1B, enolase (EC 4.1.1.11) and polynucleotide phosphorylase (EC 2.7.7.8).  
**References:** [754, 683, 488, 3051, 2758, 356]

[EC 3.1.26.12 created 2008]

#### EC 3.1.26.13

**Accepted name:** retroviral ribonuclease H  
**Reaction:** Endohydrolysis of RNA in RNA/DNA hybrids. Three different cleavage modes: 1. sequence-specific internal cleavage of RNA [1-4]. Human immunodeficiency virus type 1 and Moloney murine leukemia virus enzymes prefer to cleave the RNA strand one nucleotide away from the RNA-DNA junction [5]. 2. RNA 5'-end directed cleavage 13-19 nucleotides from the RNA end [6,7]. 3. DNA 3'-end directed cleavage 15-20 nucleotides away from the primer terminus [8-10].  
**Other name(s):** RT/RNase H; retroviral reverse transcriptase RNaseH; HIV RNase H  
**Comments:** Retroviral reverse transcriptase is a multifunctional enzyme responsible for viral replication. To perform this task the enzyme combines two distinct activities. The polymerase domain (EC 2.7.7.49, RNA-directed DNA polymerase) occupies the N-terminal two-thirds of the reverse transcriptase whereas the ribonuclease H domain comprises the C-terminal remaining one-third [390, 2589]. The RNase H domain of Moloney murine leukemia virus and Human immunodeficiency virus display two metal binding sites [937, 548, 2233]  
**References:** [2590, 2527, 2378, 286, 2591, 591, 1396, 2225, 828, 185, 1203, 1540, 390, 2589, 937, 548, 2233]



[EC 3.1.26.13 created 2009]

### EC 3.1.27 Endoribonucleases producing 3'-phosphomonoesters

[3.1.27.1 *Transferred entry. ribonuclease T<sub>2</sub>. Now EC 4.6.1.19, ribonuclease T<sub>2</sub>, since the primary reaction is that of a lyase*]

[EC 3.1.27.1 created 1972 as EC 3.1.4.23, transferred 1978 to EC 3.1.27.1, modified 1981, deleted 2018]

[3.1.27.2 *Transferred entry. Bacillus subtilis ribonuclease. Now EC 4.6.1.22, Bacillus subtilis ribonuclease, since the reaction catalysed is that of a lyase*]

[EC 3.1.27.2 created 1978, deleted 2018]

#### EC 3.1.27.3

**Accepted name:** ribonuclease T<sub>1</sub>

**Reaction:** Two-stage endonucleolytic cleavage to nucleoside 3'-phosphates and 3'-phosphooligonucleotides ending in Gp with 2',3'-cyclic phosphate intermediates

**Other name(s):** guanyloribonuclease; *Aspergillus oryzae* ribonuclease; RNase N<sub>1</sub>; RNase N<sub>2</sub>; ribonuclease N<sub>3</sub>; ribonuclease U<sub>1</sub>; ribonuclease F1; ribonuclease Ch; ribonuclease PP1; ribonuclease SA; RNase F1; ribonuclease C2; binase; RNase Sa; guanyl-specific RNase; RNase G; RNase T<sub>1</sub>; ribonuclease guaninenucleotido-2'-transferase (cyclizing); ribonuclease N<sub>3</sub>; ribonuclease N<sub>1</sub>

**Comments:** Formerly EC 2.7.7.26 and EC 3.1.4.8.

**References:** [1392, 2837]

[EC 3.1.27.3 created 1961 as EC 3.1.4.8, transferred 1965 to EC 2.7.7.26, reinstated 1972 as EC 3.1.4.8, transferred 1978 to EC 3.1.27.3]

[3.1.27.4 *Transferred entry. ribonuclease U<sub>2</sub>. Now EC 4.6.1.20, ribonuclease U<sub>2</sub>, since the primary reaction is that of a lyase*]

[EC 3.1.27.4 created 1978, modified 1981, deleted 2018]

[3.1.27.5 *Transferred entry. pancreatic ribonuclease. Now EC 4.6.1.18, pancreatic ribonuclease. This reaction is now known to involve an internal-transfer (lyase) process to produce the cyclic derivative, followed by a reversal of that step with water in the "hydrolytic step"*]

[EC 3.1.27.5 created 1972 as EC 3.1.4.22, transferred 1978 to EC 3.1.27.5, modified 1981, deleted 2018]

[3.1.27.6 *Transferred entry. Enterobacter ribonuclease. Now EC 4.6.1.21, Enterobacter ribonuclease, since the primary reaction is that of a lyase*]

[EC 3.1.27.6 created 1978, modified 1981, deleted 2018]

#### EC 3.1.27.7

**Accepted name:** ribonuclease F

**Reaction:** Endonucleolytic cleavage of RNA precursor into two, leaving 5'-hydroxy and 3'-phosphate groups

**Other name(s):** ribonuclease F (*E. coli*)

**References:** [1004, 3138]

[EC 3.1.27.7 created 1984]

#### EC 3.1.27.8

**Accepted name:** ribonuclease V

**Reaction:** Hydrolysis of poly(A), forming oligoribonucleotides and ultimately 3'-AMP

**Other name(s):** endoribonuclease V

**Comments:** Also hydrolyses poly(U).

**References:** [2585]

[EC 3.1.27.8 created 1984]

[3.1.27.9 Transferred entry. tRNA-intron endonuclease. Now EC 4.6.1.16, tRNA-intron lyase]

[EC 3.1.27.9 created 1992, deleted 2014]

#### EC 3.1.27.10

**Accepted name:** rRNA endonuclease

**Reaction:** Hydrolysis of the phosphodiester linkage between guanosine and adenosine residues at one specific position in 28S rRNA from rat ribosomes

**Other name(s):**  $\alpha$ -sarcin

**Comments:** Also acts on bacterial rRNA.

**References:** [704]

[EC 3.1.27.10 created 1992]

### EC 3.1.30 Endoribonucleases that are active with either ribo- or deoxyribonucleic acids and produce 5'-phosphomonoesters

#### EC 3.1.30.1

**Accepted name:** *Aspergillus* nuclease S<sub>1</sub>

**Reaction:** Endonucleolytic cleavage to 5'-phosphomononucleotide and 5'-phosphooligonucleotide end-products

**Other name(s):** endonuclease S<sub>1</sub> (*Aspergillus*); single-stranded-nucleate endonuclease; deoxyribonuclease S<sub>1</sub>; deoxyribonuclease S<sub>1</sub>; nuclease S<sub>1</sub>; *Neurospora crassa* single-strand specific endonuclease; S1 nuclease; single-strand endodeoxyribonuclease; single-stranded DNA specific endonuclease; single-strand-specific endodeoxyribonuclease; single strand-specific DNase; *Aspergillus oryzae* S1 nuclease

**References:** [55, 2809, 3076]

[EC 3.1.30.1 created 1972 as EC 3.1.4.21, transferred 1978 to EC 3.1.30.1, modified 1981]

#### EC 3.1.30.2

**Accepted name:** *Serratia marcescens* nuclease

**Reaction:** Endonucleolytic cleavage to 5'-phosphomononucleotide and 5'-phosphooligonucleotide end-products

**Other name(s):** endonuclease (*Serratia marcescens*); barley nuclease; plant nuclease I; nucleate endonuclease

**Comments:** Hydrolyses double- or single-stranded substrate.

**References:** [1909, 2767, 2768, 3145]

[EC 3.1.30.2 created 1965 as EC 3.1.4.9, transferred 1978 to EC 3.1.30.2, modified 1981]

### EC 3.1.31 Endoribonucleases that are active with either ribo- or deoxyribonucleic acids and produce 3'-phosphomonoesters

#### EC 3.1.31.1

**Accepted name:** micrococcal nuclease

**Reaction:** Endonucleolytic cleavage to nucleoside 3'-phosphates and 3'-phosphooligonucleotide end-products

**Other name(s):** spleen endonuclease; thermonuclease; nuclease T; micrococcal endonuclease; nuclease T'; staphylococcal nuclease; spleen phosphodiesterase; *Staphylococcus aureus* nuclease; *Staphylococcus aureus* nuclease B; ribonucleate (deoxynucleate) 3'-nucleotidohydrolase

**Comments:** Hydrolyses double- or single-stranded substrate.

**References:** [31, 57, 2393, 2798]

[EC 3.1.31.1 created 1961 as EC 3.1.4.7, transferred 1978 to EC 3.1.31.1, modified 1981]

## EC 3.2 Glycosylases

This subclass contains the glycosylases, which are classified as hydrolases, although some of them can also transfer glycosyl residues to oligosaccharides, polysaccharides and other alcoholic acceptors. The glycosylases are subdivided into glycosidases, i.e., enzymes that hydrolyse *O*- and *S*-glycosyl compounds (EC 3.2.1) and those that hydrolyse *N*-glycosyl compounds (EC 3.2.2). Common names for enzymes acting on D-sugars or their derivatives do not normally contain 'D', unless ambiguity would result from the common existence of the corresponding L-sugar. Enzymes that hydrolyse a terminal, non-reducing-end glucose (or a well-defined di-, tri- or oligosaccharide) from a glycan, i.e. exoenzymes, are given systematic names based on 'glycohydrolase'; enzymes that hydrolyse internal glycosidic bonds, i.e. endoenzymes, are given systematic names based on 'glycanohydrolase'. The same structure is often used when providing accepted names for these enzymes.

### EC 3.2.1 Glycosidases, i.e. enzymes that hydrolyse *O*- and *S*-glycosyl compounds

#### EC 3.2.1.1

- Accepted name:**  $\alpha$ -amylase  
**Reaction:** Endohydrolysis of (1 $\rightarrow$ 4)- $\alpha$ -D-glucosidic linkages in polysaccharides containing three or more (1 $\rightarrow$ 4)- $\alpha$ -linked D-glucose units  
**Other name(s):** glycogenase;  $\alpha$  amylase,  $\alpha$ -amylase; endoamylase; Taka-amylase A; 1,4- $\alpha$ -D-glucan glucohydrolase  
**Systematic name:** 4- $\alpha$ -D-glucan glucohydrolase  
**Comments:** Acts on starch, glycogen and related polysaccharides and oligosaccharides in a random manner; reducing groups are liberated in the  $\alpha$ -configuration. The term " $\alpha$ " relates to the initial anomeric configuration of the free sugar group released and not to the configuration of the linkage hydrolysed.  
**References:** [771, 1800, 2598]

[EC 3.2.1.1 created 1961]

#### EC 3.2.1.2

- Accepted name:**  $\beta$ -amylase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 4)- $\alpha$ -D-glucosidic linkages in polysaccharides so as to remove successive maltose units from the non-reducing ends of the chains  
**Other name(s):** saccharogen amylase; glycogenase;  $\beta$  amylase,  $\beta$ -amylase; 1,4- $\alpha$ -D-glucan maltohydrolase  
**Systematic name:** 4- $\alpha$ -D-glucan maltohydrolase  
**Comments:** Acts on starch, glycogen and related polysaccharides and oligosaccharides producing  $\beta$ -maltose by an inversion. The term " $\beta$ " relates to the initial anomeric configuration of the free sugar group released and not to the configuration of the linkage hydrolysed.  
**References:** [129, 807, 1800]

[EC 3.2.1.2 created 1961]

#### EC 3.2.1.3

- Accepted name:** glucan 1,4- $\alpha$ -glucosidase  
**Reaction:** Hydrolysis of terminal (1 $\rightarrow$ 4)-linked  $\alpha$ -D-glucose residues successively from non-reducing ends of the chains with release of  $\beta$ -D-glucose  
**Other name(s):** glucoamylase; amyloglucosidase;  $\gamma$ -amylase; lysosomal  $\alpha$ -glucosidase; acid maltase; exo-1,4- $\alpha$ -glucosidase; glucose amylase;  $\gamma$ -1,4-glucan glucohydrolase; acid maltase; 1,4- $\alpha$ -D-glucan glucohydrolase  
**Systematic name:** 4- $\alpha$ -D-glucan glucohydrolase

**Comments:** Most forms of the enzyme can rapidly hydrolyse 1,6- $\alpha$ -D-glucosidic bonds when the next bond in the sequence is 1,4, and some preparations of this enzyme hydrolyse 1,6- and 1,3- $\alpha$ -D-glucosidic bonds in other polysaccharides. This entry covers all such enzymes acting on polysaccharides more rapidly than on oligosaccharides. EC 3.2.1.20  $\alpha$ -glucosidase, from mammalian intestine, can catalyse similar reactions.

**References:** [808, 307, 1314, 1417, 1916, 2980]

[EC 3.2.1.3 created 1961]

#### EC 3.2.1.4

**Accepted name:** cellulase

**Reaction:** Endohydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-glucosidic linkages in cellulose, lichenin and cereal  $\beta$ -D-glucans

**Other name(s):** endo-1,4- $\beta$ -D-glucanase;  $\beta$ -1,4-glucanase;  $\beta$ -1,4-endoglucan hydrolase; cellulase A; cellulose AP; endoglucanase D; alkali cellulase; cellulase A 3; celludextrinase; 9.5 cellulase; avicelase; pancellase SS; 1,4-(1,3;1,4)- $\beta$ -D-glucan 4-glucanohydrolase

**Systematic name:** 4- $\beta$ -D-glucan 4-glucanohydrolase

**Comments:** Will also hydrolyse 1,4-linkages in  $\beta$ -D-glucans also containing 1,3-linkages.

**References:** [540, 1603, 2015, 2096, 3169, 1077, 1078, 1252]

[EC 3.2.1.4 created 1961, modified 2001]

#### [3.2.1.5 Deleted entry. licheninase]

[EC 3.2.1.5 created 1961, deleted 1964]

#### EC 3.2.1.6

**Accepted name:** endo-1,3(4)- $\beta$ -glucanase

**Reaction:** Endohydrolysis of (1 $\rightarrow$ 3)- or (1 $\rightarrow$ 4)-linkages in  $\beta$ -D-glucans when the glucose residue whose reducing group is involved in the linkage to be hydrolysed is itself substituted at C-3

**Other name(s):** endo-1,3- $\beta$ -D-glucanase; laminarinase; laminaranase;  $\beta$ -1,3-glucanase;  $\beta$ -1,3-1,4-glucanase; endo-1,3- $\beta$ -glucanase; endo- $\beta$ -1,3(4)-glucanase; endo- $\beta$ -1,3-1,4-glucanase; endo- $\beta$ -(1 $\rightarrow$ 3)-D-glucanase; endo-1,3-1,4- $\beta$ -D-glucanase; endo- $\beta$ -(1-3)-D-glucanase; endo- $\beta$ -1,3-glucanase IV; endo-1,3- $\beta$ -D-glucanase; 1,3-(1,3;1,4)- $\beta$ -D-glucan 3(4)-glucanohydrolase

**Systematic name:** 3(or 4)- $\beta$ -D-glucan 3(4)-glucanohydrolase

**Comments:** Substrates include laminarin, lichenin and cereal D-glucans; different from EC 3.2.1.52  $\beta$ -N-acetylhexosaminidase.

**References:** [152, 153, 519, 2397, 2734]

[EC 3.2.1.6 created 1961, modified 1976]

#### EC 3.2.1.7

**Accepted name:** inulinase

**Reaction:** Endohydrolysis of (2 $\rightarrow$ 1)- $\beta$ -D-fructosidic linkages in inulin

**Other name(s):** inulase; indoinulinase; endo-inulinase; exoinulinase; 2,1- $\beta$ -D-fructan fructanohydrolase

**Systematic name:** 1- $\beta$ -D-fructan fructanohydrolase

**References:** [16]

[EC 3.2.1.7 created 1961]

#### EC 3.2.1.8

**Accepted name:** endo-1,4- $\beta$ -xylanase

**Reaction:** Endohydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-xylosidic linkages in xylans

**Other name(s):** endo-(1→4)- $\beta$ -xylan 4-xylanohydrolase; endo-1,4-xylanase; xylanase;  $\beta$ -1,4-xylanase; endo-1,4-xylanase; endo- $\beta$ -1,4-xylanase; endo-1,4- $\beta$ -D-xylanase; 1,4- $\beta$ -xylan xylanohydrolase;  $\beta$ -xylanase;  $\beta$ -1,4-xylan xylanohydrolase; endo-1,4- $\beta$ -xylanase;  $\beta$ -D-xylanase  
**Systematic name:** 4- $\beta$ -D-xylan xylanohydrolase  
**References:** [1192, 3168]

[EC 3.2.1.8 created 1961]

[3.2.1.9 Deleted entry. amylopectin-1,6-glucosidase]

[EC 3.2.1.9 created 1961, deleted 1972]

#### EC 3.2.1.10

**Accepted name:** oligo-1,6-glucosidase  
**Reaction:** Hydrolysis of (1→6)- $\alpha$ -D-glucosidic linkages in some oligosaccharides produced from starch and glycogen by EC 3.2.1.1 ( $\alpha$ -amylase), and in isomaltose  
**Other name(s):** limit dextrinase (erroneous); isomaltase; sucrase-isomaltase; exo-oligo-1,6-glucosidase; dextrin 6 $\alpha$ -glucanohydrolase;  $\alpha$ -limit dextrinase; dextrin 6-glucanohydrolase; oligosaccharide  $\alpha$ -1,6-glucohydrolase;  $\alpha$ -methylglucosidase  
**Systematic name:** oligosaccharide 6- $\alpha$ -glucohydrolase  
**Comments:** This enzyme, like EC 3.2.1.33 (amylo- $\alpha$ -1,6-glucosidase), can release an  $\alpha$ -1→6-linked glucose, whereas the shortest chain that can be released by EC 3.2.1.41 (pullulanase), EC 3.2.1.142 (limit dextrinase), and EC 3.2.1.68 (isoamylase) is maltose. It also hydrolyses isomaltulose (palatinose), isomaltotriose and panose, but has no action on glycogen or phosphorylase limit dextrin. The enzyme from intestinal mucosa is a single polypeptide chain that also catalyses the reaction of EC 3.2.1.48 (sucrose  $\alpha$ -glucosidase). Differs from EC 3.2.1.33 (amylo- $\alpha$ -1,6-glucosidase) in its preference for short-chain substrates and in its not requiring the 6-glucosylated residue to be at a branch point, i.e. linked at both C-1 and C-4.  
**References:** [1080, 2678, 2449, 1433, 3248]

[EC 3.2.1.10 created 1961, modified 2000, modified 2013]

#### EC 3.2.1.11

**Accepted name:** dextranase  
**Reaction:** Endohydrolysis of (1→6)- $\alpha$ -D-glucosidic linkages in dextran  
**Other name(s):** dextran hydrolase; endodextranase; dextranase DL 2; DL 2; endo-dextranase;  $\alpha$ -D-1,6-glucan-6-glucanohydrolase; 1,6- $\alpha$ -D-glucan 6-glucanohydrolase  
**Systematic name:** 6- $\alpha$ -D-glucan 6-glucanohydrolase  
**References:** [121, 593, 772, 2474]

[EC 3.2.1.11 created 1961]

[3.2.1.12 Deleted entry. cycloheptagluconase. Now included with EC 3.2.1.54 cyclomaltodextrinase]

[EC 3.2.1.12 created 1961, deleted 1976]

[3.2.1.13 Deleted entry. cyclohexagluconase. Now included with EC 3.2.1.54 cyclomaltodextrinase]

[EC 3.2.1.13 created 1961, deleted 1976]

#### EC 3.2.1.14

**Accepted name:** chitinase  
**Reaction:** Random endo-hydrolysis of *N*-acetyl- $\beta$ -D-glucosaminide (1→4)- $\beta$ -linkages in chitin and chitodextrins  
**Other name(s):** ChiC; chitodextrinase (ambiguous); 1,4- $\beta$ -poly-*N*-acetylglucosaminidase; poly- $\beta$ -glucosaminidase;  $\beta$ -1,4-poly-*N*-acetyl glucosaminidase; poly[1,4-(*N*-acetyl- $\beta$ -D-glucosaminide)] glycanohydrolase

**Systematic name:** (1→4)-2-acetamido-2-deoxy-β-D-glucan glycanohydrolase  
**Comments:** The enzyme binds to chitin and randomly cleaves glycosidic linkages in chitin and chitodextrins in a non-processive mode, generating chitooligosaccharides and free ends on which exo-chitinases and exo-chitodextrinases can act. Activity is greatly stimulated in the presence of EC 1.14.99.53, lytic chitin monooxygenase, which attacks the crystalline structure of chitin and makes the polymer more accessible to the chitinase. *cf.* EC 3.2.1.202, endo-chitodextrinase.  
**References:** [3319, 2958, 772, 477, 802, 3352, 2465]

[EC 3.2.1.14 created 1961, modified 2017]

#### EC 3.2.1.15

**Accepted name:** polygalacturonase  
**Reaction:** Random hydrolysis of (1→4)-α-D-galactosiduronic linkages in pectate and other galacturonans  
**Other name(s):** pectin depolymerase; pectinase; endopolygalacturonase; pectolase; pectin hydrolase; pectin polygalacturonase; endo-polygalacturonase; poly-α-1,4-galacturonide glycanohydrolase; endogalacturonase; endo-D-galacturonase; poly(1,4-α-D-galacturonide) glycanohydrolase  
**Systematic name:** (1→4)-α-D-galacturonan glycanohydrolase  
**References:** [593, 1704, 1864, 1911, 2272]

[EC 3.2.1.15 created 1961]

[3.2.1.16 Deleted entry. *alginate*]

[EC 3.2.1.16 created 1961, deleted 1972]

#### EC 3.2.1.17

**Accepted name:** lysozyme  
**Reaction:** Hydrolysis of (1→4)-β-linkages between *N*-acetylmuramic acid and *N*-acetyl-D-glucosamine residues in a peptidoglycan and between *N*-acetyl-D-glucosamine residues in chitodextrins  
**Other name(s):** muramidase; globulin G; mucopeptide glucohydrolase; globulin G1; *N,O*-diacetylmuramidase; lysozyme g; L-7001; 1,4-*N*-acetylmuramidase; mucopeptide *N*-acetylmuramoylhydrolase; PR1-lysozyme  
**Systematic name:** peptidoglycan *N*-acetylmuramoylhydrolase  
**Comments:** *cf.* also EC 3.2.1.14 chitinase.  
**References:** [232, 234, 1334]

[EC 3.2.1.17 created 1961]

#### EC 3.2.1.18

**Accepted name:** exo-α-sialidase  
**Reaction:** Hydrolysis of α-(2→3)-, α-(2→6)-, α-(2→8)- glycosidic linkages of terminal sialic acid residues in oligosaccharides, glycoproteins, glycolipids, colominic acid and synthetic substrates  
**Other name(s):** neuraminidase; sialidase; α-neuraminidase; acetylneuraminidase  
**Systematic name:** acetylneuraminyl hydrolase  
**Comments:** The enzyme does not act on 4-*O*-acetylated sialic acids. endo-α-Sialidase activity is listed as EC 3.2.1.129, endo-α-sialidase. See also EC 4.2.2.15 anhydrosialidase.  
**References:** [2556, 347]

[EC 3.2.1.18 created 1961, modified 1999]

[3.2.1.19 Deleted entry. *heparinase*]

[EC 3.2.1.19 created 1961, deleted 1978]

#### EC 3.2.1.20

**Accepted name:**  $\alpha$ -glucosidase  
**Reaction:** Hydrolysis of terminal, non-reducing (1 $\rightarrow$ 4)-linked  $\alpha$ -D-glucose residues with release of D-glucose  
**Other name(s):** maltase; glucoinvertase; glucosidosucrase; maltase-glucoamylase;  $\alpha$ -glucopyranosidase; glucosidoinvertase;  $\alpha$ -D-glucosidase;  $\alpha$ -glucoside hydrolase;  $\alpha$ -1,4-glucosidase  
**Systematic name:**  $\alpha$ -D-glucoside glucohydrolase  
**Comments:** This single entry covers a group of enzymes whose specificity is directed mainly towards the exo-hydrolysis of (1 $\rightarrow$ 4)- $\alpha$ -glucosidic linkages, and that hydrolyse oligosaccharides rapidly, relative to polysaccharide, which are hydrolysed relatively slowly, or not at all. The intestinal enzyme also hydrolyses polysaccharides, catalysing the reactions of EC 3.2.1.3 glucan 1,4- $\alpha$ -glucosidase and, more slowly, hydrolyses (1 $\rightarrow$ 6)- $\alpha$ -D-glucose links.  
**References:** [319, 781, 1603, 2675, 2726]

[EC 3.2.1.20 created 1961]

#### EC 3.2.1.21

**Accepted name:**  $\beta$ -glucosidase  
**Reaction:** Hydrolysis of terminal, non-reducing  $\beta$ -D-glucosyl residues with release of  $\beta$ -D-glucose  
**Other name(s):** gentiobiase; cellobiase; emulsin; elaterase; aryl- $\beta$ -glucosidase;  $\beta$ -D-glucosidase;  $\beta$ -glucoside glucohydrolase; arbutinase; amygdalinase; *p*-nitrophenyl  $\beta$ -glucosidase; primeverosidase; amygdalase; linamarase; salicilinase;  $\beta$ -1,6-glucosidase  
**Systematic name:**  $\beta$ -D-glucoside glucohydrolase  
**Comments:** Wide specificity for  $\beta$ -D-glucosides. Some examples also hydrolyse one or more of the following:  $\beta$ -D-galactosides,  $\alpha$ -L-arabinosides,  $\beta$ -D-xylosides,  $\beta$ -D-fucosides.  
**References:** [435, 471, 529, 1135, 1603, 2520]

[EC 3.2.1.21 created 1961]

#### EC 3.2.1.22

**Accepted name:**  $\alpha$ -galactosidase  
**Reaction:** Hydrolysis of terminal, non-reducing  $\alpha$ -D-galactose residues in  $\alpha$ -D-galactosides, including galactose oligosaccharides, galactomannans and galactolipids  
**Other name(s):** melibiase;  $\alpha$ -D-galactosidase;  $\alpha$ -galactosidase A;  $\alpha$ -galactoside galactohydrolase  
**Systematic name:**  $\alpha$ -D-galactoside galactohydrolase  
**Comments:** Also hydrolyses  $\alpha$ -D-fucosides.  
**References:** [2811, 3181]

[EC 3.2.1.22 created 1961]

#### EC 3.2.1.23

**Accepted name:**  $\beta$ -galactosidase  
**Reaction:** Hydrolysis of terminal non-reducing  $\beta$ -D-galactose residues in  $\beta$ -D-galactosides  
**Other name(s):** lactase (ambiguous);  $\beta$ -lactosidase; maxilact; hydrolact;  $\beta$ -D-lactosidase; S 2107; lactozym; trilactase;  $\beta$ -D-galactanase; oryzatym; sumiklat  
**Systematic name:**  $\beta$ -D-galactoside galactohydrolase  
**Comments:** Some enzymes in this group hydrolyse  $\alpha$ -L-arabinosides; some animal enzymes also hydrolyse  $\beta$ -D-fucosides and  $\beta$ -D-glucosides; *cf.* EC 3.2.1.108 lactase.  
**References:** [235, 1542, 1559, 1596, 1729, 1953, 3099, 89]

[EC 3.2.1.23 created 1961, modified 1980]

#### EC 3.2.1.24

**Accepted name:**  $\alpha$ -mannosidase  
**Reaction:** Hydrolysis of terminal, non-reducing  $\alpha$ -D-mannose residues in  $\alpha$ -D-mannosides

**Other name(s):**  $\alpha$ -D-mannosidase; *p*-nitrophenyl- $\alpha$ -mannosidase;  $\alpha$ -D-mannopyranosidase; 1,2- $\alpha$ -mannosidase; 1,2- $\alpha$ -D-mannosidase; exo- $\alpha$ -mannosidase  
**Systematic name:**  $\alpha$ -D-mannoside mannohydrolase  
**Comments:** Also hydrolyses  $\alpha$ -D-lyxosides and heptopyranosides with the same configuration at C-2, C-3 and C-4 as mannose.  
**References:** [1675, 3194]

[EC 3.2.1.24 created 1961]

#### EC 3.2.1.25

**Accepted name:**  $\beta$ -mannosidase  
**Reaction:** Hydrolysis of terminal, non-reducing  $\beta$ -D-mannose residues in  $\beta$ -D-mannosides  
**Other name(s):** mannanase; mannase;  $\beta$ -D-mannosidase;  $\beta$ -mannoside mannohydrolase; exo- $\beta$ -D-mannanase  
**Systematic name:**  $\beta$ -D-mannoside mannohydrolase  
**References:** [16, 165, 592, 1215]

[EC 3.2.1.25 created 1961]

#### EC 3.2.1.26

**Accepted name:**  $\beta$ -fructofuranosidase  
**Reaction:** Hydrolysis of terminal non-reducing  $\beta$ -D-fructofuranoside residues in  $\beta$ -D-fructofuranosides  
**Other name(s):** invertase; saccharase; glucosucrase;  $\beta$ -h-fructosidase;  $\beta$ -fructosidase; invertin; sucrase; maxinvert L 1000; fructosylinvertase; alkaline invertase; acid invertase  
**Systematic name:**  $\beta$ -D-fructofuranoside fructohydrolase  
**Comments:** Substrates include sucrose; also catalyses fructotransferase reactions.  
**References:** [2017, 2071]

[EC 3.2.1.26 created 1961]

[3.2.1.27 Deleted entry.  $\alpha$ -1,3-glucosidase]

[EC 3.2.1.27 created 1961, deleted 1972]

#### EC 3.2.1.28

**Accepted name:**  $\alpha,\alpha$ -trehalase  
**Reaction:**  $\alpha,\alpha$ -trehalose + H<sub>2</sub>O =  $\beta$ -D-glucose +  $\alpha$ -D-glucose  
**Other name(s):** trehalase  
**Systematic name:**  $\alpha,\alpha$ -trehalose glucohydrolase  
**Comments:** The enzyme is an anomer-inverting glucosidase that catalyses the hydrolysis of the  $\alpha$ -glucosidic *O*-linkage of  $\alpha,\alpha$ -trehalose, releasing initially equimolar amounts of  $\alpha$ - and  $\beta$ -D-glucose. It is widely distributed in microorganisms, plants, invertebrates and vertebrates.  
**References:** [2018, 1363, 1099, 1965]

[EC 3.2.1.28 created 1961, modified 2012]

[3.2.1.29 Deleted entry. chitobiase. Now included with EC 3.2.1.52,  $\beta$ -N-acetylhexosaminidase]

[EC 3.2.1.29 created 1961, deleted 1972]

[3.2.1.30 Deleted entry.  $\beta$ -D-acetylglucosaminidase. Now included with EC 3.2.1.52,  $\beta$ -N-acetylhexosaminidase]

[EC 3.2.1.30 created 1961, deleted 1992]

#### EC 3.2.1.31

**Accepted name:**  $\beta$ -glucuronidase



**Reaction:** a  $\beta$ -D-glucuronoside + H<sub>2</sub>O = D-glucuronate + an alcohol  
**Other name(s):**  $\beta$ -glucuronide glucuronohydrolase glucuronidase; exo- $\beta$ -D-glucuronidase; ketodase  
**Systematic name:**  $\beta$ -D-glucuronoside glucuronosohydrolase  
**References:** [612, 644, 777, 1657, 3085]

[EC 3.2.1.31 created 1961]

#### EC 3.2.1.32

**Accepted name:** *endo*-1,3- $\beta$ -xylanase  
**Reaction:** Random endohydrolysis of (1 $\rightarrow$ 3)- $\beta$ -D-glycosidic linkages in (1 $\rightarrow$ 3)- $\beta$ -D-xylans  
**Other name(s):** xylanase (ambiguous); *endo*-1,3- $\beta$ -xylosidase (misleading); 1,3- $\beta$ -xylanase; 1,3-xylanase;  $\beta$ -1,3-xylanase; *endo*- $\beta$ -1,3-xylanase; 1,3- $\beta$ -D-xylan xylanohydrolase; xylan *endo*-1,3- $\beta$ -xylosidase  
**Systematic name:** 3- $\beta$ -D-xylan xylanohydrolase  
**Comments:** This enzyme is found mostly in marine bacteria, which break down the  $\beta$ (1,3)-xylan found in the cell wall of some green and red algae. The enzyme produces mainly xylobiose, xylotriose and xylotetraose.  
**References:** [417, 61, 68, 66, 2186]

[EC 3.2.1.32 created 1965, modified 2011]

#### EC 3.2.1.33

**Accepted name:** amylo- $\alpha$ -1,6-glucosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 6)- $\alpha$ -D-glucosidic branch linkages in glycogen phosphorylase limit dextrin  
**Other name(s):** amylo-1,6-glucosidase; dextrin 6- $\alpha$ -D-glucosidase; amylopectin 1,6-glucosidase; dextrin-1,6-glucosidase; glycogen phosphorylase-limit dextrin  $\alpha$ -1,6-glucohydrolase  
**Systematic name:** glycogen phosphorylase-limit dextrin 6- $\alpha$ -glucohydrolase  
**Comments:** This enzyme hydrolyses an unsubstituted glucose unit linked by an  $\alpha$ (1 $\rightarrow$ 6) bond to an  $\alpha$ (1 $\rightarrow$ 4) glucose chain. The enzyme activity found in mammals and yeast is in a polypeptide chain containing two active centres. The other activity is similar to that of EC 2.4.1.25 (4- $\alpha$ -glucanotransferase), which acts on the glycogen phosphorylase limit dextrin chains to expose the single glucose residues, which the 6- $\alpha$ -glucosidase activity can then hydrolyse. Together, these two activities constitute the glycogen debranching system.  
**References:** [308, 1621, 2068]

[EC 3.2.1.33 created 1965, modified 2000]

[3.2.1.34 Deleted entry. *chondroitinase*. Now included with EC 3.2.1.35 *hyaluronoglucosaminidase*]

[EC 3.2.1.34 created 1965, deleted 1972]

#### EC 3.2.1.35

**Accepted name:** hyaluronoglucosaminidase  
**Reaction:** Random hydrolysis of (1 $\rightarrow$ 4)-linkages between *N*-acetyl- $\beta$ -D-glucosamine and D-glucuronate residues in hyaluronate  
**Other name(s):** hyaluronidase; hyaluronoglucosidase; chondroitinase; chondroitinase I  
**Systematic name:** hyaluronate 4-glycanohydrolase  
**Comments:** Also hydrolyses 1,4- $\beta$ -D-glycosidic linkages between *N*-acetyl-galactosamine or *N*-acetylgalactosamine sulfate and glucuronic acid in chondroitin, chondroitin 4- and 6-sulfates, and dermatan.  
**References:** [1896, 2375, 3155]

[EC 3.2.1.35 created 1965, modified 1976, modified 2001 (EC 3.2.1.34 created 1965, incorporated 1972)]

#### EC 3.2.1.36

**Accepted name:** hyaluronoglucuronidase  
**Reaction:** Random hydrolysis of (1→3)-linkages between β-D-glucuronate and *N*-acetyl-D-glucosamine residues in hyaluronate  
**Other name(s):** hyaluronidase; glucuronoglucosaminoglycan hyaluronate lyase; orgelase  
**Systematic name:** hyaluronate 3-glycanohydrolase  
**References:** [1707, 1896]

[EC 3.2.1.36 created 1965, modified 1980]

#### EC 3.2.1.37

**Accepted name:** xylan 1,4-β-xylosidase  
**Reaction:** Hydrolysis of (1→4)-β-D-xylans, to remove successive D-xylose residues from the non-reducing termini  
**Other name(s):** xylobiase; β-xylosidase; exo-1,4-β-xylosidase; β-D-xylopyranosidase; β-xylosidase; β-xylosidase; exo-1,4-xylosidase; exo-1,4-β-D-xylosidase; 1,4-β-D-xylan xylohydrolase  
**Systematic name:** 4-β-D-xylan xylohydrolase  
**Comments:** Also hydrolyses xylobiose. Some other exoglycosidase activities have been found associated with this enzyme in sheep liver.  
**References:** [435, 1192]

[EC 3.2.1.37 created 1965]

#### EC 3.2.1.38

**Accepted name:** β-D-fucosidase  
**Reaction:** Hydrolysis of terminal non-reducing β-D-fucose residues in β-D-fucosides  
**Other name(s):** β-fucosidase  
**Systematic name:** β-D-fucoside fucohydrolase  
**Comments:** Enzymes from some sources also hydrolyse β-D-galactosides and/or β-D-glucosides and/or α-L-arabinosides. The activity of EC 3.2.1.37 xylan 1,4-β-xylosidase, is an associated activity found in some sources (e.g. liver).  
**References:** [434, 435, 2450, 3182, 3183]

[EC 3.2.1.38 created 1965, deleted 1972, reinstated 1978]

#### EC 3.2.1.39

**Accepted name:** glucan endo-1,3-β-D-glucosidase  
**Reaction:** Hydrolysis of (1→3)-β-D-glucosidic linkages in (1→3)-β-D-glucans  
**Other name(s):** endo-1,3-β-glucanase; laminarinase; laminaranase; oligo-1,3-glucosidase; endo-1,3-β-glucanase; callase; β-1,3-glucanase; kitalase; 1,3-β-D-glucan 3-glucanohydrolase; endo-(1,3)-β-D-glucanase; (1→3)-β-glucan 3-glucanohydrolase; endo-1,3-β-D-glucanase; endo-1,3-β-glucosidase; 1,3-β-D-glucan glucanohydrolase  
**Systematic name:** 3-β-D-glucan glucanohydrolase  
**Comments:** Different from EC 3.2.1.6 endo-1,3(4)-β-glucanase. Very limited action on mixed-link (1→3,1→4)-β-D-glucans. Hydrolyses laminarin, paramylon and pachyman.  
**References:** [428, 2397]

[EC 3.2.1.39 created 1965]

#### EC 3.2.1.40

**Accepted name:** α-L-rhamnosidase  
**Reaction:** Hydrolysis of terminal non-reducing α-L-rhamnose residues in α-L-rhamnosides  
**Other name(s):** α-L-rhamnosidase T; α-L-rhamnosidase N  
**Systematic name:** α-L-rhamnoside rhamnhydrolase

**Comments:** The enzyme, found in animal tissues, plants, yeasts, fungi and bacteria, utilizes an inverting mechanism of hydrolysis, releasing  $\beta$ -L-rhamnose. Substrates include naringin, rutin, quercitrin, hesperidin, dioscin, terpenyl glycosides and many other natural glycosides containing terminal  $\alpha$ -L-rhamnose.

**References:** [2461, 1575, 3353, 3255, 514, 2346]

[EC 3.2.1.40 created 1972]

#### EC 3.2.1.41

**Accepted name:** pullulanase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 6)- $\alpha$ -D-glucosidic linkages in pullulan, amylopectin and glycogen, and in the  $\alpha$ - and  $\beta$ -limit dextrins of amylopectin and glycogen  
**Other name(s):** limit dextrinase (erroneous); amylopectin 6-glucanohydrolase; bacterial debranching enzyme; debranching enzyme;  $\alpha$ -dextrin endo-1,6- $\alpha$ -glucosidase; *R*-enzyme; pullulan  $\alpha$ -1,6-glucanohydrolase  
**Systematic name:** pullulan 6- $\alpha$ -glucanohydrolase  
**Comments:** Different from EC 3.2.1.142 (limit dextrinase) in its action on glycogen, and its rate of hydrolysis of limit dextrins. Its action on amylopectin is complete. Maltose is the smallest sugar that it can release from an  $\alpha$ -(1 $\rightarrow$ 6)-linkage.  
**References:** [1622, 192, 1801]

[EC 3.2.1.41 created 1972, modified 1976, modified 2000 (EC 3.2.1.69 created 1972, incorporated 1976)]

#### EC 3.2.1.42

**Accepted name:** GDP-glucosidase  
**Reaction:** GDP-glucose + H<sub>2</sub>O = D-glucose + GDP  
**Other name(s):** guanosine diphosphoglucosidase; guanosine diphosphate D-glucose glucohydrolase  
**Systematic name:** GDP-glucose glucohydrolase  
**References:** [2723]

[EC 3.2.1.42 created 1972]

#### EC 3.2.1.43

**Accepted name:**  $\beta$ -L-rhamnosidase  
**Reaction:** Hydrolysis of terminal, non-reducing  $\beta$ -L-rhamnose residues in  $\beta$ -L-rhamnosides  
**Systematic name:**  $\beta$ -L-rhamnoside rhamnohydrolase  
**References:** [142]

[EC 3.2.1.43 created 1972]

#### EC 3.2.1.44

**Accepted name:** fucoidanase  
**Reaction:** Endohydrolysis of (1 $\rightarrow$ 2)- $\alpha$ -L-fucoside linkages in fucoidan without release of sulfate  
**Other name(s):**  $\alpha$ -L-fucosidase; poly(1,2- $\alpha$ -L-fucoside-4-sulfate) glycanohydrolase  
**Systematic name:** poly[(1 $\rightarrow$ 2)- $\alpha$ -L-fucoside-4-sulfate] glycanohydrolase  
**References:** [2906]

[EC 3.2.1.44 created 1972]

#### EC 3.2.1.45

**Accepted name:** glucosylceramidase  
**Reaction:** a D-glucosyl-*N*-acylsphingosine + H<sub>2</sub>O = D-glucose + a ceramide  
**Other name(s):** psychosine hydrolase; glucosylsphingosine glucosylhydrolase; GlcCer- $\beta$ -glucosidase;  $\beta$ -D-glucocerebrosidase; glucosylcerebrosidase;  $\beta$ -glucosylceramidase; ceramide glucosidase; glucocerebrosidase; glucosylsphingosine  $\beta$ -glucosidase; glucosylsphingosine  $\beta$ -D-glucosidase

**Systematic name:** D-glucosyl-*N*-acylsphingosine glucohydrolase  
**Comments:** Also acts on glucosylsphingosine (*cf.* EC 3.2.1.62 glycosylceramidase).  
**References:** [275, 3025]

[EC 3.2.1.45 created 1972]

#### EC 3.2.1.46

**Accepted name:** galactosylceramidase  
**Reaction:** a D-galactosyl-*N*-acylsphingosine + H<sub>2</sub>O = D-galactose + a ceramide  
**Other name(s):** cerebroside galactosidase; galactocerebroside.β-galactosidase; galactosylcerebroside; galactocerebroside; ceramide galactosidase; galactocerebroside galactosidase; galactosylceramide.β-galactosidase; cerebroside β-galactosidase; galactosylceramidase I; β-galactosylceramidase; galactocerebroside-β-D-galactosidase; lactosylceramidase I; β-galactocerebroside; lactosylceramidase  
**Systematic name:** D-galactosyl-*N*-acylsphingosine galactohydrolase  
**Comments:** *cf.* EC 3.2.1.62 glycosylceramidase.  
**References:** [274]

[EC 3.2.1.46 created 1972]

#### EC 3.2.1.47

**Accepted name:** galactosylgalactosylglucosylceramidase  
**Reaction:** α-D-galactosyl-(1→4)-β-D-galactosyl-(1→4)-β-D-glucosyl-(1↔1)-ceramide + H<sub>2</sub>O = D-galactose + β-D-galactosyl-(1→4)-β-D-glucosyl-(1↔1)-ceramide  
**Other name(s):** trihexosyl ceramide galactosidase; ceramide trihexosidase; ceramidetrihexoside α-galactosidase; trihexosylceramide α-galactosidase; ceramidetrihexosidase  
**Systematic name:** α-D-galactosyl-(1→4)-β-D-galactosyl-(1→4)-β-D-glucosyl-(1↔1)-ceramide galactohydrolase  
**References:** [273, 1804]

[EC 3.2.1.47 created 1972, modified 2011]

#### EC 3.2.1.48

**Accepted name:** sucrose α-glucosidase  
**Reaction:** Hydrolysis of sucrose and maltose by an α-D-glucosidase-type action  
**Other name(s):** sucrose α-glucohydrolase; sucrase; sucrase-isomaltase; sucrose.α.-glucohydrolase; intestinal sucrase; sucrase(invertase)  
**Systematic name:** sucrose-α-D-glucohydrolase  
**Comments:** This enzyme is isolated from intestinal mucosa as a single polypeptide chain that also displays activity towards isomaltose (EC 3.2.1.10 oligo-1,6-glucosidase).  
**References:** [473, 1080, 1499, 2657, 2678, 2860]

[EC 3.2.1.48 created 1972]

#### EC 3.2.1.49

**Accepted name:** α-*N*-acetylgalactosaminidase  
**Reaction:** Cleavage of non-reducing α-(1→3)-*N*-acetylgalactosamine residues from human blood group A and AB mucin glycoproteins, Forssman hapten and blood group A lacto series glycolipids  
**Other name(s):** α-acetylgalactosaminidase; *N*-acetyl-α-D-galactosaminidase; *N*-acetyl-α-galactosaminidase; α-NAGAL; α-NAGA; α-GalNAcase  
**Systematic name:** α-*N*-acetyl-D-galactosaminide *N*-acetylgalactosaminohydrolase  
**Comments:** The human lysosomal enzyme is involved in the degradation of blood type A epitope.  
**References:** [84, 3339, 458, 1185, 1064, 3151, 87]

[EC 3.2.1.49 created 1972, modified 2011]

#### EC 3.2.1.50

**Accepted name:**  $\alpha$ -*N*-acetylglucosaminidase  
**Reaction:** Hydrolysis of terminal non-reducing *N*-acetyl-D-glucosamine residues in *N*-acetyl- $\alpha$ -D-glucosaminides  
**Other name(s):**  $\alpha$ -acetylglucosaminidase; *N*-acetyl- $\alpha$ -D-glucosaminidase; *N*-acetyl- $\alpha$ -glucosaminidase;  $\alpha$ -D-2-acetamido-2-deoxyglucosidase  
**Systematic name:**  $\alpha$ -*N*-acetyl-D-glucosaminide *N*-acetylglucosaminohydrolase  
**Comments:** Hydrolyses UDP-*N*-acetylglucosamine.  
**References:** [3078, 3079, 3157, 3163]

[EC 3.2.1.50 created 1972]

#### EC 3.2.1.51

**Accepted name:**  $\alpha$ -L-fucosidase  
**Reaction:** an  $\alpha$ -L-fucoside + H<sub>2</sub>O = L-fucose + an alcohol  
**Other name(s):**  $\alpha$ -fucosidase  
**Systematic name:**  $\alpha$ -L-fucoside fucohydrolase  
**References:** [1658, 2400, 2870]

[EC 3.2.1.51 created 1972]

#### EC 3.2.1.52

**Accepted name:**  $\beta$ -*N*-acetylhexosaminidase  
**Reaction:** Hydrolysis of terminal non-reducing *N*-acetyl-D-hexosamine residues in *N*-acetyl- $\beta$ -D-hexosaminides  
**Other name(s):** hexosaminidase;  $\beta$ -acetylaminodeoxyhexosidase; *N*-acetyl- $\beta$ -D-hexosaminidase; *N*-acetyl- $\beta$ -hexosaminidase;  $\beta$ -hexosaminidase;  $\beta$ -acetylhexosaminidase;  $\beta$ -D-*N*-acetylhexosaminidase;  $\beta$ -*N*-acetyl-D-hexosaminidase;  $\beta$ -*N*-acetylglucosaminidase; hexosaminidase A; *N*-acetylhexosaminidase;  $\beta$ -D-hexosaminidase  
**Systematic name:**  $\beta$ -*N*-acetyl-D-hexosaminide *N*-acetylhexosaminohydrolase  
**Comments:** Acts on *N*-acetylglucosides and *N*-acetylgalactosides.  
**References:** [346, 357, 823, 1667]

[EC 3.2.1.52 created 1972 (EC 3.2.1.30 created 1961, incorporated 1992 [EC 3.2.1.29 created 1961, incorporated 1972])]

#### EC 3.2.1.53

**Accepted name:**  $\beta$ -*N*-acetylgalactosaminidase  
**Reaction:** Hydrolysis of terminal non-reducing *N*-acetyl-D-galactosamine residues in *N*-acetyl- $\beta$ -D-galactosaminides  
**Other name(s):** *N*-acetyl- $\beta$ -galactosaminidase; *N*-acetyl- $\beta$ -D-galactosaminidase;  $\beta$ -acetylgalactosaminidase;  $\beta$ -D-*N*-acetylgalactosaminidase; *N*-acetylgalactosaminidase  
**Systematic name:**  $\beta$ -*N*-acetyl-D-galactosaminide *N*-acetylgalactosaminohydrolase  
**References:** [823, 1170]

[EC 3.2.1.53 created 1972]

#### EC 3.2.1.54

**Accepted name:** cyclomaltodextrinase  
**Reaction:** cyclomaltodextrin + H<sub>2</sub>O = linear maltodextrin  
**Other name(s):** cycloheptagluconase; cyclohexagluconase; cyclodextrinase; cyclomaltodextrin dextrin-hydrolase (de-cyclizing)

**Systematic name:** cyclomaltodextrin dextrin-hydrolase (ring-opening)  
**Comments:** Also hydrolyses linear maltodextrin.  
**References:** [585]

[EC 3.2.1.54 created 1972 (EC 3.2.1.12 and EC 3.2.1.13 both created 1961 and incorporated 1976)]

#### EC 3.2.1.55

**Accepted name:** non-reducing end  $\alpha$ -L-arabinofuranosidase  
**Reaction:** Hydrolysis of terminal non-reducing  $\alpha$ -L-arabinofuranoside residues in  $\alpha$ -L-arabinosides.  
**Other name(s):** arabinosidase (ambiguous);  $\alpha$ -arabinosidase;  $\alpha$ -L-arabinosidase;  $\alpha$ -arabinofuranosidase; polysaccharide  $\alpha$ -L-arabinofuranosidase;  $\alpha$ -L-arabinofuranoside hydrolase; L-arabinosidase (ambiguous);  $\alpha$ -L-arabinanase  
**Systematic name:**  $\alpha$ -L-arabinofuranoside non-reducing end  $\alpha$ -L-arabinofuranosidase  
**Comments:** The enzyme acts on  $\alpha$ -L-arabinofuranosides,  $\alpha$ -L-arabinans containing (1,3)- and/or (1,5)-linkages, arabinoxylans and arabinogalactans. Some  $\beta$ -galactosidases (EC 3.2.1.23) and  $\beta$ -D-fucosidases (EC 3.2.1.38) also hydrolyse  $\alpha$ -L-arabinosides. *cf.* EC 3.2.1.185, non-reducing end  $\beta$ -L-arabinofuranosidase.  
**References:** [2829, 1355, 1356, 1810, 1244]

[EC 3.2.1.55 created 1972, modified 1976 (EC 3.2.1.79 created 1972, incorporated 1976), modified 2013]

#### EC 3.2.1.56

**Accepted name:** glucuronosyl-disulfoglucosamine glucuronidase  
**Reaction:** 3-D-glucuronosyl- $N^2,6$ -disulfo- $\beta$ -D-glucosamine + H<sub>2</sub>O = D-glucuronate +  $N^2,6$ -disulfo-D-glucosamine  
**Other name(s):** glycuronidase; 3-D-glucuronosyl-2- $N,6$ -disulfo- $\beta$ -D-glucosamine glucuronohydrolase  
**Systematic name:** 3-D-glucuronosyl- $N^2,6$ -disulfo- $\beta$ -D-glucosamine glucuronohydrolase  
**References:** [608]

[EC 3.2.1.56 created 1972]

#### EC 3.2.1.57

**Accepted name:** isopullulanase  
**Reaction:** Hydrolysis of pullulan to isopanose (6- $\alpha$ -maltosylglucose)  
**Systematic name:** pullulan 4-gluconohydrolase (isopanose-forming)  
**Comments:** The enzyme has practically no action on starch. Panose (4- $\alpha$ -isomaltosylglucose) is hydrolysed to isomaltose and glucose. *cf.* EC 3.2.1.41 (pullulanase) and EC 3.2.1.135 (neopullulanase).  
**References:** [2504]

[EC 3.2.1.57 created 1972]

#### EC 3.2.1.58

**Accepted name:** glucan 1,3- $\beta$ -glucosidase  
**Reaction:** Successive hydrolysis of  $\beta$ -D-glucose units from the non-reducing ends of (1 $\rightarrow$ 3)- $\beta$ -D-glucans, releasing  $\alpha$ -glucose  
**Other name(s):** exo-1,3- $\beta$ -glucosidase;  $\beta$ -1,3-glucan exo-hydrolase; exo (1 $\rightarrow$ 3)-glucanohydrolase; 1,3- $\beta$ -glucan glucohydrolase  
**Systematic name:** 3- $\beta$ -D-glucan glucohydrolase  
**Comments:** Acts on oligosaccharides, but very slowly on laminaribiose.  
**References:** [152, 153]

[EC 3.2.1.58 created 1972]

### EC 3.2.1.59

**Accepted name:** glucan endo-1,3- $\alpha$ -glucosidase  
**Reaction:** Endohydrolysis of (1 $\rightarrow$ 3)- $\alpha$ -D-glucosidic linkages in isolichenin, pseudonigeran and nigeran  
**Other name(s):** endo-1,3- $\alpha$ -glucanase; mutanase; endo-(1 $\rightarrow$ 3)- $\alpha$ -glucanase; cariogenase; cariogenanase; endo-1,3- $\alpha$ -D-glucanase; 1,3(1,3;1,4)- $\alpha$ -D-glucan 3-glucanohydrolase  
**Systematic name:** 3- $\alpha$ -D-glucan 3-glucanohydrolase  
**Comments:** Products from pseudonigeran (1,3- $\alpha$ -D-glucan) are nigerose and  $\alpha$ -D-glucose.  
**References:** [1068]

[EC 3.2.1.59 created 1972]

### EC 3.2.1.60

**Accepted name:** glucan 1,4- $\alpha$ -maltotetraohydrolase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 4)- $\alpha$ -D-glucosidic linkages in amylaceous polysaccharides, to remove successive maltotetraose residues from the non-reducing chain ends  
**Other name(s):** exo-maltotetraohydrolase; 1,4- $\alpha$ -D-glucan maltotetraohydrolase  
**Systematic name:** 4- $\alpha$ -D-glucan maltotetraohydrolase  
**Comments:** Compare EC 3.2.1.2  $\beta$ -amylase, which removes successive maltose residues, and EC 3.2.1.98 (glucan 1,4- $\alpha$ -maltohexaosidase) and EC 3.2.1.116 (glucan 1,4- $\alpha$ -maltotriohydrolase).  
**References:** [2043, 2443]

[EC 3.2.1.60 created 1972]

### EC 3.2.1.61

**Accepted name:** mycodextranase  
**Reaction:** Endohydrolysis of (1 $\rightarrow$ 4)- $\alpha$ -D-glucosidic linkages in  $\alpha$ -D-glucans containing both (1 $\rightarrow$ 3)- and (1 $\rightarrow$ 4)-bonds  
**Other name(s):** 1,3-1,4- $\alpha$ -D-glucan 4-glucanohydrolase  
**Systematic name:** (1 $\rightarrow$ 3)-(1 $\rightarrow$ 4)- $\alpha$ -D-glucan 4-glucanohydrolase  
**Comments:** Products are nigerose and 4- $\alpha$ -D-nigerosylglucose. No hydrolysis of  $\alpha$ -D-glucans containing only 1,3- or 1,4-bonds.  
**References:** [2997]

[EC 3.2.1.61 created 1972]

### EC 3.2.1.62

**Accepted name:** glycosylceramidase  
**Reaction:** a glycosyl-*N*-acylsphingosine + H<sub>2</sub>O = a ceramide + a sugar  
**Other name(s):** phlorizin hydrolase; phloretin-glucosidase; glycosyl ceramide glycosylhydrolase; cerebrosidase; phloridzin  $\beta$ -glucosidase; lactase-phlorizin hydrolase; phloridzin glucosidase  
**Systematic name:** glycosyl-*N*-acylsphingosine glycohydrolase  
**Comments:** Broad specificity [*cf.* EC 3.2.1.45 (glycosylceramidase) and EC 3.2.1.46 (galactosylceramidase)]. Also hydrolyses phlorizin to phloretin and glucose. The intestinal enzyme is a complex that also catalyses the reaction of EC 3.2.1.108 lactase.  
**References:** [1639, 1736, 1786]

[EC 3.2.1.62 created 1972, modified 1976]

### EC 3.2.1.63

**Accepted name:** 1,2- $\alpha$ -L-fucosidase  
**Reaction:** methyl-2- $\alpha$ -L-fucopyranosyl- $\beta$ -D-galactoside + H<sub>2</sub>O = L-fucose + methyl  $\beta$ -D-galactoside  
**Other name(s):** almond emulsin fucosidase;  $\alpha$ -(1 $\rightarrow$ 2)-L-fucosidase  
**Systematic name:** 2- $\alpha$ -L-fucopyranosyl- $\beta$ -D-galactoside fucohydrolase

**Comments:** Highly specific for non-reducing terminal L-fucose residues linked to D-galactose residues by a 1,2- $\alpha$ -linkage. Not identical with EC 3.2.1.111 1,3- $\alpha$ -L-fucosidase.

**References:** [118, 2144, 2400]

[EC 3.2.1.63 created 1972]

#### EC 3.2.1.64

**Accepted name:** 2,6- $\beta$ -fructan 6-levanbiohydrolase

**Reaction:** Hydrolysis of (2 $\rightarrow$ 6)- $\beta$ -D-fructofuranan, to remove successive disaccharide residues as levanbiose, i.e. 6-( $\beta$ -D-fructofuranosyl)-D-fructose, from the end of the chain

**Other name(s):**  $\beta$ -2,6-fructan-6-levanbiohydrolase; 2,6- $\beta$ -D-fructan 6-levanbiohydrolase; levanbiose-producing lev-anase; 2,6- $\beta$ -D-fructan 6- $\beta$ -D-fructofuranosylfructohydrolase

**Systematic name:** (2 $\rightarrow$ 6)- $\beta$ -D-fructofuranan 6-( $\beta$ -D-fructosyl)-D-fructose-hydrolase

**References:** [105, 2498, 2499, 2715, 1385]

[EC 3.2.1.64 created 1972, modified 2004]

#### EC 3.2.1.65

**Accepted name:** levanase

**Reaction:** Random hydrolysis of (2 $\rightarrow$ 6)- $\beta$ -D-fructofuranosidic linkages in (2 $\rightarrow$ 6)- $\beta$ -D-fructans (levans) contain-ing more than 3 fructose units

**Other name(s):** levan hydrolase; 2,6- $\beta$ -D-fructan fructanohydrolase

**Systematic name:** (2 $\rightarrow$ 6)- $\beta$ -D-fructan fructanohydrolase

**References:** [104]

[EC 3.2.1.65 created 1972]

#### EC 3.2.1.66

**Accepted name:** quercitrinase

**Reaction:** quercitrin + H<sub>2</sub>O = L-rhamnose + quercetin

**Systematic name:** quercitrin 3-L-rhamnohydrolase

**Comments:** Quercitrin is quercetin 3-L-rhamnoside.

**References:** [3164]

[EC 3.2.1.66 created 1972]

#### EC 3.2.1.67

**Accepted name:** galacturan 1,4- $\alpha$ -galacturonidase

**Reaction:** [(1 $\rightarrow$ 4)- $\alpha$ -D-galacturonide]<sub>n</sub> + H<sub>2</sub>O = [(1 $\rightarrow$ 4)- $\alpha$ -D-galacturonide]<sub>n-1</sub> + D-galacturonate

**Other name(s):** exopolygalacturonase; poly(galacturonate) hydrolase; exo-D-galacturonase; exo-D-galacturonanase; exopoly-D-galacturonase; poly(1,4- $\alpha$ -D-galacturonide) galacturonohydrolase

**Systematic name:** poly[(1 $\rightarrow$ 4)- $\alpha$ -D-galacturonide] galacturonohydrolase

**References:** [1067]

[EC 3.2.1.67 created 1972]

#### EC 3.2.1.68

**Accepted name:** isoamylase

**Reaction:** Hydrolysis of (1 $\rightarrow$ 6)- $\alpha$ -D-glucosidic branch linkages in glycogen, amylopectin and their  $\beta$ -limit dex-trins

**Other name(s):** debranching enzyme; glycogen  $\alpha$ -1,6-glucanohydrolase

**Systematic name:** glycogen 6- $\alpha$ -D-glucanohydrolase



**Comments:** Also readily hydrolyses amylopectin. Differs from EC 3.2.1.41 (pullulanase) and EC 3.2.1.142 (limit dextrinase) by its inability to hydrolyse pullulan, and by limited action on  $\alpha$ -limit dextrans. Maltose is the smallest sugar it can release from an  $\alpha$ -(1 $\rightarrow$ 6)-linkage.

**References:** [3286]

[EC 3.2.1.68 created 1972, modified 1976, modified 2000]

[3.2.1.69 Deleted entry. amylopectin 6-glucanohydrolase. Now included with EC 3.2.1.41 pullulanase]

[EC 3.2.1.69 created 1972, deleted 1976]

#### EC 3.2.1.70

**Accepted name:** glucan 1,6- $\alpha$ -glucosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 6)- $\alpha$ -D-glucosidic linkages in (1 $\rightarrow$ 6)- $\alpha$ -D-glucans and derived oligosaccharides  
**Other name(s):** exo-1,6- $\beta$ -glucosidase; glucodextrinase; glucan  $\alpha$ -1,6-D-glucohydrolase  
**Systematic name:** glucan 6- $\alpha$ -D-glucohydrolase  
**Comments:** Hydrolysis is accompanied by inversion at C-1, so that new reducing ends are released in the  $\beta$ -configuration. Dextrans and isomaltosaccharides are hydrolysed, as is isomaltose, but very slowly. The enzyme from some sources also possesses the activity of EC 3.2.1.59 (glucan endo-1,3- $\alpha$ -glucosidase).  
**References:** [2176, 2547, 3095]

[EC 3.2.1.70 created 1972, modified 2001]

#### EC 3.2.1.71

**Accepted name:** glucan endo-1,2- $\beta$ -glucosidase  
**Reaction:** Random hydrolysis of (1 $\rightarrow$ 2)-glucosidic linkages in (1 $\rightarrow$ 2)- $\beta$ -D-glucans  
**Other name(s):** endo-1,2- $\beta$ -glucanase;  $\beta$ -D-1,2-glucanase; endo-(1 $\rightarrow$ 2)- $\beta$ -D-glucanase; 1,2- $\beta$ -D-glucan glucanohydrolase  
**Systematic name:** 2- $\beta$ -D-glucan glucanohydrolase  
**References:** [2398]

[EC 3.2.1.71 created 1972]

#### EC 3.2.1.72

**Accepted name:** xylan 1,3- $\beta$ -xylosidase  
**Reaction:** Hydrolysis of successive xylose residues from the non-reducing termini of (1 $\rightarrow$ 3)- $\beta$ -D-xylans  
**Other name(s):** 1,3- $\beta$ -D-xylosidase, exo-1,3- $\beta$ -xylosidase;  $\beta$ -1,3'-xylanase; exo- $\beta$ -1,3'-xylanase; 1,3- $\beta$ -D-xylan xylohydrolase  
**Systematic name:** 3- $\beta$ -D-xylan xylohydrolase  
**References:** [859]

[EC 3.2.1.72 created 1972]

#### EC 3.2.1.73

**Accepted name:** licheninase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-glucosidic linkages in  $\beta$ -D-glucans containing (1 $\rightarrow$ 3)- and (1 $\rightarrow$ 4)-bonds  
**Other name(s):** lichenase;  $\beta$ -(1 $\rightarrow$ 4)-D-glucan 4-glucanohydrolase; 1,3;1,4- $\beta$ -glucan endohydrolase; 1,3;1,4- $\beta$ -glucan 4-glucanohydrolase; 1,3-1,4- $\beta$ -D-glucan 4-glucanohydrolase  
**Systematic name:** (1 $\rightarrow$ 3)-(1 $\rightarrow$ 4)- $\beta$ -D-glucan 4-glucanohydrolase  
**Comments:** Acts on lichenin and cereal  $\beta$ -D-glucans, but not on  $\beta$ -D-glucans containing only 1,3- or 1,4-bonds.  
**References:** [151]

[EC 3.2.1.73 created 1972]

#### EC 3.2.1.74

**Accepted name:** glucan 1,4- $\beta$ -glucosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 4)-linkages in (1 $\rightarrow$ 4)- $\beta$ -D-glucans, to remove successive glucose units  
**Other name(s):** exo-1,4- $\beta$ -glucosidase; exocellulase; exo- $\beta$ -1,4-glucosidase; exo- $\beta$ -1,4-glucanase;  $\beta$ -1,4- $\beta$ -glucanase;  $\beta$ -glucosidase; exo-1,4- $\beta$ -glucanase; 1,4- $\beta$ -D-glucan glucohydrolase  
**Systematic name:** 4- $\beta$ -D-glucan glucohydrolase  
**Comments:** Acts on 1,4- $\beta$ -D-glucans and related oligosaccharides. Cellobiose is hydrolysed, but very slowly.  
**References:** [151]

[EC 3.2.1.74 created 1972]

#### EC 3.2.1.75

**Accepted name:** glucan endo-1,6- $\beta$ -glucosidase  
**Reaction:** Random hydrolysis of (1 $\rightarrow$ 6)-linkages in (1 $\rightarrow$ 6)- $\beta$ -D-glucans  
**Other name(s):** endo-1,6- $\beta$ -glucanase;  $\beta$ -1 $\rightarrow$ 6)- $\beta$ -D-glucanase;  $\beta$ -1,6-glucanase-pustulanase;  $\beta$ -1,6-glucan hydrolase;  $\beta$ -1,6-glucan 6-glucanohydrolase; 1,6- $\beta$ -D-glucan glucanohydrolase  
**Systematic name:** 6- $\beta$ -D-glucan glucanohydrolase  
**Comments:** Acts on lutean, pustulan and 1,6-oligo- $\beta$ -D-glucosides.  
**References:** [2399]

[EC 3.2.1.75 created 1972]

#### EC 3.2.1.76

**Accepted name:** L-iduronidase  
**Reaction:** Hydrolysis of unsulfated  $\alpha$ -L-iduronosidic linkages in dermatan sulfate  
**Other name(s):**  $\alpha$ -L-iduronidase  
**Systematic name:** glycosaminoglycan  $\alpha$ -L-iduronohydrolase  
**References:** [1834, 2454, 2750]

[EC 3.2.1.76 created 1972]

#### EC 3.2.1.77

**Accepted name:** mannan 1,2-(1,3)- $\alpha$ -mannosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 2)- and (1 $\rightarrow$ 3)-linkages in yeast mannan, releasing mannose  
**Other name(s):** exo-1,2-1,3- $\alpha$ -mannosidase; 1,2-1,3- $\alpha$ -D-mannan mannohydrolase  
**Systematic name:** (1 $\rightarrow$ 2)-(1 $\rightarrow$ 3)- $\alpha$ -D-mannan mannohydrolase  
**Comments:** A 1,6- $\alpha$ -D-mannan backbone remains after action on yeast mannan. This is further attacked, but slowly.  
**References:** [1336, 1337]

[EC 3.2.1.77 created 1972]

#### EC 3.2.1.78

**Accepted name:** mannan endo-1,4- $\beta$ -mannosidase  
**Reaction:** Random hydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-mannosidic linkages in mannans, galactomannans and glucomannans  
**Other name(s):** endo-1,4- $\beta$ -mannanase; endo- $\beta$ -1,4-mannase;  $\beta$ -mannanase B;  $\beta$ -1, 4-mannan 4-mannanohydrolase; endo- $\beta$ -mannanase;  $\beta$ -D-mannanase; 1,4- $\beta$ -D-mannan mannanohydrolase  
**Systematic name:** 4- $\beta$ -D-mannan mannanohydrolase  
**References:** [712, 2396]

[EC 3.2.1.78 created 1972]

[3.2.1.79 Deleted entry.  $\alpha$ -L-arabinofuranoside hydrolase. Now included with EC 3.2.1.55  $\alpha$ -N-arabinofuranosidase]

[EC 3.2.1.79 created 1972, deleted 1976]

#### EC 3.2.1.80

**Accepted name:** fructan  $\beta$ -fructosidase  
**Reaction:** Hydrolysis of terminal, non-reducing (2 $\rightarrow$ 1)- and (2 $\rightarrow$ 6)-linked  $\beta$ -D-fructofuranose residues in fructans  
**Other name(s):** exo- $\beta$ -D-fructosidase; exo- $\beta$ -fructosidase; polysaccharide  $\beta$ -fructofuranosidase; fructan exohydrolase  
**Systematic name:**  $\beta$ -D-fructan fructohydrolase  
**Comments:** Hydrolyses inulin and levan, and also sucrose.  
**References:** [526, 1292]

[EC 3.2.1.80 created 1972]

#### EC 3.2.1.81

**Accepted name:**  $\beta$ -agarase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-galactosidic linkages in agarose, giving the tetramer as the predominant product  
**Other name(s):** agarase (ambiguous); AgaA; AgaB; endo- $\beta$ -agarase; agarose 3-glycanohydrolase (incorrect)  
**Systematic name:** agarose 4-glycanohydrolase  
**Comments:** Also acts on porphyran, but more slowly [659]. This enzyme cleaves the  $\beta$ -(1 $\rightarrow$ 4) linkages of agarose in a random manner with retention of the anomeric-bond configuration, producing  $\beta$ -anomers that give rise progressively to  $\alpha$ -anomers when mutarotation takes place [1303]. The end products of hydrolysis are neoagarotetraose and neoagarohexaose in the case of AgaA from the marine bacterium *Zobellia galactanivorans*, and neoagarotetraose and neoagarobiose in the case of AgaB [1303].  
**References:** [659, 37, 2172, 2171, 2792, 1303]

[EC 3.2.1.81 created 1972, modified 2006]

#### EC 3.2.1.82

**Accepted name:** exo-poly- $\alpha$ -galacturonosidase  
**Reaction:** Hydrolysis of pectic acid from the non-reducing end, releasing digalacturonate  
**Other name(s):** exopolygalacturonosidase; exopolygalacturanosidase; poly(1,4- $\alpha$ -D-galactosiduronate) digalacturonohydrolase  
**Systematic name:** poly[(1 $\rightarrow$ 4)- $\alpha$ -D-galactosiduronate] digalacturonohydrolase  
**References:** [1067, 1075, 1076]

[EC 3.2.1.82 created 1972]

#### EC 3.2.1.83

**Accepted name:**  $\kappa$ -carrageenase  
**Reaction:** Endohydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-linkages between D-galactose 4-sulfate and 3,6-anhydro-D-galactose in  $\kappa$ -carrageenans  
**Other name(s):**  $\kappa$ -carrageenan 4- $\beta$ -D-glycanohydrolase  
**Systematic name:**  $\kappa$ -carrageenan 4- $\beta$ -D-glycanohydrolase (configuration-retaining)  
**Comments:** The main products of hydrolysis are neocarrabiose-sulfate and neocarratetraose-sulfate [1903]. Unlike EC 3.2.1.157 (1-carrageenase), but similar to EC 3.2.1.81 ( $\beta$ -agarase), this enzyme proceeds with retention of the anomeric configuration.  
**References:** [3150, 2318, 2316, 1902, 1903]

[EC 3.2.1.83 created 1972, modified 2006]

#### EC 3.2.1.84

**Accepted name:** glucan 1,3- $\alpha$ -glucosidase  
**Reaction:** Hydrolysis of terminal (1 $\rightarrow$ 3)- $\alpha$ -D-glucosidic links in (1 $\rightarrow$ 3)- $\alpha$ -D-glucans  
**Other name(s):** exo-1,3- $\alpha$ -glucanase; glucosidase II; 1,3- $\alpha$ -D-glucan 3-glucohydrolase  
**Systematic name:** 3- $\alpha$ -D-glucan 3-glucohydrolase  
**Comments:** Does not act on nigeran.  
**References:** [3348]

[EC 3.2.1.84 created 1972]

#### EC 3.2.1.85

**Accepted name:** 6-phospho- $\beta$ -galactosidase  
**Reaction:** a 6-phospho- $\beta$ -D-galactoside + H<sub>2</sub>O = 6-phospho-D-galactose + an alcohol  
**Other name(s):** phospho- $\beta$ -galactosidase;  $\beta$ -D-phosphogalactoside galactohydrolase; phospho- $\beta$ -D-galactosidase; 6-phospho- $\beta$ -D-galactosidase  
**Systematic name:** 6-phospho- $\beta$ -D-galactoside 6-phosphogalactohydrolase  
**References:** [1111]

[EC 3.2.1.85 created 1976]

#### EC 3.2.1.86

**Accepted name:** 6-phospho- $\beta$ -glucosidase  
**Reaction:** 6-phospho- $\beta$ -D-glucosyl-(1 $\rightarrow$ 4)-D-glucose + H<sub>2</sub>O = D-glucose + D-glucose 6-phosphate  
**Other name(s):** phospho- $\beta$ -glucosidase A; phospho- $\beta$ -glucosidase; phosphocellobiase; 6-phospho- $\beta$ -D-glucosyl-(1,4)-D-glucose glucohydrolase  
**Systematic name:** 6-phospho- $\beta$ -D-glucosyl-(1 $\rightarrow$ 4)-D-glucose glucohydrolase  
**Comments:** Also hydrolyses several other phospho- $\beta$ -D-glucosides, but not their non-phosphorylated forms.  
**References:** [2230]

[EC 3.2.1.86 created 1976]

#### EC 3.2.1.87

**Accepted name:** capsular-polysaccharide endo-1,3- $\alpha$ -galactosidase  
**Reaction:** Random hydrolysis of (1 $\rightarrow$ 3)- $\alpha$ -D-galactosidic linkages in *Aerobacter aerogenes* capsular polysaccharide  
**Other name(s):** polysaccharide depolymerase; capsular polysaccharide galactohydrolase  
**Systematic name:** Aerobacter-capsular-polysaccharide galactohydrolase  
**Comments:** Hydrolyses the galactosyl- $\alpha$ -1,3-D-galactose linkages only in the complex substrate, bringing about depolymerization.  
**References:** [3313, 3314]

[EC 3.2.1.87 created 1976]

#### EC 3.2.1.88

**Accepted name:** non-reducing end  $\beta$ -L-arabinopyranosidase  
**Reaction:** Removal of a terminal  $\beta$ -L-arabinopyranose residue from the non-reducing end of its substrate.  
**Other name(s):** vicianosidase;  $\beta$ -L-arabinosidase (ambiguous);  $\beta$ -L-arabinoside arabinohydrolase (ambiguous)  
**Systematic name:**  $\beta$ -L-arabinopyranoside non-reducing end  $\beta$ -L-arabinopyranosidase  
**Comments:** The enzyme, which was characterized from dormant seeds of the plant *Cajanus cajan* (pigeon pea), has been shown to remove the terminal non-reducing  $\beta$ -L-arabinopyranoside residue from the artificial substrate *p*-nitrophenyl- $\beta$ -L-arabinopyranose [599]. In the presence of methanol the enzyme demonstrates transglycosylase activity, transferring the arabinose moiety to methanol while retaining the anomeric configuration, generating 1-*O*-methyl- $\beta$ -L-arabinopyranose [598].  
**References:** [599, 598]

[EC 3.2.1.88 created 1976, modified 2013]

#### EC 3.2.1.89

**Accepted name:** arabinogalactan endo- $\beta$ -1,4-galactanase  
**Reaction:** The enzyme specifically hydrolyses (1 $\rightarrow$ 4)- $\beta$ -D-galactosidic linkages in type I arabinogalactans.  
**Other name(s):** endo-1,4- $\beta$ -galactanase; galactanase (ambiguous); arabinogalactanase; *ganB* (gene name)  
**Systematic name:** arabinogalactan 4- $\beta$ -D-galactanohydrolase  
**Comments:** This enzyme, isolated from the bacterium *Bacillus subtilis*, hydrolyses the  $\beta$ (1 $\rightarrow$ 4) bonds found in type I plant arabinogalactans, which are a component of the primary cell walls of dicots. The predominant product is a tetrasaccharide. *cf.* EC 3.2.1.181, galactan endo- $\beta$ -1,3-galactanase.  
**References:** [699, 1583, 2642]

[EC 3.2.1.89 created 1976, modified 2012]

[3.2.1.90 Deleted entry. arabinogalactan endo-1,3- $\beta$ -galactosidase. The enzyme was not sufficiently characterized to warrant an EC number]

[EC 3.2.1.90 created 1976, deleted 2001]

#### EC 3.2.1.91

**Accepted name:** cellulose 1,4- $\beta$ -cellobiosidase (non-reducing end)  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-glucosidic linkages in cellulose and cellotetraose, releasing cellobiose from the non-reducing ends of the chains  
**Other name(s):** exo-cellobiohydrolase;  $\beta$ -1,4-glucan cellobiohydrolase;  $\beta$ -1,4-glucan cellobiosylhydrolase; 1,4- $\beta$ -glucan cellobiosidase; exoglucanase; avicelase; CBH 1; C<sub>1</sub> cellulase; cellobiohydrolase I; cellobiohydrolase; exo- $\beta$ -1,4-glucan cellobiohydrolase; 1,4- $\beta$ -D-glucan cellobiohydrolase; cellobiosidase  
**Systematic name:** 4- $\beta$ -D-glucan cellobiohydrolase (non-reducing end)  
**References:** [195, 713, 1030]

[EC 3.2.1.91 created 1976, modified 2011]

#### EC 3.2.1.92

**Accepted name:** peptidoglycan  $\beta$ -*N*-acetylmuramidase  
**Reaction:** Hydrolysis of terminal, non-reducing *N*-acetylmuramic residues  
**Other name(s):** exo- $\beta$ -*N*-acetylmuramidase; exo- $\beta$ -acetylmuramidase;  $\beta$ -2-acetamido-3-*O*-(*D*-1-carboxyethyl)-2-deoxy-*D*-glucoside acetamidodeoxyglucohydrolase  
**Systematic name:** peptidoglycan  $\beta$ -*N*-acetylmuramoyloxohydrolase  
**References:** [2428]

[EC 3.2.1.92 created 1976]

#### EC 3.2.1.93

**Accepted name:**  $\alpha,\alpha$ -phosphotrehalase  
**Reaction:**  $\alpha,\alpha$ -trehalose 6-phosphate + H<sub>2</sub>O = *D*-glucose + *D*-glucose 6-phosphate  
**Other name(s):** phosphotrehalase  
**Systematic name:**  $\alpha,\alpha$ -trehalose-6-phosphate phosphoglucohydrolase  
**References:** [212]

[EC 3.2.1.93 created 1976]

#### EC 3.2.1.94

**Accepted name:** glucan 1,6- $\alpha$ -isomaltosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 6)- $\alpha$ -*D*-glucosidic linkages in polysaccharides, to remove successive isomaltose units from the non-reducing ends of the chains

**Other name(s):** exo-isomaltohydrolase; isomalto-dextranase; isomaltodextranase; G2-dextranase; 1,6- $\alpha$ -D-glucan isomaltohydrolase  
**Systematic name:** 6- $\alpha$ -D-glucan isomaltohydrolase  
**Comments:** Optimum activity is on those 1,6- $\alpha$ -D-glucans containing 6, 7 and 8 glucose units; those containing 3, 4 and 5 glucose units are hydrolysed at slower rates.  
**References:** [2546, 2545]

[EC 3.2.1.94 created 1976]

#### EC 3.2.1.95

**Accepted name:** dextran 1,6- $\alpha$ -isomaltotriosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 6)- $\alpha$ -D-glucosidic linkages in dextrans, to remove successive isomaltotriose units from the non-reducing ends of the chains  
**Other name(s):** exo-isomaltotriohydrolase; 1,6- $\alpha$ -D-glucan isomaltotriohydrolase  
**Systematic name:** 6- $\alpha$ -D-glucan isomaltotriohydrolase  
**References:** [2795]

[EC 3.2.1.95 created 1978]

#### EC 3.2.1.96

**Accepted name:** mannosyl-glycoprotein endo- $\beta$ -*N*-acetylglucosaminidase  
**Reaction:** Endohydrolysis of the *N,N'*-diacetylchitobiosyl unit in high-mannose glycopeptides and glycoproteins containing the -[Man(GlcNAc)<sub>2</sub>Asn]- structure. One *N*-acetyl-D-glucosamine residue remains attached to the protein; the rest of the oligosaccharide is released intact  
**Other name(s):** *N,N'*-diacetylchitobiosyl  $\beta$ -*N*-acetylglucosaminidase; endo- $\beta$ -*N*-acetylglucosaminidase; mannosyl-glycoprotein endo- $\beta$ -*N*-acetylglucosaminidase; di-*N*-acetylchitobiosyl  $\beta$ -*N*-acetylglucosaminidase; endo- $\beta$ -acetylglucosaminidase; endo- $\beta$ -(1 $\rightarrow$ 4)-*N*-acetylglucosaminidase; mannosyl-glycoprotein 1,4-*N*-acetamidodeoxy- $\beta$ -D-glycohydrolase; endoglycosidase S; endo-*N*-acetyl- $\beta$ -D-glucosaminidase; endo-*N*-acetyl- $\beta$ -glucosaminidase; endo- $\beta$ -*N*-acetylglucosaminidase D; endo- $\beta$ -*N*-acetylglucosaminidase F; endo- $\beta$ -*N*-acetylglucosaminidase H; endo- $\beta$ -*N*-acetylglucosaminidase L; glycopeptide-D-mannosyl-4-*N*-(*N*-acetyl-D-glucosaminyloxy)-asparagine 1,4-*N*-acetyl- $\beta$ -glucosaminohydrolase; endoglycosidase H  
**Systematic name:** glycopeptide-D-mannosyl-*N*<sup>4</sup>-(*N*-acetyl-D-glucosaminyloxy)-asparagine 1,4-*N*-acetyl- $\beta$ -glucosaminohydrolase  
**Comments:** A group of related enzymes.  
**References:** [431, 1498, 2277, 2278, 2831, 2888]

[EC 3.2.1.96 created 1978]

#### EC 3.2.1.97

**Accepted name:** endo- $\alpha$ -*N*-acetylgalactosaminidase  
**Reaction:**  $\beta$ -D-galactosyl-(1 $\rightarrow$ 3)-*N*-acetyl- $\alpha$ -D-galactosaminyloxy-[glycoprotein]-L-serine/L-threonine + H<sub>2</sub>O =  $\beta$ -D-galactosyl-(1 $\rightarrow$ 3)-*N*-acetyl-D-galactosamine + [glycoprotein]-L-serine/L-threonine  
**Other name(s):** endo- $\alpha$ -acetylgalactosaminidase; endo- $\alpha$ -*N*-acetyl-D-galactosaminidase; mucinaminyserine mucinaminidase; D-galactosyl-3-(*N*-acetyl- $\alpha$ -D-galactosaminyloxy)-L-serine mucinaminohydrolase; endo- $\alpha$ -GalNAc-ase; glycopeptide  $\alpha$ -*N*-acetylgalactosaminidase; D-galactosyl-*N*-acetyl- $\alpha$ -D-galactosamine D-galactosyl-*N*-acetyl-galactosaminohydrolase  
**Systematic name:** glycopeptide-D-galactosyl-*N*-acetyl- $\alpha$ -D-galactosamine D-galactosyl-*N*-acetyl-galactosaminohydrolase

**Comments:** The enzyme catalyses the liberation of Gal-(1→3)-β-GalNAc α-linked to serine or threonine residues of mucin-type glycoproteins. EngBF from the bacterium *Bifidobacterium longum* specifically acts on core 1-type *O*-glycan to release the disaccharide Gal-(1→3)-β-GalNAc. The enzymes from the bacteria *Clostridium perfringens*, *Enterococcus faecalis*, *Propionibacterium acnes* and *Alcaligenes faecalis* show broader specificity (e.g. they can also release the core 2 trisaccharide Gal-(1→3)-β-(GlcNAc-(1→6)-β)-GalNAc or the core 3 disaccharide GlcNAc-(1→3)-β-GalNAc) [85, 1523]. The enzyme may play an important role in the degradation and utilization of mucins having core 1 *O*-glycan.

**References:** [85, 1523, 850, 2814, 976, 88, 936]

[EC 3.2.1.97 created 1978 (EC 3.2.1.110 created 1984, incorporated 2008), modified 2008, modified 2011]

#### EC 3.2.1.98

**Accepted name:** glucan 1,4-α-maltohexaosidase  
**Reaction:** Hydrolysis of (1→4)-α-D-glucosidic linkages in amylaceous polysaccharides, to remove successive maltohexaose residues from the non-reducing chain ends  
**Other name(s):** exo-maltohexaohydrolase; 1,4-α-D-glucan maltohexaohydrolase  
**Systematic name:** 4-α-D-glucan maltohexaohydrolase  
**Comments:** cf. EC 3.2.1.3 glucan 1,4-α-glucosidase, which removes successive glucose residues; EC 3.2.1.2 β-amylase, which removes successive maltose residues; EC 3.2.1.116 glucan 1,4-α-maltotriohydrolase, which removes successive maltotriose units and EC 3.2.1.60 glucan 1,4-α-maltotetrahydrolase, which removes successive maltotetraose residues. The products have the α-configuration.  
**References:** [1353, 2043]

[EC 3.2.1.98 created 1978]

#### EC 3.2.1.99

**Accepted name:** arabinan endo-1,5-α-L-arabinanase  
**Reaction:** Endohydrolysis of (1→5)-α-arabinofuranosidic linkages in (1→5)-arabinans  
**Other name(s):** endo-1,5-α-L-arabinanase; endo-α-1,5-arabanase; endo-arabanase; 1,5-α-L-arabinan 1,5-α-L-arabinanohydrolase; arabinan endo-1,5-α-L-arabinosidase (misleading)  
**Systematic name:** 5-α-L-arabinan 5-α-L-arabinanohydrolase  
**Comments:** Acts best on linear 1,5-α-L-arabinan. Also acts on branched arabinan, but more slowly.  
**References:** [1354, 3154, 785, 1612]

[EC 3.2.1.99 created 1981, modified 2011]

#### EC 3.2.1.100

**Accepted name:** mannan 1,4-mannobiosidase  
**Reaction:** Hydrolysis of (1→4)-β-D-mannosidic linkages in (1→4)-β-D-mannans, to remove successive manno-biose residues from the non-reducing chain ends  
**Other name(s):** 1,4-β-D-mannan mannobiohydrolase; exo-β-mannanase; exo-1,4-β-mannobiohydrolase  
**Systematic name:** 4-β-D-mannan mannobiohydrolase  
**References:** [67]

[EC 3.2.1.100 created 1983]

#### EC 3.2.1.101

**Accepted name:** mannan endo-1,6-α-mannosidase  
**Reaction:** Random hydrolysis of (1→6)-α-D-mannosidic linkages in unbranched (1→6)-mannans  
**Other name(s):** endo-α-1→6-D-mannanase; endo-1,6-β-mannanase; mannan endo-1,6-β-mannosidase; 1,6-α-D-mannan mannanohydrolase  
**Systematic name:** 6-α-D-mannan mannanohydrolase  
**References:** [2041, 289, 2040]

[EC 3.2.1.101 created 1984, modified 2001]

#### EC 3.2.1.102

**Accepted name:** blood-group-substance endo-1,4- $\beta$ -galactosidase  
**Reaction:** Endohydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-galactosidic linkages in blood group A and B substances  
**Other name(s):** endo- $\beta$ -galactosidase (ambiguous); blood-group-substance 1,4- $\beta$ -D-galactanohydrolase  
**Systematic name:** blood-group-substance 4- $\beta$ -D-galactanohydrolase  
**Comments:** Hydrolyses the 1,4- $\beta$ -D-galactosyl linkages adjacent to a 1,3- $\alpha$ -D-galactosyl or *N*-acetylgalactosaminy l residues and a 1,2- $\alpha$ -D-fucosyl residue.  
**References:** [858, 2054, 2852]

[EC 3.2.1.102 created 1984]

#### EC 3.2.1.103

**Accepted name:** keratan-sulfate endo-1,4- $\beta$ -galactosidase  
**Reaction:** Endohydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-galactosidic linkages in keratan sulfate  
**Other name(s):** endo- $\beta$ -galactosidase (ambiguous); keratan sulfate endogalactosidase; keratanase; keratan-sulfate 1,4- $\beta$ -D-galactanohydrolase  
**Systematic name:** keratan-sulfate 4- $\beta$ -D-galactanohydrolase  
**Comments:** Hydrolyses the 1,4- $\beta$ -D-galactosyl linkages adjacent to 1,3-*N*-acetyl- $\alpha$ -D-glucosaminy l residues. Also acts on some non-sulfated oligosaccharides, but only acts on blood group substances when the 1,2-linked fucosyl residues have been removed (*cf.* EC 3.2.1.102 blood-group-substance endo-1,4- $\beta$ -galactosidase).  
**References:** [858]

[EC 3.2.1.103 created 1984]

#### EC 3.2.1.104

**Accepted name:** steryl- $\beta$ -glucosidase  
**Reaction:** cholesteryl- $\beta$ -D-glucoside + H<sub>2</sub>O = D-glucose + cholesterol  
**Systematic name:** cholesteryl- $\beta$ -D-glucoside glucohydrolase  
**Comments:** Acts on glucosides of cholesterol and sitosterol, but not on some related sterols such as coprostanol.  
**References:** [1364]

[EC 3.2.1.104 created 1984]

#### EC 3.2.1.105

**Accepted name:** 3 $\alpha$ (*S*)-strictosidine  $\beta$ -glucosidase  
**Reaction:** strictosidine + H<sub>2</sub>O = D-glucose + strictosidine aglycone  
**Systematic name:** strictosidine  $\beta$ -D-glucohydrolase  
**Comments:** Does not act on a number of closely related glycosides. Strictosidine is a precursor of indole alkaloids.  
**References:** [1109, 144]

[EC 3.2.1.105 created 1984]

#### EC 3.2.1.106

**Accepted name:** mannosyl-oligosaccharide glucosidase  
**Reaction:** Glc<sub>3</sub>Man<sub>9</sub>GlcNAc<sub>2</sub>-[protein] + H<sub>2</sub>O = Glc<sub>2</sub>Man<sub>9</sub>GlcNAc<sub>2</sub>-[protein] +  $\beta$ -D-glucopyranose  
**Other name(s):** Glc<sub>3</sub>Man<sub>9</sub>NAc<sub>2</sub> oligosaccharide glucosidase; trimming glucosidase I; CWH41 (gene name); MOGS (gene name); mannosyl-oligosaccharide glucohydrolase  
**Systematic name:** Glc<sub>3</sub>Man<sub>9</sub>GlcNAc<sub>2</sub>-[protein] glucohydrolase (configuration-inverting)



**Comments:** This enzyme catalyses the first step in the processing of the *N*-glycan tetradecasaccharide precursor Glc<sub>3</sub>Man<sub>9</sub>GlcNAc<sub>2</sub>, which takes place in the endoplasmic reticulum, by removing the distal  $\alpha$ -1,2-linked glucose residue. This and subsequent processing steps are required before complex *N*-glycans can be synthesized.

**References:** [696, 981, 1441, 982, 1812]

[EC 3.2.1.106 created 1984, modified 2018]

#### EC 3.2.1.107

**Accepted name:** protein-glucosylgalactosylhydroxylysine glucosidase  
**Reaction:** [collagen]-(5*R*)-5-*O*-[ $\alpha$ -D-glucosyl-(1 $\rightarrow$ 2)- $\beta$ -D-galactosyl]-5-hydroxy-L-lysine + H<sub>2</sub>O = D-glucose + [collagen]-(5*R*)-5-*O*-( $\beta$ -D-galactosyl)-5-hydroxy-L-lysine  
**Other name(s):** PGGHG (gene name); 2-*O*- $\alpha$ -D-glucopyranosyl-5-*O*- $\alpha$ -D-galactopyranosylhydroxy-L-lysine glucohydrolase; protein- $\alpha$ -D-glucosyl-1,2- $\beta$ -D-galactosyl-L-hydroxylysine glucohydrolase; protein- $\alpha$ -D-glucosyl-(1 $\rightarrow$ 2)- $\beta$ -D-galactosyl-L-hydroxylysine glucohydrolase  
**Systematic name:** [collagen]-(5*R*)-5-*O*-[ $\alpha$ -D-glucosyl-(1 $\rightarrow$ 2)- $\beta$ -D-galactosyl]-5-hydroxy-L-lysine glucohydrolase  
**Comments:** The enzyme specifically hydrolyses glucose from  $\alpha$ -D-glucosyl-(1 $\rightarrow$ 2)- $\beta$ -D-galactosyl disaccharide units that are linked to hydroxylysine residues of collagen and collagen-like proteins. Acetylation of the  $\epsilon$ -amino group of the glycosylated hydroxylysine abolishes activity.  
**References:** [1035, 1036, 2766, 1034]

[EC 3.2.1.107 created 1984]

#### EC 3.2.1.108

**Accepted name:** lactase  
**Reaction:** lactose + H<sub>2</sub>O = D-galactose + D-glucose  
**Other name(s):** lactase-phlorizin hydrolase  
**Systematic name:** lactose galactohydrolase  
**Comments:** The enzyme from intestinal mucosa is isolated as a complex that also catalyses the reaction of EC 3.2.1.62 glycosylceramidase. *cf.* EC 3.2.1.33 amylo- $\alpha$ -1,6-glucosidase.  
**References:** [1736, 2365, 2567, 2685, 2686, 89]

[EC 3.2.1.108 created 1984]

#### EC 3.2.1.109

**Accepted name:** endogalactosaminidase  
**Reaction:** Endohydrolysis of (1 $\rightarrow$ 4)- $\alpha$ -D-galactosaminidic linkages in poly(D-galactosamine)  
**Systematic name:** galactosaminoglycan glycanohydrolase  
**References:** [2403, 2867]

[EC 3.2.1.109 created 1984]

[3.2.1.110 Deleted entry. mucinaminylserine mucinaminidase. The enzyme is identical to EC 3.2.1.97, glycopeptide  $\alpha$ -*N*-acetylgalactosaminidase]

[EC 3.2.1.110 created 1984, deleted 2008]

#### EC 3.2.1.111

**Accepted name:** 1,3- $\alpha$ -L-fucosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 3)-linkages between  $\alpha$ -L-fucose and *N*-acetylglucosamine residues in glycoproteins  
**Other name(s):** almond emulsin fucosidase I  
**Systematic name:** 3- $\alpha$ -L-fucosyl-*N*-acetylglucosaminyl-glycoprotein fucohydrolase  
**Comments:** Not identical with EC 3.2.1.63 1,2- $\alpha$ -L-fucosidase.  
**References:** [1243, 2144, 3303]

[EC 3.2.1.111 created 1986]

#### EC 3.2.1.112

**Accepted name:** 2-deoxyglucosidase  
**Reaction:** a 2-deoxy- $\alpha$ -D-glucoside + H<sub>2</sub>O = 2-deoxy-D-glucose + an alcohol  
**Other name(s):** 2-deoxy- $\alpha$ -glucosidase; 2-deoxy- $\alpha$ -D-glucosidase  
**Systematic name:** 2-deoxy- $\alpha$ -D-glucoside deoxyglucohydrolase  
**References:** [366]

[EC 3.2.1.112 created 1986]

#### EC 3.2.1.113

**Accepted name:** mannosyl-oligosaccharide 1,2- $\alpha$ -mannosidase  
**Reaction:** Hydrolysis of the terminal (1 $\rightarrow$ 2)-linked  $\alpha$ -D-mannose residues in the oligo-mannose oligosaccharide Man<sub>9</sub>(GlcNAc)<sub>2</sub>  
**Other name(s):** mannosidase 1A; mannosidase 1B; 1,2- $\alpha$ -mannosidase; exo- $\alpha$ -1,2-mannanase; mannose-9 processing  $\alpha$ -mannosidase; glycoprotein processing mannosidase I; mannosidase I; Man<sub>9</sub>-mannosidase; ManI; 1,2- $\alpha$ -mannosyl-oligosaccharide  $\alpha$ -D-mannohydrolase  
**Systematic name:** 2- $\alpha$ -mannosyl-oligosaccharide  $\alpha$ -D-mannohydrolase  
**Comments:** Involved in the synthesis of glycoproteins.  
**References:** [2826, 2994]

[EC 3.2.1.113 created 1986]

#### EC 3.2.1.114

**Accepted name:** mannosyl-oligosaccharide 1,3-1,6- $\alpha$ -mannosidase  
**Reaction:** Man<sub>5</sub>GlcNAc<sub>3</sub>-[protein] + 2 H<sub>2</sub>O = Man<sub>3</sub>GlcNAc<sub>3</sub>-[protein] + 2  $\alpha$ -D-mannopyranose  
**Other name(s):** MAN2A1 (gene name); MAN2A2 (gene name); mannosidase II; exo-1,3-1,6- $\alpha$ -mannosidase;  $\alpha$ -D-mannosidase II;  $\alpha$ -mannosidase II;  $\alpha$ 1-3,6-mannosidase; GlcNAc transferase I-dependent  $\alpha$ 1,3[ $\alpha$ 1,6]mannosidase; Golgi  $\alpha$ -mannosidase II; ManII; 1,3(1,6)- $\alpha$ -D-mannosidase; 1,3-(1,6-)mannosyl-oligosaccharide  $\alpha$ -D-mannohydrolase; (1 $\rightarrow$ 3)-(1 $\rightarrow$ 6)-mannosyl-oligosaccharide  $\alpha$ -D-mannohydrolase  
**Systematic name:** (1 $\rightarrow$ 3)-(1 $\rightarrow$ 6)-mannosyl-oligosaccharide  $\alpha$ -D-mannohydrolase (configuration-retaining)  
**Comments:** The enzyme, found in plants and animals, participates in the processing of *N*-glycans in the Golgi apparatus. It removes two mannosyl residues, one linked by  $\alpha$ 1,3 linkage, and the other linked by  $\alpha$ 1,6 linkage, both of which are removed by the same catalytic site. The enzyme is sensitive to swainsonine.  
**References:** [2995, 2825, 1057, 2994, 1964, 1926, 3036, 92, 2615, 2458]

[EC 3.2.1.114 created 1986, modified 2018]

#### EC 3.2.1.115

**Accepted name:** branched-dextran exo-1,2- $\alpha$ -glucosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 2)- $\alpha$ -D-glucosidic linkages at the branch points of dextrans and related polysaccharides, producing free D-glucose  
**Other name(s):** dextran 1,2- $\alpha$ -glucosidase; dextran  $\alpha$ -1,2 debranching enzyme1 1,2- $\alpha$ -D-glucosyl-branched-dextran 2-glucohydrolase  
**Systematic name:** (1 $\rightarrow$ 2)- $\alpha$ -D-glucosyl-branched-dextran 2-glucohydrolase  
**Comments:** Does not hydrolyse disaccharides or oligosaccharides containing linear 1,2- $\alpha$ -glucosidic linkages.  
**References:** [1933, 1934]

[EC 3.2.1.115 created 1989]

#### EC 3.2.1.116

**Accepted name:** glucan 1,4- $\alpha$ -maltotriohydrolase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 4)- $\alpha$ -D-glucosidic linkages in amylaceous polysaccharides, to remove successive maltotriose residues from the non-reducing chain ends  
**Other name(s):** exo-maltotriohydrolase; maltotriohydrolase; 1,4- $\alpha$ -D-glucan maltotriohydrolase  
**Systematic name:** 4- $\alpha$ -D-glucan maltotriohydrolase  
**Comments:** *cf.* EC 3.2.1.2 ( $\beta$ -amylase), EC 3.2.1.60 (glucan 1,4- $\alpha$ -maltotetrahydrolase) and EC 3.2.1.98 (glucan 1,4- $\alpha$ -maltohexaosidase). The products have the  $\alpha$ -configuration.  
**References:** [2043]

[EC 3.2.1.116 created 1989]

#### EC 3.2.1.117

**Accepted name:** amygdalin  $\beta$ -glucosidase  
**Reaction:** (*R*)-amygdalin + H<sub>2</sub>O = (*R*)-prunasin + D-glucose  
**Other name(s):** amygdalase; amygdalinase; amygdalin hydrolase; amygdalin glucosidase  
**Systematic name:** amygdalin  $\beta$ -D-glucohydrolase  
**Comments:** Highly specific; does not act on prunasin, linamarin, gentiobiose or cellobiose (*cf.* EC 3.2.1.21  $\beta$ -glucosidase).  
**References:** [1572]

[EC 3.2.1.117 created 1989]

#### EC 3.2.1.118

**Accepted name:** prunasin  $\beta$ -glucosidase  
**Reaction:** (*R*)-prunasin + H<sub>2</sub>O = D-glucose + mandelonitrile  
**Other name(s):** prunasin hydrolase  
**Systematic name:** prunasin  $\beta$ -D-glucohydrolase  
**Comments:** Highly specific; does not act on amygdalin, linamarin or gentiobiose. (*cf.* EC 3.2.1.21  $\beta$ -glucosidase).  
**References:** [1572]

[EC 3.2.1.118 created 1989]

#### EC 3.2.1.119

**Accepted name:** vicianin  $\beta$ -glucosidase  
**Reaction:** (*R*)-vicianin + H<sub>2</sub>O = mandelonitrile + vicianose  
**Other name(s):** vicianin hydrolase  
**Systematic name:** (*R*)-vicianin  $\beta$ -D-glucohydrolase  
**Comments:** Also hydrolyses, more slowly, (*R*)-amygdalin and (*R*)-prunasin, but not gentiobiose, linamarin or cellobiose.  
**References:** [1572]

[EC 3.2.1.119 created 1989]

#### EC 3.2.1.120

**Accepted name:** oligoxyloglucan  $\beta$ -glycosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-glucosidic links in oligoxyloglucans so as to remove successive isoprimeverose [i.e.  $\alpha$ -xylo-(1 $\rightarrow$ 6)- $\beta$ -D-glucosyl-] residues from the non-reducing chain ends  
**Other name(s):** isoprimeverose-producing oligoxyloglucan hydrolase; oligoxyloglucan hydrolase  
**Systematic name:** oligoxyloglucan xyloglucohydrolase  
**References:** [1402]

[EC 3.2.1.120 created 1989]

#### EC 3.2.1.121

**Accepted name:** polymannuronate hydrolase  
**Reaction:** Endohydrolysis of the D-mannuronide linkages of polymannuronate  
**Other name(s):** polymannuronic acid polymerase  
**Systematic name:** poly(mannuronide) manuronohydrolase  
**Comments:** Does not act on alginic acid, which is a copolymer of polymannuronate.  
**References:** [668]

[EC 3.2.1.121 created 1989]

#### EC 3.2.1.122

**Accepted name:** maltose-6'-phosphate glucosidase  
**Reaction:**  $\alpha$ -maltose 6'-phosphate + H<sub>2</sub>O = D-glucose + D-glucose 6-phosphate  
**Other name(s):** phospho- $\alpha$ -glucosidase; maltose-6'-phosphate 6-phosphoglucohydrolase  
**Systematic name:**  $\alpha$ -maltose-6'-phosphate 6-phosphoglucohydrolase  
**Comments:** Hydrolyses a variety of 6-phospho- $\alpha$ -D-glucosides, including  $\alpha$ -maltose 6'-phosphate,  $\alpha,\alpha$ -trehalose 6-phosphate, sucrose 6-phosphate and *p*-nitrophenyl- $\alpha$ -D-glucopyranoside 6-phosphate (as a chromogenic substrate). The enzyme is activated by Fe<sup>II</sup>, Mn<sup>II</sup>, Co<sup>II</sup> and Ni<sup>II</sup>. It is rapidly inactivated in air.  
**References:** [2916]

[EC 3.2.1.122 created 1989, modified 1999]

#### EC 3.2.1.123

**Accepted name:** endoglycosylceramidase  
**Reaction:** oligoglycosylglucosyl-(1 $\leftrightarrow$ 1)-ceramide + H<sub>2</sub>O = ceramide + oligoglycosylglucose  
**Other name(s):** endoglycoceramidase; EGCase; glycosyl-*N*-acetyl-sphingosine 1,1- $\beta$ -D-glucanohydrolase, oligoglycosylglucosylceramide glycohydrolase; oligoglycosylglucosyl(1 $\leftrightarrow$ 1)ceramide glycohydrolase  
**Systematic name:** oligoglycosylglucosyl-(1 $\leftrightarrow$ 1)-ceramide glycohydrolase  
**Comments:** An enzyme from *Rhodococcus* sp. that degrades various acidic and neutral glycosphingolipids to oligosaccharides and ceramides, by cleaving a glucosyl bond. Does not act on monoglycosylceramides. *cf.* EC 3.2.1.62 glycosylceramidase.  
**References:** [1266]

[EC 3.2.1.123 created 1989]

#### EC 3.2.1.124

**Accepted name:** 3-deoxy-2-octulosonidase  
**Reaction:** Endohydrolysis of the  $\beta$ -ketopyranosidic linkages of 3-deoxy-D-*manno*-2-octulosonate in capsular polysaccharides  
**Other name(s):** 2-keto-3-deoxyoctonate hydrolase; octulosylono hydrolase; octulofuranosylono hydrolase; octulopyranosylonohydrolase  
**Systematic name:** capsular-polysaccharide 3-deoxy-D-*manno*-2-octulosonohydrolase  
**Comments:** The enzyme from a bacteriophage catalyses the depolymerization of capsular polysaccharides containing 3-deoxy-2-octulosonide in the cell wall of *Escherichia coli*.  
**References:** [42]

[EC 3.2.1.124 created 1989]

#### EC 3.2.1.125

**Accepted name:** raucaffricine  $\beta$ -glucosidase  
**Reaction:** raucaffricine + H<sub>2</sub>O = D-glucose + vomilenine  
**Other name(s):** raucaffricine  $\beta$ -D-glucosidase; raucaffricine glucosidase

**Systematic name:** raucaffricine  $\beta$ -D-glucohydrolase  
**Comments:** Highly specific; some other ajmalan glucoside alkaloids are hydrolysed, but more slowly.  
**References:** [2587]

[EC 3.2.1.125 created 1989]

#### EC 3.2.1.126

**Accepted name:** coniferin  $\beta$ -glucosidase  
**Reaction:** coniferin + H<sub>2</sub>O = D-glucose + coniferol  
**Other name(s):** coniferin-hydrolyzing  $\beta$ -glucosidase  
**Systematic name:** coniferin  $\beta$ -D-glucohydrolase  
**Comments:** Also hydrolyses syringin, 4-cinnamyl alcohol  $\beta$ -glucoside and, more slowly, some other aryl  $\beta$ -glycosides. A plant cell-wall enzyme involved in the biosynthesis of lignin.  
**References:** [1182, 1807]

[EC 3.2.1.126 created 1989]

#### EC 3.2.1.127

**Accepted name:** 1,6- $\alpha$ -L-fucosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 6)-linkages between  $\alpha$ -L-fucose and *N*-acetyl-D-glucosamine in glycopeptides such as immunoglobulin G glycopeptide and fucosyl-asialo-agalacto-fetuin  
**Other name(s):**  $\alpha$ -L-fucosidase; 1,6-L-fucosyl-*N*-acetyl-D-glucosaminylglycopeptide fucohydrolase  
**Systematic name:** 6-L-fucosyl-*N*-acetyl-D-glucosaminylglycopeptide fucohydrolase  
**Comments:** The enzyme from *Aspergillus niger* does not act on 1,2-, 1,3-, or 1,4-L-fucosyl linkages.  
**References:** [3280]

[EC 3.2.1.127 created 1989]

#### EC 3.2.1.128

**Accepted name:** glycyrrhizin hydrolase  
**Reaction:** glycyrrhizin + H<sub>2</sub>O =  $\beta$ -D-glucuronosyl-(1 $\rightarrow$ 2)-D-glucuronate + glycyrrhetinate  
**Other name(s):** glycyrrhizinate  $\beta$ -glucuronidase; glycyrrhizin  $\beta$ -hydrolase; glycyrrhizinic acid hydrolase  
**Systematic name:** glycyrrhizinate glucuronosylhydrolase  
**Comments:** The enzyme from *Aspergillus niger* is specific for the hydrolysis of the triterpenoid glycoside glycyrrhizin from roots of *Glycyrrhiza* sp.  
**References:** [2008]

[EC 3.2.1.128 created 1989]

#### EC 3.2.1.129

**Accepted name:** endo- $\alpha$ -sialidase  
**Reaction:** Endohydrolysis of (2 $\rightarrow$ 8)- $\alpha$ -sialosyl linkages in oligo- or poly(sialic) acids  
**Other name(s):** endo-*N*-acetylneuraminidase; endoneuraminidase; endo-*N*-acetylneuraminidase; poly( $\alpha$ -2,8-sialosyl) endo-*N*-acetylneuraminidase; poly( $\alpha$ -2,8-sialoside)  $\alpha$ -2,8-sialosylhydrolase; endosialidase; endo-*N*-polysialoside (2 $\rightarrow$ 8)- $\alpha$ -sialosylhydrolase  
**Systematic name:** polysialoside (2 $\rightarrow$ 8)- $\alpha$ -sialosylhydrolase  
**Comments:** Although the name endo-*N*-acetylneuraminidase has also been used for this enzyme, this is misleading since its activity is not restricted to acetylated substrates. An exo- $\alpha$ -sialidase activity is listed as EC 3.2.1.18 exo- $\alpha$ -sialidase. See also EC 4.2.2.15 anhydrosialidase.  
**References:** [769, 1029, 1470, 1580, 2253, 2932, 347]

[EC 3.2.1.129 created 1990, modified 1999]

### EC 3.2.1.130

- Accepted name:** glycoprotein endo- $\alpha$ -1,2-mannosidase  
**Reaction:**  $\text{GlcMan}_9\text{GlcNAc}_2\text{-[protein]} + \text{H}_2\text{O} = \text{Man}_8\text{GlcNAc}_2\text{-[protein]} (\text{isomer } 8\text{A}_{1,2,3}\text{B}_{1,2}) + \alpha\text{-D-glucosyl-}(1\rightarrow3)\text{-}\alpha\text{-D-mannopyranose}$   
**Other name(s):** glucosylmannosidase; endo- $\alpha$ -D-mannosidase; endo- $\alpha$ -mannosidase; endomannosidase; glucosyl mannosidase; MANEA (gene name); glycoprotein glucosylmannohydrolase  
**Systematic name:** glycoprotein glucosylmannohydrolase (configuration-retaining)  
**Comments:** The enzyme catalyses the hydrolysis of the terminal  $\alpha$ -D-glucosyl-(1 $\rightarrow$ 3)-D-mannosyl unit from the  $\text{GlcMan}_9(\text{GlcNAc})_2$  oligosaccharide component of *N*-glucosylated proteins during their processing in the Golgi apparatus. The name for the isomer is based on a nomenclature proposed by Prien et al [2325].  
**References:** [1749, 2993, 1144, 2741, 1037, 1052, 2325]

[EC 3.2.1.130 created 1990, modified 2017]

### EC 3.2.1.131

- Accepted name:** xylan  $\alpha$ -1,2-glucuronosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 2)- $\alpha$ -D-(4-*O*-methyl)glucuronosyl links in the main chain of hardwood xylans  
**Other name(s):** 1,2- $\alpha$ -glucuronidase;  $\alpha$ -(1 $\rightarrow$ 2)-glucuronidase; xylan  $\alpha$ -D-1,2-(4-*O*-methyl)glucuronohydrolase  
**Systematic name:** xylan 2- $\alpha$ -D-(4-*O*-methyl)glucuronohydrolase  
**References:** [1261]

[EC 3.2.1.131 created 1990]

### EC 3.2.1.132

- Accepted name:** chitosanase  
**Reaction:** Endohydrolysis of  $\beta$ -(1 $\rightarrow$ 4)-linkages between D-glucosamine residues in a partly acetylated chitosan  
**Systematic name:** chitosan *N*-acetylglucosaminohydrolase  
**Comments:** A whole spectrum of chitosanases are now known (for more details, see <http://rbrzezinski.recherche.usherbrooke.ca/>). They can hydrolyse various types of links in chitosan. The only constant property is the endohydrolysis of GlcN-GlcN links, which is common to all known chitosanases. One known chitosanase is limited to this link recognition [1808], while the majority can also recognize GlcN-GlcNAc links or GlcNAc-GlcN links but not both. They also do not recognize GlcNAc-GlcNAc links in partly acetylated chitosan.  
**References:** [755, 2496, 1286, 1808]

[EC 3.2.1.132 created 1990, modified 2004]

### EC 3.2.1.133

- Accepted name:** glucan 1,4- $\alpha$ -maltohydrolase  
**Reaction:** hydrolysis of (1 $\rightarrow$ 4)- $\alpha$ -D-glucosidic linkages in polysaccharides so as to remove successive  $\alpha$ -maltose residues from the non-reducing ends of the chains  
**Other name(s):** maltogenic  $\alpha$ -amylase; 1,4- $\alpha$ -D-glucan  $\alpha$ -maltohydrolase  
**Systematic name:** 4- $\alpha$ -D-glucan  $\alpha$ -maltohydrolase  
**Comments:** Acts on starch and related polysaccharides and oligosaccharides. The product is  $\alpha$ -maltose; *cf.* EC 3.2.1.2  $\beta$ -amylase.  
**References:** [607, 2218]

[EC 3.2.1.133 created 1992, modified 1999]

### EC 3.2.1.134

- Accepted name:** difructose-anhydride synthase  
**Reaction:** bis-D-fructose 2',1:2,1'-dianhydride + H<sub>2</sub>O = inulobiose

**Other name(s):** inulobiose hydrolase  
**Systematic name:** bis-D-fructose 2',1:2,1'-dianhydride fructohydrolase  
**Comments:** Produces difructose anhydride by the reverse reaction of partial hydrolysis, forming an  $\alpha$ -fructosidic linkage.  
**References:** [1841, 1842]

[EC 3.2.1.134 created 1992]

#### EC 3.2.1.135

**Accepted name:** neopullulanase  
**Reaction:** Hydrolysis of pullulan to panose (6- $\alpha$ -D-glucosylmaltose)  
**Other name(s):** pullulanase II  
**Systematic name:** pullulan 4-D-glucanohydrolase (panose-forming)  
**Comments:** cf. EC 3.2.1.41 (pullulanase ) and EC 3.2.1.57 (isopullulanase).  
**References:** [1242]

[EC 3.2.1.135 created 1992]

#### EC 3.2.1.136

**Accepted name:** glucuronoarabinoxylan endo-1,4- $\beta$ -xylanase  
**Reaction:** Endohydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-xylosyl links in some glucuronoarabinoxylans  
**Other name(s):** feraxan endoxylanase; feraxanase; endoarabinoxylanase; glucuronoxylan xylohydrolase; glucuronoxylanase; glucuronoxylan xylanohydrolase; glucuronoarabinoxylan 1,4- $\beta$ -D-xylanohydrolase  
**Systematic name:** glucuronoarabinoxylan 4- $\beta$ -D-xylanohydrolase  
**Comments:** High activity towards feruloylated arabinoxylans from cereal plant cell walls.  
**References:** [2095]

[EC 3.2.1.136 created 1992]

#### EC 3.2.1.137

**Accepted name:** mannan exo-1,2-1,6- $\alpha$ -mannosidase  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 2)- $\alpha$ -D- and (1 $\rightarrow$ 6)- $\alpha$ -D- linkages in yeast mannan, releasing D-mannose  
**Other name(s):** exo-1,2-1,6- $\alpha$ -mannosidase; 1,2-1,6- $\alpha$ -D-mannan D-mannohydrolase  
**Systematic name:** (1 $\rightarrow$ 2)-(1 $\rightarrow$ 6)- $\alpha$ -D-mannan D-mannohydrolase  
**Comments:** Mannose residues linked  $\alpha$ -D-1,3- are also released, but very slowly.  
**References:** [2856]

[EC 3.2.1.137 created 1992]

[3.2.1.138 *Transferred entry. anhydrosialidase. Now EC 4.2.2.15, anhydrosialidase*]

[EC 3.2.1.138 created 1992, deleted 2003]

#### EC 3.2.1.139

**Accepted name:**  $\alpha$ -glucuronidase  
**Reaction:** an  $\alpha$ -D-glucuronoside + H<sub>2</sub>O = an alcohol + D-glucuronate  
**Other name(s):**  $\alpha$ -glucosiduronase  
**Systematic name:**  $\alpha$ -D-glucosiduronate glucuronohydrolase  
**Comments:** Considerable differences in the specificities of the enzymes from different fungi for  $\alpha$ -D-glucosiduronates have been reported. Activity is also found in the snail.  
**References:** [2329, 3006]

[EC 3.2.1.139 created 1999]

#### EC 3.2.1.140

- Accepted name:** lacto-*N*-biosidase  
**Reaction:**  $\beta$ -D-Gal-(1→3)- $\beta$ -D-GlcNAc-(1→3)- $\beta$ -D-Gal-(1→4)-D-Glc + H<sub>2</sub>O =  $\beta$ -D-Gal-(1→3)-D-GlcNAc +  $\beta$ -D-Gal-(1→4)-D-Glc  
**Systematic name:** oligosaccharide lacto-*N*-biosylhydrolase  
**Comments:** The enzyme from *Streptomyces* specifically hydrolyses the terminal lacto-*N*-biosyl residue ( $\beta$ -D-Gal-(1→3)-D-GlcNAc) from the non-reducing end of oligosaccharides with the structure  $\beta$ -D-Gal-(1→3)- $\beta$ -D-GlcNAc-(1→3)- $\beta$ -D-Gal-(1→R). Lacto-*N*-hexaose ( $\beta$ -D-Gal-(1→3)- $\beta$ -D-GlcNAc-(1→3)- $\beta$ -D-Gal-(1→3)- $\beta$ -D-GlcNAc-(1→3)- $\beta$ -D-Gal-(1→4)-D-Glc) is hydrolysed to form first lacto-*N*-tetraose plus lacto-*N*-biose, with the subsequent formation of lactose. Oligosaccharides in which the non-reducing terminal Gal or the penultimate GlcNAc are replaced by fucose or sialic acid are not substrates. Asialo GM1 tetraose ( $\beta$ -D-Gal-(1→3)- $\beta$ -D-GalNAc-(1→3)- $\beta$ -D-Gal-(1→4)-D-Glc) is hydrolysed very slowly, but lacto-*N*-neotetraose ( $\beta$ -D-Gal-(1→4)- $\beta$ -D-GalNAc-(1→3)- $\beta$ -D-Gal-(1→4)-D-Glc) is not a substrate  
**References:** [2521, 2522]

[EC 3.2.1.140 created 1999]

#### EC 3.2.1.141

- Accepted name:** 4- $\alpha$ -D-(1→4)- $\alpha$ -D-glucanotrehalose trehalohydrolase  
**Reaction:** hydrolysis of (1→4)- $\alpha$ -D-glucosidic linkage in 4- $\alpha$ -D-[(1→4)- $\alpha$ -D-glucanosyl]<sub>n</sub> trehalose to yield trehalose and (1→4)- $\alpha$ -D-glucan  
**Other name(s):** malto-oligosyltrehalose trehalohydrolase  
**Systematic name:** 4- $\alpha$ -D-[(1→4)- $\alpha$ -D-glucano]trehalose glucanohydrolase (trehalose-producing)  
**References:** [1825, 2032, 2031]

[EC 3.2.1.141 created 1999]

#### EC 3.2.1.142

- Accepted name:** limit dextrinase  
**Reaction:** Hydrolysis of (1→6)- $\alpha$ -D-glucosidic linkages in  $\alpha$ - and  $\beta$ -limit dextrans of amylopectin and glycogen, and in amylopectin and pullulan  
**Other name(s):** *R*-enzyme; amylopectin-1,6-glucosidase; dextrin  $\alpha$ -1,6-glucanohydrolase  
**Systematic name:** dextrin 6- $\alpha$ -glucanohydrolase  
**Comments:** Plant enzymes with little or no action on glycogen. Action on amylopectin is incomplete, but action on  $\alpha$ -limit dextrans is complete. Maltose is the smallest sugar it can release from an  $\alpha$ -(1→6)-linkage.  
**References:** [958, 1801]

[EC 3.2.1.142 created 2000]

#### EC 3.2.1.143

- Accepted name:** poly(ADP-ribose) glycohydrolase  
**Reaction:** hydrolyses poly(ADP-D-ribose) at glycosidic (1''-2') linkage of ribose-ribose bond to produce free ADP-D-ribose  
**Comments:** Specific to (1''-2') linkage of ribose-ribose bond of poly(ADP-D-ribose).  
**References:** [1935, 1694]

[EC 3.2.1.143 created 2000]

#### EC 3.2.1.144

- Accepted name:** 3-deoxyoctulosonase  
**Reaction:** 3-deoxyoctulosonyl-lipopolysaccharide + H<sub>2</sub>O = 3-deoxyoctulosonic acid + lipopolysaccharide  
**Other name(s):**  $\alpha$ -Kdo-ase



**Systematic name:** 3-deoxyoctulosonyl-lipopolysaccharide hydrolase  
**Comments:** Releases Kdo ( $\alpha$ - and  $\beta$ -linked 3-deoxy-D-manno-octulosonic acid) from different lipopolysaccharides, including *Re*-LPS from *Escherichia coli* and *Salmonella*, Rd-LPS from *S. minnesota*, and de-*O*-acyl-re-LPS. 4-Methylumbelliferyl- $\alpha$ -Kdo ( $\alpha$ -Kdo-OMec) is also a substrate.  
**References:** [1676]

[EC 3.2.1.144 created 2000]

#### EC 3.2.1.145

**Accepted name:** galactan 1,3- $\beta$ -galactosidase  
**Reaction:** Hydrolysis of terminal, non-reducing  $\beta$ -D-galactose residues in (1 $\rightarrow$ 3)- $\beta$ -D-galactopyranans  
**Other name(s):** galactan (1 $\rightarrow$ 3)- $\beta$ -D-galactosidase  
**Systematic name:** galactan 3- $\beta$ -D-galactosidase  
**Comments:** This enzyme removes not only free galactose, but also 6-glycosylated residues, e.g., (1 $\rightarrow$ 6)- $\beta$ -D-galactobiose, and galactose bearing oligosaccharide chains on O-6. Hence, it releases branches from [*arabino*-galacto-(1 $\rightarrow$ 6)]-(1 $\rightarrow$ 3)- $\beta$ -D-galactans.  
**References:** [2983, 2254]

[EC 3.2.1.145 created 2001]

#### EC 3.2.1.146

**Accepted name:**  $\beta$ -galactofuranosidase  
**Reaction:** Hydrolysis of terminal non-reducing  $\beta$ -D-galactofuranosides, releasing galactose  
**Other name(s):** exo- $\beta$ -galactofuranosidase; exo- $\beta$ -D-galactofuranosidase;  $\beta$ -D-galactofuranosidase  
**Systematic name:**  $\beta$ -D-galactofuranoside hydrolase  
**Comments:** The enzyme from *Helminthosporium sacchari* detoxifies helminthosporoside, a bis(digalactosyl)terpene produced by this fungus, by releasing its four molecules of bound galactose.  
**References:** [2425, 530, 498, 1910]

[EC 3.2.1.146 created 2001]

#### EC 3.2.1.147

**Accepted name:** thioglucosidase  
**Reaction:** a thioglucoside + H<sub>2</sub>O = a sugar + a thiol  
**Other name(s):** myrosinase; sinigrinase; sinigrase  
**Systematic name:** thioglucoside glucohydrolase  
**Comments:** Has a wide specificity for thioglycosides.  
**References:** [956, 2281]

[EC 3.2.1.147 created 1972 as EC 3.2.3.1, transferred 2001 to EC 3.2.1.147]

[3.2.1.148 Deleted entry. ribosylhomocysteinase. This enzyme was transferred to EC 3.13.1.2, 5-deoxyribos-5-ylhomocysteinase, which has since been deleted. The activity is most probably attributable to EC 4.4.1.21, S-ribosylhomocysteine lyase]

[EC 3.2.1.148 created 1972 as EC 3.3.1.3, transferred 2001 to EC 3.2.1.148, deleted 2004]

#### EC 3.2.1.149

**Accepted name:**  $\beta$ -primeverosidase  
**Reaction:** a 6-*O*-( $\beta$ -D-xylopyranosyl)- $\beta$ -D-glucopyranoside + H<sub>2</sub>O = 6-*O*-( $\beta$ -D-xylopyranosyl)- $\beta$ -D-glucopyranose + an alcohol  
**Systematic name:** 6-*O*-( $\beta$ -D-xylopyranosyl)- $\beta$ -D-glucopyranoside 6-*O*-( $\beta$ -D-xylosyl)- $\beta$ -D-glucohydrolase

**Comments:** The enzyme is responsible for the formation of the alcoholic aroma in oolong and black tea. In addition to  $\beta$ -primeverosides [i.e. 6-*O*-( $\beta$ -D-xylopyranosyl)- $\beta$ -D-glucopyranosides], it also hydrolyses 6-*O*-( $\beta$ -D-apiofuranosyl)- $\beta$ -D-glucopyranosides and, less rapidly,  $\beta$ -vicianosides and 6-*O*-( $\alpha$ -L-arabinofuranosyl)- $\beta$ -D-glucopyranosides, but not  $\beta$ -glucosides. Geranyl-, linaloyl-, benzyl- and *p*-nitrophenol glycosides are all hydrolysed.

**References:** [1227, 2149]

[EC 3.2.1.149 created 2001]

#### EC 3.2.1.150

**Accepted name:** oligoxyloglucan reducing-end-specific cellobiohydrolase

**Reaction:** Hydrolysis of cellobiose from the reducing end of xyloglucans consisting of a (1 $\rightarrow$ 4)- $\beta$ -linked glucan carrying  $\alpha$ -D-xylosyl groups on O-6 of the glucose residues. To be a substrate, the first residue must be unsubstituted, the second residue may bear a xylosyl group, whether further glycosylated or not, and the third residue, which becomes the new terminus by the action of the enzyme, is preferably xylosylated, but this xylose residue must not be further substituted.

**Systematic name:** oligoxyloglucan reducing-end cellobiohydrolase

**Comments:** The enzyme is found in the fungus *Geotrichum* sp. M128. The substrate is a hemicellulose found in plant cell walls.

**References:** [3266]

[EC 3.2.1.150 created 2003]

#### EC 3.2.1.151

**Accepted name:** xyloglucan-specific endo- $\beta$ -1,4-glucanase

**Reaction:** xyloglucan + H<sub>2</sub>O = xyloglucan oligosaccharides

**Other name(s):** XEG; xyloglucan endo- $\beta$ -1,4-glucanase; xyloglucanase; xyloglucanendohydrolase; XH; 1,4- $\beta$ -D-glucan glucanohydrolase

**Systematic name:** [(1 $\rightarrow$ 6)- $\alpha$ -D-xylo]-(1 $\rightarrow$ 4)- $\beta$ -D-glucan glucanohydrolase

**Comments:** The enzyme from *Aspergillus aculeatus* is specific for xyloglucan and does not hydrolyse other cell-wall components. The reaction involves endohydrolysis of 1,4- $\beta$ -D-glucosidic linkages in xyloglucan with retention of the  $\beta$ -configuration of the glycosyl residues.

**References:** [2248, 985]

[EC 3.2.1.151 created 2003]

#### EC 3.2.1.152

**Accepted name:** mannosylglycoprotein endo- $\beta$ -mannosidase

**Reaction:** Hydrolysis of the  $\alpha$ -D-mannosyl-(1 $\rightarrow$ 6)- $\beta$ -D-mannosyl-(1 $\rightarrow$ 4)-*N*-acetyl- $\beta$ -D-glucosaminyl-(1 $\rightarrow$ 4)-*N*-acetyl- $\beta$ -D-glucosaminyl sequence of glycoprotein to  $\alpha$ -D-mannosyl-(1 $\rightarrow$ 6)-D-mannose and *N*-acetyl- $\beta$ -D-glucosaminyl-(1 $\rightarrow$ 4)-*N*-acetyl- $\beta$ -D-glucosaminyl sequences

**Other name(s):** endo- $\beta$ -mannosidase

**Comments:** The substrate group is a substituent on N-4 of an asparagine residue in the glycoprotein. The mannose residue at the non-reducing end of the sequence may carry further  $\alpha$ -D-mannosyl groups on O-3 or O-6, but such a substituent on O-3 of the  $\beta$ -D-mannosyl group prevents the action of the enzyme. The enzyme was obtained from the lily, *Lilium longiflorum*.

**References:** [1263, 2533]

[EC 3.2.1.152 created 2005]

#### EC 3.2.1.153

**Accepted name:** fructan  $\beta$ -(2,1)-fructosidase

**Reaction:** Hydrolysis of terminal, non-reducing (2 $\rightarrow$ 1)-linked  $\beta$ -D-fructofuranose residues in fructans

**Other name(s):**  $\beta$ -(2-1)-D-fructan fructohydrolase;  $\beta$ -(2-1)fructan exohydrolase; inulinase; 1-FEH II; 1-fructan exohydrolase; 1-FEH w1; 1-FEH w2;  $\beta$ -(2-1)-linkage-specific fructan- $\beta$ -fructosidase;  $\beta$ -(2,1)-D-fructan fructohydrolase

**Systematic name:**  $\beta$ -(2 $\rightarrow$ 1)-D-fructan fructohydrolase

**Comments:** Possesses one of the activities of EC 3.2.1.80, fructan  $\beta$ -fructosidase. While the best substrates are the inulin-type fructans, such as 1-kestose [ $\beta$ -D-fructofuranosyl-(2 $\rightarrow$ 1)- $\beta$ -D-fructofuranosyl  $\alpha$ -D-glucopyranoside] and 1,1-nystose [ $\beta$ -D-fructofuranosyl-(2 $\rightarrow$ 1)- $\beta$ -D-fructofuranosyl-(2 $\rightarrow$ 1)- $\beta$ -D-fructofuranosyl  $\alpha$ -D-glucopyranoside], some (but not all) levan-type fructans can also be hydrolysed, but more slowly [see EC 3.2.1.154, fructan  $\beta$ -(2,6)-fructosidase]. Sucrose, while being a very poor substrate, can substantially inhibit enzyme activity in some cases.

**References:** [2457, 579]

[EC 3.2.1.153 created 2005]

#### EC 3.2.1.154

**Accepted name:** fructan  $\beta$ -(2,6)-fructosidase

**Reaction:** Hydrolysis of terminal, non-reducing (2 $\rightarrow$ 6)-linked  $\beta$ -D-fructofuranose residues in fructans

**Other name(s):**  $\beta$ -(2-6)-fructan exohydrolase; levanase; 6-FEH;  $\beta$ -(2,6)-D-fructan fructohydrolase

**Systematic name:** (2 $\rightarrow$ 6)- $\beta$ -D-fructan fructohydrolase

**Comments:** Possesses one of the activities of EC 3.2.1.80, fructan  $\beta$ -fructosidase. While the best substrates are the levan-type fructans such as 6-kestotriose [ $\beta$ -D-fructofuranosyl-(2 $\rightarrow$ 6)- $\beta$ -D-fructofuranosyl  $\alpha$ -D-glucopyranoside] and 6,6-kestotetraose [ $\beta$ -D-fructofuranosyl-(2 $\rightarrow$ 6)- $\beta$ -D-fructofuranosyl-(2 $\rightarrow$ 6)- $\beta$ -D-fructofuranosyl  $\alpha$ -D-glucopyranoside], some (but not all) inulin-type fructans can also be hydrolysed, but more slowly [*cf.* EC 3.2.1.153, fructan  $\beta$ -(2,1)-fructosidase]. Sucrose, while being a very poor substrate, can substantially inhibit enzyme activity in some cases.

**References:** [1829, 580, 1119]

[EC 3.2.1.154 created 2005]

#### EC 3.2.1.155

**Accepted name:** xyloglucan-specific exo- $\beta$ -1,4-glucanase

**Reaction:** Hydrolysis of (1 $\rightarrow$ 4)-D-glucosidic linkages in xyloglucans so as to successively remove oligosaccharides from the chain end.

**Other name(s):** Cel74A

**Systematic name:** [(1 $\rightarrow$ 6)- $\alpha$ -D-xylo]-(1 $\rightarrow$ 4)- $\beta$ -D-glucan exo-glucohydrolase

**Comments:** The enzyme removes XXXG heptasaccharides, XXLG/XLXG octasaccharides and XLLG nonasaccharides from the end of tamarind seed xyloglucan polymers in a processive manner. Hydrolysis occurs at the unsubstituted D-glucopyranose residue in the main backbone. It is not known whether the cleavage takes place at the reducing or non-reducing end of the polymer. Very low activity with  $\beta$ -D-glucans. The enzyme from *Chrysosporium lucknowense* shifts to an endoglucanase mode when acting on linear substrates without bulky substituents on the polymeric backbone such as barley  $\beta$ -glucan.

**References:** [985]

[EC 3.2.1.155 created 2005, withdrawn at public-review stage, modified and reinstated 2006, modified 2011]

#### EC 3.2.1.156

**Accepted name:** oligosaccharide reducing-end xylanase

**Reaction:** Hydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-xylose residues from the reducing end of oligosaccharides

**Other name(s):** Rex; reducing end xylose-releasing exo-oligoxylanase

**Systematic name:**  $\beta$ -D-xylopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-xylopyranose reducing-end xylanase

**Comments:** The enzyme, originally isolated from the bacterium *Bacillus halodurans* C-125, releases the xylose unit at the reducing end of oligosaccharides ending with the structure  $\beta$ -D-xylopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-xylopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-xylopyranose, leaving the new reducing end in the  $\alpha$  configuration. It is specific for the  $\beta$  anomers of xylooligosaccharides whose degree of polymerization is equal to or greater than 3. The penultimate residue must be  $\beta$ -D-xylopyranose, but replacing either of the flanking residues with glucose merely slows the rate greatly.

**References:** [1166, 865]

[EC 3.2.1.156 created 2005]

#### EC 3.2.1.157

**Accepted name:**  $\iota$ -carrageenase

**Reaction:** Endohydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-linkages between D-galactose 4-sulfate and 3,6-anhydro-D-galactose-2-sulfate in  $\iota$ -carrageenans

**Systematic name:**  $\iota$ -carrageenan 4- $\beta$ -D-glycanohydrolase (configuration-inverting)

**Comments:** The main products of hydrolysis are  $\iota$ -neocarratetraose sulfate and  $\iota$ -neocarrahexaose sulfate.  $\iota$ -Neocarraoctaose is the shortest substrate oligomer that can be cleaved. Unlike EC 3.2.1.81,  $\beta$ -agarase and EC 3.2.1.83,  $\kappa$ -carrageenase, this enzyme proceeds with inversion of the anomeric configuration.  $\iota$ -Carrageenan differs from  $\kappa$ -carrageenan by possessing a sulfo group on O-2 of the 3,6-anhydro-D-galactose residues, in addition to that present in the  $\kappa$ -compound on O-4 of the D-galactose residues.

**References:** [137, 1904, 1905]

[EC 3.2.1.157 created 2006]

#### EC 3.2.1.158

**Accepted name:**  $\alpha$ -agarase

**Reaction:** Endohydrolysis of (1 $\rightarrow$ 3)- $\alpha$ -L-galactosidic linkages in agarose, yielding agarotetraose as the major product

**Other name(s):** agarase (ambiguous); agaraseA33

**Systematic name:** agarose 3-glycanohydrolase

**Comments:** Requires  $\text{Ca}^{2+}$ . The enzyme from *Thalassomonas* sp. can use agarose, agarohexaose and neoagarohexaose as substrate. The products of agarohexaose hydrolysis are dimers and tetramers, with agarotetraose being the predominant product, whereas hydrolysis of neoagarohexaose gives rise to two types of trimer. While the enzyme can also hydrolyse the highly sulfated agarose porphyran very efficiently, it cannot hydrolyse the related compounds  $\kappa$ -carrageenan (see EC 3.2.1.83) and  $\iota$ -carrageenan (see EC 3.2.1.157) [2170]. See also EC 3.2.1.81,  $\beta$ -agarase.

**References:** [2317, 2170]

[EC 3.2.1.158 created 2006]

#### EC 3.2.1.159

**Accepted name:**  $\alpha$ -neoagaro-oligosaccharide hydrolase

**Reaction:** Hydrolysis of the (1 $\rightarrow$ 3)- $\alpha$ -L-galactosidic linkages of neoagaro-oligosaccharides that are smaller than a hexamer, yielding 3,6-anhydro-L-galactose and D-galactose

**Other name(s):**  $\alpha$ -neoagaro-oligosaccharide hydrolase;  $\alpha$ -NAOS hydrolase

**Systematic name:**  $\alpha$ -neoagaro-oligosaccharide 3-glycohydrolase

**Comments:** When neoagarohexaose is used as a substrate, the oligosaccharide is cleaved at the non-reducing end to produce 3,6-anhydro-L-galactose and agaropentaose, which is further hydrolysed to agarobiose and agarotriose. With neoagarotetraose as substrate, the products are predominantly agarotriose and 3,6-anhydro-L-galactose. In *Vibrio* sp. the actions of EC 3.2.1.81,  $\beta$ -agarase and EC 3.2.1.159 can be used to degrade agarose to 3,6-anhydro-L-galactose and D-galactose.

**References:** [2791]

[EC 3.2.1.159 created 2006]

[3.2.1.160 Deleted entry. xyloglucan-specific  $\alpha$ -D-glucanase. The enzyme was shown to be identical to EC 3.2.1.155, xyloglucan-specific  $\alpha$ -D-glucanase, during the public-review process so was withdrawn before being made official]

[EC 3.2.1.160 created 2006, deleted 2006]

#### EC 3.2.1.161

**Accepted name:**  $\beta$ -apiosyl- $\beta$ -glucosidase  
**Reaction:** 7-[ $\beta$ -D-apiofuranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranosyloxy]isoflavonoid + H<sub>2</sub>O = a 7-hydroxyisoflavonoid +  $\beta$ -D-apiofuranosyl-(1 $\rightarrow$ 6)-D-glucose  
**Other name(s):** isoflavonoid-7-*O*- $\beta$ [D-apiosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucoside] disaccharidase; isoflavonoid 7-*O*- $\beta$ -apiosyl-glucoside  $\beta$ -glucosidase; furcatin hydrolase  
**Systematic name:** 7-[ $\beta$ -D-apiofuranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranosyloxy]isoflavonoid  $\beta$ -D-apiofuranosyl-(1 $\rightarrow$ 6)-D-glucohydrolase  
**Comments:** The enzyme from the tropical tree *Dalbergia nigrescens* Kurz belongs in glycosyl hydrolase family 1. The enzyme removes disaccharides from the natural substrates dalpatein 7-*O*- $\beta$ -D-apiofuranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside and 7-hydroxy-2',4',5',6-tetramethoxy-7-*O*- $\beta$ -D-apiofuranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (dalnigrein 7-*O*- $\beta$ -D-apiofuranosyl-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside) although it can also remove a single glucose residue from isoflavonoid 7-*O*-glucosides [445]. Daidzin and genistin are also substrates.  
**References:** [1181, 445, 23]

[EC 3.2.1.161 created 2006]

#### EC 3.2.1.162

**Accepted name:**  $\lambda$ -carrageenase  
**Reaction:** Endohydrolysis of (1 $\rightarrow$ 4)- $\beta$ -linkages in the backbone of  $\lambda$ -carrageenan, resulting in the tetrasaccharide  $\alpha$ -D-Galp2,6S<sub>2</sub>-(1 $\rightarrow$ 3)- $\beta$ -D-Galp2S-(1 $\rightarrow$ 4)- $\alpha$ -D-Galp2,6S<sub>2</sub>-(1 $\rightarrow$ 3)-D-Galp2S  
**Other name(s):** endo- $\beta$ -1,4-carrageenase 2,6,2'-trisulfate-hydrolase  
**Systematic name:** endo-(1 $\rightarrow$ 4)- $\beta$ -carrageenase 2,6,2'-trisulfate-hydrolase  
**Comments:** The enzyme from *Pseudoalteromonas* sp. is specific for  $\lambda$ -carrageenan.  $\iota$ -Carrageenan (see EC 3.2.1.157,  $\iota$ -carrageenase),  $\kappa$ -carrageenan (see EC 3.2.1.83,  $\kappa$ -carrageenase), agarose and porphyran are not substrates.  
**References:** [2169]

[EC 3.2.1.162 created 2007]

#### EC 3.2.1.163

**Accepted name:** 1,6- $\alpha$ -D-mannosidase  
**Reaction:** Hydrolysis of the (1 $\rightarrow$ 6)-linked  $\alpha$ -D-mannose residues in  $\alpha$ -D-Man $p$ -(1 $\rightarrow$ 6)-D-Man $p$   
**Systematic name:** (1 $\rightarrow$ 6)- $\alpha$ -mannosyl  $\alpha$ -D-mannohydrolase  
**Comments:** The enzyme is specific for (1 $\rightarrow$ 6)-linked mannobiose and has no activity towards any other linkages, or towards *p*-nitrophenyl- $\alpha$ -D-mannopyranoside or baker's yeast mannan. It is strongly inhibited by Mn<sup>2+</sup> but does not require Ca<sup>2+</sup> or any other metal cofactor for activity.  
**References:** [92]

[EC 3.2.1.163 created 2007]

#### EC 3.2.1.164

**Accepted name:** galactan endo-1,6- $\beta$ -galactosidase  
**Reaction:** Endohydrolysis of (1 $\rightarrow$ 6)- $\beta$ -D-galactosidic linkages in arabinogalactan proteins and (1 $\rightarrow$ 3):(1 $\rightarrow$ 6)- $\beta$ -galactans to yield galactose and (1 $\rightarrow$ 6)- $\beta$ -galactobiose as the final products  
**Other name(s):** endo-1,6- $\beta$ -galactanase  
**Systematic name:** endo- $\beta$ -(1 $\rightarrow$ 6)-galactanase

**Comments:** The enzyme specifically hydrolyses 1,6- $\beta$ -D-galactooligosaccharides with a degree of polymerization (DP) higher than 3, and their acidic derivatives with 4-*O*-methylglucosyluronate or glucosyluronate groups at the non-reducing terminals [2188]. 1,3- $\beta$ -D- and 1,4- $\beta$ -D-galactosyl residues cannot act as substrates. The enzyme can also hydrolyse  $\alpha$ -L-arabinofuranosidase-treated arabinogalactan protein (AGP) extracted from radish roots [2188, 1519]. AGPs are thought to be involved in many physiological events, such as cell division, cell expansion and cell death [1519].

**References:** [293, 2188, 1519]

[EC 3.2.1.164 created 2007]

#### EC 3.2.1.165

**Accepted name:** exo-1,4- $\beta$ -D-glucosaminidase

**Reaction:** Hydrolysis of chitosan or chitosan oligosaccharides to remove successive D-glucosamine residues from the non-reducing termini

**Other name(s):** CsxA; GlcNase; exochitosanase; GlmA; exo- $\beta$ -D-glucosaminidase; chitosan exo-1,4- $\beta$ -D-glucosaminidase

**Systematic name:** chitosan exo-(1 $\rightarrow$ 4)- $\beta$ -D-glucosaminidase

**Comments:** Chitosan is a partially or totally *N*-deacetylated chitin derivative that is found in the cell walls of some phytopathogenic fungi and comprises D-glucosamine residues with a variable content of GlcNAc residues [496]. Acts specifically on chitooligosaccharides and chitosan, having maximal activity on chitotetraose, chitopentaose and their corresponding alcohols [2055]. The enzyme can degrade GlcN-GlcNAc but not GlcNAc-GlcNAc [853]. A member of the glycoside hydrolase family 2 (GH-2) [496].

**References:** [2055, 2104, 853, 496, 1228]

[EC 3.2.1.165 created 2008]

#### EC 3.2.1.166

**Accepted name:** heparanase

**Reaction:** endohydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-glycosidic bonds of heparan sulfate chains in heparan sulfate proteoglycan

**Other name(s):** Hpa1 heparanase; Hpa1; heparanase 1; heparanase-1; C1A heparanase; HPSE

**Systematic name:** heparan sulfate *N*-sulfo-D-glucosamine endoglucanase

**Comments:** Heparanase cleaves the linkage between a glucuronic acid unit and an *N*-sulfo glucosamine unit carrying either a 3-*O*-sulfo or a 6-*O*-sulfo group [2265]. Heparanase-1 cuts macromolecular heparin into fragments of 5000–20000 Da [3080]. The enzyme cleaves the heparan sulfate glycosaminoglycans from proteoglycan core proteins and degrades them to small oligosaccharides. Inside cells, the enzyme is important for the normal catabolism of heparan sulfate proteoglycans, generating glycosaminoglycan fragments that are then transported to lysosomes and completely degraded. When secreted, heparanase degrades basement membrane heparan sulfate glycosaminoglycans at sites of injury or inflammation, allowing extravasion of immune cells into nonvascular spaces and releasing factors that regulate cell proliferation and angiogenesis [133].

**References:** [133, 2265, 2282, 2183, 3080, 955, 2956, 1900, 1038]

[EC 3.2.1.166 created 2010]

#### EC 3.2.1.167

**Accepted name:** baicalin- $\beta$ -D-glucuronidase

**Reaction:** baicalin + H<sub>2</sub>O = baicalein + D-glucuronate

**Other name(s):** baicalinase

**Systematic name:** 5,6,7-trihydroxyflavone-7-*O*- $\beta$ -D-glucopyranosiduronate glucuronosylhydrolase

**Comments:** The enzyme also hydrolyses wogonin 7-*O*- $\beta$ -D-glucuronide and oroxylin 7-*O*- $\beta$ -D-glucuronide with lower efficiency [1976]. Negligible activity with *p*-nitrophenyl- $\beta$ -D-glucuronide [3324].

**References:** [1229, 3324, 2534, 1976]

[EC 3.2.1.167 created 2011]

#### EC 3.2.1.168

**Accepted name:** hesperidin 6-*O*- $\alpha$ -L-rhamnosyl- $\beta$ -D-glucosidase  
**Reaction:** hesperidin + H<sub>2</sub>O = hesperetin + rutinose  
**Systematic name:** hesperetin 7-(6-*O*- $\alpha$ -L-rhamnopyranosyl- $\beta$ -D-glucopyranoside) 6-*O*- $\alpha$ -rhamnopyranosyl- $\beta$ -glucohydrolase  
**Comments:** The enzyme exhibits high specificity towards 7-O-linked flavonoid  $\beta$ -rutinosides.  
**References:** [1859, 1860]

[EC 3.2.1.168 created 2011]

#### EC 3.2.1.169

**Accepted name:** protein *O*-GlcNAcase  
**Reaction:** (1) [protein]-3-*O*-(*N*-acetyl- $\beta$ -D-glucosaminy)-L-serine + H<sub>2</sub>O = [protein]-L-serine + *N*-acetyl-D-glucosamine  
(2) [protein]-3-*O*-(*N*-acetyl- $\beta$ -D-glucosaminy)-L-threonine + H<sub>2</sub>O = [protein]-L-threonine + *N*-acetyl-D-glucosamine  
**Other name(s):** OGA; glycoside hydrolase *O*-GlcNAcase; *O*-GlcNAcase; BtGH84; *O*-GlcNAc hydrolase  
**Systematic name:** [protein]-3-*O*-(*N*-acetyl- $\beta$ -D-glucosaminy)-L-serine/threonine *N*-acetylglucosaminy hydrolase  
**Comments:** Within higher eukaryotes post-translational modification of protein serines/threonines with *N*-acetylglucosamine (*O*-GlcNAc) is dynamic, inducible and abundant, regulating many cellular processes by interfering with protein phosphorylation. EC 2.4.1.255 (protein *O*-GlcNAc transferase) transfers GlcNAc onto substrate proteins and EC 3.2.1.169 (protein *O*-GlcNAcase) cleaves GlcNAc from the modified proteins.  
**References:** [873, 3160, 387, 584, 1447, 633]

[EC 3.2.1.169 created 2011]

#### EC 3.2.1.170

**Accepted name:** mannosylglycerate hydrolase  
**Reaction:** 2-*O*-( $\alpha$ -D-mannopyranosyl)-D-glycerate + H<sub>2</sub>O = D-mannopyranose + D-glycerate  
**Other name(s):** MgH  
**Systematic name:** 2-*O*-( $\alpha$ -D-mannopyranosyl)-D-glycerate D-mannohydrolase  
**Comments:** The enzyme occurs in thermophilic bacteria and has been characterized in *Thermus thermophilus* and *Rubrobacter radiotolerans*. It also has been identified in the moss *Selaginella moellendorffii*.  
**References:** [28, 2102]

[EC 3.2.1.170 created 2011, modified 2018]

#### EC 3.2.1.171

**Accepted name:** rhamnogalacturonan hydrolase  
**Reaction:** Endohydrolysis of  $\alpha$ -D-GalA-(1 $\rightarrow$ 2)- $\alpha$ -L-Rha glycosidic bond in the rhamnogalacturonan I backbone with initial inversion of anomeric configuration releasing oligosaccharides with  $\beta$ -D-GalA at the reducing end.  
**Other name(s):** rhamnogalacturonase A; RGase A; RG-hydrolase  
**Systematic name:** rhamnogalacturonan  $\alpha$ -D-GalA-(1 $\rightarrow$ 2)- $\alpha$ -L-Rha hydrolase  
**Comments:** The enzyme is part of the degradation system for rhamnogalacturonan I in *Aspergillus aculeatus*.  
**References:** [2264, 1494, 107, 2263, 2289]

[EC 3.2.1.171 created 2011]

#### EC 3.2.1.172



**Accepted name:** unsaturated rhamnogalacturonyl hydrolase  
**Reaction:** 2-*O*-(4-deoxy- $\beta$ -L-*threo*-hex-4-enopyranuronosyl)- $\alpha$ -L-rhamnopyranose + H<sub>2</sub>O = 5-dehydro-4-deoxy-D-glucuronate + L-rhamnopyranose  
**Other name(s):** YteR; YesR  
**Systematic name:** 2-*O*-(4-deoxy- $\beta$ -L-*threo*-hex-4-enopyranuronosyl)- $\alpha$ -L-rhamnopyranose hydrolase  
**Comments:** The enzyme is part of the degradation system for rhamnogalacturonan I in *Bacillus subtilis* strain 168.  
**References:** [1274, 3330, 1275]

[EC 3.2.1.172 created 2011, modified 2012]

#### EC 3.2.1.173

**Accepted name:** rhamnogalacturonan galacturonohydrolase  
**Reaction:** Exohydrolysis of the  $\alpha$ -D-GalA-(1 $\rightarrow$ 2)- $\alpha$ -L-Rha bond in rhamnogalacturonan oligosaccharides with initial inversion of configuration releasing D-galacturonic acid from the non-reducing end of rhamnogalacturonan oligosaccharides.  
**Other name(s):** RG-galacturonohydrolase  
**Systematic name:** rhamnogalacturonan oligosaccharide  $\alpha$ -D-GalA-(1 $\rightarrow$ 2)- $\alpha$ -L-Rha galacturonohydrolase  
**Comments:** The enzyme is part of the degradation system for rhamnogalacturonan I in *Aspergillus aculeatus*.  
**References:** [2012]

[EC 3.2.1.173 created 2011]

#### EC 3.2.1.174

**Accepted name:** rhamnogalacturonan rhamnohydrolase  
**Reaction:** Exohydrolysis of the  $\alpha$ -L-Rha-(1 $\rightarrow$ 4)- $\alpha$ -D-GalA bond in rhamnogalacturonan oligosaccharides with initial inversion of configuration releasing  $\beta$ -L-rhamnose from the non-reducing end of rhamnogalacturonan oligosaccharides.  
**Other name(s):** RG-rhamnohydrolase; RG  $\alpha$ -L-rhamnopyranohydrolase  
**Systematic name:** rhamnogalacturonan oligosaccharide  $\alpha$ -L-Rha-(1 $\rightarrow$ 4)- $\alpha$ -D-GalA rhamnohydrolase  
**Comments:** The enzyme is part of the degradation system for rhamnogalacturonan I in *Aspergillus aculeatus*.  
**References:** [2289, 2013]

[EC 3.2.1.174 created 2011]

#### EC 3.2.1.175

**Accepted name:**  $\beta$ -D-glucopyranosyl abscisate  $\beta$ -glucosidase  
**Reaction:** D-glucopyranosyl abscisate + H<sub>2</sub>O = D-glucose + abscisate  
**Other name(s):** AtBG1; ABA- $\beta$ -D-glucosidase; ABA-specific  $\beta$ -glucosidase; ABA-GE hydrolase;  $\beta$ -D-glucopyranosyl abscisate hydrolase  
**Systematic name:**  $\beta$ -D-glucopyranosyl abscisate glucohydrolase  
**Comments:** The enzyme hydrolyzes the biologically inactive  $\beta$ -D-glucopyranosyl ester of abscisic acid to produce active abscisate. Abscisate is a phytohormone critical for plant growth, development and adaptation to various stress conditions. The enzyme does not hydrolyse  $\beta$ -D-glucopyranosyl zeatin [1624].  
**References:** [1624, 1403, 610]

[EC 3.2.1.175 created 2011]

#### EC 3.2.1.176

**Accepted name:** cellulose 1,4- $\beta$ -cellobiosidase (reducing end)  
**Reaction:** Hydrolysis of (1 $\rightarrow$ 4)- $\beta$ -D-glucosidic linkages in cellulose and similar substrates, releasing cellobiose from the reducing ends of the chains.  
**Other name(s):** CelS; CelSS; endoglucanase SS; cellulase SS; cellobiohydrolase CelS; Cel48A  
**Systematic name:** 4- $\beta$ -D-glucan cellobiohydrolase (reducing end)



**Comments:** Some exocellulases, most of which belong to the glycoside hydrolase family 48 (GH48, formerly known as cellulase family L), act at the reducing ends of cellulose and similar substrates. The CelS enzyme from *Clostridium thermocellum* is the most abundant subunit of the cellulosome formed by the organism. It liberates cellobiose units from the reducing end by hydrolysis of the glycosidic bond, employing an inverting reaction mechanism [2493]. Different from EC 3.2.1.91, which attacks cellulose from the non-reducing end.

**References:** [150, 2493]

[EC 3.2.1.176 created 2011]

#### EC 3.2.1.177

**Accepted name:**  $\alpha$ -D-xyloside xylohydrolase

**Reaction:** Hydrolysis of terminal, non-reducing  $\alpha$ -D-xylose residues with release of  $\alpha$ -D-xylose.

**Other name(s):**  $\alpha$ -xylosidase

**Systematic name:**  $\alpha$ -D-xyloside xylohydrolase

**Comments:** The enzyme catalyses hydrolysis of a terminal, unsubstituted xyloside at the extreme reducing end of a xylogluco-oligosaccharide. Representative  $\alpha$ -xylosidases from glycoside hydrolase family 31 utilize a two-step (double-displacement) mechanism involving a covalent glycosyl-enzyme intermediate, and retain the anomeric configuration of the product.

**References:** [1961, 2514, 505, 1741, 1222, 2190, 1604]

[EC 3.2.1.177 created 2011]

#### EC 3.2.1.178

**Accepted name:**  $\beta$ -porphyranase

**Reaction:** Hydrolysis of  $\beta$ -D-galactopyranose-(1 $\rightarrow$ 4)- $\alpha$ -L-galactopyranose-6-sulfate linkages in porphyran

**Other name(s):** porphyranase; PorA; PorB; endo- $\beta$ -porphyranase

**Systematic name:** porphyran  $\beta$ -D-galactopyranose-(1 $\rightarrow$ 4)- $\alpha$ -L-galactopyranose-6-sulfate 4-glycanohydrolase

**Comments:** The backbone of porphyran consists largely (70%) of (1 $\rightarrow$ 3)-linked  $\beta$ -D-galactopyranose followed by (1 $\rightarrow$ 4)-linked  $\alpha$ -L-galactopyranose-6-sulfate [the other 30% are mostly agarobiose repeating units of (1 $\rightarrow$ 3)-linked  $\beta$ -D-galactopyranose followed by (1 $\rightarrow$ 4)-linked 3,6-anhydro- $\alpha$ -L-galactopyranose] [489]. This enzyme cleaves the (1 $\rightarrow$ 4) linkages between  $\beta$ -D-galactopyranose and  $\alpha$ -L-galactopyranose-6-sulfate, forming mostly the disaccharide  $\alpha$ -L-galactopyranose-6-sulfate-(1 $\rightarrow$ 3)- $\beta$ -D-galactose, although some longer oligosaccharides of even number of residues are also observed. Since the enzyme is inactive on the non-sulfated agarose portion of the porphyran backbone, some agarose fragments are also included in the products [1098]. Methylation of the D-galactose prevents its binding at position -1 [489].

**References:** [1098, 489]

[EC 3.2.1.178 created 2011]

#### EC 3.2.1.179

**Accepted name:** gellan tetrasaccharide unsaturated glucuronosyl hydrolase

**Reaction:**  $\beta$ -D-4-deoxy- $\Delta^4$ -GlcAp-(1 $\rightarrow$ 4)- $\beta$ -D-Glcp-(1 $\rightarrow$ 4)- $\alpha$ -L-Rhap-(1 $\rightarrow$ 3)-D-Glcp + H<sub>2</sub>O = 5-dehydro-4-deoxy-D-glucuronate +  $\beta$ -D-Glcp-(1 $\rightarrow$ 4)- $\alpha$ -L-Rhap-(1 $\rightarrow$ 3)-D-Glcp

**Other name(s):** UGL (ambiguous); unsaturated glucuronyl hydrolase (ambiguous); gellan tetrasaccharide unsaturated glucuronyl hydrolase

**Systematic name:**  $\beta$ -D-4-deoxy- $\Delta^4$ -GlcAp-(1 $\rightarrow$ 4)- $\beta$ -D-Glcp-(1 $\rightarrow$ 4)- $\alpha$ -L-Rhap-(1 $\rightarrow$ 3)-D-Glcp  $\beta$ -D-4-deoxy- $\Delta^4$ -GlcAp hydrolase

**Comments:** The enzyme releases 4-deoxy-4(5)-unsaturated D-glucuronic acid from oligosaccharides produced by polysaccharide lyases, e.g. the tetrasaccharide  $\beta$ -D-4-deoxy- $\Delta^4$ -GlcAp-(1 $\rightarrow$ 4)- $\beta$ -D-Glcp-(1 $\rightarrow$ 4)- $\alpha$ -L-Rhap-(1 $\rightarrow$ 3)-D-Glcp produced by EC 4.2.2.25, gellan lyase. The enzyme can also hydrolyse unsaturated chondroitin and hyaluronate disaccharides ( $\beta$ -D-4-deoxy- $\Delta^4$ -GlcAp-(1 $\rightarrow$ 3)-D-GalNAc,  $\beta$ -D-4-deoxy- $\Delta^4$ -GlcAp-(1 $\rightarrow$ 3)-D-GalNAc6S,  $\beta$ -D-4-deoxy- $\Delta^4$ -GlcAp2S-(1 $\rightarrow$ 3)-D-GalNAc,  $\beta$ -D-4-deoxy- $\Delta^4$ -GlcAp-(1 $\rightarrow$ 3)-D-GlcNAc), preferring the unsulfated disaccharides to the sulfated disaccharides.

**References:** [1272, 1070, 1273]

[EC 3.2.1.179 created 2011, modified 2016]

#### EC 3.2.1.180

**Accepted name:** unsaturated chondroitin disaccharide hydrolase

**Reaction:**  $\beta$ -D-4-deoxy- $\Delta^4$ -GlcAp-(1 $\rightarrow$ 3)- $\beta$ -D-GalNAc6S + H<sub>2</sub>O = 5-dehydro-4-deoxy-D-glucuronate + N-acetyl- $\beta$ -D-galactosamine-6-O-sulfate

**Other name(s):** UGL (ambiguous); unsaturated glucuronyl hydrolase (ambiguous)

**Systematic name:**  $\beta$ -D-4-deoxy- $\Delta^4$ -GlcAp-(1 $\rightarrow$ 3)- $\beta$ -D-GalNAc6S hydrolase

**Comments:** The enzyme releases 4-deoxy-4,5-didehydro D-glucuronic acid or 4-deoxy-4,5-didehydro L-iduronic acid from chondroitin disaccharides, hyaluronan disaccharides and heparin disaccharides and cleaves both glycosidic (1 $\rightarrow$ 3) and (1 $\rightarrow$ 4) bonds. It prefers the sulfated disaccharides to the unsulfated disaccharides.

**References:** [1828, 2044]

[EC 3.2.1.180 created 2011]

#### EC 3.2.1.181

**Accepted name:** galactan endo- $\beta$ -1,3-galactanase

**Reaction:** The enzyme specifically hydrolyses  $\beta$ -1,3-galactan and  $\beta$ -1,3-galactooligosaccharides

**Other name(s):** endo- $\beta$ -1,3-galactanase

**Systematic name:** arabinogalactan 3- $\beta$ -D-galactanohydrolase

**Comments:** The enzyme from the fungus *Flammulina velutipes* (winter mushroom) hydrolyses the  $\beta$ (1 $\rightarrow$ 3) bonds found in type II plant arabinogalactans, which occur in cell walls of dicots and cereals. The enzyme is an endohydrolase, and requires at least 3 contiguous  $\beta$ -1,3-residues. *cf.* EC 3.2.1.89, arabinogalactan endo- $\beta$ -1,4-galactanase and EC 3.2.1.145, galactan 1,3- $\beta$ -galactosidase.

**References:** [1518]

[EC 3.2.1.181 created 2012]

#### EC 3.2.1.182

**Accepted name:** 4-hydroxy-7-methoxy-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-2-yl glucoside  $\beta$ -D-glucosidase

**Reaction:** (1) (2R)-4-hydroxy-7-methoxy-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-2-yl  $\beta$ -D-glucopyranoside + H<sub>2</sub>O = 2,4-dihydroxy-7-methoxy-2H-1,4-benzoxazin-3(4H)-one + D-glucose

(2) (2R)-4-hydroxy-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-2-yl  $\beta$ -D-glucopyranoside + H<sub>2</sub>O = 2,4-dihydroxy-2H-1,4-benzoxazin-3(4H)-one + D-glucose

**Other name(s):** DIMBOAGlc hydrolase; DIMBOA glucosidase

**Systematic name:** (2R)-4-hydroxy-7-methoxy-3-oxo-3,4-dihydro-2H-1,4-benzoxazin-2-yl  $\beta$ -D-glucopyranoside  $\beta$ -D-glucosidase

**Comments:** The enzyme from *Triticum aestivum* (wheat) has a higher affinity for DIMBOA glucoside than DIMBOA glucoside. With *Secale cereale* (rye) the preference is reversed.

**References:** [2788, 2787, 525, 2087, 2790, 2789]

[EC 3.2.1.182 created 2012]

#### EC 3.2.1.183

**Accepted name:** UDP-*N*-acetylglucosamine 2-epimerase (hydrolysing)  
**Reaction:** UDP-*N*-acetyl- $\alpha$ -D-glucosamine + H<sub>2</sub>O = *N*-acetyl-D-mannosamine + UDP  
**Other name(s):** UDP-*N*-acetylglucosamine 2-epimerase (ambiguous); GNE (gene name); *siaA* (gene name); *neuC* (gene name)  
**Systematic name:** UDP-*N*-acetyl- $\alpha$ -D-glucosamine hydrolase (2-epimerising)  
**Comments:** The enzyme is found in mammalian liver, as well as in some pathogenic bacteria including *Neisseria meningitidis* and *Staphylococcus aureus*. It catalyses the first step of sialic acid (*N*-acetylneuraminic acid) biosynthesis. The initial product formed is the  $\alpha$  anomer, which rapidly mutarotates to a mixture of anomers [440]. The mammalian enzyme is bifunctional and also catalyses EC 2.7.1.60, *N*-acetylmannosamine kinase. *cf.* EC 5.1.3.14, UDP-*N*-acetylglucosamine 2-epimerase (non-hydrolysing).  
**References:** [2756, 440, 247, 2007]

[EC 3.2.1.183 created 2012]

#### EC 3.2.1.184

**Accepted name:** UDP-*N,N'*-diacetylbacillosamine 2-epimerase (hydrolysing)  
**Reaction:** UDP-*N,N'*-diacetylbacillosamine + H<sub>2</sub>O = UDP + 2,4-diacetamido-2,4,6-trideoxy-D-mannopyranose  
**Other name(s):** UDP-Bac2Ac<sub>4</sub>Ac 2-epimerase; NeuC  
**Systematic name:** UDP-*N,N'*-diacetylbacillosamine hydrolase (2-epimerising)  
**Comments:** Requires Mg<sup>2+</sup>. Involved in biosynthesis of legionaminic acid, a nonulosonate derivative that is incorporated by some bacteria into assorted virulence-associated cell surface glycoconjugates. The initial product formed by the enzyme from *Legionella pneumophila*, which incorporates legionaminic acid into the O-antigen moiety of its lipopolysaccharide, is 2,4-diacetamido-2,4,6-trideoxy- $\alpha$ -D-mannopyranose, which rapidly mutarotates to a mixture of anomers [928]. The enzyme from *Campylobacter jejuni*, which incorporates legionaminic acid into flagellin, prefers GDP-*N,N'*-diacetylbacillosamine [2582].  
**References:** [928, 2582]

[EC 3.2.1.184 created 2012]

#### EC 3.2.1.185

**Accepted name:** non-reducing end  $\beta$ -L-arabinofuranosidase  
**Reaction:**  $\beta$ -L-arabinofuranosyl-(1 $\rightarrow$ 2)- $\beta$ -L-arabinofuranose + H<sub>2</sub>O = 2  $\beta$ -L-arabinofuranose  
**Other name(s):** HypBA1  
**Systematic name:**  $\beta$ -L-arabinofuranoside non-reducing end  $\beta$ -L-arabinofuranosidase  
**Comments:** The enzyme, which was identified in the bacterium *Bifidobacterium longum* JCM1217, removes the  $\beta$ -L-arabinofuranose residue from the non-reducing end of multiple substrates, including  $\beta$ -L-arabinofuranosyl-hydroxyproline (Ara-Hyp), Ara<sub>2</sub>-Hyp, Ara<sub>3</sub>-Hyp, and  $\beta$ -L-arabinofuranosyl-(1 $\rightarrow$ 2)-1-*O*-methyl- $\beta$ -L-arabinofuranose. In the presence of 1-alkanols, the enzyme demonstrates transglycosylation activity, retaining the anomeric configuration of the arabinofuranose residue. *cf.* EC 3.2.1.55, non-reducing end  $\alpha$ -L-arabinofuranosidase  
**References:** [852]

[EC 3.2.1.185 created 2013]

#### EC 3.2.1.186

**Accepted name:** protodioscin 26-*O*- $\beta$ -D-glucosidase  
**Reaction:** protodioscin + H<sub>2</sub>O = 26-deglucoprotodioscin + D-glucose  
**Other name(s):** F26G; torvosidase; CSF26G1; furostanol glycoside 26-*O*- $\beta$ -D-glucosidase; furostanol 26-*O*- $\beta$ -D-glucoside glucohydrolase  
**Systematic name:** protodioscin glucohydrolase  
**Comments:** The enzyme has been characterized from the plants *Cheilocostus speciosus* and *Solanum torvum*. It also hydrolyses the 26- $\beta$ -D-glucose group from related steroid glucosides such as protogracillin, torvoside A and torvoside H.

**References:** [1254, 77]

[EC 3.2.1.186 created 2013]

#### EC 3.2.1.187

**Accepted name:** (Ara-f)<sub>3</sub>-Hyp β-L-arabinobiosidase  
**Reaction:** 4-*O*-(β-L-arabinofuranosyl-(1→2)-β-L-arabinofuranosyl-(1→2)-β-L-arabinofuranosyl)-(2*S*,4*S*)-4-hydroxyproline + H<sub>2</sub>O = 4-*O*-(β-L-arabinofuranosyl)-(2*S*,4*S*)-4-hydroxyproline + β-L-arabinofuranosyl-(1→2)-β-L-arabinofuranose  
**Other name(s):** *hypBA2* (gene name); β-L-arabinobiosidase  
**Systematic name:** 4-*O*-(β-L-arabinofuranosyl-(1→2)-β-L-arabinofuranosyl-(1→2)-β-L-arabinofuranosyl)-(2*S*,4*S*)-4-hydroxyproline β-L-arabinofuranosyl-(1→2)-β-L-arabinofuranose hydrolase  
**Comments:** The enzyme, which was identified in the bacterium *Bifidobacterium longum* JCM1217, is specific for (Ara-f)<sub>3</sub>-Hyp, a sugar chain found in hydroxyproline-rich glycoproteins such as extensin and lectin. The enzyme was not able to accept (Ara-f)<sub>2</sub>-Hyp or (Ara-f)<sub>4</sub>-Hyp as substrates. In the presence of 1-alkanols, the enzyme demonstrates transglycosylation activity, retaining the anomeric configuration of the arabinofuranose residue.  
**References:** [851]

[EC 3.2.1.187 created 2013]

#### EC 3.2.1.188

**Accepted name:** avenacosidase  
**Reaction:** avenacoside B + H<sub>2</sub>O = 26-desgluco-avenacoside B + D-glucose  
**Other name(s):** As-P60  
**Systematic name:** avenacoside B 26-β-D-glucohydrolase  
**Comments:** Isolated from oat (*Avena sativa*) seedlings. The product acts as a defense system against fungal infection. Also acts on avenacoside A.  
**References:** [1006, 1005]

[EC 3.2.1.188 created 2013]

#### EC 3.2.1.189

**Accepted name:** dioscin glycosidase (diosgenin-forming)  
**Reaction:** 3-*O*-[α-L-Rha-(1→4)-[α-L-Rha-(1→2)]-β-D-Glc]diosgenin + 3 H<sub>2</sub>O = D-glucose + 2 L-rhamnose + diosgenin  
**Other name(s):** dioscin glycosidase (aglycone-forming)  
**Systematic name:** 3-*O*-[α-L-Rha-(1→4)-[α-L-Rha-(1→2)]-β-D-Glc]diosgenin hydrolase (diosgenin-forming)  
**Comments:** The enzyme is involved in degradation of the steroid saponin dioscin by some fungi of the *Absidia* genus. The enzyme can also hydrolyse 3-*O*-[α-L-Ara-(1→4)-[α-L-Rha-(1→2)]-β-D-Glc]diosgenin into diosgenin and free sugars as the final products. *cf.* EC 3.2.1.190, dioscin glycosidase (3-*O*-β-D-Glc-diosgenin-forming).  
**References:** [829]

[EC 3.2.1.189 created 2013]

#### EC 3.2.1.190

**Accepted name:** dioscin glycosidase (3-*O*-β-D-Glc-diosgenin-forming)  
**Reaction:** 3-*O*-[α-L-Rha-(1→4)-[α-L-Rha-(1→2)]-β-D-Glc]diosgenin + 2 H<sub>2</sub>O = 2 L-rhamnopyranose + diosgenin 3-*O*-β-D-glucofuranoside  
**Other name(s):** dioscin-α-L-rhamnosidase  
**Systematic name:** 3-*O*-[α-L-Rha-(1→4)-[α-L-Rha-(1→2)]-β-D-Glc]diosgenin (3-*O*-β-D-Glc-diosgenin-forming)

**Comments:** The enzyme is involved in the hydrolysis of the steroid saponin dioscin by the digestive system of *Sus scrofa* (pig). cf. EC 3.2.1.189, dioscin glycosidase (diosgenin-forming).

**References:** [2336]

[EC 3.2.1.190 created 2013]

#### EC 3.2.1.191

**Accepted name:** ginsenosidase type III

**Reaction:** a protopanaxadiol-type ginsenoside with two glucosyl residues at position 3 + 2 H<sub>2</sub>O = a protopanaxadiol-type ginsenoside with no glycosidic modification at position 3 + 2 D-glucopyranose (overall reaction)  
(1a) a protopanaxadiol-type ginsenoside with two glucosyl residues at position 3 + H<sub>2</sub>O = a protopanaxadiol-type ginsenoside with one glucosyl residue at position 3 + D-glucopyranose  
(1b) a protopanaxadiol-type ginsenoside with one glucosyl residue at position 3 + H<sub>2</sub>O = a protopanaxadiol-type ginsenoside with no glycosidic modification at position 3 + D-glucopyranose

**Systematic name:** protopanaxadiol-type ginsenoside 3-β-D-hydrolase

**Comments:** Ginsenosidase type III catalyses the sequential hydrolysis of the 3-O-β-D-(1→2)-glucopyranosyl bond followed by hydrolysis of the 3-O-β-D-glucopyranosyl bond of protopanaxadiol ginsenosides. When acting for example on ginsenoside Rb1 the enzyme first generates ginsenoside XVII, and subsequently ginsenoside LXXV.

**References:** [1323, 48, 1167]

[EC 3.2.1.191 created 2014]

#### EC 3.2.1.192

**Accepted name:** ginsenoside Rb1 β-glucosidase

**Reaction:** ginsenoside Rb1 + 2 H<sub>2</sub>O = ginsenoside Rg3 + 2 D-glucopyranose (overall reaction)  
(1a) ginsenoside Rb1 + H<sub>2</sub>O = ginsenoside Rd + D-glucopyranose  
(1b) ginsenoside Rd + H<sub>2</sub>O = ginsenoside Rg3 + D-glucopyranose

**Systematic name:** ginsenoside Rb1 glucohydrolase

**Comments:** Ginsenosidases catalyse the hydrolysis of glycosyl moieties attached to the C-3, C-6 or C-20 position of ginsenosides. They are specific with respect to the nature of the glycosidic linkage, the position and the order in which the linkages are cleaved. Ginsenoside Rb1 β-glucosidase specifically and sequentially hydrolyses the 20-[β-D-glucopyranosyl-(1→6)-β-D glucopyranosyloxy] residues attached to position 20 by first hydrolysing the (1→6)-glucosidic bond to generate ginsenoside Rd as an intermediate, followed by hydrolysis of the remaining 20-O-β-D-glucosidic bond.

**References:** [3252]

[EC 3.2.1.192 created 2014]

#### EC 3.2.1.193

**Accepted name:** ginsenosidase type I

**Reaction:** (1) a protopanaxadiol-type ginsenoside with two glucosyl residues at position 3 + H<sub>2</sub>O = a protopanaxadiol-type ginsenoside with one glucosyl residue at position 3 + D-glucopyranose  
(2) a protopanaxadiol-type ginsenoside with one glucosyl residue at position 3 + H<sub>2</sub>O = a protopanaxadiol-type ginsenoside with no glycosidic modifications at position 3 + D-glucopyranose  
(3) a protopanaxadiol-type ginsenoside with two glycosyl residues at position 20 + H<sub>2</sub>O = a protopanaxadiol-type ginsenoside with a single glucosyl residue at position 20 + a monosaccharide

**Systematic name:** ginsenoside glucohydrolase

**Comments:** Ginsenosidase type I is slightly activated by  $Mg^{2+}$  or  $Ca^{2+}$  [3309]. The enzyme hydrolyses the 3-*O*- $\beta$ -D-(1→2)-glucosidic bond, the 3-*O*- $\beta$ -D-glucopyranosyl bond and the 20-*O*- $\beta$ -D-(1→6)-glycosidic bond of protopanaxadiol-type ginsenosides. It usually leaves a single glucosyl residue attached at position 20 and one or no glucosyl residues at position 3. Starting with a ginsenoside that is glycosylated at both positions (e.g. ginsenoside Rb1, Rb2, Rb3, Rc or Rd), the most common products are ginsenoside F2 and ginsenoside C-K, with low amounts of ginsenoside Rh2.

**References:** [3309]

[EC 3.2.1.193 created 2014]

#### EC 3.2.1.194

**Accepted name:** ginsenosidase type IV

**Reaction:** a protopanaxatriol-type ginsenoside with two glycosyl residues at position 6 + 2 H<sub>2</sub>O = a protopanaxatriol-type ginsenoside with no glycosidic modification at position 6 + D-glucopyranose + a monosaccharide (overall reaction)  
(1a) a protopanaxatriol-type ginsenoside with two glycosyl residues at position 6 + H<sub>2</sub>O = a protopanaxatriol-type ginsenoside with a single glucosyl at position 6 + a monosaccharide  
(1b) a protopanaxatriol-type ginsenoside with a single glucosyl at position 6 + H<sub>2</sub>O = a protopanaxatriol-type ginsenoside with no glycosidic modification at position 6 + D-glucopyranose

**Systematic name:** protopanaxatriol-type ginsenoside 6- $\beta$ -D-glucohydrolase

**Comments:** Ginsenosidase type IV catalyses the sequential hydrolysis of the 6-*O*- $\beta$ -D-(1→2)-glycosidic bond or the 6-*O*- $\alpha$ -D-(1→2)-glycosidic bond in protopanaxatriol-type ginsenosides with a disacchride attached to the C<sub>6</sub> position, followed by the hydrolysis of the remaining 6-*O*- $\beta$ -D-glycosidic bond (e.g. ginsenoside Re → ginsenoside Rg1 → ginsenoside F1).

**References:** [3107, 3106]

[EC 3.2.1.194 created 2014]

#### EC 3.2.1.195

**Accepted name:** 20-*O*-multi-glycoside ginsenosidase

**Reaction:** a protopanaxadiol-type ginsenoside with two glycosyl residues at position 20 + H<sub>2</sub>O = a protopanaxadiol-type ginsenoside with a single glucosyl residue at position 20 + a monosaccharide  
ginsenosidase type II (erroneous)

**Other name(s):**

**Systematic name:** protopanaxadiol-type ginsenoside 20- $\beta$ -D-glucohydrolase

**Comments:** The 20-*O*-multi-glycoside ginsenosidase catalyses the hydrolysis of the 20-*O*- $\alpha$ -(1→6)-glycosidic bond and the 20-*O*- $\beta$ -(1→6)-glycosidic bond of protopanaxadiol-type ginsenosides. The enzyme usually leaves a single glucosyl residue attached at position 20, although it can cleave the remaining glucosyl residue with a lower efficiency. Starting with a ginsenoside that is glycosylated at positions 3 and 20, such as ginsenosides Rb1, Rb2, Rb3 and Rc, the most common product is ginsenoside Rd, with a low amount of ginsenoside Rg3 also formed.

**References:** [3308]

[EC 3.2.1.195 created 2014]

#### EC 3.2.1.196

**Accepted name:** limit dextrin  $\alpha$ -1,6-maltotetraose-hydrolase

**Reaction:** Hydrolysis of (1→6)- $\alpha$ -D-glucosidic linkages to branches with degrees of polymerization of three or four glucose residues in limit dextrin.

**Other name(s):** *glgX* (gene name); glycogen debranching enzyme (ambiguous)

**Systematic name:** glycogen phosphorylase-limit dextrin maltotetraose-hydrolase

**Comments:** This bacterial enzyme catalyses a reaction similar to EC 3.2.1.33, amylo- $\alpha$ -1,6-glucosidase (one of the activities of the eukaryotic glycogen debranching enzyme). However, while EC 3.2.1.33 removes single glucose residues linked by 1,6- $\alpha$ -linkage, and thus requires the additional activity of 4- $\alpha$ -glucanotransferase (EC 2.4.1.25) to act on limit dextrans formed by glycogen phosphorylase (EC 2.4.1.1), this enzyme removes maltotriose and maltotetraose chains that are attached by 1,6- $\alpha$ -linkage to the limit dextrin main chain, generating a debranched limit dextrin without a need for another enzyme.

**References:** [1313, 541, 2716]

[EC 3.2.1.196 created 2016]

#### EC 3.2.1.197

**Accepted name:**  $\beta$ -1,2-mannosidase  
**Reaction:**  $\beta$ -D-mannopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-mannopyranosyl-(1 $\rightarrow$ 2)-D-mannopyranose + H<sub>2</sub>O =  $\beta$ -D-mannopyranosyl-(1 $\rightarrow$ 2)-D-mannopyranose +  $\alpha$ -D-mannopyranose  
**Systematic name:**  $\beta$ -1,2-D-mannoside mannohydrolase  
**Comments:** The enzyme, characterized from multiple bacterial species, catalyses the hydrolysis of terminal, non-reducing D-mannose residues from  $\beta$ -1,2-mannotriose and  $\beta$ -1,2-mannobiose. The mechanism involves anomeric inversion, resulting in the release of  $\alpha$ -D-mannopyranose. Activity with  $\beta$ -1,2-mannotriose or higher oligosaccharides is higher than that with  $\beta$ -1,2-mannobiose.  
**References:** [523, 2083]

[EC 3.2.1.197 created 2016]

#### EC 3.2.1.198

**Accepted name:**  $\alpha$ -mannan endo-1,2- $\alpha$ -mannanase  
**Reaction:** Hydrolysis of the terminal  $\alpha$ -D-mannosyl-(1 $\rightarrow$ 3)- $\alpha$ -D-mannose disaccharide from  $\alpha$ -D-mannosyl-(1 $\rightarrow$ 3)- $\alpha$ -D-mannosyl-(1 $\rightarrow$ 2)- $\alpha$ -D-mannosyl-(1 $\rightarrow$ 2)- $\alpha$ -D-mannosyl side chains in fungal cell wall  $\alpha$ -mannans.  
**Systematic name:**  $\alpha$ -mannan 1,2-[ $\alpha$ -D-mannosyl-(1 $\rightarrow$ 3)- $\alpha$ -D-mannose] hydrolase  
**Comments:** The enzyme, characterized from the gut bacteria *Bacteroides thetaiotaomicron* and *Bacteroides xylinisolvens*, can also catalyse the reaction of EC 3.2.1.130, glycoprotein endo- $\alpha$ -1,2-mannosidase.  
**References:** [1023, 524]

[EC 3.2.1.198 created 2016]

#### EC 3.2.1.199

**Accepted name:** sulfoquinovosidase  
**Reaction:** an 6-sulfo- $\alpha$ -D-quinovosyl diacylglycerol + H<sub>2</sub>O = 6-sulfo-D-quinovose + a 1,2-diacylglycerol  
**Other name(s):** *yihQ* (gene name)  
**Systematic name:** 6-sulfo- $\alpha$ -D-quinovosyl diacylglycerol 6-sulfo-D-quinovohydrolase  
**Comments:** The enzyme, characterized from the bacteria *Escherichia coli* and *Pseudomonas putida*, hydrolyses terminal non-reducing  $\alpha$ -sulfoquinovoside residues in  $\alpha$ -sulfoquinovosyl diacylglycerides and  $\alpha$ -sulfoquinovosyl glycerol.  
**References:** [2629, 2738]

[EC 3.2.1.199 created 2016]

#### EC 3.2.1.200

**Accepted name:** exo-chitinase (non-reducing end)  
**Reaction:** Hydrolysis of *N,N'*-diacetylchitobiose from the non-reducing end of chitin and chitodextrans.  
**Other name(s):** *chiB* (gene name)  
**Systematic name:** (1 $\rightarrow$ 4)-2-acetamido-2-deoxy- $\beta$ -D-glucan diacetylchitobiohydrolase (non-reducing end)



**Comments:** The enzyme hydrolyses the second glycosidic (1→4) linkage from non-reducing ends of chitin and chitodextrin molecules, liberating *N,N'*-diacetylchitobiose disaccharides. *cf.* EC 3.2.1.201, exo-chitinase (reducing end).

**References:** [2873, 1207, 2166, 1007]

[EC 3.2.1.200 created 2017]

#### EC 3.2.1.201

**Accepted name:** exo-chitinase (reducing end)

**Reaction:** Hydrolysis of *N,N'*-diacetylchitobiose from the reducing end of chitin and chitodextrins.

**Other name(s):** *chiA* (gene name)

**Systematic name:** (1→4)-2-acetamido-2-deoxy-β-D-glucan diacetylchitobiohydrolase (reducing end)

**Comments:** The enzyme hydrolyses the second glycosidic (1→4) linkage from reducing ends of chitin and chitodextrin molecules, liberating *N,N'*-diacetylchitobiose disaccharides. *cf.* EC 3.2.1.200, exo-chitinase (non-reducing end).

**References:** [1207, 2036, 1007, 321]

[EC 3.2.1.201 created 2017]

#### EC 3.2.1.202

**Accepted name:** endo-chitodextrinase

**Reaction:** Hydrolysis of chitodextrins, releasing *N,N'*-diacetylchitobiose and small amounts of *N,N',N''*-triacetylchitotriose.

**Other name(s):** endo I (gene name); chitodextrinase (ambiguous); endolytic chitodextrinase; periplasmic chitodextrinase

**Systematic name:** (1→4)-2-acetamido-2-deoxy-β-D-glucan diacetylchitobiohydrolase (endo-cleaving)

**Comments:** The enzyme, characterized from the bacterium *Vibrio furnissii*, is an endo-cleaving chitodextrinase that participates in the chitin catabolic pathway found in members of the *Vibrionaceae*. Unlike EC 3.2.1.14, chitinase, it has no activity on chitin. The smallest substrate is a tetrasaccharide, and the final products are *N,N'*-diacetylchitobiose and small amounts of *N,N',N''*-triacetylchitotriose. *cf.* EC 3.2.1.200, exo-chitinase (non-reducing end), and EC 3.2.1.201, exo-chitinase (reducing end).

**References:** [170, 1432]

[EC 3.2.1.202 created 2017]

#### EC 3.2.1.203

**Accepted name:** carboxymethylcellulase

**Reaction:** Endohydrolysis of (1→4)-β-D-glucosidic linkages in (carboxymethyl)cellulose.

**Other name(s):** CMCase

**Systematic name:** 4-β-D-(carboxymethyl)glucan 4-(carboxymethyl)glucanohydrolase

**Comments:** The enzyme from the acidophilic bacterium *Alicyclobacillus acidocaldarius* is an endo-cleaving hydrolase that cleaves β(1→4)-linked residues. However, it is specific for (carboxymethyl)cellulose and does not act on cellulosic substrates such as avicel.

**References:** [1962]

[EC 3.2.1.203 created 2017]

#### EC 3.2.1.204

**Accepted name:** 1,3-α-isomaltosidase

**Reaction:** cyclobis-(1→6)-α-nigerosyl + 2 H<sub>2</sub>O = 2 isomaltose (overall reaction)

(1a) cyclobis-(1→6)-α-nigerosyl + H<sub>2</sub>O = α-isomaltosyl-(1→3)-isomaltose

(1b) α-isomaltosyl-(1→3)-isomaltose + H<sub>2</sub>O = 2 isomaltose

**Systematic name:** 1,3-α-isomaltohydrolase (configuration-retaining)



**Comments:** The enzyme, characterized from the bacteria *Bacillus* sp. NRRL B-21195 and *Kribbella flavida*, participates in the degradation of starch. The cyclic tetrasaccharide cyclobis-(1→6)- $\alpha$ -nigerosyl is formed from starch extracellularly and imported into the cell, where it is degraded to glucose.

**References:** [1457, 2828]

[EC 3.2.1.204 created 2017]

#### EC 3.2.1.205

**Accepted name:** isomaltose glucohydrolase

**Reaction:** isomaltose + H<sub>2</sub>O =  $\beta$ -D-glucose + D-glucose

**Systematic name:** isomaltose 6- $\alpha$ -glucohydrolase (configuration-inverting)

**Comments:** The enzyme catalyses the hydrolysis of  $\alpha$ -1,6-glycosidic linkages from the non-reducing end of its substrate. Unlike EC 3.2.1.10, oligo-1,6-glycosidase, the enzyme inverts the anomeric configuration of the released residue. The enzyme can also act on panose and maltotriose at a lower rate.

**References:** [2828]

[EC 3.2.1.205 created 2017]

#### EC 3.2.1.206

**Accepted name:** oleuropein  $\beta$ -glucosidase

**Reaction:** oleuropein + H<sub>2</sub>O = oleuropein aglycone + D-glucopyranose

**Other name(s):** *OeGLU* (gene name)

**Systematic name:** oleuropein 2- $\beta$ -D-glucohydrolase

**Comments:** Oleuropein is a glycosylated secoiridoid exclusively biosynthesized by members of the Oleaceae plant family where it is part of a defence system against herbivores. The enzyme also hydrolyses ligstroside and demethyloleuropein.

**References:** [451, 2455, 1008, 1520, 1521]

[EC 3.2.1.206 created 2017]

#### EC 3.2.1.207

**Accepted name:** mannosyl-oligosaccharide  $\alpha$ -1,3-glycosidase

**Reaction:** (1) Glc<sub>2</sub>Man<sub>9</sub>GlcNAc<sub>2</sub>-[protein] + H<sub>2</sub>O = GlcMan<sub>9</sub>GlcNAc<sub>2</sub>-[protein] +  $\beta$ -D-glucopyranose

(2) GlcMan<sub>9</sub>GlcNAc<sub>2</sub>-[protein] + H<sub>2</sub>O = Man<sub>9</sub>GlcNAc<sub>2</sub>-[protein] +  $\beta$ -D-glucopyranose

**Other name(s):** ER glucosidase II;  $\alpha$ -glucosidase II; trimming glucosidase II; ROT2 (gene name); GTB1 (gene name); GANAB (gene name); PRKCSH (gene name)

**Systematic name:** Glc<sub>2</sub>Man<sub>9</sub>GlcNAc<sub>2</sub>-[protein] 3- $\alpha$ -glucohydrolase (configuration-inverting)

**Comments:** This eukaryotic enzyme cleaves off sequentially the two  $\alpha$ -1,3-linked glucose residues from the Glc<sub>2</sub>Man<sub>9</sub>GlcNAc<sub>2</sub> oligosaccharide precursor of immature *N*-glycosylated proteins.

**References:** [2965, 3341, 3190, 1960]

[EC 3.2.1.207 created 2018]

#### EC 3.2.1.208

**Accepted name:** glucosylglycerate hydrolase

**Reaction:** 2-*O*-( $\alpha$ -D-glucopyranosyl)-D-glycerate + H<sub>2</sub>O = D-glucopyranose + D-glycerate

**Other name(s):** GG hydrolase; GgH

**Systematic name:** 2-*O*-( $\alpha$ -D-glucopyranosyl)-D-glycerate D-glucohydrolase

**Comments:** The enzyme has been isolated from nontuberculous mycobacteria (e.g. *Mycobacterium hassiacum*), which accumulate 2-*O*-( $\alpha$ -D-glucopyranosyl)-D-glycerate during growth under nitrogen deprivation.

**References:** [27, 386]

[EC 3.2.1.208 created 2018]

## EC 3.2.2 Hydrolysing *N*-glycosyl compounds

### EC 3.2.2.1

- Accepted name:** purine nucleosidase  
**Reaction:** a purine nucleoside + H<sub>2</sub>O = D-ribose + a purine base  
**Other name(s):** nucleosidase (misleading); purine β-ribosidase; purine nucleoside hydrolase; purine ribonucleosidase; ribonucleoside hydrolase (misleading); nucleoside hydrolase (misleading); *N*-ribosyl purine ribohydrolase; nucleosidase g; *N*-D-ribosylpurine ribohydrolase; inosine-adenosine-guanosine preferring nucleoside hydrolase; purine-specific nucleoside *N*-ribohydrolase; IAG-nucleoside hydrolase; IAG-NH  
**Systematic name:** purine-nucleoside ribohydrolase  
**Comments:** The enzyme from the bacterium *Ochrobactrum anthropi* specifically catalyses the irreversible *N*-riboside hydrolysis of purine nucleosides. Pyrimidine nucleosides, purine and pyrimidine nucleotides, NAD<sup>+</sup>, NADP<sup>+</sup> and nicotinamide mononucleotide are not substrates [2148].  
**References:** [1122, 1362, 2834, 2890, 2237, 2148, 3063, 1858]

[EC 3.2.2.1 created 1961, modified 2006, modified 2011]

### EC 3.2.2.2

- Accepted name:** inosine nucleosidase  
**Reaction:** inosine + H<sub>2</sub>O = D-ribose + hypoxanthine  
**Other name(s):** inosinase; inosine-guanosine nucleosidase  
**Systematic name:** inosine ribohydrolase  
**References:** [1492, 2890]

[EC 3.2.2.2 created 1961]

### EC 3.2.2.3

- Accepted name:** uridine nucleosidase  
**Reaction:** uridine + H<sub>2</sub>O = D-ribose + uracil  
**Other name(s):** uridine hydrolase  
**Systematic name:** uridine ribohydrolase  
**References:** [377]

[EC 3.2.2.3 created 1961]

### EC 3.2.2.4

- Accepted name:** AMP nucleosidase  
**Reaction:** AMP + H<sub>2</sub>O = D-ribose 5-phosphate + adenine  
**Other name(s):** adenylate nucleosidase; adenosine monophosphate nucleosidase  
**Systematic name:** AMP phosphoribohydrolase  
**References:** [1211]

[EC 3.2.2.4 created 1961]

### EC 3.2.2.5

- Accepted name:** NAD<sup>+</sup> glycohydrolase  
**Reaction:** NAD<sup>+</sup> + H<sub>2</sub>O = ADP-D-ribose + nicotinamide  
**Other name(s):** NAD glycohydrolase; nicotinamide adenine dinucleotide glycohydrolase; β-NAD<sup>+</sup> glycohydrolase; DPNase (ambiguous); NAD hydrolase (ambiguous); diphosphopyridine nucleosidase (ambiguous); nicotinamide adenine dinucleotide nucleosidase (ambiguous); NAD nucleosidase (ambiguous); DPN hydrolase (ambiguous); NADase (ambiguous); *nga* (gene name); NAD<sup>+</sup> nucleosidase

**Systematic name:** NAD<sup>+</sup> glycohydrolase  
**Comments:** This enzyme catalyses the hydrolysis of NAD<sup>+</sup>, without associated ADP-ribosyl cyclase activity (unlike the metazoan enzyme EC 3.2.2.6, bifunctional ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase). The enzyme from Group A streptococci has been implicated in the pathogenesis of diseases such as streptococcal toxic shock-like syndrome (STSS) and necrotizing fasciitis. The enzyme from the venom of the snake *Agkistrodon acutus* also catalyses EC 3.6.1.5, apyrase [3329].  
**References:** [750, 995, 3329, 905, 2696]

[EC 3.2.2.5 created 1961, modified 2013]

#### EC 3.2.2.6

**Accepted name:** ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase  
**Reaction:** NAD<sup>+</sup> + H<sub>2</sub>O = ADP-D-ribose + nicotinamide (overall reaction)  
(1a) NAD<sup>+</sup> = cyclic ADP-ribose + nicotinamide  
(1b) cyclic ADP-ribose + H<sub>2</sub>O = ADP-D-ribose  
**Other name(s):** NAD<sup>+</sup> nucleosidase; NADase (ambiguous); DPNase (ambiguous); DPN hydrolase (ambiguous); NAD hydrolase (ambiguous); nicotinamide adenine dinucleotide nucleosidase (ambiguous); NAD glycohydrolase (misleading); NAD nucleosidase (ambiguous); nicotinamide adenine dinucleotide glycohydrolase (misleading); CD38 (gene name); BST1 (gene name)  
**Systematic name:** NAD<sup>+</sup> glycohydrolase (cyclic ADP-ribose-forming)  
**Comments:** This multiunctional enzyme acts on NAD<sup>+</sup>, catalysing both the synthesis and hydrolysis of cyclic ADP-ribose, a calcium messenger that can mobilize intracellular Ca<sup>2+</sup> stores and activate Ca<sup>2+</sup> influx to regulate a wide range of physiological processes. In addition, the enzyme also catalyses EC 2.4.99.20, 2'-phospho-ADP-ribosyl cyclase/2'-phospho-cyclic-ADP-ribose transferase. It is also able to act on β-nicotinamide D-ribonucleotide. *cf.* EC 3.2.2.5, NAD<sup>+</sup> glycohydrolase.  
**References:** [1238, 1193, 2853, 2928, 827, 3250, 1724]

[EC 3.2.2.6 created 1961, modified 2004, modified 2014, modified 2018]

#### EC 3.2.2.7

**Accepted name:** adenosine nucleosidase  
**Reaction:** adenosine + H<sub>2</sub>O = D-ribose + adenine  
**Other name(s):** adenosinase; *N*-ribosyladenine ribohydrolase; adenosine hydrolase; ANase  
**Systematic name:** adenosine ribohydrolase  
**Comments:** Also acts on adenosine *N*-oxide.  
**References:** [1856]

[EC 3.2.2.7 created 1972]

#### EC 3.2.2.8

**Accepted name:** ribosylpyrimidine nucleosidase  
**Reaction:** a pyrimidine nucleoside + H<sub>2</sub>O = D-ribose + a pyrimidine base  
**Other name(s):** *N*-ribosylpyrimidine nucleosidase; pyrimidine nucleosidase; *N*-ribosylpyrimidine ribohydrolase; pyrimidine nucleoside hydrolase; RihB; YeiK; nucleoside ribohydrolase  
**Systematic name:** pyrimidine-nucleoside ribohydrolase  
**Comments:** Also hydrolyses purine D-ribonucleosides, but much more slowly. 2'-, 3'- and 5'-deoxynucleosides are not substrates [910].  
**References:** [2900, 2260, 910, 911]

[EC 3.2.2.8 created 1972]

#### EC 3.2.2.9

**Accepted name:** adenosylhomocysteine nucleosidase

**Reaction:**  $S\text{-adenosyl-L-homocysteine} + \text{H}_2\text{O} = S\text{-(5-deoxy-D-ribose-5-yl)-L-homocysteine} + \text{adenine}$   
**Other name(s):** *S*-adenosylhomocysteine hydrolase (ambiguous); *S*-adenosylhomocysteine nucleosidase; 5'-methyladenosine nucleosidase; *S*-adenosylhomocysteine/5'-methylthioadenosine nucleosidase; Ado-Hcy/MTA nucleosidase  
**Systematic name:** *S*-adenosyl-L-homocysteine homocysteinylribohydrolase  
**Comments:** Also acts on *S*-methyl-5'-thioadenosine to give adenine and *S*-methyl-5-thioribose (*cf.* EC 3.2.2.16, methylthioadenosine nucleosidase).  
**References:** [662, 761]

[EC 3.2.2.9 created 1972, modified 2004]

#### EC 3.2.2.10

**Accepted name:** pyrimidine-5'-nucleotide nucleosidase  
**Reaction:** a pyrimidine 5'-nucleotide +  $\text{H}_2\text{O} = \text{D-ribose 5-phosphate} + \text{a pyrimidine base}$   
**Other name(s):** pyrimidine nucleotide *N*-ribosidase; Pyr5N  
**Systematic name:** pyrimidine-5'-nucleotide phosphoribo(deoxyribo)hydrolase  
**Comments:** Also acts on dUMP, dTMP and dCMP.  
**References:** [1233, 1234]

[EC 3.2.2.10 created 1972]

#### EC 3.2.2.11

**Accepted name:**  $\beta$ -aspartyl-*N*-acetylglucosaminidase  
**Reaction:**  $1\text{-}\beta\text{-aspartyl-}N\text{-acetyl-D-glucosaminylamine} + \text{H}_2\text{O} = \text{L-asparagine} + N\text{-acetyl-D-glucosamine}$   
**Other name(s):**  $\beta$ -aspartylacetylglucosaminidase  
**Systematic name:** 1- $\beta$ -aspartyl-*N*-acetyl-D-glucosaminylamine L-asparaginohydrolase  
**References:** [728]

[EC 3.2.2.11 created 1972]

#### EC 3.2.2.12

**Accepted name:** inosinate nucleosidase  
**Reaction:**  $\text{IMP} + \text{H}_2\text{O} = \text{D-ribose 5-phosphate} + \text{hypoxanthine}$   
**Other name(s):** 5'-inosinate phosphoribohydrolase  
**Systematic name:** IMP phosphoribohydrolase  
**References:** [1554]

[EC 3.2.2.12 created 1972]

#### EC 3.2.2.13

**Accepted name:** 1-methyladenosine nucleosidase  
**Reaction:**  $1\text{-methyladenosine} + \text{H}_2\text{O} = 1\text{-methyladenine} + \text{D-ribose}$   
**Other name(s):** 1-methyladenosine hydrolase  
**Systematic name:** 1-methyladenosine ribohydrolase  
**References:** [2891]

[EC 3.2.2.13 created 1976]

#### EC 3.2.2.14

**Accepted name:** NMN nucleosidase  
**Reaction:**  $\beta\text{-nicotinamide D-ribonucleotide} + \text{H}_2\text{O} = \text{D-ribose 5-phosphate} + \text{nicotinamide}$   
**Other name(s):** NMNase; nicotinamide mononucleotide nucleosidase; nicotinamide mononucleotidase; NMN glycohydrolase; NMNGhase

**Systematic name:** nicotinamide-nucleotide phosphoribohydrolase  
**Comments:** The enzyme is thought to participate in an NAD<sup>+</sup>-salvage pathway. In eukaryotic organisms this activity has been attributed to EC 3.2.2.6, ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase.  
**References:** [56, 1236, 1237]

[EC 3.2.2.14 created 1976, modified 2018]

#### EC 3.2.2.15

**Accepted name:** DNA-deoxyinosine glycosylase  
**Reaction:** Hydrolyses DNA and polynucleotides, releasing free hypoxanthine  
**Other name(s):** DNA(hypoxanthine) glycohydrolase; deoxyribonucleic acid glycosylase; hypoxanthine-DNA glycosylase  
**Systematic name:** DNA-deoxyinosine deoxyribohydrolase  
**References:** [1391]

[EC 3.2.2.15 created 1980, modified 1982, modified 2000]

#### EC 3.2.2.16

**Accepted name:** methylthioadenosine nucleosidase  
**Reaction:** *S*-methyl-5'-thioadenosine + H<sub>2</sub>O = *S*-methyl-5-thio-D-ribose + adenine  
**Other name(s):** 5'-methylthioadenosine nucleosidase; MTA nucleosidase; MeSAdo nucleosidase; methylthioadenosine methylthioribohydrolase  
**Systematic name:** *S*-methyl-5'-thioadenosine adeninehydrolase  
**Comments:** Does not act on *S*-adenosylhomocysteine. *cf.* EC 3.2.2.9 adenosylhomocysteine nucleosidase.  
**References:** [1003]

[EC 3.2.2.16 created 1983, modified 2004]

#### EC 3.2.2.17

**Accepted name:** deoxyribodipyrimidine endonucleosidase  
**Reaction:** Cleaves the *N*-glycosidic bond between the 5'-pyrimidine residue in cyclobutadipyrimidine (in DNA) and the corresponding deoxy-D-ribose residue  
**Other name(s):** pyrimidine dimer DNA-glycosylase; endonuclease V; deoxyribonucleate pyrimidine dimer glycosidase; pyrimidine dimer DNA glycosylase; T<sub>4</sub>-induced UV endonuclease; PD-DNA glycosylase  
**Systematic name:** deoxy-D-ribocyclobutadipyrimidine polynucleotidodeoxyribohydrolase  
**References:** [1069]

[EC 3.2.2.17 created 1983]

[3.2.2.18 Deleted entry. glycopeptide *N*-glycosidase. Now included with EC 3.5.1.52, peptide-*N*<sup>4</sup>-(*N*-acetyl-β-glucosaminyl)asparagine amidase]

[EC 3.2.2.18 created 1984, deleted 1989]

#### EC 3.2.2.19

**Accepted name:** [protein ADP-ribosylarginine] hydrolase  
**Reaction:** (1) protein-*N*<sup>ω</sup>-(ADP-D-ribosyl)-L-arginine + H<sub>2</sub>O = ADP-D-ribose + protein-L-arginine  
(2) *N*<sup>ω</sup>-(ADP-D-ribosyl)-L-arginine + H<sub>2</sub>O = ADP-D-ribose + L-arginine  
**Other name(s):** ADP-ribose-L-arginine cleavage enzyme; ADP-ribosylarginine hydrolase; *N*<sup>ω</sup>-(ADP-D-ribosyl)-L-arginine ADP-ribosylhydrolase; protein-ω-*N*-(ADP-D-ribosyl)-L-arginine ADP-ribosylhydrolase  
**Systematic name:** protein-*N*<sup>ω</sup>-(ADP-D-ribosyl)-L-arginine ADP-ribosylhydrolase  
**Comments:** The enzyme will remove ADP-D-ribose from arginine residues in ADP-ribosylated proteins.  
**References:** [1985, 1986, 1507, 2832, 2164]

[EC 3.2.2.19 created 1989, modified 2004]

#### EC 3.2.2.20

**Accepted name:** DNA-3-methyladenine glycosylase I  
**Reaction:** Hydrolysis of alkylated DNA, releasing 3-methyladenine  
**Other name(s):** deoxyribonucleate 3-methyladenine glycosidase I; 3-methyladenine DNA glycosylase I; DNA-3-methyladenine glycosidase I  
**Systematic name:** alkylated-DNA glycohydrolase (releasing methyladenine and methylguanine)  
**Comments:** Involved in the removal of alkylated bases from DNA in *Escherichia coli* (cf. EC 2.1.1.63 methylated-DNA—[protein]-cysteine *S*-methyltransferase).  
**References:** [725, 1390, 2914]

[EC 3.2.2.20 created 1990, modified 2000]

#### EC 3.2.2.21

**Accepted name:** DNA-3-methyladenine glycosylase II  
**Reaction:** Hydrolysis of alkylated DNA, releasing 3-methyladenine, 3-methylguanine, 7-methylguanine and 7-methyladenine  
**Other name(s):** deoxyribonucleate 3-methyladenine glycosidase II; 3-methyladenine DNA glycosylase II; DNA-3-methyladenine glycosidase II; AlkA  
**Systematic name:** alkylated-DNA glycohydrolase (releasing methyladenine and methylguanine)  
**Comments:** Involved in the removal of alkylated bases from DNA in *Escherichia coli* (cf. EC 2.1.1.63 methylated-DNA—[protein]-cysteine *S*-methyltransferase).  
**References:** [725, 1390, 2417, 2914]

[EC 3.2.2.21 created 1990, modified 2000]

#### EC 3.2.2.22

**Accepted name:** rRNA *N*-glycosylase  
**Reaction:** Hydrolysis of the *N*-glycosylic bond at A-4324 in 28S rRNA from rat ribosomes  
**Other name(s):** ribosomal ribonucleate *N*-glycosidase; nigrin b; RNA *N*-glycosidase; rRNA *N*-glycosidase; ricin; momorcochin-*S*; Mirabilis antiviral protein; momorcochin-*S*; gelonin; saporins  
**Systematic name:** rRNA *N*-glycohydrolase  
**Comments:** Ricin A-chain and related toxins show this activity. Naked rRNA is attacked more slowly than rRNA in intact ribosomes. Naked rRNA from *Escherichia coli* is cleaved at a corresponding position.  
**References:** [704]

[EC 3.2.2.22 created 1990, modified 2000]

#### EC 3.2.2.23

**Accepted name:** DNA-formamidopyrimidine glycosylase  
**Reaction:** Hydrolysis of DNA containing ring-opened 7-methylguanine residues, releasing 2,6-diamino-4-hydroxy-5-(*N*-methyl)formamidopyrimidine  
**Other name(s):** Fapy-DNA glycosylase; deoxyribonucleate glycosidase; 2,6-diamino-4-hydroxy-5-*N*-formamidopyrimidine-DNA glycosylase; 2,6-diamino-4-hydroxy-5(*N*-methyl)formamidopyrimidine-DNA glycosylase; formamidopyrimidine-DNA glycosylase; DNA-formamidopyrimidine glycosidase; Fpg protein  
**Systematic name:** DNA glycohydrolase [2,6-diamino-4-hydroxy-5-(*N*-methyl)formamidopyrimidine releasing]  
**Comments:** May play a significant role in processes leading to recovery from mutagenesis and/or cell death by alkylating agents. Also involved in the GO system responsible for removing an oxidatively damaged form of guanine (7,8-dihydro-8-oxoguanine) from DNA.  
**References:** [255]

[EC 3.2.2.23 created 1990, modified 2000]

#### EC 3.2.2.24

- Accepted name:** ADP-ribosyl-[dinitrogen reductase] hydrolase  
**Reaction:** [dinitrogen reductase]- $N^{10}$ - $\alpha$ -(ADP-D-ribosyl)-L-arginine = ADP-D-ribose + [dinitrogen reductase]-L-arginine  
**Other name(s):** azoferredoxin glycosidase; azoferredoxin-activating enzymes; dinitrogenase reductase-activating glycohydrolase; ADP-ribosyl glycohydrolase; *draG* (gene name)  
**Systematic name:** ADP-D-ribosyl-[dinitrogen reductase] ADP-ribosylhydrolase  
**Comments:** The enzyme restores the activity of EC 1.18.6.1, nitrogenase, by catalysing the removal of ADP-ribose from an arginine residue of the dinitrogenase reductase component of nitrogenase. This activity occurs only when the nitrogenase product, ammonium, is not available. The combined activity of this enzyme and EC 2.4.2.37,  $NAD^+$ -dinitrogen-reductase ADP-D-ribosyltransferase, controls the level of activity of nitrogenase.  
**References:** [779, 1674, 206]

[EC 3.2.2.24 created 1992]

#### EC 3.2.2.25

- Accepted name:** *N*-methyl nucleosidase  
**Reaction:** 7-methylxanthosine +  $H_2O$  = 7-methylxanthine + D-ribose  
**Other name(s):** 7-methylxanthosine nucleosidase; *N*-MeNase; *N*-methyl nucleoside hydrolase; methylpurine nucleosidase  
**Systematic name:** 7-methylxanthosine ribohydrolase  
**Comments:** The enzyme preferentially hydrolyses 3- and 7-methylpurine nucleosides, such as 3-methylxanthosine, 3-methyladenosine and 7-methylguanosine. Hydrolysis of 7-methylxanthosine to form 7-methylxanthine is the second step in the caffeine-biosynthesis pathway.  
**References:** [2066]

[EC 3.2.2.25 created 2007]

#### EC 3.2.2.26

- Accepted name:** fufalosine hydrolase  
**Reaction:** fufalosine +  $H_2O$  = dehypoxanthine fufalosine + hypoxanthine  
**Other name(s):** fufalosine nucleosidase; MqnB  
**Systematic name:** fufalosine ribohydrolase  
**Comments:** This enzyme, which is specific for fufalosine, catalyses the second step of a novel menaquinone biosynthetic pathway that is found in some prokaryotes.  
**References:** [1149]

[EC 3.2.2.26 created 2008]

#### EC 3.2.2.27

- Accepted name:** uracil-DNA glycosylase  
**Reaction:** Hydrolyses single-stranded DNA or mismatched double-stranded DNA and polynucleotides, releasing free uracil  
**Other name(s):** UdgB (ambiguous); uracil-DNA *N*-glycosylase; UDG (ambiguous); uracil DNA glycohydrolase  
**Systematic name:** uracil-DNA deoxyribohydrolase (uracil-releasing)  
**Comments:** Uracil-DNA glycosylases are widespread enzymes that are found in all living organisms. EC 3.2.2.27 and double-stranded uracil-DNA glycosylase (EC 3.2.2.28) form a central part of the DNA-repair machinery since they initiate the DNA base-excision repair pathway by hydrolysing the *N*-glycosidic bond between uracil and the deoxyribose sugar thereby catalysing the removal of mis-incorporated uracil from DNA.  
**References:** [1626, 1448, 2234, 2773]



[EC 3.2.2.27 created 2009]

#### EC 3.2.2.28

- Accepted name:** double-stranded uracil-DNA glycosylase  
**Reaction:** Specifically hydrolyses mismatched double-stranded DNA and polynucleotides, releasing free uracil  
**Other name(s):** Mug; double-strand uracil-DNA glycosylase; Dug; dsUDG; double-stranded DNA specific UDG; dsDNA specific UDG; UdgB (ambiguous); G:T/U mismatch-specific DNA glycosylase; UDG (ambiguous)  
**Systematic name:** uracil-double-stranded DNA deoxyribohydrolase (uracil-releasing)  
**Comments:** No activity on DNA containing a T/G mismatch or single-stranded DNA containing either a site-specific uracil or 3,*N*<sup>4</sup>-ethenocytosine residue [2804], significant role for double-stranded uracil-DNA glycosylase in mutation avoidance in non-dividing *E. coli* [1948]. Uracil-DNA glycosylases are widespread enzymes that are found in all living organisms. Uracil-DNA glycosylase (EC 3.2.2.27) and EC 3.2.2.28 form a central part of the DNA-repair machinery since they initiate the DNA base-excision repair pathway by hydrolysing the *N*-glycosidic bond between uracil and the deoxyribose sugar thereby catalysing the removal of mis-incorporated uracil from DNA.  
**References:** [161, 2804, 1948]

[EC 3.2.2.28 created 2009]

#### EC 3.2.2.29

- Accepted name:** thymine-DNA glycosylase  
**Reaction:** Hydrolyses mismatched double-stranded DNA and polynucleotides, releasing free thymine.  
**Other name(s):** mismatch-specific thymine-DNA glycosylase; mismatch-specific thymine-DNA *N*-glycosylase; hTDG; hsTDG; TDG; thymine DNA glycosylase; G/T glycosylase; uracil/thymine DNA glycosylase; T:G mismatch-specific thymidine-DNA glycosylase; G:T mismatch-specific thymine DNA-glycosylase  
**Systematic name:** thymine-DNA deoxyribohydrolase (thymine-releasing)  
**Comments:** Thymine-DNA glycosylase is part of the DNA-repair machinery. Thymine removal is fastest when it is from a G/T mismatch with a 5'-flanking C/G pair. The glycosylase removes uracil from G/U, C/U, and T/U base pairs faster than it removes thymine from G/T [3134].  
**References:** [3135, 2062, 3134]

[EC 3.2.2.29 created 2009]

#### EC 3.2.2.30

- Accepted name:** aminodeoxyfutilosine nucleosidase  
**Reaction:** 6-amino-6-deoxyfutilosine + H<sub>2</sub>O = dehypoxanthine futilosine + adenine  
**Other name(s):** AFL nucleosidase; aminofutilosine nucleosidase; methylthioadenosine nucleosidase; MqnB  
**Systematic name:** 6-amino-6-deoxyfutilosine ribohydrolase  
**Comments:** The enzyme, found in several bacterial species, catalyses a step in a modified futilosine pathway for menaquinone biosynthesis. While the enzyme from some organisms also has the activity of EC 3.2.2.9, adenosylhomocysteine nucleosidase, the enzyme from *Chlamydia trachomatis* is specific for 6-amino-6-deoxyfutilosine [164].  
**References:** [1149, 1672, 65, 3117, 1927, 1454, 164]

[EC 3.2.2.30 created 2014]

#### EC 3.2.2.31

- Accepted name:** adenine glycosylase  
**Reaction:** Hydrolyses free adenine bases from 7,8-dihydro-8-oxoguanine:adenine mismatched double-stranded DNA, leaving an apurinic site.  
**Other name(s):** *mutY* (gene name); A/G-specific adenine glycosylase



**Systematic name:** adenine-DNA deoxyribohydrolase (adenine-releasing)  
**Comments:** The enzyme serves as a mismatch repair enzyme that works to correct 7,8-dihydro-8-oxoguanine:adenine mispairs that arise in DNA when error-prone synthesis occurs past 7,8-dihydro-8-oxoguanine (GO) lesions in DNA. The enzyme excises the adenine of the mispair, producing an apurinic site sensitive to AP endonuclease activity. After removing the undamaged adenine the enzyme remains bound to the site to prevent EC 3.2.2.23 (MutM) from removing the GO lesion, which could lead to a double strand break. *In vitro* the enzyme is also active with adenine:guanine, adenine:cytosine, and adenine:7,8-dihydro-8-oxoadenine (AO) mispairs, removing the adenine in all cases.  
**References:** [96, 1901]

[EC 3.2.2.31 created 2018]

### EC 3.2.3 Hydrolysing S-glycosyl compounds (deleted sub-subclass)

[3.2.3.1 *Transferred entry. thioglucosidase. Now EC 3.2.1.147, thioglucosidase*]

[EC 3.2.3.1 created 1972, deleted 2001]

## EC 3.3 Acting on ether bonds

This subclass contains enzymes that act on ether bonds. It is subdivided into those hydrolysing thioether and trialkylsulfonium compounds (EC 3.3.1) and those acting on ethers (EC 3.3.2).

### EC 3.3.1 Thioether and trialkylsulfonium hydrolases

#### EC 3.3.1.1

**Accepted name:** adenosylhomocysteinase  
**Reaction:**  $S$ -adenosyl-L-homocysteine + H<sub>2</sub>O = L-homocysteine + adenosine  
**Other name(s):**  $S$ -adenosylhomocysteine synthase;  $S$ -adenosylhomocysteine hydrolase (ambiguous); adenosylhomocysteine hydrolase;  $S$ -adenosylhomocysteinase; SAHase; AdoHcyase  
**Systematic name:**  $S$ -adenosyl-L-homocysteine hydrolase  
**Comments:** The enzyme contains one tightly bound NAD<sup>+</sup> per subunit. This appears to bring about a transient oxidation at C-3' of the 5'-deoxyadenosine residue, thus labilizing the thioether bond [2229] (for mechanism, click here), *cf.* EC 5.5.1.4, inositol-3-phosphate synthase.  
**References:** [559, 2229]

[EC 3.3.1.1 created 1961, modified 2004]

#### EC 3.3.1.2

**Accepted name:**  $S$ -adenosyl-L-methionine hydrolase (L-homoserine-forming)  
**Reaction:**  $S$ -adenosyl-L-methionine + H<sub>2</sub>O = L-homoserine +  $S$ -methyl-5'-thioadenosine  
**Other name(s):**  $S$ -adenosylmethionine cleaving enzyme; methylmethionine-sulfonium-salt hydrolase; adenosylmethionine lyase; adenosylmethionine hydrolase;  $S$ -adenosylmethionine hydrolase;  $S$ -adenosyl-L-methionine hydrolase  
**Systematic name:**  $S$ -adenosyl-L-methionine hydrolase (L-homoserine-forming)  
**Comments:** Also hydrolyses  $S$ -methyl-L-methionine to dimethyl sulfide and homoserine. *cf.* EC 3.13.1.8,  $S$ -adenosyl-L-methionine hydrolase (adenosine-forming).  
**References:** [1857]

[EC 3.3.1.2 created 1972, modified 1976, modified 2018]

[3.3.1.3 Deleted entry. ribosylhomocysteinase. This enzyme was transferred to EC 3.2.1.148, ribosylhomocysteinase, which has since been deleted. The activity is most probably attributable to EC 4.4.1.21, S-ribosylhomocysteine lyase]

[EC 3.3.1.3 created 1972, deleted 2001]

## EC 3.3.2 Ether hydrolases

### EC 3.3.2.1

**Accepted name:** isochorismatase  
**Reaction:** isochorismate + H<sub>2</sub>O = (2*S*,3*S*)-2,3-dihydroxy-2,3-dihydrobenzoate + pyruvate  
**Other name(s):** 2,3-dihydro-2,3-dihydroxybenzoate synthase; 2,3-dihydroxy-2,3-dihydrobenzoate synthase; 2,3-dihydroxy-2,3-dihydrobenzoic synthase  
**Systematic name:** isochorismate pyruvate-hydrolase  
**Comments:** The enzyme is involved in the biosynthesis of several siderophores, such as 2,3-dihydroxybenzoylglycine, enterobactin, bacillibactin, and vibriobactin.  
**References:** [3306]

[EC 3.3.2.1 created 1972]

### EC 3.3.2.2

**Accepted name:** lysoplasmalogenase  
**Reaction:** (1) 1-(1-alkenyl)-*sn*-glycero-3-phosphocholine + H<sub>2</sub>O = an aldehyde + *sn*-glycero-3-phosphocholine  
(2) 1-(1-alkenyl)-*sn*-glycero-3-phosphoethanolamine + H<sub>2</sub>O = an aldehyde + *sn*-glycero-3-phosphoethanolamine  
**Other name(s):** alkenylglycerophosphocholine hydrolase; alkenylglycerophosphoethanolamine hydrolase; 1-(1-alkenyl)-*sn*-glycero-3-phosphocholine aldehydohydrolase  
**Systematic name:** lysoplasmalogen aldehydohydrolase  
**Comments:** Lysoplasmalogenase is specific for the *sn*-2-deacylated (lyso) form of plasmalogen and catalyses hydrolytic cleavage of the vinyl ether bond, releasing a fatty aldehyde and *sn*-glycero-3-phosphocholine or *sn*-glycero-3-phosphoethanolamine.  
**References:** [3129, 691, 1000, 78, 3219]

[EC 3.3.2.2 created 1972, modified 1976, (EC 3.3.2.5 created 1984, incorporated 2016), modified 2016]

[3.3.2.3 Transferred entry. epoxide hydrolase. Now known to comprise two enzymes, microsomal epoxide hydrolase (EC 3.3.2.9) and soluble epoxide hydrolase (EC 3.3.2.10)]

[EC 3.3.2.3 created 1978, modified 1999, deleted 2006]

### EC 3.3.2.4

**Accepted name:** *trans*-epoxysuccinate hydrolase  
**Reaction:** *trans*-2,3-epoxysuccinate + H<sub>2</sub>O = *meso*-tartrate  
**Other name(s):** *trans*-epoxysuccinate hydratase; tartrate epoxydase  
**Systematic name:** *trans*-2,3-epoxysuccinate hydrolase  
**Comments:** Acts on both optical isomers of the substrate.  
**References:** [36]

[EC 3.3.2.4 created 1984]

[3.3.2.5 Transferred entry. alkenylglycerophosphoethanolamine hydrolase. Now included in EC 3.3.2.2, lysoplasmalogenase.]

[EC 3.3.2.5 created 1984, deleted 2016]

### EC 3.3.2.6

- Accepted name:** leukotriene-A<sub>4</sub> hydrolase  
**Reaction:** leukotriene A<sub>4</sub> + H<sub>2</sub>O = leukotriene B<sub>4</sub>  
**Other name(s):** LTA<sub>4</sub> hydrolase; LTA<sub>4</sub>H; leukotriene A<sub>4</sub> hydrolase  
**Systematic name:** (7E,9E,11Z,14Z)-(5S,6S)-5,6-epoxycosa-7,9,11,14-tetraenoate hydrolase  
**Comments:** This is a bifunctional zinc metalloprotease that displays both epoxide hydrolase and aminopeptidase activities [2073, 2201]. It preferentially cleaves tripeptides at an arginyl bond, with dipeptides and tetrapeptides being poorer substrates [2201] (see EC 3.4.11.6, aminopeptidase B). It also converts leukotriene A<sub>4</sub> into leukotriene B<sub>4</sub>, unlike EC 3.3.2.10, soluble epoxide hydrolase, which converts leukotriene A<sub>4</sub> into 5,6-dihydroxy-7,9,11,14-icosatetraenoic acid [1017, 2073]. In vertebrates, five epoxide-hydrolase enzymes have been identified to date: EC 3.3.2.6 (leukotriene A<sub>4</sub> hydrolase), EC 3.3.2.7 (hepoxilin-epoxide hydrolase), EC 3.3.2.9 (microsomal epoxide hydrolase), EC 3.3.2.10 (soluble epoxide hydrolase) and EC 3.3.2.11 (cholesterol-5,6-oxide hydrolase) [812].  
**References:** [724, 1922, 1017, 2073, 812, 2201, 2161]

[EC 3.3.2.6 created 1989, modified 2006]

### EC 3.3.2.7

- Accepted name:** hepoxilin-epoxide hydrolase  
**Reaction:** hepoxilin A<sub>3</sub> + H<sub>2</sub>O = trioxilin A<sub>3</sub>  
**Other name(s):** hepoxilin epoxide hydrolase; hepoxilin hydrolase; hepoxilin A<sub>3</sub> hydrolase  
**Systematic name:** (5Z,9E,14Z)-(8ξ,11R,12S)-11,12-epoxy-8-hydroxyicoso-5,9,14-trienoate hydrolase  
**Comments:** Converts hepoxilin A<sub>3</sub> into trioxilin A<sub>3</sub>. Highly specific for the substrate, having only slight activity with other epoxides such as leukotriene A<sub>4</sub> and styrene oxide [2222]. Hepoxilin A<sub>3</sub> is an hydroxy-epoxide derivative of arachidonic acid that is formed via the 12-lipoxygenase pathway [2222]. It is probable that this enzyme plays a modulatory role in inflammation, vascular physiology, systemic glucose metabolism and neurological function [2073]. In vertebrates, five epoxide-hydrolase enzymes have been identified to date: EC 3.3.2.6 (leukotriene-A<sub>4</sub> hydrolase), EC 3.3.2.7 (hepoxilin-epoxide hydrolase), EC 3.3.2.9 (microsomal epoxide hydrolase), EC 3.3.2.10 (soluble epoxide hydrolase) and EC 3.3.2.11 (cholesterol 5,6-oxide hydrolase) [812].  
**References:** [2221, 2222, 812, 2073]

[EC 3.3.2.7 created 1992, modified 2006]

### EC 3.3.2.8

- Accepted name:** limonene-1,2-epoxide hydrolase  
**Reaction:** 1,2-epoxymenth-8-ene + H<sub>2</sub>O = menth-8-ene-1,2-diol  
**Other name(s):** limonene oxide hydrolase  
**Systematic name:** 1,2-epoxymenth-8-ene hydrolase  
**Comments:** Involved in the monoterpene degradation pathway of the actinomycete *Rhodococcus erythropolis*. The enzyme hydrolyses several alicyclic and 1-methyl-substituted epoxides, such as 1-methylcyclohexene oxide, indene oxide and cyclohexene oxide. It differs from the previously described epoxide hydrolases [EC 3.3.2.4 (*trans*-epoxysuccinate hydrolase), EC 3.3.2.6 (leukotriene-A<sub>4</sub> hydrolase), EC 3.3.2.7 (hepoxilin-epoxide hydrolase), EC 3.3.2.9 (microsomal epoxide hydrolase) and EC 3.3.2.10 (soluble epoxide hydrolase)] as it is not inhibited by 2-bromo-4'-nitroacetophenone, diethyl dicarbonate, 4-fluorochalcone oxide or 1,10-phenanthroline. Both enantiomers of menth-8-ene-1,2-diol [i.e. (1R,2R,4S)-menth-8-ene-1,2-diol and (1S,2S,4R)-menth-8-ene-1,2-diol] are metabolized.  
**References:** [3041, 138, 3042]

[EC 3.3.2.8 created 2001]

### EC 3.3.2.9

- Accepted name:** microsomal epoxide hydrolase  
**Reaction:** (1) *cis*-stilbene oxide + H<sub>2</sub>O = (1R,2R)-1,2-diphenylethane-1,2-diol

(2) 1-(4-methoxyphenyl)-*N*-methyl-*N*-[(3-methyloxetan-3-yl)methyl]methanamine + H<sub>2</sub>O = 2-[(4-methoxyphenyl)methyl](methyl)aminomethyl)-2-methylpropane-1,3-diol

- Other name(s):** microsomal oxirane/oxetane hydrolase; epoxide hydratase (ambiguous); microsomal epoxide hydratase (ambiguous); epoxide hydrase; microsomal epoxide hydrase; arene-oxide hydratase (ambiguous); benzo[*a*]pyrene-4,5-oxide hydratase; benzo(a)pyrene-4,5-epoxide hydratase; aryl epoxide hydrase (ambiguous); *cis*-epoxide hydrolase; mEH; EPHX1 (gene name)
- Systematic name:** *cis*-stilbene-oxide hydrolase
- Comments:** This is a key hepatic enzyme that catalyses the hydrolytic ring opening of oxiranes (epoxides) and oxetanes to give the corresponding diols. The enzyme is involved in the metabolism of numerous substrates including the stereoselective hydrolytic ring opening of 7-oxabicyclo[4.1.0]hepta-2,4-dienes (arene oxides) to the corresponding *trans*-dihydrodiols. The reaction proceeds via a triad mechanism and involves the formation of an hydroxyalkyl-enzyme intermediate. Five epoxide-hydrolase enzymes have been identified in vertebrates to date: EC 3.3.2.6 (leukotriene-A<sub>4</sub> hydrolase), EC 3.3.2.7 (hepoxilin-epoxide hydrolase), EC 3.3.2.9 (microsomal epoxide hydrolase), EC 3.3.2.10 (soluble epoxide hydrolase) and EC 3.3.2.11 (cholesterol-5,6-oxide hydrolase).
- References:** [2142, 1301, 2140, 2141, 1743, 188, 812, 1977, 2073, 2950]

[EC 3.3.2.9 created 2006 (EC 3.3.2.3 created 1978, modified 1999, part incorporated 2006), modified 2017]

### EC 3.3.2.10

- Accepted name:** soluble epoxide hydrolase
- Reaction:** an epoxide + H<sub>2</sub>O = a glycol
- Other name(s):** epoxide hydrase (ambiguous); epoxide hydratase (ambiguous); arene-oxide hydratase (ambiguous); aryl epoxide hydrase (ambiguous); *trans*-stilbene oxide hydrolase; sEH; cytosolic epoxide hydrolase
- Systematic name:** epoxide hydrolase
- Comments:** Catalyses the hydrolysis of *trans*-substituted epoxides, such as *trans*-stilbene oxide, as well as various aliphatic epoxides derived from fatty-acid metabolism [812]. It is involved in the metabolism of arachidonic epoxides (epoxyicosatrienoic acids; EETs) and linoleic acid epoxides. The EETs, which are endogenous chemical mediators, act at the vascular, renal and cardiac levels to regulate blood pressure [1977, 3312]. The enzyme from mammals is a bifunctional enzyme: the C-terminal domain exhibits epoxide-hydrolase activity and the N-terminal domain has the activity of EC 3.1.3.76, lipid-phosphate phosphatase [2074, 507]. Like EC 3.3.2.9, microsomal epoxide hydrolase, it is probable that the reaction involves the formation of an hydroxyalkyl—enzyme intermediate [1977, 1587]. The enzyme can also use leukotriene A<sub>4</sub>, the substrate of EC 3.3.2.6, leukotriene-A<sub>4</sub> hydrolase, but it forms 5,6-dihydroxy-7,9,11,14-icosatetraenoic acid rather than leukotriene B<sub>4</sub> as the product [1017, 2073]. In vertebrates, five epoxide-hydrolase enzymes have been identified to date: EC 3.3.2.6 (leukotriene-A<sub>4</sub> hydrolase), EC 3.3.2.7 (hepoxilin-epoxide hydrolase), EC 3.3.2.9 (microsomal epoxide hydrolase), EC 3.3.2.10 (soluble epoxide hydrolase) and EC 3.3.2.11 (cholesterol 5,6-oxide hydrolase) [812].
- References:** [2074, 507, 2140, 1977, 3312, 1587, 812, 3320, 1017, 2073]

[EC 3.3.2.10 created 2006 (EC 3.3.2.3 created 1978, part incorporated 2006)]

### EC 3.3.2.11

- Accepted name:** cholesterol-5,6-oxide hydrolase
- Reaction:** (1) 5,6 $\alpha$ -epoxy-5 $\alpha$ -cholestan-3 $\beta$ -ol + H<sub>2</sub>O = 5 $\alpha$ -cholestane-3 $\beta$ ,5 $\alpha$ ,6 $\beta$ -triol  
(2) 5,6 $\beta$ -epoxy-5 $\beta$ -cholestan-3 $\beta$ -ol + H<sub>2</sub>O = 5 $\alpha$ -cholestane-3 $\beta$ ,5 $\alpha$ ,6 $\beta$ -triol
- Other name(s):** cholesterol-epoxide hydrolase; ChEH
- Systematic name:** 5,6 $\alpha$ -epoxy-5 $\alpha$ -cholestan-3 $\beta$ -ol hydrolase
- Comments:** The enzyme appears to work equally well with either epoxide as substrate [2614]. The product is a competitive inhibitor of the reaction. In vertebrates, five epoxide-hydrolase enzymes have been identified to date: EC 3.3.2.6 (leukotriene-A<sub>4</sub> hydrolase), EC 3.3.2.7 (hepoxilin-epoxide hydrolase), EC 3.3.2.9 (microsomal epoxide hydrolase), EC 3.3.2.10 (soluble epoxide hydrolase) and EC 3.3.2.11 (cholesterol 5,6-oxide hydrolase) [2614].
- References:** [1655, 2143, 2614, 812, 2073]

[EC 3.3.2.11 created 2006]

#### EC 3.3.2.12

- Accepted name:** oxepin-CoA hydrolase  
**Reaction:** 2-oxepin-2(3*H*)-ylideneacetyl-CoA + H<sub>2</sub>O = 3-oxo-5,6-dehydrosuberyl-CoA semialdehyde  
**Other name(s):** *paaZ* (gene name)  
**Systematic name:** 2-oxepin-2(3*H*)-ylideneacetyl-CoA hydrolase  
**Comments:** The enzyme from *Escherichia coli* is a bifunctional fusion protein that also catalyses EC 1.17.1.7, 3-oxo-5,6-dehydrosuberyl-CoA semialdehyde dehydrogenase. Combined the two activities result in a two-step conversion of oxepin-CoA to 3-oxo-5,6-dehydrosuberyl-CoA, part of an aerobic phenylacetate degradation pathway [1,3,4]. The enzyme from *Escherichia coli* also exhibits enoyl-CoA hydratase activity utilizing crotonyl-CoA as a substrate [2235].  
**References:** [758, 2235, 1265, 2905]

[EC 3.3.2.12 created 2011 as EC 3.7.1.16, transferred 2013 to EC 3.3.2.12]

#### EC 3.3.2.13

- Accepted name:** chorismatase  
**Reaction:** chorismate + H<sub>2</sub>O = (4*R*,5*R*)-4,5-dihydroxycyclohexa-1(6),2-diene-1-carboxylate + pyruvate  
**Other name(s):** chorismate/3,4-dihydroxycyclohexa-1,5-dienoate synthase; *fkbO* (gene name); *rapK* (gene name)  
**Systematic name:** chorismate pyruvate-hydrolase  
**Comments:** The enzyme found in several bacterial species is involved in the biosynthesis of macrocyclic polyketides.  
**References:** [54, 1348]

[EC 3.3.2.13 created 2013]

#### EC 3.3.2.14

- Accepted name:** 2,4-dinitroanisole *O*-demethylase  
**Reaction:** 2,4-dinitroanisole + H<sub>2</sub>O = methanol + 2,4-dinitrophenol  
**Other name(s):** 2,4-dinitroanisole ether hydrolase; *dnhA* (gene name); *dnhB* (gene name); DNAN demethylase  
**Systematic name:** 2,4-dinitroanisole methanol hydrolase  
**Comments:** The enzyme, characterized from the bacterium *Nocardioides* sp. JS1661, is involved in the degradation of 2,4-dinitroanisole. Unlike other known *O*-demethylases, such as EC 1.14.99.15, 4-methoxybenzoate monooxygenase (*O*-demethylating), or EC 1.14.11.32, codeine 3-*O*-demethylase, it does not require oxygen or electron donors, and produces methanol rather than formaldehyde.  
**References:** [762]

[EC 3.3.2.14 created 2015]

#### EC 3.3.2.15

- Accepted name:** *trans*-2,3-dihydro-3-hydroxyanthranilic acid synthase  
**Reaction:** (2*S*)-2-amino-4-deoxychorismate + H<sub>2</sub>O = (5*S*,6*S*)-6-amino-5-hydroxycyclohexa-1,3-diene-1-carboxylate + pyruvate  
**Other name(s):** isochorismatase (ambiguous); *phzD* (gene name)  
**Systematic name:** (2*S*)-2-amino-4-deoxychorismate pyruvate-hydrolase  
**Comments:** Isolated from the bacterium *Pseudomonas aeruginosa*. Involved in phenazine biosynthesis.  
**References:** [1852, 2240]

[EC 3.3.2.15 created 2016]

## EC 3.4 Acting on peptide bonds (peptidases)

It is recommended that the term "peptidase" be used as being synonymous with "peptide hydrolase" for any enzyme that hydrolyses peptide bonds. Peptidases are recommended to be further divided into "exo-peptidases" that act only near a terminus of a polypeptide chain and "endo-peptidases" that act internally in polypeptide chains. The types of exo-peptidases and endo-peptidases are described more fully below. The usage of "peptidase", which is now recommended, is synonymous with "protease" as it was originally used [1] as a general term for both exo-peptidases and endo-peptidases, but it should be noted that previously, in Enzyme Nomenclature (1984), "peptidase" was restricted to the enzymes included in sub-subclasses EC 3.4.11 and EC 3.4.13-19, the exo-peptidases. Also, the term "proteinase" used previously for the enzymes included in sub-subclasses EC 3.4.21-25 carried the same meaning as "endo-peptidase", and has been replaced by "endo-peptidase", for consistency.¶

The nomenclature of the peptidases is troublesome. Their specificity is commonly difficult to define, depending upon the nature of several amino-acid residues around the peptide bond to be hydrolysed and also on the conformation of the substrate's polypeptide chain. A classification involving the additional criterion of catalytic mechanism is therefore used.¶

Two sets of sub-subclasses of peptidases are recognized, those of the exo-peptidases (EC 3.4.11 and EC 3.4.13-19) and those of the endo-peptidases (EC 3.4.21-25). The exo-peptidases act only near the ends of polypeptide chains, and those acting at a free N-terminus liberate a single amino-acid residue (amino-peptidases; EC 3.4.11), or a dipeptide or a tripeptide (dipeptidyl-peptidases and tripeptidyl-peptidases; EC 3.4.14). The exo-peptidases that act at a free C-terminus liberate a single residue (carboxypeptidases, EC 3.4.16-18), or a dipeptide (peptidyl-dipeptidases; EC 3.4.15). The carboxypeptidases are allocated to three groups on the basis of catalytic mechanism: the serine-type carboxypeptidases (EC 3.4.16), the metallo-carboxypeptidases (EC 3.4.17) and the cysteine-type carboxypeptidases (EC 3.4.18). Other exo-peptidases are specific for dipeptides (dipeptidases, EC 3.4.13), or for removal of terminal residues that are substituted, cyclized or linked by isopeptide bonds (peptide linkages other than those of alpha-carboxyl to alpha-amino groups) (omega peptidases; EC 3.4.19).¶

The endo-peptidases are divided into sub-subclasses on the basis of catalytic mechanism, and specificity is used only to identify individual enzymes within the groups. The sub-subclasses are: serine endo-peptidases (EC 3.4.21), cysteine endo-peptidases (EC 3.4.22), aspartic endo-peptidases (EC 3.4.23), metallo-endo-peptidases (EC 3.4.24) and threonine endo-peptidases (EC 3.4.25). ¶

There are characteristic inhibitors of the members of each catalytic type of endo-peptidase; to save space, these have not been listed separately for each individual enzyme but are reviewed in [2] and [3]. A general source of information on peptidases that similarly has not been cited for each individual enzyme is reference [4].¶

In describing the specificity of peptidases, use is made of a model in which the catalytic site is considered to be flanked on one or both sides by specificity subsites, each able to accommodate the sidechain of a single amino-acid residue (based on [5]). These sites are numbered from the catalytic site, S1...Sn towards the N-terminus of the substrate, and S1'...Sn' towards the C-terminus. The residues they accommodate are numbered P1...Pn, and P1'...Pn', respectively, as follows:¶

Substrate: - P3 - P2 - P1 — P1' - P2' - P3' - ;¶

Enzyme: - S3 - S2 - S1 \* S1' - S2' - S3' - ;¶

In this representation, the catalytic site of the enzyme is marked by an asterisk (\*). The peptide bond cleaved (the scissile bond) is indicated by the symbol '—' or a hyphen in the structural formula of the substrate, or a hyphen in the name of the enzyme.¶

Finally, in describing the specificity of endo-peptidases, the term oligopeptidase' is used to refer to those that act optimally on substrates smaller than proteins.¶

Families of peptidases are referred to by use of the numbering system of Rawlings & Barrett [6,7].¶

### References;¶

1. Grassmann, W. & Dyckerhoff, H. Über die Proteinase und die Polypeptidase der Hefe. 13. Abhandlung über Pflanzenproteasen in der von R. Willstätter und Mitarbeitern begonnenen Untersuchungsreihe. Hoppe-Seyler's Z. Physiol. Chem. 179 (1928) 41-78.

2. Barrett, A. J. & Salvesen, G. S. (eds) Proteinase Inhibitors, Elsevier Science Publishers, Amsterdam (1986).

3. Beynon, R. J. & Bond, J. S. Proteolytic Enzymes. A Practical Approach. IRL Press, Oxford (1989).

4. Barrett, A. J., Rawlings, N. D. & Woessner, J. F. (eds) Handbook of Proteolytic Enzymes, Academic Press, London (1998).

5. Berger, A. and Schechter, I. Mapping the active site of papain with the aid of peptide substrates and inhibitors. Philos. Trans. R. Soc. London, Ser. B Biol. Sci. 257 (1970) 249-264.

6. Rawlings, N.D. and Barrett, A.J. In: Methods Enzymol. 244 (1994) 19-61 and 461-486; Methods Enzymol. 248 (1995) 105-120 and 183-228.

7. Rawlings, N. D. and Barrett, A. J. MEROPS: the peptidase database. *Nucleic Acids Res.* 27 (1999) 325-331.

### **EC 3.4.1 $\alpha$ -Amino-acyl-peptide hydrolases (deleted sub-subclass)**

[3.4.1.1 *Transferred entry. leucyl aminopeptidase. Now EC 3.4.11.1, leucyl aminopeptidase*]

[EC 3.4.1.1 created 1961, deleted 1972]

[3.4.1.2 *Transferred entry. aminopeptidase. Now EC 3.4.11.2, membrane alanyl aminopeptidase*]

[EC 3.4.1.2 created 1961, deleted 1972]

[3.4.1.3 *Transferred entry. aminotripeptidase. Now EC 3.4.11.4, tripeptide aminopeptidase*]

[EC 3.4.1.3 created 1961, deleted 1972]

[3.4.1.4 *Transferred entry. proline iminopeptidase. Now EC 3.4.11.5, prolyl aminopeptidase*]

[EC 3.4.1.4 created 1965, deleted 1972]

### **EC 3.4.2 Peptidyl-amino-acid hydrolases (deleted sub-subclass)**

[3.4.2.1 *Transferred entry. carboxypeptidase A. Now EC 3.4.17.1, carboxypeptidase A*]

[EC 3.4.2.1 created 1961, deleted 1972]

[3.4.2.2 *Transferred entry. carboxypeptidase B. Now EC 3.4.17.2, carboxypeptidase B*]

[EC 3.4.2.2 created 1961, deleted 1972]

[3.4.2.3 *Transferred entry. yeast carboxypeptidase. Now EC 3.4.17.4, Gly-Xaa carboxypeptidase*]

[EC 3.4.2.3 created 1961, deleted 1972]

### **EC 3.4.3 Dipeptide hydrolases (deleted sub-subclass)**

[3.4.3.1 *Transferred entry. glycyl-glycine dipeptidase. Now EC 3.4.13.18, cytosol nonspecific dipeptidase*]

[EC 3.4.3.1 created 1961, deleted 1972]

[3.4.3.2 *Transferred entry. glycyl-leucine dipeptidase. Now EC 3.4.13.18, cytosol nonspecific dipeptidase*]

[EC 3.4.3.2 created 1961, deleted 1972]

[3.4.3.3 *Transferred entry. aminoacyl-histidine dipeptidase. Now EC 3.4.13.3, Xaa-His dipeptidase*]

[EC 3.4.3.3 created 1961, deleted 1972]

[3.4.3.4 *Transferred entry. aminoacyl-methylhistidine dipeptidase. Now EC 3.4.13.5, Xaa-methyl-His dipeptidase*]

[EC 3.4.3.4 created 1961, deleted 1972]

[3.4.3.5 *Transferred entry. cysteinylglycine dipeptidase. Now EC 3.4.11.2, membrane alanyl aminopeptidase*]

[EC 3.4.3.5 created 1961, deleted 1972]

[3.4.3.6 *Transferred entry. iminodipeptidase. Now EC 3.4.13.18, cytosol nonspecific dipeptidase*]

[EC 3.4.3.6 created 1961, deleted 1972]

[3.4.3.7 *Transferred entry. iminodipeptidase. Now EC 3.4.13.9, Xaa-Pro dipeptidase*]

[EC 3.4.3.7 created 1961, deleted 1972]

### EC 3.4.4 Peptidyl peptide hydrolases (deleted sub-subclass)

- [3.4.4.1 *Transferred entry. pepsin. Now EC 3.4.23.1, pepsin A*  
[EC 3.4.4.1 created 1961, deleted 1972]
- [3.4.4.2 *Transferred entry. pepsin B. Now EC 3.4.23.2, pepsin B*  
[EC 3.4.4.2 created 1961, deleted 1972]
- [3.4.4.3 *Transferred entry. rennin. Now EC 3.4.23.4, chymosin*  
[EC 3.4.4.3 created 1961, deleted 1972]
- [3.4.4.4 *Transferred entry. trypsin. Now EC 3.4.21.4, trypsin*  
[EC 3.4.4.4 created 1961, deleted 1972]
- [3.4.4.5 *Transferred entry. chymotrypsin. Now EC 3.4.21.1, chymotrypsin*  
[EC 3.4.4.5 created 1961, deleted 1972]
- [3.4.4.6 *Transferred entry. chymotrypsin B. Now EC 3.4.21.1, chymotrypsin*  
[EC 3.4.4.6 created 1961, deleted 1972]
- [3.4.4.7 *Transferred entry. elastase. Now covered by EC 3.4.21.36, pancreatic elastase and EC 3.4.21.37, leukocyte elastase*  
[EC 3.4.4.7 created 1961, deleted 1972]
- [3.4.4.8 *Transferred entry. enteropeptidase. Now EC 3.4.21.9, enteropeptidase*  
[EC 3.4.4.8 created 1961, deleted 1972]
- [3.4.4.9 *Transferred entry. cathepsin C. Now EC 3.4.14.1, dipeptidyl-peptidase I*  
[EC 3.4.4.9 created 1961, deleted 1972]
- [3.4.4.10 *Transferred entry. papain. Now EC 3.4.22.2, papain*  
[EC 3.4.4.10 created 1961, deleted 1972]
- [3.4.4.11 *Transferred entry. chymopapain. Now EC 3.4.22.6, chymopapain*  
[EC 3.4.4.11 created 1961, deleted 1972]
- [3.4.4.12 *Transferred entry. ficin. Now EC 3.4.22.3, ficain*  
[EC 3.4.4.12 created 1961, deleted 1972]
- [3.4.4.13 *Transferred entry. thrombin. Now EC 3.4.21.5, thrombin*  
[EC 3.4.4.13 created 1961, deleted 1972]
- [3.4.4.14 *Transferred entry. plasmin. Now EC 3.4.21.7, plasmin*  
[EC 3.4.4.14 created 1961, deleted 1972]
- [3.4.4.15 *Transferred entry. renin. Now EC 3.4.23.15, renin*  
[EC 3.4.4.15 created 1961, deleted 1972]
- [3.4.4.16 *Transferred entry. subtiloepitidase A. Now covered by the microbial serine proteinases EC 3.4.21.62 (subtilisin), EC 3.4.21.63 (oryzin), EC 3.4.21.64 (endopeptidase K), EC 3.4.21.65 (thermomycolin), EC 3.4.21.66 (thermitase) and EC 3.4.21.67 (ndopeptidase So)*  
[EC 3.4.4.16 created 1961, deleted 1972]



[3.4.4.17 *Transferred entry. aspergillopeptidase A. Now covered by the microbial aspartic proteinases EC 3.4.23.20 (penicillopepsin), EC 3.4.23.21 (rhizopuspepsin), EC 3.4.23.22 (endothiapepsin), EC 3.4.23.23 (mucorpepsin), EC 3.4.23.24 (candidapepsin), EC 3.4.23.25 (saccharopepsin), EC 3.4.23.26 (rhodotorulapepsin), EC 3.4.21.103 (physarolisin), EC 3.4.23.28 (acrocyllindropepsin), EC 3.4.23.29 (polyporopepsin) and EC 3.4.23.30 (pyncporopepsin)]*

[EC 3.4.4.17 created 1961, deleted 1972]

[3.4.4.18 *Transferred entry. streptococcus peptidase A. Now EC 3.4.22.10, streptopain]*

[EC 3.4.4.18 created 1961, deleted 1972]

[3.4.4.19 *Transferred entry. clostridiopeptidase A. Now EC 3.4.24.3, microbial collagenase]*

[EC 3.4.4.19 created 1961, deleted 1972]

[3.4.4.20 *Transferred entry. clostridiopeptidase B. Now EC 3.4.22.8, clostripain]*

[EC 3.4.4.20 created 1961, deleted 1972]

[3.4.4.21 *Transferred entry. kallikrein. Now EC 3.4.21.34 (plasma kallikrein) and EC 3.4.21.35 (tissue kallikrein)]*

[EC 3.4.4.21 created 1965, deleted 1972]

[3.4.4.22 *Transferred entry. now EC 3.4.23.3, gastricsin]*

[EC 3.4.4.22 created 1965, deleted 1972]

[3.4.4.23 *Transferred entry. now EC 3.4.23.5, cathepsin D]*

[EC 3.4.4.23 created 1965, deleted 1972]

[3.4.4.24 *Transferred entry. now covered by EC 3.4.22.32 (stem bromelain) and EC 3.4.22.33 (fruit bromelain)]*

[EC 3.4.4.24 created 1965, deleted 1972]

[3.4.4.25 *Deleted entry. Streptomyces alkalophilic keratinase]*

[EC 3.4.4.25 created 1965, deleted 1972]

## EC 3.4.11 Aminopeptidases

### EC 3.4.11.1

**Accepted name:** leucyl aminopeptidase

**Reaction:** Release of an N-terminal amino acid, Xaa—Yaa-, in which Xaa is preferably Leu, but may be other amino acids including Pro although not Arg or Lys, and Yaa may be Pro. Amino acid amides and methyl esters are also readily hydrolysed, but rates on arylamides are exceedingly low

**Other name(s):** leucine aminopeptidase; leucyl peptidase; peptidase S; cytosol aminopeptidase; cathepsin III; L-leucine aminopeptidase; leucinaminopeptidase; leucinamide aminopeptidase; FTBL proteins; pro-teinates FTBL; aminopeptidase II; aminopeptidase III; aminopeptidase I

**Comments:** A zinc enzyme isolated from pig kidney and cattle lens; activated by heavy metal ions. Type example of peptidase family M17.

**References:** [1142, 574, 3047]

[EC 3.4.11.1 created 1961 as EC 3.4.1.1, transferred 1972 to EC 3.4.11.1]

### EC 3.4.11.2

**Accepted name:** membrane alanyl aminopeptidase

**Reaction:** Release of an N-terminal amino acid, Xaa—Yaa- from a peptide, amide or arylamide. Xaa is preferably Ala, but may be most amino acids including Pro (slow action). When a terminal hydrophobic residue is followed by a prolyl residue, the two may be released as an intact Xaa-Pro dipeptide

**Other name(s):** microsomal aminopeptidase; aminopeptidase M; aminopeptidase N; particle-bound aminopeptidase; amino-oligopeptidase; alanine aminopeptidase; membrane aminopeptidase I; pseudo leucine aminopeptidase; alanyl aminopeptidase; alanine-specific aminopeptidase; cysteinylglycine dipeptidase; cysteinylglycinase; L-alanine aminopeptidase; CD13

**Comments:** A zinc enzyme, not activated by heavy metal ions. Type example of peptidase family M1.

**References:** [3081, 1458, 968, 2679, 756]

[EC 3.4.11.2 created 1961 as EC 3.4.1.2, transferred 1972 to EC 3.4.11.2 (EC 3.4.13.6 created 1961 as EC 3.4.3.5, transferred 1972 to EC 3.4.13.6, incorporated 1997)]

### EC 3.4.11.3

**Accepted name:** cystinyl aminopeptidase

**Reaction:** Release of an N-terminal amino acid, Cys—Xaa-, in which the half-cystine residue is involved in a disulfide loop, notably in oxytocin or vasopressin. Hydrolysis rates on a range of aminoacyl arylamides exceed that for the cystinyl derivative, however [4]

**Other name(s):** cystyl-aminopeptidase; oxytocinase; cystine aminopeptidase; L-cystine aminopeptidase; oxytocin peptidase; vasopressinase

**Comments:** A zinc-containing sialoglycoprotein in peptidase family M1 (membrane alanyl aminopeptidase family)

**References:** [2676, 2677, 3285, 2507]

[EC 3.4.11.3 created 1972]

### EC 3.4.11.4

**Accepted name:** tripeptide aminopeptidase

**Reaction:** Release of the N-terminal residue from a tripeptide

**Other name(s):** tripeptidase; aminotripeptidase; aminoexotripeptidase; lymphopeptidase; imidoendopeptidase; peptidase B; alanine-phenylalanine-proline arylamidase; peptidase T

**Comments:** A zinc enzyme, widely distributed in mammalian tissues. Formerly EC 3.4.1.3

**References:** [641, 2486]

[EC 3.4.11.4 created 1961 as EC 3.4.1.3, transferred 1972 to EC 3.4.11.4]

### EC 3.4.11.5

**Accepted name:** prolyl aminopeptidase

**Reaction:** Release of N-terminal proline from a peptide

**Other name(s):** proline aminopeptidase; Pro-X aminopeptidase; cytosol aminopeptidase V; proline iminopeptidase

**Comments:** A Mn<sup>2+</sup>-requiring enzyme present in the cytosol of mammalian and microbial cells. In contrast to the mammalian form, the bacterial form of the enzyme (type example of peptidase family S33) hydrolyses both polyproline and prolyl-2-naphthylamide. The mammalian enzyme, which is not specific for prolyl bonds, is possibly identical with EC 3.4.11.1, leucyl aminopeptidase.

**References:** [2530, 2112, 3004]

[EC 3.4.11.5 created 1965 as EC 3.4.1.4, transferred 1972 to EC 3.4.11.5]

### EC 3.4.11.6

**Accepted name:** aminopeptidase B

**Reaction:** Release of N-terminal Arg and Lys from oligopeptides when P1' is not Pro. Also acts on arylamides of Arg and Lys

**Other name(s):** arylamidase II; arginine aminopeptidase; arginyl aminopeptidase; Cl<sup>-</sup>-activated arginine aminopeptidase; cytosol aminopeptidase IV; L-arginine aminopeptidase

**Comments:** Cytosolic or membrane-associated enzyme from mammalian tissues, activated by chloride ions and low concentrations of thiol compounds. This is one of the activities of the bifunctional enzyme EC 3.3.2.6 (membrane alanyl aminopeptidase family) [855, 349].

**References:** [869, 187, 350, 855, 349, 2201]

[EC 3.4.11.6 created 1972, modified 1997]

#### EC 3.4.11.7

**Accepted name:** glutamyl aminopeptidase

**Reaction:** Release of N-terminal glutamate (and to a lesser extent aspartate) from a peptide

**Other name(s):** aminopeptidase A; aspartate aminopeptidase; angiotensinase A; glutamyl peptidase; Ca<sup>2+</sup>-activated glutamate aminopeptidase; membrane aminopeptidase II; antigen BP-1/6C3 of mouse B lymphocytes; L-aspartate aminopeptidase; angiotensinase A2

**Comments:** Ca<sup>2+</sup>-activated and generally membrane-bound. A zinc-metallopeptidase in family M1 (membrane alanyl aminopeptidase family)

**References:** [929, 446, 536, 2926, 3221]

[EC 3.4.11.7 created 1972]

[3.4.11.8 *Transferred entry. pyroglutamyl aminopeptidase. Now EC 3.4.19.3, pyroglutamyl-peptidase I*]

[EC 3.4.11.8 created 1972, deleted 1981]

#### EC 3.4.11.9

**Accepted name:** Xaa-Pro aminopeptidase

**Reaction:** Release of any N-terminal amino acid, including proline, that is linked to proline, even from a dipeptide or tripeptide

**Other name(s):** proline aminopeptidase; aminopeptidase P; aminoacylproline aminopeptidase; X-Pro aminopeptidase

**Comments:** A Mn<sup>2+</sup>-dependent, generally membrane-bound enzyme present in both mammalian and bacterial cells. In peptidase family M24 (methionyl aminopeptidase family)

**References:** [3269, 3268, 782, 2199, 1172]

[EC 3.4.11.9 created 1972]

#### EC 3.4.11.10

**Accepted name:** bacterial leucyl aminopeptidase

**Reaction:** Release of an N-terminal amino acid, preferentially leucine, but not glutamic or aspartic acids

**Other name(s):** *Aeromonas proteolytica* aminopeptidase

**Comments:** A zinc enzyme. Forms of the enzyme have been isolated from *Aeromonas proteolytica*, *Escherichia coli* and *Streptococcus thermophilus*. Examples are known from peptidase families M17 and M28 (of leucyl aminopeptidase and aminopeptidase Y, respectively)

**References:** [2323, 604, 2347]

[EC 3.4.11.10 created 1972]

[3.4.11.11 *Deleted entry. aminopeptidase*]

[EC 3.4.11.11 created 1978, deleted 1992]

[3.4.11.12 *Deleted entry. thermophilic aminopeptidase*]

[EC 3.4.11.12 created 1978, deleted 1997]

#### EC 3.4.11.13

**Accepted name:** clostridial aminopeptidase

**Reaction:** Release of any N-terminal amino acid, including proline and hydroxyproline, but no cleavage of Xaa-Pro-

**Other name(s):** *Clostridium histolyticum* aminopeptidase

**Comments:** A secreted enzyme from *Clostridium histolyticum*, requiring Mn<sup>2+</sup> or Co<sup>2+</sup>

**References:** [1427, 1428, 1429]

[EC 3.4.11.13 created 1978]

**EC 3.4.11.14**

**Accepted name:** cytosol alanyl aminopeptidase

**Reaction:** Release of an N-terminal amino acid, preferentially alanine, from a wide range of peptides, amides and arylamides

**Other name(s):** arylamidase; aminopolypeptidase; thiol-activated aminopeptidase; human liver aminopeptidase; puromycin-sensitive aminopeptidase; soluble alanyl aminopeptidase; cytosol aminopeptidase III; alanine aminopeptidase

**Comments:** A puromycin-sensitive,  $\text{Co}^{2+}$ -activated zinc-sialoglycoprotein that is generally cytosolic. Multiple forms are widely distributed in mammalian tissues and body fluids. In peptidase family M1 (membrane alanyl aminopeptidase family)

**References:** [2755, 1387, 2654]

[EC 3.4.11.14 created 1978]

**EC 3.4.11.15**

**Accepted name:** aminopeptidase Y

**Reaction:** Preferentially, release of N-terminal lysine

**Other name(s):** aminopeptidase Co; aminopeptidase (cobalt-activated); lysyl aminopeptidase

**Comments:** Requires  $\text{Co}^{2+}$ ; inhibited by  $\text{Zn}^{2+}$  and  $\text{Mn}^{2+}$ . An enzyme best known from *Saccharomyces cerevisiae* that hydrolyses Lys-NHPhNO<sub>2</sub> and, more slowly, Arg-NHPhNO<sub>2</sub>. Type example of peptidase family M28

**References:** [8, 3270, 2097]

[EC 3.4.11.15 created 1989, modified 1997]

**EC 3.4.11.16**

**Accepted name:** Xaa-Trp aminopeptidase

**Reaction:** Release of a variety of N-terminal residues (especially glutamate and leucine) from peptides, provided tryptophan (or at least phenylalanine or tyrosine) is the penultimate residue. Also acts on Glu-Trp, Leu-Trp and a number of other dipeptides

**Other name(s):** aminopeptidase W; aminopeptidase X-Trp; X-Trp aminopeptidase

**Comments:** A glycoprotein containing  $\text{Zn}^{2+}$ , from renal and intestinal brush border membranes

**References:** [885, 886]

[EC 3.4.11.16 created 1989]

**EC 3.4.11.17**

**Accepted name:** tryptophanyl aminopeptidase

**Reaction:** Preferential release of N-terminal tryptophan

**Other name(s):** tryptophan aminopeptidase; L-tryptophan aminopeptidase

**Comments:** From *Trichosporon cutaneum*. Also acts on L-tryptophanamide. Requires  $\text{Mn}^{2+}$

**References:** [1283]

[EC 3.4.11.17 created 1989]

**EC 3.4.11.18**

**Accepted name:** methionyl aminopeptidase

**Reaction:** Release of N-terminal amino acids, preferentially methionine, from peptides and arylamides

**Other name(s):** methionine aminopeptidase; peptidase M; L-methionine aminopeptidase; MAP

**Comments:** This membrane-bound enzyme, which is present in both prokaryotes and eukaryotes, releases the initiator methionine from nascent peptides. The activity is dependent on the identity of the second, third and fourth amino acid residues of the target protein, but in general the enzyme acts only when the penultimate residue is small and uncharged (e.g. Gly, Ala, Cys, Ser, Thr, and Val).

**References:** [3298, 2985, 806, 189, 2446]

[EC 3.4.11.18 created 1990]

#### EC 3.4.11.19

**Accepted name:** D-stereospecific aminopeptidase

**Reaction:** Release of an N-terminal D-amino acid from a peptide, Xaa-Yaa-, in which Xaa is preferably D-Ala, D-Ser or D-Thr. D-Amino acid amides and methyl esters also are hydrolysed, as is glycine amide

**Other name(s):** D-aminopeptidase

**Comments:** Known from the bacterium *Ochrobactrum anthropi*. In peptidase family S12 (D-Ala-D-Ala carboxypeptidase family) [82]

**References:** [83, 82]

[EC 3.4.11.19 created 1993]

#### EC 3.4.11.20

**Accepted name:** aminopeptidase Ey

**Reaction:** Differs from other aminopeptidases in broad specificity for amino acids in the P1 position and the ability to hydrolyse peptides of four or five residues that contain Pro in the P1' position

**Comments:** A zinc glycoprotein in peptidase family M1 (membrane alanyl aminopeptidase family), composed of two 150 kDa subunits. From the plasma fraction of hen egg yolk

**References:** [1220, 2875, 2874]

[EC 3.4.11.20 created 1995]

#### EC 3.4.11.21

**Accepted name:** aspartyl aminopeptidase

**Reaction:** Release of an N-terminal aspartate or glutamate from a peptide, with a preference for aspartate

**Comments:** Aminoacyl-arylamides are poor substrates. This is an abundant cytosolic enzyme in mammalian cells, in peptidase family M18 of aminopeptidase I

**References:** [1416, 3186]

[EC 3.4.11.21 created 2000]

#### EC 3.4.11.22

**Accepted name:** aminopeptidase I

**Reaction:** Release of an N-terminal amino acid, preferably a neutral or hydrophobic one, from a polypeptide. Aminoacyl-arylamides are poor substrates

**Other name(s):** aminopeptidase III; aminopeptidase yscI; leucine aminopeptidase IV; yeast aminopeptidase I

**Comments:** A 640-kDa, dodecameric enzyme best known as the major vacuolar aminopeptidase of yeast, *Saccharomyces cerevisiae*, in which species it was first given the name aminopeptidase I (one), amongst others. Activity is stimulated by both  $Zn^{2+}$  and  $Cl^{-}$  ions. Type example of peptidase family M18

**References:** [1330, 1892, 398, 2137]

[EC 3.4.11.22 created 1997]

#### EC 3.4.11.23

**Accepted name:** PepB aminopeptidase

**Reaction:** Release of an N-terminal amino acid, Xaa, from a peptide or arylamide. Xaa is preferably Glu or Asp but may be other amino acids, including Leu, Met, His, Cys and Gln  
**Other name(s):** *Salmonella enterica* serovar Typhimurium peptidase B  
**Comments:** A 270-kDa protein composed of six 46.3-kDa subunits. The pH optimum is in the alkaline range and activity is stimulated by KCl. In peptidase family M17.  
**References:** [1835]

[EC 3.4.11.23 created 2003]

#### EC 3.4.11.24

**Accepted name:** aminopeptidase S  
**Reaction:** Release of an N-terminal amino acid with a preference for large hydrophobic amino-terminus residues  
**Other name(s):** Mername-AA022 peptidase; SGAP; aminopeptidase (*Streptomyces griseus*); *Streptomyces griseus* aminopeptidase; *S. griseus* AP; double-zinc aminopeptidase  
**Comments:** Aminopeptidases are associated with many biological functions, including protein maturation, protein degradation, cell-cycle control and hormone-level regulation [72, 863]. This enzyme contains two zinc molecules in its active site and is activated by Ca<sup>2+</sup> [863]. In the presence of Ca<sup>2+</sup>, the best substrates are Leu-Phe, Leu-Ser, Leu-pNA (aminoacyl-*p*-nitroanilide), Phe-Phe-Phe and Phe-Phe [72]. Peptides with proline in the P1' position are not substrates [72]. Belongs in peptidase family M28.  
**References:** [2744, 190, 72, 863, 921]

[EC 3.4.11.24 created 2008]

#### EC 3.4.11.25

**Accepted name:** β-peptidyl aminopeptidase  
**Reaction:** Cleaves N-terminal β-homoamino acids from peptides composed of 2 to 6 amino acids  
**Other name(s):** BapA (ambiguous)  
**Comments:** *Sphingosinicella xenopeptidilytica* strain 3-2W4 is able to utilize the β-peptides β-homoVal-β-homoAla-β-homoLeu and β-homoAla-β-homoLeu as sole carbon and energy sources [898].  
**References:** [1095, 898, 897, 1094]

[EC 3.4.11.25 created 2011]

#### EC 3.4.11.26

**Accepted name:** intermediate cleaving peptidase 55  
**Reaction:** The enzyme cleaves the Pro<sup>36</sup>-Pro<sup>37</sup> bond of cysteine desulfurase (EC 2.8.1.7) removing three amino acid residues (Tyr-Ser-Pro) from the N-terminus after cleavage by mitochondrial processing peptidase.  
**Other name(s):** Icp55; mitochondrial intermediate cleaving peptidase 55 kDa  
**Comments:** Icp55 removes the destabilizing N-terminal amino acid residues that are left after cleavage by the mitochondrial processing peptidase, leading to the stabilisation of the substrate. The enzyme can remove single amino acids or a short peptide, as in the case of cysteine desulfurase (EC 2.8.1.7), where three amino acids are removed.  
**References:** [2020, 3077]

[EC 3.4.11.26 created 2011]

### EC 3.4.12 Peptidylamino-acid hydrolases or acylamino-acid hydrolases (deleted sub-subclass)

[3.4.12.1 Transferred entry. now EC 3.4.16.5 (carboxypeptidase C) and EC 3.4.16.6 (carboxypeptidase D)]

[EC 3.4.12.1 created 1972, deleted 1978]

[3.4.12.2 Transferred entry. now EC 3.4.17.1, carboxypeptidase A]

[EC 3.4.12.2 created 1972, deleted 1978]

[3.4.12.3 *Transferred entry. now EC 3.4.17.2, carboxypeptidase B*]

[EC 3.4.12.3 created 1972, deleted 1978]

[3.4.12.4 *Transferred entry. now EC 3.4.16.2, lysosomal Pro-Xaa carboxypeptidase*]

[EC 3.4.12.4 created 1972, modified 1976, deleted 1978]

[3.4.12.5 *Transferred entry. now EC 3.5.1.28, N-acetylmuramoyl-L-alanine amidase*]

[EC 3.4.12.5 created 1972, deleted 1978]

[3.4.12.6 *Transferred entry. now EC 3.4.17.8, muramoyl-pentapeptidase carboxypeptidase*]

[EC 3.4.12.6 created 1972, deleted 1978]

[3.4.12.7 *Transferred entry. now EC 3.4.17.3, lysine carboxypeptidase*]

[EC 3.4.12.7 created 1972, deleted 1978]

[3.4.12.8 *Transferred entry. now EC 3.4.17.4, Gly-Xaa carboxypeptidase*]

[EC 3.4.12.8 created 1972, deleted 1978]

[3.4.12.9 *Deleted entry. aspartate carboxypeptidase*]

[EC 3.4.12.9 created 1972, deleted 1978]

[3.4.12.10 *Transferred entry. now EC 3.4.19.9,  $\gamma$ -glutamyl hydrolase*]

[EC 3.4.12.10 created 1972, modified 1976, deleted 1978]

[3.4.12.11 *Transferred entry. now EC 3.4.17.6, alanine carboxypeptidase*]

[EC 3.4.12.11 created 1972, deleted 1978]

[3.4.12.12 *Transferred entry. now EC 3.4.16.5 (carboxypeptidase C) and EC 3.4.16.6 (carboxypeptidase D)*]

[EC 3.4.12.12 created 1972, deleted 1978]

[3.4.12.13 *Deleted entry.  $\gamma$ -glutamylglutamate carboxypeptidase*]

[EC 3.4.12.13 created 1975, modified 1976, deleted 1978]

### EC 3.4.13 Dipeptidases

[3.4.13.1 *Transferred entry. glycyl-glycine dipeptidase. Now EC 3.4.13.18, cytosol nonspecific dipeptidase*]

[EC 3.4.13.1 created 1972, deleted 1978 [transferred to EC 3.4.13.11, deleted 1992]]

[3.4.13.2 *Transferred entry. glycyl-leucine dipeptidase. Now EC 3.4.13.18, cytosol nonspecific dipeptidase*]

[EC 3.4.13.2 created 1972, deleted 1978 [transferred to EC 3.4.13.11, deleted 1992]]

[3.4.13.3 *Deleted entry. Xaa-His dipeptidase. The activity is covered by EC 3.4.13.18, cytosol nonspecific dipeptidase and EC 3.4.13.20,  $\beta$ -Ala-His dipeptidase.*]

[EC 3.4.13.3 created 1961 as EC 3.4.3.3, transferred 1972 to EC 3.4.13.3, modified 1989 (EC 3.4.13.13 created 1981, incorporated 1992), deleted 2011]

#### EC 3.4.13.4

**Accepted name:** Xaa-Arg dipeptidase

**Reaction:** Preferential hydrolysis of Xaa-Arg, Xaa-Lys or Xaa-ornithine dipeptides

**Other name(s):** aminoacyl-lysine dipeptidase;  $N^2$ -(4-amino-butyl)-L-lysine hydrolase; X-Arg dipeptidase

**Comments:** Widely distributed in mammals

**References:** [1552]

[EC 3.4.13.4 created 1972]

#### EC 3.4.13.5

**Accepted name:** Xaa-methyl-His dipeptidase

**Reaction:** Hydrolysis of anserine ( $\beta$ -alanyl- $\mid$ - $N^{\pi}$ -methyl-L-histidine), carnosine, homocarnosine, glycyl- $\mid$ -leucine and other dipeptides with broad specificity

**Other name(s):** anserinase; aminoacyl-methylhistidine dipeptidase; acetylhistidine deacetylase; *N*-acetylhistidine deacetylase;  $\alpha$ -*N*-acetyl-L-histidine aminohydrolase; X-methyl-His dipeptidase

**References:** [1338, 168, 1645]

[EC 3.4.13.5 created 1961 as EC 3.4.3.4, transferred 1972 to EC 3.4.13.5, modified 1981 (EC 3.5.1.34 created 1972, incorporated 1981)]

[3.4.13.6 *Transferred entry. Cys-Gly dipeptidase. Now EC 3.4.11.2, membrane alanyl aminopeptidase*]

[EC 3.4.13.6 created 1961 as EC 3.4.3.5, transferred 1972 to EC 3.4.13.6]

#### EC 3.4.13.7

**Accepted name:** Glu-Glu dipeptidase

**Reaction:** Hydrolysis of the Glu- $\mid$ -Glu dipeptide

**Other name(s):**  $\alpha$ -glutamyl-glutamate dipeptidase; glutamylglutamic arylamidase

**Comments:** It is unclear whether the specificity of this enzyme extends to other  $\alpha$ -glutamyl dipeptides

**References:** [2322]

[EC 3.4.13.7 created 1972]

[3.4.13.8 *Transferred entry. Pro-X dipeptidase. Now EC 3.4.13.18, cytosol nonspecific dipeptidase*]

[EC 3.4.13.8 created 1961 as EC 3.4.3.6, transferred 1972 to EC 3.4.13.8]

#### EC 3.4.13.9

**Accepted name:** Xaa-Pro dipeptidase

**Reaction:** Hydrolysis of Xaa- $\mid$ -Pro dipeptides; also acts on aminoacyl-hydroxyproline analogs. No action on Pro-Pro

**Other name(s):** prolidase; imidodipeptidase; proline dipeptidase; peptidase D;  $\gamma$ -peptidase; X-Pro dipeptidase

**Comments:** A  $Mn^{2+}$ -activated enzyme, in peptidase family M24 (methionyl aminopeptidase family); cytosolic from most animal tissues.

**References:** [549, 2680, 127, 311]

[EC 3.4.13.9 created 1961 as EC 3.4.3.7, transferred 1972 to EC 3.4.13.9]

[3.4.13.10 *Transferred entry.  $\beta$ -aspartyl dipeptidase. Now EC 3.4.19.5,  $\beta$ -aspartyl-peptidase*]

[EC 3.4.13.10 created 1972, deleted 1992]

[3.4.13.11 *Transferred entry. dipeptidase. Now EC 3.4.13.19, membrane dipeptidase*]

[EC 3.4.13.11 created 1972, deleted 1992]

#### EC 3.4.13.12

**Accepted name:** Met-Xaa dipeptidase

**Reaction:** Hydrolysis of Met- $\mid$ -Xaa dipeptides

**Other name(s):** methionyl dipeptidase; dipeptidase M; Met-X dipeptidase

**Comments:** A  $Mn^{2+}$ -activated *Escherichia coli* enzyme with thiol dependence

**References:** [309]



[EC 3.4.13.12 created 1976]

[3.4.13.13 *Transferred entry. homocarnosinase. Now EC 3.4.13.3, X-His dipeptidase*]

[EC 3.4.13.13 created 1981, deleted 1992]

[3.4.13.14 *Deleted entry.  $\gamma$ -glutamyl dipeptidase*]

[EC 3.4.13.14 created 1989, deleted 1992]

[3.4.13.15 *Transferred entry.  $N^2$ - $\beta$ -alanylarginine dipeptidase. Now EC 3.4.13.18, cytosol nonspecific dipeptidase*]

[EC 3.4.13.15 created 1989, deleted 1992]

[3.4.13.16 *Deleted entry. aspartylphenylalanine dipeptidase*]

[EC 3.4.13.16 created 1989, deleted 1992]

#### EC 3.4.13.17

**Accepted name:** non-stereospecific dipeptidase

**Reaction:** Hydrolysis of dipeptides containing either D- or L-amino acids or both

**Other name(s):** peptidyl-D-amino acid hydrolase; D-(or L-)aminoacyl-dipeptidase

**Comments:** A digestive enzyme of cephalopods

**References:** [535]

[EC 3.4.13.17 created 1990]

#### EC 3.4.13.18

**Accepted name:** cytosol nonspecific dipeptidase

**Reaction:** Hydrolysis of dipeptides, preferentially hydrophobic dipeptides including prolyl amino acids

**Other name(s):**  $N^2$ - $\beta$ -alanylarginine dipeptidase; glycyl-glycine dipeptidase; glycyl-leucine dipeptidase; iminodipeptidase; peptidase A; Pro-X dipeptidase; prolinase; prolyl dipeptidase; prolylglycine dipeptidase; iminodipeptidase; prolinase; L-prolylglycine dipeptidase; prolylglycine dipeptidase; diglycinase; Gly-Leu hydrolase; glycyl-L-leucine dipeptidase; glycyl-L-leucine hydrolase; glycyl-L-leucine peptidase; L-amino-acyl-L-amino-acid hydrolase; glycyllucine peptidase; glycyllucine hydrolase; glycyllucine dipeptide hydrolase; non-specific dipeptidase; human cytosolic non-specific dipeptidase; glycyl-L-leucine hydrolase; glycyl-glycine dipeptidase

**Comments:** A zinc enzyme with broad specificity, varying somewhat with source species. Activated and stabilized by dithiothreitol and  $Mn^{2+}$ . Inhibited by bestatin and leucine.

**References:** [173]

[EC 3.4.13.18 created 1961 as EC 3.4.3.1 and EC 3.4.3.2, transferred 1972 to EC 3.4.13.1 and EC 3.4.13.2, transferred 1978 to EC 3.4.13.11, part transferred 1992 to EC 3.4.13.18, modified 2000 (EC 3.4.13.15 created 1989, incorporated 1992)]

#### EC 3.4.13.19

**Accepted name:** membrane dipeptidase

**Reaction:** Hydrolysis of dipeptides

**Other name(s):** renal dipeptidase; dehydropeptidase I (DPH I); dipeptidase (ambiguous); aminodipeptidase; dipeptide hydrolase (ambiguous); dipeptidyl hydrolase (ambiguous); nonspecific dipeptidase; glycosyl-phosphatidylinositol-anchored renal dipeptidase; MDP

**Comments:** A membrane-bound, zinc enzyme with broad specificity. Abundant in the kidney cortex. Inhibited by bestatin and cilastatin. Type example of peptidase family M19.

**References:** [359, 360, 1539, 1173]

[EC 3.4.13.19 created 1961 as EC 3.4.3.1 and EC 3.4.3.2, transferred 1972 to EC 3.4.13.1 and EC 3.4.13.2, transferred 1978 to EC 3.4.13.11, part transferred 1992 to EC 3.4.13.19, modified 2011]

#### EC 3.4.13.20

- Accepted name:**  $\beta$ -Ala-His dipeptidase  
**Reaction:** Preferential hydrolysis of the  $\beta$ -Ala—His dipeptide (carnosine), and also anserine, Xaa—His dipeptides and other dipeptides including homocarnosine  
**Other name(s):** serum carnosinase  
**Comments:** Present in the serum of humans and higher primates, but not in the serum of other mammals. Activated by  $\text{Cd}^{2+}$  and citrate. Belongs in peptidase family M20.  
**References:** [1646, 1290]

[EC 3.4.13.20 created 1992]

#### EC 3.4.13.21

- Accepted name:** dipeptidase E  
**Reaction:** Dipeptidase E catalyses the hydrolysis of dipeptides Asp—Xaa. It does not act on peptides with N-terminal Glu, Asn or Gln, nor does it cleave isoaspartyl peptides  
**Other name(s):** aspartyl dipeptidase; peptidase E; PepE gene product (*Salmonella typhimurium*)  
**Comments:** A free carboxy group is not absolutely required in the substrate since Asp-Phe-NH<sub>2</sub> and Asp-Phe-OMe are hydrolysed somewhat more slowly than dipeptides with free C-termini. No peptide larger than a C-blocked dipeptide is known to be a substrate. Asp-NH-Np is hydrolysed and is a convenient substrate for routine assay. The enzyme is most active near pH 7.0, and is not inhibited by diisopropylfluorophosphate or phenylmethanesulfonyl fluoride. Belongs in peptidase family S51.  
**References:** [1012, 1609]

[EC 3.4.13.21 created 2001]

#### EC 3.4.13.22

- Accepted name:** D-Ala-D-Ala dipeptidase  
**Reaction:** D-Ala-D-Ala + H<sub>2</sub>O = 2 D-Ala  
**Other name(s):** D-alanyl-D-alanine dipeptidase; *vanX* D-Ala-D-Ala dipeptidase; VanX  
**Comments:** A Zn<sup>2+</sup>-dependent enzyme [335]. The enzyme protects *Enterococcus faecium* from the antibiotic vancomycin, which can bind to the -D-Ala-D-Ala sequence at the C-terminus of the peptidoglycan pentapeptide (see diagram). This enzyme reduces the availability of the free dipeptide D-Ala-D-Ala, which is the precursor for this pentapeptide sequence, allowing D-Ala-(R)-lactate (for which vancomycin has much less affinity) to be added to the cell wall instead [3224, 1861]. The enzyme is stereospecific, as L-Ala-L-Ala, D-Ala-L-Ala and L-Ala-D-Ala are not substrates [3224]. Belongs in peptidase family M15.  
**References:** [2413, 3224, 1861, 335, 2869, 1847]

[EC 3.4.13.22 created 2006]

### EC 3.4.14 Dipeptidyl-peptidases and tripeptidyl-peptidases

#### EC 3.4.14.1

- Accepted name:** dipeptidyl-peptidase I  
**Reaction:** Release of an N-terminal dipeptide, Xaa-Yaa—Zaa-, except when Xaa is Arg or Lys, or Yaa or Zaa is Pro  
**Other name(s):** cathepsin C; dipeptidyl aminopeptidase I; dipeptidyl transferase; cathepsin C; dipeptidyl transferase; dipeptide arylamidase I; DAP I  
**Comments:** A Cl<sup>-</sup>-dependent, lysosomal cysteine-type peptidase maximally active at acidic pH. Also polymerizes dipeptide amides, arylamides and esters at neutral pH. In peptidase family C1 (papain family).  
**References:** [2291, 1891, 1870, 1869]

[EC 3.4.14.1 created 1961 as EC 3.4.4.9, transferred 1972 to EC 3.4.14.1]

#### EC 3.4.14.2

- Accepted name:** dipeptidyl-peptidase II  
**Reaction:** Release of an N-terminal dipeptide, Xaa-Yaa—, preferentially when Yaa is Ala or Pro. Substrates are oligopeptides, preferentially tripeptides  
**Other name(s):** dipeptidyl aminopeptidase II; dipeptidyl arylamidase II; carboxytripeptidase; dipeptidyl peptidase II; dipeptidyl arylamidase II; DAP II; dipeptidyl(amino)peptidase II; dipeptidylarylamidase  
**Comments:** A lysosomal serine-type peptidase in family S28 (Pro-X carboxypeptidase family); maximally active at acidic pH  
**References:** [1868, 1869]

[EC 3.4.14.2 created 1978]

[3.4.14.3 Transferred entry. acylamino-acid-releasing enzyme. Now EC 3.4.19.1, acylaminoacyl-peptidase]

[EC 3.4.14.3 created 1978, deleted 1981]

#### EC 3.4.14.4

- Accepted name:** dipeptidyl-peptidase III  
**Reaction:** Release of an N-terminal dipeptide from a peptide comprising four or more residues, with broad specificity. Also acts on dipeptidyl 2-naphthylamides.  
**Other name(s):** dipeptidyl aminopeptidase III; dipeptidyl arylamidase III; enkephalinase B; red cell angiotensinase  
**Comments:** A cytosolic peptidase that is active at neutral pH. It has broad activity on peptides, although it is highly selective for Arg-Arg-2-naphthylamide, at pH 9.2. Active in the hydrolysis of enkephalins. A metallopeptidase, the type example of peptidase family M49.  
**References:** [1866, 854]

[EC 3.4.14.4 created 1981, modified 2001]

#### EC 3.4.14.5

- Accepted name:** dipeptidyl-peptidase IV  
**Reaction:** Release of an N-terminal dipeptide, Xaa-Yaa—Zaa-, from a polypeptide, preferentially when Yaa is Pro, provided Zaa is neither Pro nor hydroxyproline  
**Other name(s):** dipeptidyl aminopeptidase IV; Xaa-Pro-dipeptidyl-aminopeptidase; Gly-Pro naphthylamidase; post-proline dipeptidyl aminopeptidase IV; lymphocyte antigen CD26; glycoprotein GP110; dipeptidyl peptidase IV; glycyloproline aminopeptidase; glycyloproline aminopeptidase; X-prolyl dipeptidyl aminopeptidase; pep X; leukocyte antigen CD26; glycyloprolyl dipeptidylaminopeptidase; dipeptidyl-peptide hydrolase; glycyloprolyl aminopeptidase; dipeptidyl-aminopeptidase IV; DPP IV/CD26; amino acyl-prolyl dipeptidyl aminopeptidase; T cell triggering molecule Tp103; X-PDAP  
**Comments:** A homodimer. An integral protein of the plasma membrane of lymphocytes and other mammalian cells, in peptidase family S9 (prolyl oligopeptidase family). The reaction is similar to that of the unrelated EC 3.4.14.11 Xaa-Pro dipeptidyl-peptidase of lactococci  
**References:** [1930, 543, 1230]

[EC 3.4.14.5 created 1981, modified 1996]

#### EC 3.4.14.6

- Accepted name:** dipeptidyl-dipeptidase  
**Reaction:** Preferential release of dipeptides from a tetrapeptide, e.g. Ala-Gly—Ala-Gly. Acts more slowly on Ala-Ala—Ala-Ala and Gly-Gly—Gly-Gly  
**Other name(s):** dipeptidyl tetrapeptide hydrolase; dipeptidyl ligase; tetrapeptide dipeptidase  
**Comments:** A thiol-activated peptidase from cabbage (*Brassica oleracea*). Tetrapeptides are formed from Ala-Ala, Gly-Gly, Ala-Gly and Gly-Ala  
**References:** [705]

[EC 3.4.14.6 created 1989]

[3.4.14.7 Deleted entry. tetralysine endopeptidase]

[EC 3.4.14.7 created 1989, deleted 1992]

[3.4.14.8 Transferred entry. tripeptidyl peptidase. Now EC 3.4.14.10, tripeptidyl-peptidase II]

[EC 3.4.14.8 created 1989, deleted 1992]

#### EC 3.4.14.9

**Accepted name:** tripeptidyl-peptidase I

**Reaction:** Release of an N-terminal tripeptide from a polypeptide, but also has endopeptidase activity.

**Other name(s):** tripeptidyl aminopeptidase; tripeptidyl peptidase

**Comments:** A lysosomal enzyme that is active at acidic pH. Deficient in classical late-infantile neuronal ceroid lipofuscinosis brain tissue. Belongs in peptidase family S53. Formerly included in EC 3.4.14.8.

**References:** [730, 2384, 729, 1347, 1692]

[EC 3.4.14.9 created 1992 (part of EC 3.4.14.8 created 1989, incorporated 1992), modified 2000, modified 2001, modified 2003]

#### EC 3.4.14.10

**Accepted name:** tripeptidyl-peptidase II

**Reaction:** Release of an N-terminal tripeptide from a polypeptide

**Other name(s):** tripeptidyl aminopeptidase; tripeptidyl peptidase; tripeptidyl aminopeptidase II; tripeptidyl peptidase II; TPP

**Comments:** A cytosolic enzyme in peptidase family S8 (subtilisin family). Active at neutral pH. Inhibited by diisopropyl fluorophosphate. Formerly included in EC 3.4.14.8

**References:** [131, 132, 2937]

[EC 3.4.14.10 created 1992 (part of EC 3.4.14.8 created 1989, incorporated 1992)]

#### EC 3.4.14.11

**Accepted name:** Xaa-Pro dipeptidyl-peptidase

**Reaction:** Hydrolyses Xaa-Pro— bonds to release unblocked, N-terminal dipeptides from substrates including Ala-Pro—*p*-nitroanilide and (sequentially) Tyr-Pro—Phe-Pro—Gly-Pro—Ile

**Other name(s):** X-prolyl dipeptidyl aminopeptidase; PepX; X-prolyl dipeptidyl peptidase; X-Pro dipeptidyl-peptidase

**Comments:** The intracellular enzyme from *Lactococcus lactis* (190-kDa) is the type example of peptidase family S15. The reaction is similar to that catalysed by dipeptidyl-peptidase IV of animals

**References:** [3322, 1897, 1015, 430, 429]

[EC 3.4.14.11 created 1996]

#### EC 3.4.14.12

**Accepted name:** Xaa-Xaa-Pro tripeptidyl-peptidase

**Reaction:** Hydrolysis of Xaa-Xaa-Pro—Yaa- releasing the N-terminal tripeptide of a peptide with Pro as the third residue (position P1) and where Yaa is not proline

**Other name(s):** prolyltri-peptidyl amino peptidase; prolyl tripeptidyl peptidase; prolyltri-peptidyl aminopeptidase; PTP-A; TPP

**Comments:** This cell-surface-associated serine exopeptidase is found in the Gram-negative, anaerobic bacterium *Porphyromonas gingivalis*, which has been implicated in adult periodontal disease [134]. The enzyme releases the N-terminal tripeptide of peptides, such as interleukin-6. It has an absolute requirement for a proline residue at the P1 position but is completely inactivated by a proline residue at the P1' position [134]. The size of the peptide does not affect the rate of reaction [134].

**References:** [134, 843]

[EC 3.4.14.12 created 2006]

#### EC 3.4.14.13

- Accepted name:**  $\gamma$ -D-glutamyl-L-lysine dipeptidyl-peptidase  
**Reaction:** The enzyme releases L-Ala- $\gamma$ -D-Glu dipeptides from cell wall peptides via cleavage of an L-Ala- $\gamma$ -D-Glu—L-Lys bond.  
**Other name(s):** YkFC  
**Comments:** The enzyme, characterized from the bacterium *Bacillus subtilis*, is involved in the recycling of the murein peptide.  
**References:** [2571, 3234]

[EC 3.4.14.13 created 2015]

### EC 3.4.15 Peptidyl-dipeptidases

#### EC 3.4.15.1

- Accepted name:** peptidyl-dipeptidase A  
**Reaction:** Release of a C-terminal dipeptide, oligopeptide—Xaa-Yaa, when Xaa is not Pro, and Yaa is neither Asp nor Glu. Thus, conversion of angiotensin I to angiotensin II, with increase in vasoconstrictor activity, but no action on angiotensin II  
**Other name(s):** dipeptidyl carboxypeptidase I; peptidase P; dipeptide hydrolase (ambiguous); peptidyl dipeptidase; angiotensin converting enzyme; kininase II; angiotensin I-converting enzyme; carboxycathepsin; dipeptidyl carboxypeptidase; peptidyl dipeptidase I; peptidyl-dipeptide hydrolase; peptidyl dipeptide hydrolase; endothelial cell peptidyl dipeptidase; ACE; peptidyl dipeptidase-4; PDH; peptidyl dipeptide hydrolase; DCP  
**Comments:** A Cl<sup>-</sup>-dependent, zinc glycoprotein that is generally membrane-bound. A potent inhibitor is captopril. Important in elevation of blood pressure, through formation of angiotensin II (vasoconstrictor) and destruction of bradykinin (vasodilator). Two molecular forms exist in mammalian tissues, a widely-distributed somatic form of 150- to 180-kDa that contains two non-identical catalytic sites, and a testicular form of 90- to 100-kDa that contains only a single catalytic site. Type example of peptidase family M2  
**References:** [2730, 682, 3147, 490]

[EC 3.4.15.1 created 1972, modified 1981, modified 1989, modified 1996, modified 2011]

[3.4.15.2 Transferred entry. *peptidyl carboxyamidase*. Now EC 3.4.19.2, *peptidyl-glycinamidase*]

[EC 3.4.15.2 created 1978, deleted 1981]

[3.4.15.3 Transferred entry. *dipeptidyl carboxypeptidase*. Now EC 3.4.15.5, *peptidyl-dipeptidase Dcp*]

[EC 3.4.15.3 created 1981, modified 1989, deleted 1996]

#### EC 3.4.15.4

- Accepted name:** peptidyl-dipeptidase B  
**Reaction:** Release of a C-terminal dipeptide or exceptionally a tripeptide  
**Other name(s):** dipeptidyl carboxyhydrolase; atriopeptin convertase; atrial di-(tri)peptidyl carboxyhydrolase; peptidyl dipeptidase B; atrial dipeptidyl carboxyhydrolase; atrial peptide convertase  
**Comments:** A membrane-bound, zinc metallopeptidase located in mammalian atrial, but not ventricular, myocytes. Although it is capable of converting the 126-residue atriopeptin III directly to atriopeptin I by releasing a C-terminal tripeptide Phe-Arg-Tyr, it is generally restricted to the release of dipeptides. In contrast to peptidyl-dipeptidase A (EC 3.4.15.1) it displays no Cl<sup>-</sup> dependence and shows no action on angiotensin I. Conversely, peptidyl-dipeptidase A is unable to release Phe-Arg from the C-terminus of atriopeptin II  
**References:** [1061, 1062, 2711, 2712]

[EC 3.4.15.4 created 1992]

#### EC 3.4.15.5

**Accepted name:** peptidyl-dipeptidase Dcp

**Reaction:** Hydrolysis of unblocked, C-terminal dipeptides from oligopeptides, with broad specificity. Does not hydrolyse bonds in which P1' is Pro, or both P1 and P1' are Gly

**Other name(s):** dipeptidyl carboxypeptidase (Dcp); dipeptidyl carboxypeptidase

**Comments:** Known from *Escherichia coli* and *Salmonella typhimurium*. A zinc metallopeptidase in peptidase family M3 (thimet oligopeptidase family). Ac-Ala—Ala-Ala is a good test substrate [474]. Inhibited by captopril, as is peptidyl-dipeptidase A. Formerly EC 3.4.15.3, and included in EC 3.4.15.1, peptidyl-dipeptidase A.

**References:** [3267, 1114, 474]

[EC 3.4.15.5 created 1981 as EC 3.4.15.3, modified 1989, transferred 1996 to EC 3.4.15.5]

#### EC 3.4.15.6

**Accepted name:** cyanophycinase

**Reaction:**  $[L\text{-Asp}(4\text{-L-Arg})]_n + \text{H}_2\text{O} = [L\text{-Asp}(4\text{-L-Arg})]_{n-1} + L\text{-Asp}(4\text{-L-Arg})$

**Other name(s):** cyanophycin degrading enzyme;  $\beta$ -Asp-Arg hydrolysing enzyme; CGPase; CphB; CphE; cyanophycin granule polypeptidase; extracellular CGPase

**Comments:** The enzyme is highly specific for the branched polypeptide cyanophycin and does not hydrolyse poly-L-aspartate or poly-L-arginine [2423]. A serine-type exopeptidase that belongs in peptidase family S51.

**References:** [2125, 2126, 2423]

[EC 3.4.15.6 created 2007]

### EC 3.4.16 Serine-type carboxypeptidases

[3.4.16.1 *Transferred entry. serine carboxypeptidase. Now EC 3.4.16.6, carboxypeptidase D*]

[EC 3.4.16.1 created 1972 as EC 3.4.12.1 and EC 3.4.21.13, both transferred 1978 to EC 3.4.16.1, deleted 1993]

#### EC 3.4.16.2

**Accepted name:** lysosomal Pro-Xaa carboxypeptidase

**Reaction:** Cleavage of a -Pro—Xaa bond to release a C-terminal amino acid

**Other name(s):** angiotensinase C; lysosomal carboxypeptidase C; peptidylprolylamino acid carboxypeptidase; aminoacylproline carboxypeptidase; prolyl carboxypeptidase; carboxypeptidase P; proline-specific carboxypeptidase P; PCP; lysosomal Pro-Xaa carboxypeptidase

**Comments:** A lysosomal peptidase active at acidic pH that inactivates angiotensin II. Inhibited by diisopropyl fluorophosphate. In peptidase family S28 (Pro-X carboxypeptidase family).

**References:** [3102, 2139]

[EC 3.4.16.2 created 1972 as EC 3.4.12.4, transferred 1978 to EC 3.4.16.2]

[3.4.16.3 *Transferred entry. tyrosine carboxypeptidase. Now included with EC 3.4.16.5, carboxypeptidase C*]

[EC 3.4.16.3 created 1972 as EC 3.4.12.12, transferred 1978 to EC 3.4.16.3, deleted 1992]

#### EC 3.4.16.4

**Accepted name:** serine-type D-Ala-D-Ala carboxypeptidase

**Reaction:** Preferential cleavage:  $(\text{Ac})_2\text{-L-Lys-D-Ala—D-Ala}$ . Also transpeptidation of peptidyl-alanyl moieties that are N-acyl substituents of D-alanine

**Other name(s):** DD-peptidase; D-alanyl-D-alanine-carboxypeptidase; D-alanyl-D-alanine-cleaving-peptidase; D-alanyl-D-alanine-cleaving peptidase; DD-transpeptidase; D-alanine carboxypeptidase; DD-carboxypeptidase; D-alanyl carboxypeptidase

**Comments:** A membrane-bound, bacterial enzyme inhibited by penicillin and other  $\beta$ -lactam antibiotics, which acylate the active site serine. Examples are known from peptidase families S11, S12 and S13. Distinct from EC 3.4.17.14, zinc D-Ala-D-Ala carboxypeptidase

**References:** [909, 810]

[EC 3.4.16.4 created 1989]

#### EC 3.4.16.5

**Accepted name:** carboxypeptidase C

**Reaction:** Release of a C-terminal amino acid with broad specificity

**Other name(s):** carboxypeptidase Y; serine carboxypeptidase I; cathepsin A; lysosomal protective protein; deamidase; lysosomal carboxypeptidase A; phaseolin

**Comments:** A carboxypeptidase with optimum pH 4.5-6.0, inhibited by diisopropyl fluorophosphate, and sensitive to thiol-blocking reagents (reviewed in [283]). Widely distributed in eukaryotes. Type example of peptidase family S10.

**References:** [283, 3030, 1287, 1914]

[EC 3.4.16.5 created 1972 as EC 3.4.12.1, transferred 1978 to EC 3.4.16.1, part transferred 1993 to EC 3.4.16.5 (EC 3.4.16.3 created 1972 as EC 3.4.12.12, transferred 1978 to EC 3.4.16.3, transferred 1992 to EC 3.4.16.1), (EC 3.4.21.13 created 1972, transferred 1978 to EC 3.4.16.1)]

#### EC 3.4.16.6

**Accepted name:** carboxypeptidase D

**Reaction:** Preferential release of a C-terminal arginine or lysine residue

**Other name(s):** cereal serine carboxypeptidase II; *Saccharomyces cerevisiae* KEX1 gene product; carboxypeptidase Kex1; gene KEX1 serine carboxypeptidase; KEX1 carboxypeptidase; KEX1 proteinase; KEX1DELTAp; CPDW-II; serine carboxypeptidase (misleading); *Phaseolus* proteinase

**Comments:** A carboxypeptidase with optimum pH 4.5-6.0, inhibited by diisopropyl fluorophosphate, and sensitive to thiol-blocking reagents (reviewed in [283]). In peptidase family S10 (carboxypeptidase C family).

**References:** [283, 285, 621, 1678]

[EC 3.4.16.6 created 1972 as EC 3.4.12.1, transferred 1978 to EC 3.4.16.1, part transferred 1993 to EC 3.4.16.6 (EC 3.4.16.3 created 1972 as EC 3.4.12.12, transferred 1978 to EC 3.4.16.3, transferred 1992 to EC 3.4.16.1), (EC 3.4.21.13 created 1972, transferred 1978 to EC 3.4.16.1), modified 2011]

### EC 3.4.17 Metallo-carboxypeptidases

#### EC 3.4.17.1

**Accepted name:** carboxypeptidase A

**Reaction:** Release of a C-terminal amino acid, but little or no action with -Asp, -Glu, -Arg, -Lys or -Pro

**Other name(s):** carboxypolypeptidase; pancreatic carboxypeptidase A; tissue carboxypeptidase A

**Comments:** A zinc enzyme formed from procarboxypeptidase A. Isolated from cattle, pig and dogfish pancreas, and other sources including mast cells [726] and skeletal muscle [252]. Type example of peptidase family M14.

**References:** [2267, 2394, 726, 252]

[EC 3.4.17.1 created 1961 as EC 3.4.2.1, transferred 1972 to EC 3.4.12.2, transferred 1978 to EC 3.4.17.1]

#### EC 3.4.17.2

**Accepted name:** carboxypeptidase B



**Reaction:** Preferential release of a C-terminal lysine or arginine amino acid  
**Other name(s):** protaminase; pancreatic carboxypeptidase B; tissue carboxypeptidase B; peptidyl-L-lysine [L-arginine]hydrolase  
**Comments:** A zinc enzyme formed from procarboxypeptidase B. Isolated from cattle, pig and dogfish pancreas and other sources, including skin fibroblasts [338] and adrenal medulla [3098]. In peptidase family M14 (carboxypeptidase A family).  
**References:** [788, 299, 338, 3098]

[EC 3.4.17.2 created 1961 as EC 3.4.2.2, transferred 1972 to EC 3.4.12.3, transferred 1978 to EC 3.4.17.2]

### EC 3.4.17.3

**Accepted name:** lysine carboxypeptidase  
**Reaction:** Release of a C-terminal basic amino acid, preferentially lysine  
**Other name(s):** carboxypeptidase N; arginine carboxypeptidase; kininase I; anaphylatoxin inactivator; plasma carboxypeptidase B; creatine kinase conversion factor; bradykinase; kininase Ia; hippuryllysine hydrolase; bradykinin-decomposing enzyme; protaminase; CPase N; creatinine kinase convertase; peptidyl-L-lysine(-L-arginine) hydrolase; CPN  
**Comments:** A zinc enzyme found in plasma. Inactivates bradykinin and anaphylatoxins in blood plasma. In peptidase family M14 (carboxypeptidase A family).  
**References:** [2294, 1656, 2681]

[EC 3.4.17.3 created 1972 as EC 3.4.12.7, transferred 1978 to EC 3.4.17.3, modified 1989]

### EC 3.4.17.4

**Accepted name:** Gly-Xaa carboxypeptidase  
**Reaction:** Release of a C-terminal amino acid from a peptide in which glycine is the penultimate amino acid, e.g. Z-Gly—Leu  
**Other name(s):** glycine carboxypeptidase; carboxypeptidase a; carboxypeptidase S; peptidase  $\alpha$ ; yeast carboxypeptidase; Gly-X carboxypeptidase  
**Comments:** From yeast. In peptidase family M20 (glutamate carboxypeptidase family).  
**References:** [751, 3206]

[EC 3.4.17.4 created 1961 as EC 3.4.2.3, transferred 1972 to EC 3.4.12.8, transferred 1978 to EC 3.4.17.4 (EC 3.4.17.9 created 1981, incorporated 1992)]

### [3.4.17.5 Deleted entry. aspartate carboxypeptidase]

[EC 3.4.17.5 created 1972 as EC 3.4.12.9, transferred 1978 to EC 3.4.17.5, deleted 1992]

### EC 3.4.17.6

**Accepted name:** alanine carboxypeptidase  
**Reaction:** Release of a C-terminal alanine from a peptide or a variety of pteroyl or acyl groups  
**Other name(s):** N-benzoyl-L-alanine-amidohydrolase  
**Comments:** From soil bacteria. The enzyme from *Corynebacterium equi* also hydrolyses N-benzoylglycine and N-benzoyl-L-aminobutyric acid.  
**References:** [1659, 1936]

[EC 3.4.17.6 created 1972 as EC 3.4.12.11, transferred 1978 to EC 3.4.17.6]

### [3.4.17.7 Transferred entry. acylmuramoyl-alanine carboxypeptidase. Now EC 3.5.1.28, N-acetylmuramoyl-L-alanine amidase]

[EC 3.4.17.7 created 1978, deleted 1992]

### EC 3.4.17.8



**Accepted name:** muramoylpentapeptide carboxypeptidase  
**Reaction:** Cleavage of the bond UDP-*N*-acetylmuramoyl-L-alanyl- $\gamma$ -D-glutamyl-6-carboxy-L-lysyl-D-alanyl-D-alanine  
**Other name(s):** D-alanine carboxypeptidase I; DD-carboxypeptidase; D-alanine carboxypeptidase; D-alanyl-D-alanine carboxypeptidase; D-alanine-D-alanine-carboxypeptidase; carboxypeptidase D-alanyl-D-alanine; carboxypeptidase I; UDP-*N*-acetylmuramoyl-tetrapeptidyl-D-alanine alanine-hydrolase; D-alanyl-D-alanine peptidase; DD-peptidase; penicillin binding protein 5; PBP5; PdcA; VanY  
**Comments:** A bacterial enzyme that requires a divalent cation for activity. Does not cleave the C-terminal D-alanine from the product of the above reaction, UDP-*N*-acetyl-muramoyl-L-alanyl- $\gamma$ -D-glutamyl-6-carboxy-L-lysyl-D-alanine. Competitively inhibited by penicillins and cephalosporins.  
**References:** [1285]

[EC 3.4.17.8 created 1972 as EC 3.4.12.6, transferred 1978 to EC 3.4.17.8]

[3.4.17.9 *Transferred entry. carboxypeptidase S. Now included with EC 3.4.17.4, Gly-Xaa carboxypeptidase*]

[EC 3.4.17.9 created 1981, deleted 1992]

#### EC 3.4.17.10

**Accepted name:** carboxypeptidase E  
**Reaction:** Release of C-terminal arginine or lysine residues from polypeptides  
**Other name(s):** carboxypeptidase H; enkephalin convertase; cobalt-stimulated chromaffin granule carboxypeptidase; insulin granule-associated carboxypeptidase; enkephalin convertase; membrane-bound carboxypeptidase; carboxypeptidase E; enkephalin-precursor endopeptidase; enkephalin precursor carboxypeptidase; peptidyl-L-lysine(-L-arginine) hydrolase  
**Comments:** A zinc enzyme, activated by Co<sup>2+</sup>. Inhibited by 1,10-phenanthroline and other chelating agents. pH optimum 5.6. Located in storage granules of secretory cells, and active in processing of protein hormones and bioactive peptides. In peptidase family M14 (carboxypeptidase A family).  
**References:** [2337, 818, 817, 1803, 816]

[EC 3.4.17.10 created 1986, modified 2000]

#### EC 3.4.17.11

**Accepted name:** glutamate carboxypeptidase  
**Reaction:** Release of C-terminal glutamate residues from a wide range of *N*-acylating moieties, including peptidyl, aminoacyl, benzoyl, benzyloxycarbonyl, foyl and pteroyl groups  
**Other name(s):** carboxypeptidase G; carboxypeptidase G<sub>1</sub>; carboxypeptidase G<sub>2</sub>; glutamyl carboxypeptidase; *N*-pteroyl-L-glutamate hydrolase  
**Comments:** A zinc enzyme produced by pseudomonads, *Flavobacterium* sp. and *Acinetobacter* sp. Its ability to hydrolyse pteroyl-L-glutamate (folic acid) has led to its use as a folate-depleting, antitumour agent. Type example of peptidase family M20  
**References:** [947, 1865, 29, 2627]

[EC 3.4.17.11 created 1992]

#### EC 3.4.17.12

**Accepted name:** carboxypeptidase M  
**Reaction:** Cleavage of C-terminal arginine or lysine residues from polypeptides  
**Other name(s):** CPM  
**Comments:** A membrane-bound enzyme optimally active at neutral pH. In peptidase family M14 (carboxypeptidase A family)  
**References:** [2682, 566, 2683]

[EC 3.4.17.12 created 1992]

### EC 3.4.17.13

- Accepted name:** muramoyltetrapeptide carboxypeptidase  
**Reaction:** Hydrolysis of the bond: *N*-acetyl-D-glucosaminyl-*N*-acetylmuramoyl-L-Ala-D-glutamyl-6-carboxy-L-lysyl-D-alanine  
**Other name(s):** carboxypeptidase IIW; carboxypeptidase II; lysyl-D-alanine carboxypeptidase; L-lysyl-D-alanine carboxypeptidase; LD-carboxypeptidase  
**Comments:** Variants are known from various microorganisms. Involved in peptidoglycan synthesis, catalysing both decarboxylation and transpeptidation. Stimulated by divalent cations such as Mg<sup>2+</sup> and Ca<sup>2+</sup>, but not by Zn<sup>2+</sup>. Inhibited by thiol-blocking reagents, but unaffected by penicillin  
**References:** [539, 2467, 1893]

[EC 3.4.17.13 created 1992]

### EC 3.4.17.14

- Accepted name:** zinc D-Ala-D-Ala carboxypeptidase  
**Reaction:** Cleavage of the bond: (Ac)<sub>2</sub>-L-lysyl-D-alanyl-D-alanine  
**Other name(s):** Zn<sup>2+</sup> G peptidase, D-alanyl-D-alanine hydrolase; D-alanyl-D-alanine-cleaving carboxypeptidase; DD-carboxypeptidase; G enzyme; DD-carboxypeptidase-transpeptidase  
**Comments:** A zinc enzyme. Catalyses carboxypeptidation but not transpeptidation reactions involved in bacterial cell wall metabolism. Weakly inhibited by β-lactams. In peptidase family M15. Distinct from EC 3.4.16.4, serine-type D-Ala-D-Ala carboxypeptidase.  
**References:** [606, 1340, 909]

[EC 3.4.17.14 created 1992]

### EC 3.4.17.15

- Accepted name:** carboxypeptidase A<sub>2</sub>  
**Reaction:** Similar to that of carboxypeptidase A (EC 3.4.17.1), but with a preference for bulkier C-terminal residues  
**Other name(s):** CPA2  
**Comments:** Isolated from rat pancreas but not present in cattle pancreas. In peptidase family M14 (carboxypeptidase A family).  
**References:** [878]

[EC 3.4.17.15 created 1992]

### EC 3.4.17.16

- Accepted name:** membrane Pro-Xaa carboxypeptidase  
**Reaction:** Release of a C-terminal residue other than proline, by preferential cleavage of a prolyl bond  
**Other name(s):** carboxypeptidase P; microsomal carboxypeptidase; membrane Pro-X carboxypeptidase  
**Comments:** One of the renal brush border exopeptidases  
**References:** [569, 263, 1096]

[EC 3.4.17.16 created 1992]

### EC 3.4.17.17

- Accepted name:** tubuliny-Tyr carboxypeptidase  
**Reaction:** Cleavage of the -Glu-Tyr bond to release the C-terminal tyrosine residue from the native tyrosinated tubulin. Inactive on Z-Glu-Tyr  
**Other name(s):** carboxypeptidase-tubulin; soluble carboxypeptidase; tubulin-tyrosine carboxypeptidase; tubulin carboxypeptidase; tubulinytyrosine carboxypeptidase; tyrosinotubulin carboxypeptidase; tyrosyltubulin carboxypeptidase; TTCPase; brain I carboxypeptidase  
**Comments:** Active at neutral pH, from brain  
**References:** [2019, 1549, 70]

[EC 3.4.17.17 created 1992]

#### EC 3.4.17.18

**Accepted name:** carboxypeptidase T  
**Reaction:** Releases a C-terminal residue, which may be hydrophobic or positively charged  
**Other name(s):** CPT  
**Comments:** Known from *Thermoactinomyces vulgaris*. In peptidase family M14 (carboxypeptidase A family)  
**References:** [2208, 2703, 2898]

[EC 3.4.17.18 created 1993]

#### EC 3.4.17.19

**Accepted name:** carboxypeptidase Taq  
**Reaction:** Release of a C-terminal amino acid with broad specificity, except for -Pro  
**Comments:** A 56-kDa enzyme from *Thermus aquaticus*. Most active at 80° C. Type example of peptidase family M32  
**References:** [1627, 1628]

[EC 3.4.17.19 created 1996]

#### EC 3.4.17.20

**Accepted name:** carboxypeptidase U  
**Reaction:** Release of C-terminal Arg and Lys from a polypeptide  
**Other name(s):** arginine carboxypeptidase; carboxypeptidase R; plasma carboxypeptidase B (misleading, since the term carboxypeptidase B is used for other enzymes); thrombin-activatable fibrinolysis inhibitor  
**Comments:** Pro-carboxypeptidase U in (human) plasma is activated by thrombin or plasmin during clotting to form the unstable carboxypeptidase U, with activity similar to that of the more stable lysine carboxypeptidase, except that no preference is shown for Lys over Arg. A zinc enzyme, in peptidase family M14 (carboxypeptidase A family)  
**References:** [674, 2640, 3121, 2868, 312]

[EC 3.4.17.20 created 1997]

#### EC 3.4.17.21

**Accepted name:** glutamate carboxypeptidase II  
**Reaction:** Release of an unsubstituted, C-terminal glutamyl residue, typically from Ac-Asp-Glu or folylpoly- $\gamma$ -glutamates  
**Other name(s):** *N*-acetylated- $\gamma$ -linked-acidic dipeptidase (NAALADase); folate hydrolase; prostate-specific membrane antigen; pteroylpoly- $\gamma$ -glutamate carboxypeptidase; microsomal  $\gamma$ -glutamyl carboxypeptidase; pteroylpolyglutamate hydrolase; folylpolyglutamate hydrolase; pteroylpoly- $\gamma$ -glutamate hydrolase; pteroylpoly $\gamma$ glutamyl hydrolase; pteroylpolyglutamate hydrolase; pteroylpolyglutamic acid hydrolase; PSM antigen; acetylaspartylglutamate dipeptidase; NAALA dipeptidase; rat NAAG peptidase; mGCP; membrane glutamate carboxypeptidase; *N*-acetylated- $\alpha$ -linked-amino dipeptidase; prostrate-specific membrane antigen; *N*-Acetylated  $\alpha$ -linked acidic dipeptidase; PSMA  
**Comments:** A metallo-carboxypeptidase that is predominantly expressed as a membrane-bound enzyme of 94-100 kDa, but also exists in a soluble form. Hydrolyses  $\alpha$ -peptide bonds in Ac-Asp-Glu, Asp-Glu, and Glu-Glu, but also  $\gamma$ -glutamyl bonds in  $\gamma$ -Glu-Glu, and folylpoly- $\gamma$ -glutamates. With folylpoly- $\gamma$ -glutamates, shows processive carboxypeptidase activity to produce pteroylmonoglutamate [1758]. Does not hydrolyse Ac- $\beta$ -Asp-Glu. Known inhibitors: quisqualic acid, Ac- $\beta$ -Asp-Glu, and 2-phosphonomethyl-pentanedioate. In peptidase family M28 of *Vibrio leucyl* aminopeptidase. The release of C-terminal glutamate from folylpoly- $\gamma$ -glutamates is also catalysed by EC 3.4.17.11 (glutamate carboxypeptidase) and EC 3.4.19.9 ( $\gamma$ -Glu-X carboxypeptidase).  
**References:** [1134, 2383, 1032, 1758]

[EC 3.4.17.21 created 1997, modified 2000 (EC 3.4.13.8 created 1972 and EC 3.4.19.8 created 1992, incorporated 2000)]

#### EC 3.4.17.22

**Accepted name:** metallocarboxypeptidase D  
**Reaction:** Releases C-terminal Arg and Lys from polypeptides  
**Other name(s):** carboxypeptidase D (cattle, human, mouse, rat); gp180 (duck)  
**Comments:** Activated by  $\text{Co}^{2+}$ ; inhibited by guanidinoethylmercaptosuccinic acid. Large molecule (180 kDa) because of presence of three copies of metallopeptidase domain. The product of the silver gene (*Drosophila*) is similar. A zinc metallopeptidase in peptidase family M14 (carboxypeptidase A family)  
**References:** [1573, 2718, 2719]

[EC 3.4.17.22 created 1997]

#### EC 3.4.17.23

**Accepted name:** angiotensin-converting enzyme 2  
**Reaction:** angiotensin II +  $\text{H}_2\text{O}$  = angiotensin-(1-7) + L-phenylalanine  
**Other name(s):** ACE-2; ACE2; hACE2; angiotensin converting enzyme 2; angiotensin converting enzyme-2; Tmem27  
**Comments:** A transmembrane glycoprotein with an extracellular catalytic domain. Angiotensin-converting enzyme 2 functions as a carboxypeptidase, cleaving a single C-terminal residue from a distinct range of substrates [1594]. Catalytic efficiency is 400-fold higher with angiotensin II (1-8) as a substrate than with angiotensin I (1-10). Angiotensin-converting enzyme 2 also efficiently hydrolyses des-Arg<sup>9</sup>-bradykinin, but it does not hydrolyse bradykinin [3066]. In peptidase family M2.  
**References:** [3066, 1594, 2954]

[EC 3.4.17.23 created 2009]

### EC 3.4.18 Cysteine-type carboxypeptidases

#### EC 3.4.18.1

**Accepted name:** cathepsin X  
**Reaction:** Release of C-terminal amino acid residues with broad specificity, but lacks action on C-terminal proline. Shows weak endopeptidase activity  
**Other name(s):** cathepsin B2; cysteine-type carboxypeptidase; cathepsin IV; cathepsin Z; acid carboxypeptidase; lysosomal carboxypeptidase B  
**Comments:** Cathepsin X is a lysosomal cysteine peptidase of family C1 (papain family). The pH optimum is dependent on the substrate and is 5.0 for the carboxypeptidase activity. Unstable above pH 7.0. Compound E-64, leupeptin and antipain are inhibitors, but not cystatin C. Cathepsin X is ubiquitously distributed in mammalian tissues. The propeptide is extremely short (38 amino acid residues) and the proenzyme is catalytically active. Human gene locus: 20q13.  
**References:** [2027, 2026, 2526, 1867, 2216, 2091]

[EC 3.4.18.1 created 1981, modified 2000]

### EC 3.4.19 Omega peptidases

#### EC 3.4.19.1

**Accepted name:** acylaminoacyl-peptidase  
**Reaction:** Cleavage of an N-acetyl or N-formyl amino acid from the N-terminus of a polypeptide  
**Other name(s):** acylamino-acid-releasing enzyme; N-acylpeptide hydrolase; N-formylmethionine (fMet) aminopeptidase;  $\alpha$ -N-acylpeptide hydrolase

**Comments:** Active at neutral pH. Several variants of this enzyme exist; the human erythrocyte enzyme is relatively specific for removal of *N*-acetylalanine from peptides. Displays dipeptidyl-peptidase activity on glycyl-peptides, perhaps as a result of mis-recognition of the glycyl residue as an uncharged *N*-acyl group. Inhibited by diisopropyl fluorophosphate. In peptidase family S9 (prolyl oligopeptidase family).

**References:** [2986, 3012, 1490]

[EC 3.4.19.1 created 1978 as EC 3.4.14.3, transferred 1981 to EC 3.4.19.1]

#### EC 3.4.19.2

**Accepted name:** peptidyl-glycinamidase

**Reaction:** Cleavage of C-terminal glycinamide from polypeptides

**Other name(s):** carboxyamidase; peptidyl carboxy-amidase; peptidyl-aminoacylamidase; carboxamidopeptidase; peptidyl amino acid amide hydrolase

**Comments:** Inactivates vasopressin and oxytocin by splitting off glycinamide. Also cleaves ester substrates of trypsin and chymotrypsin. Although glycinamide is by far the preferred leaving group, other aminoacylamides may also be released, e.g. phenylalaninamide. The toad skin enzyme is inhibited by diisopropyl fluorophosphate.

**References:** [825, 2057, 2666]

[EC 3.4.19.2 created 1978 as EC 3.4.15.2, transferred 1981 to EC 3.4.19.2]

#### EC 3.4.19.3

**Accepted name:** pyroglutamyl-peptidase I

**Reaction:** Release of an N-terminal pyroglutamyl group from a polypeptide, the second amino acid generally not being Pro

**Other name(s):** 5-oxoprolyl-peptidase; pyrased; pyroglutamate aminopeptidase; pyroglutamyl aminopeptidase; L-pyroglutamyl peptide hydrolase; pyrrolidone-carboxyl peptidase; pyrrolidone-carboxylate peptidase; pyrrolidonyl peptidase; L-pyrrolidonecarboxylate peptidase; pyroglutamidase; pyrrolidonecarboxyl peptidase

**Comments:** A cysteine peptidase, known from bacteria, plants and animals. The enzyme from bacterial sources is used in protein sequencing, and is the type example of peptidase family C15.

**References:** [2988, 106, 2247, 2536]

[EC 3.4.19.3 created 1972 as EC 3.4.11.8, transferred 1981 to EC 3.4.19.3, modified 1997]

[3.4.19.4 Deleted entry. *N*-acetylmethionylpeptide peptidase]

[EC 3.4.19.4 created 1989, deleted 1992]

#### EC 3.4.19.5

**Accepted name:**  $\beta$ -aspartyl-peptidase

**Reaction:** Cleavage of a  $\beta$ -linked Asp residue from the N-terminus of a polypeptide

**Other name(s):**  $\beta$ -aspartyl dipeptidase;  $\beta$ -aspartyl peptidase;  $\beta$ -aspartyldipeptidase

**Comments:** Other isopeptide bonds, e.g.  $\gamma$ -glutamyl and  $\beta$ -alanyl, are not hydrolysed. A mammalian, cytosolic enzyme.

**References:** [1025]

[EC 3.4.19.5 created 1972 as EC 3.4.13.10, transferred 1992 to EC 3.4.19.5, modified 1997]

#### EC 3.4.19.6

**Accepted name:** pyroglutamyl-peptidase II

**Reaction:** Release of the N-terminal pyroglutamyl group from pGlu—His-Xaa tripeptides and pGlu—His-Xaa-Gly tetrapeptides

**Other name(s):** thyroliberinase; pyroglutamyl aminopeptidase II; , thyrotropin-releasing factor pyroglutamate aminopeptidase; pyroglutamate aminopeptidase II; pyroglutamyl peptidase II; thyroliberin-hydrolyzing pyroglutamate aminopeptidase; thyrotropin-releasing hormone-degrading pyroglutamate aminopeptidase; thyrotropin-releasing hormone-degrading peptidase; TRH aminopeptidase  
**Comments:** Highly specific for thyrotropin releasing hormone (pyroglutamyl-histidyl-prolylamide). Will not cleave the pyroglutamyl-histidyl bond of luteinizing hormone releasing hormone. Found in serum and brain. Inhibited by metal chelators. In peptidase family M1 (membrane alanyl aminopeptidase family)  
**References:** [174, 2128, 3187]

[EC 3.4.19.6 created 1992]

#### EC 3.4.19.7

**Accepted name:** *N*-formylmethionyl-peptidase  
**Reaction:** Release of an N-terminal, formyl-methionyl residue from a polypeptide  
**Other name(s):** (fMet)-releasing enzyme; formylmethionine aminopeptidase  
**Comments:** Highly specific for *N*-formylmethionyl peptides. Will not cleave methionyl peptides or *N*-formyl derivatives of amino acids other than methionine. Isolated from rat liver. Inhibited by heavy metals and activated by Cl<sup>-</sup>  
**References:** [2786]

[EC 3.4.19.7 created 1992]

[3.4.19.8 Transferred entry. now EC 3.4.17.21, glutamate carboxypeptidase II]

[EC 3.4.19.8 created 1992, deleted 2000]

#### EC 3.4.19.9

**Accepted name:** folate  $\gamma$ -glutamyl hydrolase  
**Reaction:** tetrahydropteroyl-( $\gamma$ -glutamyl)<sub>*n*</sub> + (*n*-1) H<sub>2</sub>O = 5,6,7,8-tetrahydrofolate + (*n*-1) L-glutamate  
**Other name(s):** GGH (gene name); conjugase; folate conjugase; lysosomal  $\gamma$ -glutamyl carboxypeptidase;  $\gamma$ -Glu-X carboxypeptidase; pteroyl-poly- $\gamma$ -glutamate hydrolase; carboxypeptidase G; folic acid conjugase; poly( $\gamma$ -glutamic acid) endohydrolase; polyglutamate hydrolase; poly(glutamic acid) hydrolase II; pteroylpoly- $\gamma$ -glutamyl hydrolase;  $\gamma$ -glutamyl hydrolase  
**Systematic name:** tetrahydropteroyl-poly- $\gamma$ -glutamyl  $\gamma$ -glutamyl hydrolase  
**Comments:** The enzyme, which occurs only in animals and plants, can be either endo- and/or exopeptidase. It acts on tetrahydropteroyl polyglutamates and their modified forms, as well as the polyglutamates of the folate breakdown product *N*-(4-aminobenzoyl)-L-glutamate (pABA-Glu). The initial cleavage may release either monoglutamate or poly- $\gamma$ -glutamate of two or more residues, depending on the specific enzyme. For example, GGH1 from the plant *Arabidopsis thaliana* cleaves pentaglutamates, mainly to di- and triglutamates, whereas GGH2 from the same organism yields mainly monoglutamates. The enzyme is lysosomal (and secreted) in animals and vacuolar in plants.  
**References:** [1873, 3124, 3264, 3265, 3263, 2204, 25]

[EC 3.4.19.9 created 1972 as EC 3.4.12.10, transferred 1978 to EC 3.4.22.12, transferred 1992 to EC 3.4.19.9, modified 1997, modified 2018]

[3.4.19.10 Transferred entry. acylmuramoyl-Ala peptidase. Now EC 3.5.1.28, N-acetylmuramoyl-L-alanine amidase]

[EC 3.4.19.10 created 1972 as EC 3.4.12.5, transferred 1978 to EC 3.4.17.7, transferred 1992 to EC 3.4.19.10, deleted 1997]

#### EC 3.4.19.11

**Accepted name:**  $\gamma$ -D-glutamyl-*meso*-diaminopimelate peptidase  
**Reaction:** Hydrolysis of  $\gamma$ -D-glutamyl bonds to the L-terminus (position 7) of *meso*-diaminopimelic acid (*meso*-A2pm) in 7-(L-Ala- $\gamma$ -D-Glu)-*meso*-A2pm and 7-(L-Ala- $\gamma$ -D-Glu)-7-(D-Ala)-*meso*-A2pm. It is required that the D-terminal amino and carboxy groups of *meso*-A2pm are unsubstituted

**Other name(s):** endopeptidase I;  $\gamma$ -D-glutamyl-diaminopimelate endopeptidase;  $\gamma$ -D-glutamyl-L-*meso*-diaminopimelate peptidoglycan hydrolase;  $\gamma$ -glutamyl-L-*meso*-diaminopimelyl endopeptidase;  $\gamma$ -D-glutamyl-*meso*-diaminopimelate endopeptidase;  $\gamma$ -D-glutamyl-*meso*-diaminopimelic peptidoglycan hydrolase;  $\gamma$ -D-glutamyl-*meso*-diaminopimelic endopeptidase;  $\gamma$ -D-glutamyl-*meso*-D-aminopimelic endopeptidase

**Comments:** A 45-kDa metallopeptidase from *Bacillus sphaericus*, the substrates being components of the bacterial spore wall. A member of peptidase family M14 (carboxypeptidase A family). Endopeptidase II has similar activity, but differs in cellular location, molecular mass and catalytic mechanism [1188]

**References:** [74, 880, 1188]

[EC 3.4.19.11 created 1996]

#### EC 3.4.19.12

**Accepted name:** ubiquitinyl hydrolase 1

**Reaction:** Thiol-dependent hydrolysis of ester, thioester, amide, peptide and isopeptide bonds formed by the C-terminal Gly of ubiquitin (a 76-residue protein attached to proteins as an intracellular targeting signal)

**Other name(s):** ubiquitin C-terminal hydrolase; yeast ubiquitin hydrolase

**Comments:** Links to polypeptides smaller than 60 residues are hydrolysed more readily than those to larger polypeptides. Isoforms exist with quantitatively different specificities, amongst the best known being UCH-L1 and UCH-L3, which are major proteins of the brain of mammals [1333]. Inhibited by ubiquitin aldehyde (in which Gly<sup>76</sup> is replaced by aminoacetaldehyde). Ubiquitinyl hydrolase 1 is the type example of peptidase family C12, with a similar protein fold to papain and catalytic amino acids Cys, His and Asp. There is a separate family (C19) of enzymes that also hydrolyse ubiquitinyl bonds, and it is thought that all the ubiquitinyl hydrolases are also ubiquitin thiolesterases (EC 3.1.2.15)

**References:** [1333, 3191]

[EC 3.4.19.12 created 2000]

#### EC 3.4.19.13

**Accepted name:** glutathione hydrolase

**Reaction:** glutathione + H<sub>2</sub>O = L-cysteinylglycine + L-glutamate

**Other name(s):** glutathionase; GGT (ambiguous);  $\gamma$ -glutamyltranspeptidase (ambiguous)

**Comments:** This protein also has EC 2.3.2.2 ( $\gamma$ -glutamyltransferase) activity. The enzyme consists of two chains that are created by the proteolytic cleavage of a single precursor polypeptide. The N-terminal L-threonine of the C-terminal subunit functions as the active site for both the cleavage and the hydrolysis reactions [2810, 2179, 248, 2180]. The human enzyme also hydrolyses oxidized glutathione and leukotriene C<sub>4</sub> with similar efficiency, while the mouse enzyme does not [3177, 376].

**References:** [1044, 2810, 2179, 248, 2180, 3177, 376]

[EC 3.4.19.13 created 2011]

#### EC 3.4.19.14

**Accepted name:** leukotriene-C<sub>4</sub> hydrolase

**Reaction:** leukotriene C<sub>4</sub> + H<sub>2</sub>O = leukotriene D<sub>4</sub> + L-glutamate

**Other name(s):**  $\gamma$ -glutamyl leukotrienase; GGT5

**Comments:** The mouse enzyme is specific for leukotriene C<sub>4</sub>, while the human enzyme also has considerable activity towards glutathione and oxidized glutathione (*cf.* EC 3.4.19.13, glutathione hydrolase) [1039, 3177].

**References:** [376, 2628, 1039, 3177]

[EC 3.4.19.14 created 2012]

#### EC 3.4.19.15

**Accepted name:** desampylase



**Reaction:** an  $N^6$ -[small archaeal modifier protein]-[protein]-L-lysine + H<sub>2</sub>O = a [protein]-L-lysine + a small archaeal modifier protein

**Other name(s):** SAMP-protein conjugate cleaving protease; HvJAMM1

**Systematic name:**  $N^6$ -[small archaeal modifier protein]-[protein]-L-lysine hydrolase

**Comments:** The enzyme, characterized from the archaeon *Haloferax volcanii*, specifically cleaves the ubiquitin-like small modifier proteins SAMP1 and SAMP2 from protein conjugates, hydrolysing the isopeptide bond between a lysine residue of the target protein and the C-terminal glycine of the modifier protein. The enzyme contains Zn<sup>2+</sup>. cf. EC 3.4.19.12, ubiquitinyl hydrolase 1. In peptidase family M67.

**References:** [1120]

[EC 3.4.19.15 created 2015 as EC 3.4.24.88, transferred 2016 to EC 3.4.19.15]

#### EC 3.4.19.16

**Accepted name:** glucosinolate  $\gamma$ -glutamyl hydrolase

**Reaction:** (1) an (*E*)-1-(glutathion-*S*-yl)-*N*-hydroxy- $\omega$ -(methylsulfanyl)alkan-1-imine + H<sub>2</sub>O = an (*E*)-1-(L-cysteinylglycin-*S*-yl)-*N*-hydroxy- $\omega$ -(methylsulfanyl)alkan-1-imine + L-glutamate  
(2) (*E*)-1-(glutathion-*S*-yl)-*N*-hydroxy-2-(1*H*-indol-3-yl)ethan-1-imine + H<sub>2</sub>O = (*E*)-1-(L-cysteinylglycin-*S*-yl)-*N*-hydroxy-2-(1*H*-indol-3-yl)ethan-1-imine + L-glutamate  
(3) (glutathion-*S*-yl)(1*H*-indol-3-yl)acetonitrile + H<sub>2</sub>O = (L-cysteinylglycin-*S*-yl)(1*H*-indol-3-yl)acetonitrile + L-glutamate  
(4) (*Z*)-1-(glutathion-*S*-yl)-*N*-hydroxy-2-phenylethan-1-imine + H<sub>2</sub>O = (*Z*)-1-(L-cysteinylglycin-*S*-yl)-*N*-hydroxy-2-phenylethan-1-imine + L-glutamate

**Other name(s):** GGP1 (gene name); GGP3 (gene name)

**Comments:** This enzyme, characterized from the plant *Arabidopsis thaliana*, participates in the biosynthesis of the plant defense compounds glucosinolates and camalexin. It is the only known plant enzyme capable of hydrolysing the  $\gamma$ -glutamyl residue of glutathione in the cytosol.

**References:** [896]

[EC 3.4.19.16 created 2017]

### EC 3.4.21 Serine endopeptidases

#### EC 3.4.21.1

**Accepted name:** chymotrypsin

**Reaction:** Preferential cleavage: Tyr—|, Trp—|, Phe—|, Leu—|

**Other name(s):** chymotrypsins A and B;  $\alpha$ -chymar ophth; avazyme; chymar; chymotest; enzeon; quimar; quimotrase;  $\alpha$ -chymar;  $\alpha$ -chymotrypsin A;  $\alpha$ -chymotrypsin

**Comments:** Chymotrypsin A is formed from cattle and pig chymotrypsinogen A, several iso-forms being produced according to the number of bonds hydrolysed in the precursor. Chymotrypsin B (formerly listed as EC 3.4.4.6), formed from chymotrypsinogen B, is homologous with chymotrypsin A. Enzymes with specificity similar to that of chymotrypsins A and B have been isolated from many species. In peptidase family S1 (trypsin family)

**References:** [3184, 245, 172, 2303, 2935]

[EC 3.4.21.1 created 1961 as EC 3.4.4.5 and EC 3.4.4.6, transferred 1972 to EC 3.4.21.1]

#### EC 3.4.21.2

**Accepted name:** chymotrypsin C

**Reaction:** Preferential cleavage: Leu—|, Tyr—|, Phe—|, Met—|, Trp—|, Gln—|, Asn—|

**Comments:** Formed from pig chymotrypsinogen C, and from cattle subunit II of procarboxypeptidase A. Reacts more readily with Tos-Leu-CH<sub>2</sub>Cl than Tos-Phe-CH<sub>2</sub>Cl in contrast to chymotrypsin. In peptidase family S1 (trypsin family)

**References:** [2251, 789, 3184]



[EC 3.4.21.2 created 1972]

### EC 3.4.21.3

**Accepted name:** metridin  
**Reaction:** Preferential cleavage: Leu—, Tyr—, Phe—, Met—, Trp—, Gln—, Asn—  
**Other name(s):** Metridium proteinase A; sea anemone protease A; sea anemone proteinase A  
**Comments:** Digestive enzyme from the sea anemone *Metridium senile*.  
**References:** [912, 2769]

[EC 3.4.21.3 created 1972]

### EC 3.4.21.4

**Accepted name:** trypsin  
**Reaction:** Preferential cleavage: Arg—, Lys—  
**Other name(s):**  $\alpha$ -trypsin;  $\beta$ -trypsin; cocoonase; parenzyme; parenzymol; tryptar; trypure; pseudotrypsin; tryptase; tripcellim; sperm receptor hydrolase  
**Comments:** The single polypeptide chain cattle  $\beta$ -trypsin is formed from trypsinogen by cleavage of one peptide bond. Further peptide bond cleavages produce  $\alpha$  and other iso-forms. Isolated as multiple cationic and anionic trypsins [783] from the pancreas of many vertebrates and from lower species including crayfish, insects (cocoonase) and microorganisms (*Streptomyces griseus*) [2389]. Type example of peptidase family S1.  
**References:** [1205, 3101, 2389, 763, 783, 2303, 2883]

[EC 3.4.21.4 created 1961 as EC 3.4.4.4, transferred 1972 to EC 3.4.21.4]

### EC 3.4.21.5

**Accepted name:** thrombin  
**Reaction:** Selective cleavage of Arg—Gly bonds in fibrinogen to form fibrin and release fibrinopeptides A and B  
**Other name(s):** fibrinogenase; thrombase; thrombofort; topical; thrombin-C; tropostasin; activated blood-coagulation factor II; blood-coagulation factor IIa; factor IIa; E thrombin;  $\beta$ -thrombin;  $\gamma$ -thrombin  
**Comments:** Formed from prothrombin. More selective than trypsin and plasmin. In peptidase family S1 (trypsin family).  
**References:** [175, 1772, 1915, 1753, 1799, 546, 437, 1764]

[EC 3.4.21.5 created 1961 as EC 3.4.4.13, transferred 1972 to EC 3.4.21.5]

### EC 3.4.21.6

**Accepted name:** coagulation factor Xa  
**Reaction:** Selective cleavage of Arg—Thr and then Arg—Ile bonds in prothrombin to form thrombin  
**Other name(s):** thrombokinase; prothrombase; prothrombinase; activated blood-coagulation factor X; autoprothrombin C; thromboplastin; plasma thromboplastin; factor Xa; activated Stuart-Prower factor; activated factor X  
**Comments:** A blood coagulation factor formed from the proenzyme factor X by limited proteolysis. Factor X is a glycoprotein composed of a heavy chain and a light chain, which are generated from a precursor protein by the excision of the tripeptide RKR and held together by one or more disulfide bonds. The activated factor Xa converts prothrombin to thrombin in the presence of factor Va, Ca<sup>2+</sup> and phospholipids. Scutellarin (EC 3.4.21.60) has similar specificity, but does not require factor Va.  
**References:** [836, 1320, 546, 1289, 1874, 437]

[EC 3.4.21.6 created 1972, modified 2011]

### EC 3.4.21.7

**Accepted name:** plasmin  
**Reaction:** Selective cleavage of Arg—Thr and then Arg—Ile bonds in prothrombin to form thrombin  
**Other name(s):** fibrinase; fibrinolysin; actase; serum tryptase; thrombolysin  
**Comments:** Formed from plasminogen by proteolysis which results in multiple forms of the active plasmin. In peptidase family S1 (trypsin family).  
**References:** [381, 380, 2433]

[EC 3.4.21.7 created 1961 as EC 3.4.4.14, transferred 1972 to EC 3.4.21.7]

[3.4.21.8 *Transferred entry. kallikrein. Now EC 3.4.21.34 (plasma kallikrein) and EC 3.4.21.35 (tissue kallikrein)*]

[EC 3.4.21.8 created 1972, deleted 1981]

#### EC 3.4.21.9

**Accepted name:** enteropeptidase  
**Reaction:** Activation of trypsinogen by selective cleavage of Lys<sup>6</sup>—Ile bond  
**Other name(s):** enterokinase  
**Comments:** Is not inhibited by protein inhibitors of trypsin. In peptidase family S1 (trypsin family).  
**References:** [1685]

[EC 3.4.21.9 created 1961 as EC 3.4.4.8, transferred 1972 to EC 3.4.21.9]

#### EC 3.4.21.10

**Accepted name:** acrosin  
**Reaction:** Preferential cleavage: Arg—, Lys—  
**Other name(s):** acrosomal proteinase; acrozonase;  $\alpha$ -acrosin;  $\beta$ -acrosin; epsilon-acrosin; acrosomal protease; acrosin amidase  
**Comments:** Occurs in spermatozoa; formed from proacrosin by limited proteolysis. Inhibited by naturally occurring trypsin inhibitors. In peptidase family S1 (trypsin family)  
**References:** [2001, 2684, 1413]

[EC 3.4.21.10 created 1972]

[3.4.21.11 *Transferred entry. elastase. Now EC 3.4.21.37, leukocyte elastase*]

[EC 3.4.21.11 created 1972, deleted 1981]

#### EC 3.4.21.12

**Accepted name:**  $\alpha$ -lytic endopeptidase  
**Reaction:** Preferential cleavage: Ala—, Val— in bacterial cell walls, elastin and other proteins  
**Other name(s):** myxobacter  $\alpha$ -lytic proteinase;  $\alpha$ -lytic proteinase;  $\alpha$ -lytic protease; *Mycobacterium* sorangium  $\alpha$ -lytic proteinase; *Myxobacter* 495  $\alpha$ -lytic proteinase;  $\alpha$ -lytic proteinase; *Myxobacter*  $\alpha$ -lytic proteinase; *Mycobacterium* sorangium  $\alpha$ -lytic proteinase  
**Comments:** From the myxobacterium *Lysobacter enzymogenes*. In peptidase family S1 (trypsin family)  
**References:** [2195, 2303, 708, 262]

[EC 3.4.21.12 created 1972]

[3.4.21.13 *Transferred entry. Phaseolus proteinase. Now EC 3.4.16.6, carboxypeptidase D*]

[EC 3.4.21.13 created 1972, deleted 1978]

[3.4.21.14 *Transferred entry. now EC 3.4.21.67 endopeptidase So*]

[EC 3.4.21.14 created 1961 as EC 3.4.4.16, transferred 1972 to EC 3.4.21.14, modified 1986, deleted 1992]

[3.4.21.15 *Transferred entry. Aspergillus alkaline proteinase. Now EC 3.4.21.63, oryzin*]

[EC 3.4.21.15 created 1972, deleted 1978 (transferred to EC 3.4.21.14, deleted 1992)]

[3.4.21.16 Deleted entry. *Alternaria serine proteinase*]

[EC 3.4.21.16 created 1972, deleted 1992]

[3.4.21.17 Deleted entry. *Arthrobacter serine proteinase*]

[EC 3.4.21.17 created 1972, deleted 1978 [transferred to EC 3.4.21.14, deleted 1992]]

[3.4.21.18 Deleted entry. *Tenebrio α-proteinase*]

[EC 3.4.21.18 created 1972 [EC 3.4.99.24 created 1972, incorporated 1978], deleted 1992]

#### EC 3.4.21.19

**Accepted name:** glutamyl endopeptidase

**Reaction:** Preferential cleavage: Glu—|, Asp—|

**Other name(s):** V8 proteinase; endoproteinase Glu-C; staphylococcal serine proteinase

**Comments:** From *Staphylococcus aureus* strain V8. In appropriate buffer the specificity is restricted to Glu—|. In peptidase family S1 (trypsin family)

**References:** [646, 648, 374]

[EC 3.4.21.19 created 1978]

#### EC 3.4.21.20

**Accepted name:** cathepsin G

**Reaction:** Specificity similar to chymotrypsin C

**Other name(s):** chymotrypsin-like proteinase; neutral proteinase

**Comments:** From azurophil granules of polymorphonuclear leukocytes. In peptidase family S1 (trypsin family)

**References:** [155, 2879, 1159]

[EC 3.4.21.20 created 1978]

#### EC 3.4.21.21

**Accepted name:** coagulation factor VIIa

**Reaction:** Selective cleavage of Arg—|Ile bond in factor X to form factor Xa

**Other name(s):** blood-coagulation factor VIIa; activated blood coagulation factor VII

**Comments:** Formed from the precursor factor VII. The cattle enzyme is more readily inhibited by diisopropyl fluorophosphate than the human [2069]. In peptidase family S1 (trypsin family)

**References:** [2069, 546, 1289, 313]

[EC 3.4.21.21 created 1978]

#### EC 3.4.21.22

**Accepted name:** coagulation factor IXa

**Reaction:** Selective cleavage of Arg—|Ile bond in factor X to form factor Xa

**Other name(s):** activated Christmas factor; blood-coagulation factor IXa; activated blood-coagulation factor IX; auto-prothrombin II; blood platelet cofactor II; activated blood coagulation factor XI

**Comments:** A chymotrypsin homologue, and one of the γ-carboxyglutamic acid-containing blood coagulation factors. The proenzyme factor IX is activated by factor XIa. In peptidase family S1 (trypsin family)

**References:** [835, 546, 1706, 437]

[EC 3.4.21.22 created 1978]

[3.4.21.23 Deleted entry. *Vipera russelli proteinase*]

[EC 3.4.21.23 created 1978, deleted 1992]

[3.4.21.24 Deleted entry. red cell neutral endopeptidase]

[EC 3.4.21.24 created 1978, deleted 1992]

#### EC 3.4.21.25

**Accepted name:** cucumisin

**Reaction:** Hydrolysis of proteins with broad specificity

**Other name(s):** euphorbain; solanain; hurain; tabernamontanain

**Comments:** From the sarcocarp of the musk melon (*Cucumis melo*). In peptidase family S8 (subtilisin family). Other endopeptidases from plants, which are less well characterized but presumably of serine-type, include euphorbain from *Euphorbia cerifera* [1759], solanain from horse-nettle *Solanum elaeagnifolium* [973], hurain from *Hura crepitans* [1296] and tabernamontanain from *Tabernamontana grandiflora* [1295].

**References:** [973, 1296, 1295, 1380, 1379, 1759, 1381]

[EC 3.4.21.25 created 1978 (EC 3.4.21.56 created 1972 as EC 3.4.99.7 transferred 1989 to EC 3.4.21.56, deleted 1992; EC 3.4.99.9 created 1972 deleted 1992; EC 3.4.99.21 created 1972 deleted 1992; EC 3.4.99.23 created 1972 deleted 1992; all covered by EC 3.4.21.25)]

#### EC 3.4.21.26

**Accepted name:** prolyl oligopeptidase

**Reaction:** Hydrolysis of —Pro— and to a lesser extent —Ala— in oligopeptides

**Other name(s):** post-proline cleaving enzyme; proline-specific endopeptidase; post-proline endopeptidase; proline endopeptidase; endoprolylpeptidase; prolyl endopeptidase

**Comments:** Found in vertebrates, plants and *Flavobacterium*. Generally cytosolic, commonly activated by thiol compounds. Type example of peptidase family S9.

**References:** [3103, 2107, 1978, 2409]

[EC 3.4.21.26 created 1978, modified 1981 (EC 3.4.22.18 created 1981, incorporated 1992)]

#### EC 3.4.21.27

**Accepted name:** coagulation factor XIa

**Reaction:** Selective cleavage of Arg—Ala and Arg—Val bonds in factor IX to form factor IXa

**Other name(s):** blood-coagulation factor XIa; activated blood-coagulation factor XI; activated plasma thromboplastin antecedent

**Comments:** In peptidase family S1 (trypsin family), and one of the  $\gamma$ -carboxyglutamic acid-containing blood coagulation factors. The proenzyme factor XI is activated by factor XIIa

**References:** [1563, 437, 834]

[EC 3.4.21.27 created 1978]

[3.4.21.28 Transferred entry. Agkistrodon serine proteinase. Now EC 3.4.21.74, venombin A]

[EC 3.4.21.28 created 1978, deleted 1992]

[3.4.21.29 Transferred entry. Bothrops atrox serine proteinase. Now EC 3.4.21.74, venombin A]

[EC 3.4.21.29 created 1978, deleted 1992]

[3.4.21.30 Transferred entry. Crotalus adamanteus serine proteinase. Now EC 3.4.21.74, venombin A]

[EC 3.4.21.30 created 1978, deleted 1992]

[3.4.21.31 Transferred entry. urokinase. Now EC 3.4.21.73, u-plasminogen activator]

[EC 3.4.21.31 created 1972 as EC 3.4.99.26, transferred 1978 to EC 3.4.21.31, deleted 1992]

#### EC 3.4.21.32

- Accepted name:** brachyurin
- Reaction:** Hydrolysis of proteins, with broad specificity for peptide bonds. Native collagen is cleaved about 75% of the length of the molecule from the N-terminus. Low activity on small molecule substrates of both trypsin and chymotrypsin
- Other name(s):** *Uca pugilator* collagenolytic proteinase; crab protease I; crab protease II
- Comments:** From hepatopancreas of the fiddler crab, *Uca pugilator*. In peptidase family S1 (trypsin family). Other serine endopeptidases that degrade collagen, but are not listed separately here, include a second endopeptidase from *Uca pugilator* [3158], digestive enzymes from other decapod crustacea [1475, 1745], and an enzyme from the fungus *Entomophthora coronata* [1210].
- References:** [1210, 965, 3159, 3158, 1475, 1745]

[EC 3.4.21.32 created 1978]

[3.4.21.33 Deleted entry. *Entomophthora collagenolytic proteinase*]

[EC 3.4.21.33 created 1978, deleted 1992]

#### EC 3.4.21.34

- Accepted name:** plasma kallikrein
- Reaction:** Selective cleavage of some Arg— and Lys— bonds, including Lys—Arg and Arg—Ser in (human) kininogen to release bradykinin
- Other name(s):** serum kallikrein; kininogenin; kallikrein I; kallikrein II; kininogenase; kallikrein; callicrein; glumorin; padreatin; padutin; kallidinogenase; bradykininogenase; pancreatic kallikrein; onokrein P; diliminal D; depot-Padutin; urokallikrein; urinary kallikrein
- Comments:** Formed from plasma prokallikrein (Fletcher factor) by factor XIIa. Activates coagulation factors XII, VII and plasminogen. Selective for Arg > Lys in P1, in small molecule substrates.
- References:** [1100, 1876, 2663, 2608, 2978]

[EC 3.4.21.34 created 1965 as EC 3.4.4.21, transferred 1972 to EC 3.4.21.8, part transferred 1981 to EC 3.4.21.34]

#### EC 3.4.21.35

- Accepted name:** tissue kallikrein
- Reaction:** Preferential cleavage of Arg— bonds in small molecule substrates. Highly selective action to release kallidin (lysyl-bradykinin) from kininogen involves hydrolysis of Met— or Leu—. The rat enzyme is unusual in liberating bradykinin directly from autologous kininogens by cleavage at two Arg— bonds [5]
- Other name(s):** glandular kallikrein; pancreatic kallikrein; submandibular kallikrein; submaxillary kallikrein; kidney kallikrein; urinary kallikrein; kallikrein; salivary kallikrein; kininogenin; kininogenase; callicrein; glumorin; padreatin; padutin; kallidinogenase; bradykininogenase; depot-padutin; urokallikrein; diliminal D; onokrein P
- Comments:** Formed from tissue prokallikrein by activation with trypsin. In peptidase family S1 (trypsin family). A large number of tissue kallikrein-related sequences have been reported for rats [3195] and mice [723], though fewer seem to exist in other mammals. The few that have been isolated and tested on substrates include mouse  $\gamma$ -renin (EC 3.4.21.54), submandibular proteinase A [60, 208], epidermal growth-factor-binding protein, nerve growth factor  $\gamma$ -subunit, rat tonin [3,4,9], submaxillary proteinases A and B [1398], T-kininogenase [3230], kallikreins k7 and k8 [694] and human prostate-specific antigen ( $\gamma$ -seminoprotein, [26])
- References:** [764, 60, 2259, 1009, 1397, 26, 723, 763, 844, 1398, 119, 226, 400, 889, 208, 3195, 694, 3230]

[EC 3.4.21.35 created 1965 as EC 3.4.4.21, transferred 1972 to EC 3.4.21.8, part transferred 1981 to EC 3.4.21.35]

#### EC 3.4.21.36

- Accepted name:** pancreatic elastase

**Reaction:** Hydrolysis of proteins, including elastin. Preferential cleavage: Ala—  
**Other name(s):** pancrotopetidase E; pancreatic elastase I; elastase; elaszym; serine elastase  
**Comments:** Formed by activation of proelastase from mammalian pancreas by trypsin. In peptidase family S1 (trypsin family). Formerly included in EC 3.4.21.11  
**References:** [2646, 1059, 1407, 214, 251]

[EC 3.4.21.36 created 1981 (EC 3.4.4.7 created 1961, transferred 1972 to EC 3.4.21.11 created 1972, part incorporated 1984)]

#### EC 3.4.21.37

**Accepted name:** leukocyte elastase  
**Reaction:** Hydrolysis of proteins, including elastin. Preferential cleavage Val— ; Ala—  
**Other name(s):** lysosomal elastase; neutrophil elastase; polymorphonuclear leukocyte elastase; elastase; elaszym; serine elastase; lysosomal elastase; granulocyte elastase  
**Comments:** Differs from pancreatic elastase in specificity on synthetic substrates and in inhibitor sensitivity. In peptidase family S1 (trypsin family). Formerly included in EC 3.4.21.11  
**References:** [156, 1059, 2760, 251]

[EC 3.4.21.37 created 1981 (EC 3.4.4.7 created 1961, transferred 1972 to EC 3.4.21.11 created 1972, part incorporated 1984)]

#### EC 3.4.21.38

**Accepted name:** coagulation factor XIIa  
**Reaction:** Selective cleavage of Arg—Ile bonds in factor VII to form factor VIIa and factor XI to form factor XIa  
**Other name(s):** Hageman factor (activated); blood-coagulation factor XIIIf; activated  $\beta$  blood-coagulation factor XII; prealbumin activator; Hageman factor  $\beta$ -fragment; Hageman factor fragment HFf; blood-coagulation factor XIIa $\beta$ ; prekallikrein activator; kallikreinogen activator  
**Comments:** Also activates plasminogen and plasma prokallikrein. Formed from the proenzyme, factor XII, by plasma kallikrein or factor XIIa. In peptidase family S1 (trypsin family)  
**References:** [837, 437, 2342, 833, 2662]

[EC 3.4.21.38 created 1981]

#### EC 3.4.21.39

**Accepted name:** chymase  
**Reaction:** Preferential cleavage: Phe— ; Tyr— ; Trp— ; Leu—  
**Other name(s):** mast cell protease I; skeletal muscle protease; skin chymotryptic proteinase; mast cell serine proteinase, chymase; skeletal muscle (SK) protease  
**Comments:** In mast cell granules. In peptidase family S1 (trypsin family)  
**References:** [3212, 2321, 1327]

[EC 3.4.21.39 created 1981]

[3.4.21.40 Deleted entry. submandibular proteinase A]

[EC 3.4.21.40 created 1981, deleted 1992]

#### EC 3.4.21.41

**Accepted name:** complement subcomponent C<sup>1r</sup>  
**Reaction:** Selective cleavage of Lys(or Arg)—Ile bond in complement subcomponent C1s to form C<sup>1s</sup> (EC 3.4.21.42)  
**Other name(s):** activated complement C1r; C<sup>1r</sup> esterase; activated complement C1r  
**Comments:** Activated from proenzyme C<sup>1r</sup> in plasma during activation of the complement system by the "classical" route. In peptidase family S1 (trypsin family)  
**References:** [2664, 1662, 2000]

[EC 3.4.21.41 created 1981]

**EC 3.4.21.42**

**Accepted name:** complement subcomponent C<sup>1s</sup>

**Reaction:** Cleavage of Arg—Ala bond in complement component C4 to form C4a and C4b, and Lys(or Arg)—Lys bond in complement component C2 to form C2a and C2b: the "classical" pathway C3 convertase

**Other name(s):** C1 esterase; activated complement C1s; complement C<sup>1r</sup>

**Comments:** Activated from proenzyme C1s in plasma by complement subcomponent C<sup>1r</sup>. In peptidase family S1 (trypsin family)

**References:** [2664, 1767, 2000, 2684]

[EC 3.4.21.42 created 1981]

**EC 3.4.21.43**

**Accepted name:** classical-complement-pathway C3/C5 convertase

**Reaction:** Selective cleavage of Arg—Ser bond in complement component C3  $\alpha$ -chain to form C3a and C3b, and Arg— bond in complement component C5  $\alpha$ -chain to form C5a and C5b

**Other name(s):** C3 convertase; C<sup>42</sup>; C4b,2a; C5 convertase; C<sup>423</sup>; C4b,2a,3b; C42; C5 convertase; C423; C4b,2a,3b; complement C.hivin.4.hivin2; complement C3 convertase

**Comments:** A complex of complement fragments C4b, C2a and C2b. C2a contains the active site, C2b the site for C4b binding. C2a and C2b are formed by cleavage of proenzyme C2 by complement subcomponent C<sup>1s</sup>. Cleavage of C5 requires complement fragment C3b which binds C5 and renders it susceptible to cleavage by the C4b,2a complex. Includes former EC 3.4.21.44. Complement component C2a is in peptidase family S1 (trypsin family)

**References:** [1420, 2000]

[EC 3.4.21.43 created 1981 (EC 3.4.21.44 created 1981, incorporated 1984)]

[3.4.21.44 Transferred entry. complement component C5 convertase. Now EC 3.4.21.43, classical-complement-pathway C3/C5 convertase]

[EC 3.4.21.44 created 1981, deleted 1984]

**EC 3.4.21.45**

**Accepted name:** complement factor I

**Reaction:** Inactivates complement subcomponents C3b, iC3b and C4b by proteolytic cleavage

**Other name(s):** complement component C3b inactivator; C3b inactivator; C3b/C4b inactivator; C3bINA; complement C3b/C4b inactivator; complement C4b inactivator; conglutinogen-activating factor C; complement C3b inactivator; factor I; complement C4bi

**Comments:** Cleavage of complement subcomponent C3b requires its binding to cofactor factor H or complement receptor CR1; cleavage of iC3b requires complement receptor CR1; cleavage of C4b requires C4b-binding protein. In peptidase family S1 (trypsin family)

**References:** [2023, 509, 2000]

[EC 3.4.21.45 created 1981]

**EC 3.4.21.46**

**Accepted name:** complement factor D

**Reaction:** Selective cleavage of Arg—Lys bond in complement factor B when in complex with complement subcomponent C3b or with cobra venom factor

**Other name(s):** C3 proactivator convertase; properdin factor D esterase; factor D; factor D (complement)

**Comments:** A component of the alternative pathway of complement activation. This reaction is analogous to the activation of complement component C2 by complement subcomponent C<sup>1s</sup>. In peptidase family S1 (trypsin family)

**References:** [2401, 2000]

[EC 3.4.21.46 created 1981]

#### EC 3.4.21.47

**Accepted name:** alternative-complement-pathway C3/C5 convertase

**Reaction:** Cleavage of Arg—Ser bond in complement component C3  $\alpha$ -chain to yield C3a and C3b, and Arg— bond in complement component C5  $\alpha$ -chain to yield C5a and C5b

**Other name(s):** complement component C3/C5 convertase (alternative); proenzyme factor B; properdin factor B; C3 proactivator; glycine-rich  $\beta$ -glycoprotein; heat-labile factor; C3 convertase; C3b,Bb,CVF,Bb,C5 convertase; (C3b)n,Bb; complement C 3(C 5) convertase (amplification); alternative complement pathway C3(C5) convertase; C5 convertase; CVF,Bb; (CVF)-dependent glycine-rich- $\beta$ -glucoprotein; cobra venom factor-dependent C3 convertase

**Comments:** A bimolecular complex of complement fragment Bb with either C3b or cobra venom factor; Bb contains the active site. Bb is formed by cleavage of proenzyme factor B by factor D. Cleavage of complement component C5 requires additional C3b which binds C5 and renders it susceptible to cleavage by C3b,Bb complex. C3b,Bb is stabilized in plasma by factor P. Complement factor B is in peptidase family S1 (trypsin family)

**References:** [1421, 1980, 2000]

[EC 3.4.21.47 created 1981]

#### EC 3.4.21.48

**Accepted name:** cerevisin

**Reaction:** Hydrolysis of proteins with broad specificity, and of Bz-Arg-OEt Ac-Tyr-OEt. Does not hydrolyse peptide amides

**Other name(s):** yeast proteinase B; proteinase *yscB*; baker's yeast proteinase B; brewer's yeast proteinase; peptidase  $\beta$

**Comments:** From *Saccharomyces cerevisiae* (baker's yeast, brewer's yeast). In peptidase family S8 (subtilisin family), but contains a Cys residue near the active site His, and is inhibited by mercurials. Proteinase *ycaB* is a similar enzyme from the yeast *Candida albicans* [738]

**References:** [751, 1505, 738, 1944]

[EC 3.4.21.48 created 1972 as EC 3.4.22.9, transferred 1981 to EC 3.4.21.48]

#### EC 3.4.21.49

**Accepted name:** hypodermin C

**Reaction:** Hydrolysis of proteins including native collagen at —Ala bond leaving an N-terminal (75%) and a C-terminal (25%) fragment

**Other name(s):** Hypoderma collagenase

**Comments:** From the larva of a warble fly, *Hypoderma lineatum*. Little action on small molecule substrates of trypsin, chymotrypsin, elastase or microbial collagenases. In peptidase family S1 (trypsin family)

**References:** [1613, 1615, 1614]

[EC 3.4.21.49 created 1981]

#### EC 3.4.21.50

**Accepted name:** lysyl endopeptidase

**Reaction:** Preferential cleavage: Lys—, including -Lys—Pro-

**Other name(s):** *Achromobacter* proteinase I (also see Comment); *Achromobacter lyticus* alkaline proteinase I; protease I; achromopeptidase; lysyl bond specific proteinase



**Comments:** From *Achromobacter lyticus* [2984]. Enzymes with similar specificity are produced by *Lysobacter enzymogenes* (Endoproteinase Lys-C; [1315]) and *Pseudomonas aeruginosa* (Ps-1; [692]). In peptidase family S1 (trypsin family)

**References:** [1831, 1830, 1315, 692, 2157, 2984]

[EC 3.4.21.50 created 1983]

[3.4.21.51 Deleted entry. *Leukocyte-membrane neutral endopeptidase*]

[EC 3.4.21.51 created 1984, deleted 1992]

[3.4.21.52 Deleted entry. *Cathepsin R*]

[EC 3.4.21.52 created 1981 as EC 3.4.99.33, transferred 1984 to EC 3.4.21.52, deleted 1992]

#### EC 3.4.21.53

**Accepted name:** endopeptidase La

**Reaction:** Hydrolysis of proteins in presence of ATP

**Other name(s):** ATP-dependent serine proteinase; lon proteinase; protease La; proteinase La; ATP-dependent lon proteinase; ATP-dependent protease La; *Escherichia coli* proteinase La; *Escherichia coli* serine proteinase La; gene lon protease; gene lon proteins; PIM1 protease; PIM1 proteinase; serine protease La

**Comments:** Product of the *lon* gene in *Escherichia coli*. ATP hydrolysis is linked with peptide bond hydrolysis; vanadate inhibits both reactions. Type example of peptidase family S16. A similar enzyme occurs in animal mitochondria

**References:** [587, 1602, 432]

[EC 3.4.21.53 created 1986]

#### EC 3.4.21.54

**Accepted name:**  $\gamma$ -renin

**Reaction:** Cleavage of the Leu—Leu bond in synthetic tetradecapeptide renin substrate (horse), to produce angiotensin I, but not active on natural angiotensinogen, unlike renin (EC 3.4.23.15). Also hydrolyses Bz-Arg-*p*-nitroanilide

**Comments:** A member of the tissue kallikrein family, from submandibular glands of male mice. In peptidase family S1 (trypsin family)

**References:** [2298, 653]

[EC 3.4.21.54 created 1986]

#### EC 3.4.21.55

**Accepted name:** venombin AB

**Reaction:** Selective cleavage at Arg— bonds in fibrinogen to form fibrin and release fibrinopeptides A and B gabonase; okinaxobin II; *Bitis gabonica* venom serine proteinase; afaâcytin

**Other name(s):** gabonase; okinaxobin II; *Bitis gabonica* venom serine proteinase; afaâcytin

**Comments:** From the venom of the Gaboon viper *Bitis gabonica*. Activates Factor XIII. Not inhibited by antithrombin III/heparin or hirudin, unlike EC 3.4.21.5, thrombin

**References:** [2286]

[EC 3.4.21.55 created 1989]

[3.4.21.56 Deleted entry. *euphorbain*. Now considered EC 3.4.21.25, *cucumis*in]

[EC 3.4.21.56 created 1972 as EC 3.4.99.7, transferred 1989 to EC 3.4.21.56, deleted 1992]

#### EC 3.4.21.57

**Accepted name:** leucyl endopeptidase

**Reaction:** Hydrolysis of proteins. Preferential cleavage: Leu— in small molecule substrates  
**Other name(s):** plant Leu-proteinase; leucine-specific serine proteinase; leucine endopeptidase; spinach serine proteinase (leucine specific); spinach leucine-specific serine proteinase; Leu-proteinase  
**Comments:** From leaves of the spinach plant (*Spinacia oleracea*)  
**References:** [19, 18]

[EC 3.4.21.57 created 1989]

[3.4.21.58 Deleted entry. prohormone serine proteinase]

[EC 3.4.21.58 created 1989, deleted 1992]

#### EC 3.4.21.59

**Accepted name:** tryptase  
**Reaction:** Preferential cleavage: Arg—, Lys—, but with more restricted specificity than trypsin  
**Other name(s):** mast cell tryptase; mast cell protease II; skin tryptase; lung tryptase; pituitary tryptase; mast cell neutral proteinase; mast cell tryptase; mast cell neutral proteinase; mast cell serine proteinase II; mast cell proteinase II; mast cell serine proteinase tryptase; rat mast cell protease II; tryptase M  
**Comments:** Occurs as a tetrameric molecule with high affinity for heparin, in mast cell granules. In peptidase family S1 (trypsin family). Not inhibited by  $\alpha_1$ -proteinase inhibitor or  $\alpha_2$ -macroglobulin  
**References:** [2878, 1436, 506, 1065, 3050]

[EC 3.4.21.59 created 1992]

#### EC 3.4.21.60

**Accepted name:** scutellarin  
**Reaction:** Selective cleavage of Arg—Thr and Arg—Ile in prothrombin to form thrombin and two inactive fragments  
**Other name(s):** taipan activator; *Oxyuranus scutellatus* prothrombin-activating proteinase  
**Comments:** From the venom of the Taipan snake (*Oxyuranus scutellatus*). Converts prothrombin to thrombin. Specificity is similar to that of Factor Xa (EC 3.4.21.6). However, unlike Factor Xa this enzyme can cleave its target in the absence of coagulation Factor Va. Activity is potentiated by phospholipid and  $\text{Ca}^{2+}$  which binds via  $\gamma$ -carboxyglutamic acid residues. Similar enzymes are known from the venom of other Australian elapid snakes, including *Pseudonaja textilis textilis*, *Oxyuranus microlepidotus* and *Demansia nuchalis affinis*.  
**References:** [3094, 2739]

[EC 3.4.21.60 created 1978 as EC 3.4.99.28, transferred 1992 to EC 3.4.21.60, modified 2010, modified 2011]

#### EC 3.4.21.61

**Accepted name:** kexin  
**Reaction:** Cleavage of -Lys-Arg— and -Arg-Arg— bonds to process yeast  $\alpha$ -factor pheromone and killer toxin precursors  
**Other name(s):** yeast KEX2 protease; proteinase yscF; prohormone-processing endoprotease; paired-basic endopeptidase; yeast cysteine proteinase F (misleading); paired-basic endopeptidase; adrenorphin-Gly-generating enzyme; endoproteinase Kex2p; gene KEX2 dibasic proteinase; Kex 2p proteinase; Kex2 endopeptidase; Kex2 endoprotease; Kex2 endoproteinase; Kex2 protease; proteinase Kex2p; Kex2-like precursor protein processing endoprotease; prohormone-processing KEX2 proteinase; prohormone-processing proteinase; proprotein convertase; protease KEX2; Kex2 proteinase; Kex2-like endoproteinase  
**Comments:** A  $\text{Ca}^{2+}$ -activated peptidase of peptidase family S8, containing Cys near the active site His, and inhibited by *p*-mercuribenzoate. Similar enzymes occur in mammals.  
**References:** [1346, 10, 1940, 862, 1941]

[EC 3.4.21.61 created 1989 as EC 3.4.22.23, transferred 1992 to EC 3.4.21.61]

#### EC 3.4.21.62

- Accepted name:** subtilisin
- Reaction:** Hydrolysis of proteins with broad specificity for peptide bonds, and a preference for a large uncharged residue in P1. Hydrolyses peptide amides
- Other name(s):** alcalase; alcalase 0.6L; alcalase 2.5L; ALK-enzyme; bacillopeptidase A; bacillopeptidase B; *Bacillus subtilis* alkaline proteinase biopraser; biopraser AL 15; biopraser APL 30; colistinase; (see also comments); subtilisin J; subtilisin S41; subtilisin Sendai; subtilisin GX; subtilisin E; subtilisin BL; genenase I; esperase; maxatase; alcalase; thermoase PC 10; protease XXVII; thermoase; superase; subtilisin DY; subtilopeptidase; SP 266; savinase 8.0L; savinase 4.0T; kazusase; protease VIII; opticlean; *Bacillus subtilis* alkaline proteinase; protin A 3L; savinase; savinase 16.0L; savinase 32.0 L EX; orientase 10B; protease S
- Comments:** Subtilisin is a serine endopeptidase, type example of peptidase family S8. It contains no cysteine residues (although these are found in homologous enzymes). Species variants include subtilisin BPN' (also subtilisin B, subtilopeptidase B, subtilopeptidase C, Nagarse, Nagarse proteinase, subtilisin Novo, bacterial proteinase Novo) and subtilisin Carlsberg (subtilisin A, subtilopeptidase A, alcalase Novo). Similar enzymes are produced by various *Bacillus subtilis* strains and other *Bacillus* species [2215, 2275]
- References:** [2215, 1814, 2275, 2063, 1232, 2303]

[EC 3.4.21.62 created 1992 (EC 3.4.21.14 created 1961 as EC 3.4.4.16, transferred 1972 to EC 3.4.21.14, modified 1986, part incorporated 1992)]

#### EC 3.4.21.63

- Accepted name:** oryzin
- Reaction:** Hydrolysis of proteins with broad specificity, and of Bz-Arg-OEt > Ac-Tyr-OEt. Does not hydrolyse peptide amides
- Other name(s):** *Aspergillus* alkaline proteinase; aspergillopeptidase B; API 21; aspergillopepsin B; aspergillopepsin F; *Aspergillus candidus* alkaline proteinase; *Aspergillus flavus* alkaline proteinase; *Aspergillus melleus* semi-alkaline proteinase; *Aspergillus oryzae* alkaline proteinase; *Aspergillus parasiticus* alkaline proteinase; *Aspergillus* serine proteinase; *Aspergillus sydowi* alkaline proteinase; *Aspergillus soya* alkaline proteinase; *Aspergillus melleus* alkaline proteinase; *Aspergillus sulphureus* alkaline proteinase; prozyme; P 5380; kyorinase; seaprose S; semi-alkaline protease; sumizyme MP; prozyme 10; onoprose; onoprose SA; protease P; promelase
- Comments:** A peptidase of family S8 (subtilisin family), not containing cysteine, that is the predominant extracellular alkaline endopeptidase of the mold *Aspergillus oryzae*. Identical or closely related enzymes are produced by *A. flavus* and *A. sojae* [2,3,4]
- References:** [2035, 1089, 2998, 1969, 2735]

[EC 3.4.21.63 created 1992 (EC 3.4.21.14 created 1961 as EC 3.4.4.16, transferred 1972 to EC 3.4.21.14, modified 1986, part incorporated 1992)]

#### EC 3.4.21.64

- Accepted name:** peptidase K
- Reaction:** Hydrolysis of keratin, and of other proteins with subtilisin-like specificity. Hydrolyses peptide amides
- Other name(s):** *Tritirachium* alkaline proteinase; *Tritirachium album* serine proteinase; proteinase K; *Tritirachium album* proteinase K; endopeptidase K
- Comments:** From the mold *Tritirachium album* Limber. A peptidase of family S8 (subtilisin family) containing two disulfide bridges and one free Cys near the active site His. Formerly included in EC 3.4.21.14
- References:** [677, 1972, 1530, 1310, 209]

[EC 3.4.21.64 created 1992 (EC 3.4.21.14 created 1961 as EC 3.4.4.16, transferred 1972 to EC 3.4.21.14, modified 1986, part incorporated 1992)]

#### EC 3.4.21.65

- Accepted name:** thermomycolin  
**Reaction:** Rather nonspecific hydrolysis of proteins. Preferential cleavage: Ala—, Tyr—, Phe— in small molecule substrates  
**Other name(s):** thermomycolase  
**Comments:** A peptidase of family S8 (subtilisin family) from the thermophilic fungus *Malbranchea pulchella* var. *sulfurea* containing Cys, but not inhibited by *p*-mercuribenzoate. Very thermostable. Formerly included in EC 3.4.21.14  
**References:** [884]

[EC 3.4.21.65 created 1992 (EC 3.4.21.14 created 1961 as EC 3.4.4.16, transferred 1972 to EC 3.4.21.14, modified 1986, part incorporated 1992)]

#### EC 3.4.21.66

- Accepted name:** thermitase  
**Reaction:** Hydrolysis of proteins, including collagen  
**Other name(s):** thermophilic *Streptomyces* serine proteinase; *Thermoactinomyces vulgaris* serine proteinase  
**Comments:** A peptidase of family S8 (subtilisin family) from *Thermoactinomyces vulgaris* containing a single Cys, near the active site His, and inhibited by *p*-mercuribenzoate. The N-terminal extension of the polypeptide chain relative to subtilisin contributes to Ca<sup>2+</sup>-binding and the high thermostability. The amino acid composition and properties of the thermostable enzyme from *Streptomyces rectus* var. *proteolyticus* (formerly included in EC 3.4.21.14) are closely similar [1942, 264].  
**References:** [1942, 264, 1474, 1883, 2899]

[EC 3.4.21.66 created 1992]

#### EC 3.4.21.67

- Accepted name:** endopeptidase So  
**Reaction:** Hydrolysis of proteins, but not Bz-Tyr-OEt, Ac-Phe-β-naphthylester, or Bz-Arg-OEt  
**Other name(s):** *E. coli* cytoplasmic proteinase; proteinase So; *Escherichia coli* serine proteinase So  
**Comments:** An *Escherichia coli* cytoplasmic endopeptidase formerly included in EC 3.4.21.14. Inhibited by Tos-Phe-CH<sub>2</sub>Cl, but not by Tos-Lys-CH<sub>2</sub>Cl  
**References:** [943, 447]

[EC 3.4.21.67 created 1992 (EC 3.4.21.14 created 1961 as EC 3.4.4.16, transferred 1972 to EC 3.4.21.14, modified 1986, part incorporated 1992)]

#### EC 3.4.21.68

- Accepted name:** *t*-plasminogen activator  
**Reaction:** Specific cleavage of Arg—Val bond in plasminogen to form plasmin  
**Other name(s):** tissue plasminogen activator; plasminogen activator, tissue-type; tissue-type plasminogen activator; tPA; t-PA  
**Comments:** A peptidase of family S1 (trypsin family) from a wide variety of mammalian tissues, especially endothelial cells. Secreted as a single chain precursor which is cleaved to a two-chain form by plasmin. Activity is considerably enhanced by fibrin. Formerly included in EC 3.4.21.31 and EC 3.4.99.26  
**References:** [2255, 1738, 2261, 3059, 891, 467]

[EC 3.4.21.68 created 1972 as EC 3.4.99.26, transferred 1978 as EC 3.4.21.31, part transferred 1992 to EC 3.4.21.68]

#### EC 3.4.21.69

- Accepted name:** protein C (activated)  
**Reaction:** Degradation of blood coagulation factors Va and VIIIa  
**Other name(s):** blood-coagulation factor XIVa; activated blood coagulation factor XIV; activated protein C; autoprothrombin II-A; protein Ca; APC; GSAPC

**Comments:** A peptidase of family S1 (trypsin family), one of the  $\gamma$ -carboxyglutamic acid-containing coagulation factors. Formed from protein C, the proenzyme that circulates in plasma, by the action of a complex of thrombin with thrombomodulin, or by serine endopeptidases present in several snake venoms

**References:** [716, 717]

[EC 3.4.21.69 created 1992]

#### EC 3.4.21.70

**Accepted name:** pancreatic endopeptidase E

**Reaction:** Preferential cleavage: Ala—|. Does not hydrolyse elastin

**Other name(s):** cholesterol-binding proteinase; proteinase E; cholesterol-binding serine proteinase; pancreatic propease E; pancreatic proteinase E; cholesterol-binding pancreatic proteinase; CBPP; pancreas E proteinase

**Comments:** A peptidase of family S1 (trypsin family) from pancreatic juice. Unlike elastases, has an acidic pI. Binds cholesterol

**References:** [1792, 2626]

[EC 3.4.21.70 created 1992]

#### EC 3.4.21.71

**Accepted name:** pancreatic elastase II

**Reaction:** Preferential cleavage: Leu—|, Met—| and Phe—|. Hydrolyses elastin

**Other name(s):** pancreatic elastase 2

**Comments:** A peptidase of family S1 (trypsin family) formed by activation of proelastase II from mammalian pancreas by trypsin. Usually, only one of the pancreatic elastases (see also EC 3.4.21.36) is expressed in a given species; human pancreatic elastase is of type II

**References:** [784, 2643]

[EC 3.4.21.71 created 1992]

#### EC 3.4.21.72

**Accepted name:** IgA-specific serine endopeptidase

**Reaction:** Cleavage of immunoglobulin A molecules at certain Pro—| bonds in the hinge region. No small molecule substrates are known

**Other name(s):** IgA protease; IgA proteinase; IgA-specific proteinase; immunoglobulin A protease; immunoglobulin A proteinase

**Comments:** Species variants differing slightly in specificity are secreted by Gram-negative bacteria *Neisseria gonorrhoeae* and *Haemophilus influenzae*. Type example of peptidase family S6. Some other bacterial endopeptidases with similar specificity are of metallo- type (see EC 3.4.24.13, IgA-specific metalloendopeptidase)

**References:** [2292, 116]

[EC 3.4.21.72 created 1992]

#### EC 3.4.21.73

**Accepted name:** u-plasminogen activator

**Reaction:** Specific cleavage of Arg—|Val bond in plasminogen to form plasmin

**Other name(s):** urokinase; urinary plasminogen activator; cellular plasminogen activator; urokinase-type plasminogen activator; double-chain urokinase-type plasminogen activator; two-chain urokinase-type plasminogen activator; urokinase plasminogen activator; uPA; u-PA; abbokinase; urinary esterase A

**Comments:** Formed from the inactive precursor by action of plasmin or plasma kallikrein. Differs in structure from *t*-plasminogen activator (EC 3.4.21.68), and does not bind to fibrin. In peptidase family S1 (trypsin family). Formerly included in EC 3.4.21.31 and EC 3.4.99.26

**References:** [1740, 1738, 2505, 467, 1686]

[EC 3.4.21.73 created 1972 as EC 3.4.99.26, transferred 1978 as EC 3.4.21.31, part transferred 1992 to EC 3.4.21.73]

#### EC 3.4.21.74

**Accepted name:** venombin A

**Reaction:** Selective cleavage of Arg— bond in fibrinogen, to form fibrin, and release fibrinopeptide A. The specificity of further degradation of fibrinogen varies with species origin of the enzyme

**Other name(s):**  $\alpha$ -fibrinogenase; habutobin; zinc metalloproteinase Cbfb1.1; zinc metalloproteinase Cbfb1.2; zinc metalloproteinase Cbfb2; ancrod; (see also Comments)

**Comments:** A somewhat thrombin-like enzyme from venoms of snakes of the viper/rattlesnake group. Species variants of the enzyme include ancrod from *Agkistrodon rhodostoma* (Malayan pit viper) (formerly EC 3.4.21.28) [2106], batroxobin from *Bothrops atrox* (South American pit viper) (formerly EC 3.4.21.29) [2774, 1269] and crotalase from *Crotalus adamanteus* (Eastern diamondback rattlesnake) (formerly EC 3.4.21.30) [1813, 2665]. In peptidase family S1 (trypsin family). Does not require activation by  $\text{Ca}^{2+}$

**References:** [2106, 2774, 1813, 2665, 1269]

[EC 3.4.21.74 created 1992 (EC 3.4.21.28, EC 3.4.21.29 and 3.4.21.30 all created 1978 and incorporated 1992)]

#### EC 3.4.21.75

**Accepted name:** furin

**Reaction:** Release of mature proteins from their proproteins by cleavage of -Arg-Xaa-Yaa-Arg— bonds, where Xaa can be any amino acid and Yaa is Arg or Lys. Releases albumin, complement component C3 and von Willebrand factor from their respective precursors

**Other name(s):** prohormone convertase; dibasic processing enzyme; PACE; paired basic amino acid cleaving enzyme; paired basic amino acid converting enzyme; serine proteinase PACE; PC1; SPC3; proprotein convertase

**Comments:** One of a group of peptidases in peptidase family S8 (subtilisin family) that is structurally and functionally similar to kexin. All are activated by  $\text{Ca}^{2+}$ , contain Cys near the active site His, and are inhibited by *p*-mercuribenzoate. At least three related enzymes are recognized in mammals: PC2, PC3 and PC4, which have somewhat different specificities

**References:** [564, 563, 1079, 2604, 2761]

[EC 3.4.21.75 created 1993]

#### EC 3.4.21.76

**Accepted name:** myeloblastin

**Reaction:** Hydrolysis of proteins, including elastin, by preferential cleavage: -Ala— > -Val—

**Other name(s):** leukocyte proteinase 3; leukocyte proteinase 4; Wegener's granulomatosis autoantigen; proteinase PR-3; proteinase-3; PMNL proteinase

**Comments:** From polymorphonuclear leukocyte granules. In peptidase family S1 (trypsin family). Not inhibited by secretory leukocyte proteinase inhibitor

**References:** [1584, 2374, 314, 1367]

[EC 3.4.21.76 created 1993]

#### EC 3.4.21.77

**Accepted name:** semenogelase

**Reaction:** Preferential cleavage: -Tyr—

**Other name(s):** prostate-specific antigen;  $\alpha$ -seminoprotein; seminin; P-30 antigen; antigen (human clone HPSA-1 prostate-specific protein moiety reduced);  $\gamma$ -seminoglycoprotein (human protein moiety reduced);  $\gamma$ -SM; antigen PSA (human prostate-specific); human glandular kallikrein; antigen PSA (human clone SP1 protein moiety reduced)

**Comments:** A peptidase of family S1 (trypsin family) from seminal plasma. Slowly inhibited by  $\alpha_1$ -antichymotrypsin

**References:** [613, 443]

[EC 3.4.21.77 created 1993]

#### EC 3.4.21.78

**Accepted name:** granzyme A

**Reaction:** Hydrolysis of proteins, including fibronectin, type IV collagen and nucleolin. Preferential cleavage: -Arg-|, -Lys-| >> -Phe-| in small molecule substrates

**Other name(s):** CTLA3; HuTPS; T-cell associated protease 1; cytotoxic T lymphocyte serine protease; TSP-1; T-cell derived serine proteinase

**Comments:** From cytotoxic T lymphocyte granules. In peptidase family S1 (trypsin family). The human enzyme does not cleave Phe-|

**References:** [2667, 895, 2138]

[EC 3.4.21.78 created 1993]

#### EC 3.4.21.79

**Accepted name:** granzyme B

**Reaction:** Preferential cleavage: -Asp-| >> -Asn-| > -Met-|, -Ser-|

**Other name(s):** CTLA1; CCPII; cytotoxic cell proteinase-1; granzyme G; granzyme H; CCP<sub>1</sub> proteinase

**Comments:** From cytotoxic T lymphocyte granules. In peptidase family S1 (trypsin family)

**References:** [2568, 2138, 2297]

[EC 3.4.21.79 created 1993]

#### EC 3.4.21.80

**Accepted name:** streptogrisin A

**Reaction:** Hydrolysis of proteins with specificity similar to chymotrypsin

**Other name(s):** *Streptomyces griseus* protease A; protease A; proteinase A; *Streptomyces griseus* proteinase A; *Streptomyces griseus* serine proteinase 3; *Streptomyces griseus* serine proteinase A

**Comments:** From *Streptomyces griseus*. A component of Pronase, in family S1 (trypsin family). Not inhibited by Tos-Phe-CH<sub>2</sub>Cl or ovomucoid

**References:** [1331, 2656, 1306, 576, 1110]

[EC 3.4.21.80 created 1993]

#### EC 3.4.21.81

**Accepted name:** streptogrisin B

**Reaction:** Hydrolysis of proteins with trypsin-like specificity

**Other name(s):** *Streptomyces griseus* protease B; pronase B; serine proteinase B; *Streptomyces griseus* proteinase B; *Streptomyces griseus* proteinase 1; *Streptomyces griseus* serine proteinase B

**Comments:** From *Streptomyces griseus*. A component of Pronase, in peptidase family S1 (trypsin family), distinct from *Streptomyces* trypsin

**References:** [1350, 845, 2390, 1110, 975]

[EC 3.4.21.81 created 1993]

#### EC 3.4.21.82

**Accepted name:** glutamyl endopeptidase II

**Reaction:** Preferential cleavage: -Glu-| >> -Asp-|. Preference for Pro or Leu at P2 and Phe at P3. Cleavage of -Glu-|Asp- and -Glu-|Pro- bonds is slow

**Other name(s):** GluSGP



**Comments:** From *Streptomyces griseus*. A peptidase of family S1 (trypsin family). Inhibited by [Leu<sup>18</sup>→Glu]-modified turkey ovomucoid third domain

**References:** [3300, 1506, 2025, 2819, 284]

[EC 3.4.21.82 created 1993]

#### EC 3.4.21.83

**Accepted name:** oligopeptidase B

**Reaction:** Hydrolysis of -Arg<sup>+</sup>, -Lys<sup>+</sup> bonds in oligopeptides, even when P1' residue is proline

**Other name(s):** protease II; *Escherichia coli* alkaline proteinase II; protease II

**Comments:** Known from *Escherichia coli*. Inhibited by Tos-Lys-CH<sub>2</sub>Cl. In peptidase family S9 (prolyl oligopeptidase family)

**References:** [1378]

[EC 3.4.21.83 created 1993]

#### EC 3.4.21.84

**Accepted name:** limulus clotting factor  $\bar{C}$

**Reaction:** Selective cleavage of -Arg<sup>103</sup><sup>+</sup>Ser- and -Ile<sup>124</sup><sup>+</sup>Ile- bonds in limulus clotting factor B to form factor  $\bar{B}$ . Cleavage of -Pro-Arg<sup>+</sup> bonds in synthetic substrates

**Other name(s):** factor C; limulus factor C

**Comments:** From the hemocyte granules of the horseshoe crabs *Limulus* and *Tachypleus*. Factor C is activated by Gram-negative bacterial lipopolysaccharides and chymotrypsin. Inhibited by antithrombin III. In peptidase family S1 (trypsin family)

**References:** [2048, 2011, 2931]

[EC 3.4.21.84 created 1993]

#### EC 3.4.21.85

**Accepted name:** limulus clotting factor  $\bar{B}$

**Reaction:** Selective cleavage of -Arg<sup>98</sup><sup>+</sup>Ile- bond in limulus proclotting enzyme to form active clotting enzyme

**Comments:** From the hemocyte granules of the horseshoe crabs *Limulus* and *Tachypleus*. Factor B is activated by limulus clotting factor  $\bar{C}$ . In peptidase family S1 (trypsin family)

**References:** [2046]

[EC 3.4.21.85 created 1993]

#### EC 3.4.21.86

**Accepted name:** limulus clotting enzyme

**Reaction:** Selective cleavage of -Arg<sup>18</sup><sup>+</sup> and -Arg<sup>47</sup><sup>+</sup> bonds in coagulogen to form coagulin and fragments

**Other name(s):** clotting enzyme

**Comments:** From the hemocyte granules of horseshoe crabs *Limulus* and *Tachypleus*. Proclotting enzyme is activated by limulus clotting factor  $\bar{C}$ . In peptidase family S1 (trypsin family)

**References:** [2010, 2931]

[EC 3.4.21.86 created 1993]

[3.4.21.87 Transferred entry. *omptin*. Now EC 3.4.23.49, *omptin*. The enzyme is not a serine protease, as thought previously, but an aspartate protease]

[EC 3.4.21.87 created 1993, deleted 2006]

#### EC 3.4.21.88



**Accepted name:** repressor LexA  
**Reaction:** Hydrolysis of Ala<sup>84</sup>—Gly bond in repressor LexA  
**Other name(s):** LexA repressor  
**Comments:** RecA protein and single-stranded DNA are required for activity, which is attributed to a Ser/Lys dyad [2690]. The LexA protein represses the SOS regulon, which regulates the genes involved in DNA repair. In the presence of single-stranded DNA, the RecA protein interacts with repressor LexA, causing it to undergo an autocatalytic cleavage which disrupts the DNA-binding part of the repressor, and inactivates it. The consequent derepression of the SOS regulon leads to DNA repair. This peptidase activity of LexA was previously attributed to the RecA protein. Type example of peptidase family S24  
**References:** [1178, 2690, 1442, 1714]

[EC 3.4.21.88 created 1995]

#### EC 3.4.21.89

**Accepted name:** signal peptidase I  
**Reaction:** Cleavage of hydrophobic, N-terminal signal or leader sequences  
**Other name(s):** leader peptidase I; signal proteinase; *Escherichia coli* leader peptidase; eukaryotic signal peptidase; eukaryotic signal proteinase; leader peptidase; leader peptide hydrolase; leader proteinase; signal peptidase; pilin leader peptidase; SPC; prokaryotic signal peptidase; prokaryotic leader peptidase; HOSP; prokaryotic signal proteinase; propeptidase; PuIO prepilin peptidase; signal peptide hydrolase; signal peptide peptidase; signalase; bacterial leader peptidase 1; pilin leader peptidase  
**Comments:** The enzyme is found in bacterial membranes and in chloroplast thylakoid membranes. Unaffected by inhibitors of most serine peptidases, but site-directed mutagenesis implicates a Ser/Lys catalytic dyad in activity [227, 2973]. Hydrolyses a single bond -Ala—Ala- in M13 phage procoat protein, producing free signal peptide and coat protein. Formerly included in EC 3.4.99.36. Eukaryote signal peptidases that may have somewhat different specificity are known from the endoplasmic reticulum membrane [1728] and mitochondrial inner membrane [2120]. Type example of peptidase family S26  
**References:** [227, 2120, 2973, 1728, 2972, 388, 1253]

[EC 3.4.21.89 created 1984 as EC 3.4.99.36, transferred 1995 to EC 3.4.21.89]

#### EC 3.4.21.90

**Accepted name:** togavirin  
**Reaction:** Autocatalytic release of the core protein from the N-terminus of the togavirus structural polyprotein by hydrolysis of a -Trp—Ser- bond  
**Other name(s):** Sindbis virus protease; Sindbis virus core protein; Nsp2 proteinase  
**Comments:** Known from the Sindbis and Semliki forest togaviruses. Once released, the core protein does not retain catalytic activity. Togavirin is the type example of peptidase family S3 and has a similar tertiary structure to chymotrypsin [2943]  
**References:** [1531, 2779, 2943]

[EC 3.4.21.90 created 1995]

#### EC 3.4.21.91

**Accepted name:** flavivirin  
**Reaction:** Selective hydrolysis of -Xaa-Xaa—Yaa- bonds in which each of the Xaa can be either Arg or Lys and Yaa can be either Ser or Ala  
**Other name(s):** Yellow fever virus (flavivirus) protease; NS2B-3 proteinase  
**Comments:** Known from classical flaviviruses (yellow fever, dengue fever). The functional viral peptidase is part of the NS2B protein. Catalytic His, Asp and Ser residues are arranged as in chymotrypsin, but flavivirin is the type example of peptidase family S7.  
**References:** [389, 353, 1690]

[EC 3.4.21.91 created 1995]

#### EC 3.4.21.92

**Accepted name:** endopeptidase Clp

**Reaction:** Hydrolysis of proteins to small peptides in the presence of ATP and  $Mg^{2+}$ .  $\alpha$ -Casein is the usual test substrate. In the absence of ATP, only oligopeptides shorter than five residues are hydrolysed (such as succinyl-Leu-Tyr-NHMec; and Leu-Tyr-Leu-Tyr-Trp, in which cleavage of the -Tyr-Leu- and -Tyr-Trp bonds also occurs)

**Other name(s):** endopeptidase Ti; caseinolytic protease; protease Ti; ATP-dependent Clp protease; endopeptidase Ti; caseinolytic protease; ClpP; Clp protease

**Comments:** An enzyme from bacteria that contains subunits of two types, ClpP, with peptidase activity, and ClpA, with ATPase activity. The ClpAP complex, which displays ATP-dependent endopeptidase activity, has the composition  $(ClpP_{14}ClpA_6)_2$  [1425]. ClpP is the type example of peptidase family S14

**References:** [960, 1850, 1851, 1425]

[EC 3.4.21.92 created 1996]

#### EC 3.4.21.93

**Accepted name:** proprotein convertase 1

**Reaction:** Release of protein hormones, neuropeptides and renin from their precursors, generally by hydrolysis of -Lys-Arg- bonds

**Other name(s):** prohormone convertase 3; neuroendocrine convertase 1; PC1

**Comments:** A  $Ca^{2+}$ -dependent enzyme, maximally active at about pH 5.5. Substrates include *pro*-opiomelanocortin, prorenin, proenkephalin, prodynorphin, prosomatostatin and proinsulin. Unlike prohormone convertase 2, does not hydrolyse proluteinizing-hormone-releasing-hormone. Unusually, processing of prodynorphin occurs at a bond in which P2 is Thr. Present in the regulated secretory pathway of neuroendocrine cells, commonly acting co-operatively with prohormone convertase 2. In peptidase family S8 (subtilisin family)

**References:** [2607, 2693, 2761, 2605, 1312]

[EC 3.4.21.93 created 1996]

#### EC 3.4.21.94

**Accepted name:** proprotein convertase 2

**Reaction:** Release of protein hormones and neuropeptides from their precursors, generally by hydrolysis of -Lys-Arg- bonds

**Other name(s):** neuroendocrine convertase 2; PC2

**Comments:** A  $Ca^{2+}$ -dependent enzyme, maximally active at about pH 5.5. Specificity is broader than that of prohormone convertase 1. Substrates include *pro*-opiomelanocortin, proenkephalin, prodynorphin, proglucagon, proinsulin and proluteinizing-hormone-releasing-hormone. Does not hydrolyse prorenin or prosomatostatin, however. Unusually, processing of prodynorphin occurs at a bond in which P2 is Thr. Present in the regulated secretory pathway of neuroendocrine cells, commonly acting co-operatively with prohormone convertase 1. In peptidase family S8 (subtilisin family)

**References:** [2607, 2694, 2466, 2605]

[EC 3.4.21.94 created 1996]

#### EC 3.4.21.95

**Accepted name:** snake venom factor V activator

**Reaction:** Fully activates human clotting factor V by a single cleavage at the Trp-Tyr-Leu-Arg<sup>1545</sup>-Ser-Asn-Asn-Gly bond. Cattle, but not rabbit, factor V is cleaved, and no other proteins of the clotting system are attacked. Esterase activity is observed on Bz-Arg-OEt and Tos-Arg-OMe, and amidase activity on Phe-pipecolyl-Arg-NHPhNO<sub>2</sub>

**Comments:** Known from venom of *Vipera russelli*. Inhibited by di-isopropyl fluorophosphate, unlike the metallopeptidase russellysin (EC 3.4.24.58) that is specific for factor X [1468]. In peptidase family S1 (trypsin family) [2930].

**References:** [1468, 2930]

[EC 3.4.21.95 created 1997]

#### EC 3.4.21.96

**Accepted name:** lactocepin

**Reaction:** Endopeptidase activity with very broad specificity, although some subsite preferences have been noted, e.g. large hydrophobic residues in the P1 and P4 positions, and Pro in the P2 position [1,2]. Best known for its action on caseins, although it has been shown to hydrolyse hemoglobin and oxidized insulin B chain

**Other name(s):** CEP; extracellular lactococcal proteinase; lactococcal cell wall-associated proteinase; lactococcal cell envelope-associated proteinase; lactococcal proteinase; PrtP

**Comments:** Associated with the cell envelope of *Lactococcus lactis* and attached via a C-terminal membrane anchor sequence. Responsible for the hydrolysis of casein in milk and the provision of peptides essential to cell growth. Important in cheese making and the production of lactic casein, being required for rapid growth to high cell densities with concomitant production of adequate levels of lactic acid. Specificity differences between lactocepins from different starter strains may be partly responsible for imparting different flavour qualities to cheese [2326]. In peptidase family S8 (subtilisin family)

**References:** [3069, 1952, 727, 2326]

[EC 3.4.21.96 created 1997]

#### EC 3.4.21.97

**Accepted name:** assemblin

**Reaction:** Cleaves -Ala-Ser- and -Ala-Ala- bonds in the scaffold protein

**Comments:** Involved in the breakdown of the scaffold protein during the late stages of assembly of the herpesvirus virion. Inhibited by diisopropyl fluorophosphate. Type example of peptidase family S21. Catalytic residues are His, Ser, His, a combination not known for any other peptidase, and the protein fold also is unique. Known from herpes viruses of several types, cytomegalovirus, Epstein-Barr virus and human herpesvirus 3

**References:** [415, 537]

[EC 3.4.21.97 created 2000]

#### EC 3.4.21.98

**Accepted name:** hepacivirin

**Reaction:** Hydrolysis of four peptide bonds in the viral precursor polyprotein, commonly with Asp or Glu in the P6 position, Cys or Thr in P1 and Ser or Ala in P1'

**Other name(s):** Cpro-2; hepatitis C virus NS3 serine proteinase; NS3-4A serine proteinase complex

**Comments:** Encoded by the genome of the viruses of the hepatitis C group, and contributes to the maturation of the precursor polyproteins. The enzyme is greatly activated by binding of the 54-residue NS4A 'co-factor' protein also derived from the viral polyprotein. Type example of peptidase family S29. The crystallographic structure shows a chymotrypsin-like fold.

**References:** [1450, 2418]

[EC 3.4.21.98 created 2000]

#### EC 3.4.21.99

**Accepted name:** spermosin

**Reaction:** Hydrolyses arginyl bonds, preferably with Pro in the P2 position

**Comments:** The enzyme from the ascidian (Prochordate) *Halocynthia roretzi* is localized in the sperm head, and released during sperm activation. A proline-rich region is involved in binding to the vitelline coat of the egg. Belongs in peptidase family S1 (trypsin family).

**References:** [2541, 2542, 2539, 2540]

[EC 3.4.21.99 created 2001]

#### EC 3.4.21.100

**Accepted name:** sedolisin

**Reaction:** Hydrolysis of the B chain of insulin at -Glu<sup>13</sup>-Ala-, -Leu<sup>15</sup>-Tyr- and -Phe<sup>25</sup>-Tyr-, and angiotensin I at -Tyr<sup>4</sup>-Ile-. A good synthetic substrate is Lys-Pro-Ile-Glu-Phe-Phe(NO<sub>2</sub>)-Arg-Leu.

**Other name(s):** *Pseudomonas* sp. pepstatin-insensitive carboxyl proteinase; pseudomonapepsin; pseudomonalisin; sedolysin

**Comments:** An enzyme secreted by *Pseudomonas* sp. No. 101. Optimum pH is 4. It is distinguished from xanthomonapepsin by its insensitivity to EPNP and from scytalidopepsin B by this property and by its unrelated amino-acid sequence. Inhibited by tyrostatin, a peptide aldehyde [2133]. Type example of peptidase family S53.

**References:** [2135, 2133, 3203, 3204]

[EC 3.4.21.100 created 1995 as EC 3.4.23.37, transferred 2001 to EC 3.4.21.100, modified 2003]

#### EC 3.4.21.101

**Accepted name:** xanthomonalisin

**Reaction:** Cleavage of casein

**Other name(s):** *Xanthomonas* aspartic proteinase; xanthomonapepsin; sedolisin-B

**Comments:** Secreted by the bacterium *Xanthomonas* sp. Belongs in peptidase family S53.

**References:** [2134, 3204]

[EC 3.4.21.101 created 1995 as EC 3.4.23.33, transferred 2001 to EC 3.4.21.101, modified 2003]

#### EC 3.4.21.102

**Accepted name:** C-terminal processing peptidase

**Reaction:** The enzyme shows specific recognition of a C-terminal tripeptide, Xaa-Yaa-Zaa, in which Xaa is preferably Ala or Leu, Yaa is preferably Ala or Tyr, and Zaa is preferably Ala, but then cleaves at a variable distance from the C-terminus. A typical cleavage is -Ala-Ala-Arg-Ala-Ala-Lys-Glu-Asn-Tyr-Ala-Leu-Ala-Ala. In the plant chloroplast, the enzyme removes the C-terminal extension of the D1 polypeptide of photosystem II

**Other name(s):** CtpA gene product (*Synechocystis* sp.); photosystem II D1 protein processing peptidase; protease Re; tail-specific protease; Tsp protease

**Comments:** Proteolytic processing of the D1 protein of photosystem II is necessary to allow the light-driven assembly of the tetranuclear manganese cluster, which is responsible for photosynthetic water oxidation. The recognition of the substrate is mediated by a PDZ domain, a small protein module that promotes protein-protein interactions by binding to internal or C-terminal sequences of their partner proteins. Type example of peptidase family S41.

**References:** [1412, 184, 1679]

[EC 3.4.21.102 created 2001]

#### EC 3.4.21.103

**Accepted name:** physarolisin

**Reaction:** Milk clotting activity. Preferential cleavage of Gly<sup>8</sup>-Ser in B chain of insulin most rapidly, followed by Leu<sup>11</sup>-Val, Cys(SO<sub>3</sub>H)<sup>19</sup>-Gly and Phe<sup>24</sup>-Phe. No action on Ac-Phe-Tyr(I)<sub>2</sub>.

**Other name(s):** *Dictyostelium discoideum* aspartic proteinase; *Dictyostelium discoideum* aspartic proteinase E; *Physarum flavicomum* aspartic proteinase; *Physarum polycephalum* acid proteinase; *Physarum* aspartic proteinase; physaropepsin

**Comments:** Belongs in peptidase family S53. From the slime mold *Physarum polycephalum*. Is not inhibited by pepstatin, but is blocked by methyl 2-diazoacetamidohexanoate. Closely similar enzymes are found in *Dictyostelium discoideum* and *P. flavicomum*. Formerly included in EC 3.4.23.6.

**References:** [1113, 2003, 2115, 3204, 2093]

[EC 3.4.21.103 created 1992 as EC 3.4.23.27 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992), transferred 2003 to EC 3.4.21.103]

#### EC 3.4.21.104

**Accepted name:** mannan-binding lectin-associated serine protease-2

**Reaction:** Selective cleavage after Arg<sup>223</sup> in complement component C2 (-Ser-Leu-Gly-Arg-|Lys-Ile-Gln-Ile) and after Arg<sup>76</sup> in complement component C4 (-Gly-Leu-Gln-Arg-|Ala-Leu-Glu-Ile)

**Other name(s):** MASP-2; MASP2; MBP-associated serine protease-2; mannan-binding lectin-associated serine protease-2; p100; mannan-binding lectin-associated serine peptidase 2

**Comments:** Mannan-binding lectin (MBL) recognizes patterns of neutral carbohydrates, such as mannose and *N*-acetylglucosamine, on a wide range of microbial surfaces and is able to initiate activation of the lectin pathway of complement [2897]. This enzyme displays C<sup>1s</sup>-like esterolytic activity (*cf.* EC 3.4.21.42, complement subcomponent C<sup>1s</sup>). It also cleaves C4 and C2 with efficiencies that are relatively higher than those of EC 3.4.21.42 [2464]. Belongs in peptidase family S1A.

**References:** [1839, 2909, 2464, 45, 1055, 410, 2897]

[EC 3.4.21.104 created 2005]

#### EC 3.4.21.105

**Accepted name:** rhomboid protease

**Reaction:** Cleaves type-1 transmembrane domains using a catalytic dyad composed of serine and histidine that are contributed by different transmembrane domains

**Comments:** These endopeptidases are multi-spanning membrane proteins. Their catalytic site is embedded within the membrane and they cleave type-1 transmembrane domains. A catalytic dyad is involved in proteolysis rather than a catalytic triad, as was thought previously [1643]. They are important for embryo development in *Drosophila melanogaster*. Rhomboid is a key regulator of EGF receptor signalling and is responsible for cleaving Spitz, the main ligand of the *Drosophila* EGF receptor pathway. Belongs in peptidase family S54. Parasite-encoded rhomboid enzymes are also important for invasion of host cells by *Toxoplasma* and the malaria parasite. Rhomboids are widely conserved from bacteria to archaea to humans [1509, 3020].

**References:** [3023, 306, 1128, 1311, 2658, 3019, 1129, 1875, 1509, 3018, 3022, 3021, 3020, 1643, 3125]

[EC 3.4.21.105 created 2005]

#### EC 3.4.21.106

**Accepted name:** hepsin

**Reaction:** Cleavage after basic amino-acid residues, with Arg strongly preferred to Lys

**Comments:** This type-II membrane-associated serine peptidase has been implicated in cell growth and development [3340, 2949]. The enzyme has been shown to activate blood coagulation factor VII by cleavage of the Arg<sup>152</sup>-|Ile<sup>153</sup> peptide bound in BHK cells, thus indicating a possible role in the initiation of blood coagulation [1408]. There is no cleavage after aromatic or aliphatic residues [3340]. The occupancy of the S2 site is an absolute requirement for catalysis and a basic residue at that site is preferred to an aliphatic residue. The nature of the residue at S3 also affects hydrolysis, with Gln being much more favourable than Ala [3340]. Belongs in peptidase family S1A.

**References:** [3340, 1408, 2949]

[EC 3.4.21.106 created 2006]

**EC 3.4.21.107**

**Accepted name:** peptidase Do

**Reaction:** Acts on substrates that are at least partially unfolded. The cleavage site P1 residue is normally between a pair of hydrophobic residues, such as Val—|—Val

**Other name(s):** DegP; DegP protease; HtrA; high temperature requirement protease A; HtrA heat shock protein; protease Do; Do protease

**Comments:** This serine endopeptidase is essential for the clearance of denatured or aggregated proteins from the inner-membrane and periplasmic space in *Escherichia coli*. Natural substrates of the enzyme include colicin A lysis protein, pilin subunits and MalS from *E. coli* [1335]. The enzyme has weak peptidase activity with casein and other non-native substrates [1335]. The peptidase acts as a chaperone at low temperatures but switches to a peptidase (heat shock protein) at higher temperatures [1710, 1537]. Molecular chaperones and peptidases control the folded state of proteins by recognizing hydrophobic stretches of polypeptide that become exposed by misfolding or unfolding. They then bind these hydrophobic substrates to prevent aggregation or assist in protein refolding. If attempts at refolding fail, then irreversibly damaged proteins are degraded by peptidases such as this enzyme [1537]. Belongs in peptidase family S1C.

**References:** [1710, 2611, 1335, 2820, 2226, 1537]

[EC 3.4.21.107 created 2006]

**EC 3.4.21.108**

**Accepted name:** HtrA2 peptidase

**Reaction:** Cleavage of non-polar aliphatic amino-acids at the P1 position, with a preference for Val, Ile and Met. At the P2 and P3 positions, Arg is selected most strongly with a secondary preference for other hydrophilic residues

**Other name(s):** high temperature requirement protein A2; HtrA2; Omi stress-regulated endoprotease; serine proteinase OMI; HtrA2 protease; OMI/HtrA2 protease; HtrA2/Omi; Omi/HtrA2

**Comments:** This enzyme is upregulated in mammalian cells in response to stress induced by both heat shock and tunicamycin treatment [967]. It can induce apoptosis in a caspase-independent manner through its peptidase activity and in a caspase-dependent manner by disrupting the interaction between caspase and the inhibitor of apoptosis (IAP) [1824]. Belongs in peptidase family S1C.

**References:** [2746, 2538, 1824, 967, 1671]

[EC 3.4.21.108 created 2006]

**EC 3.4.21.109**

**Accepted name:** matriptase

**Reaction:** Cleaves various synthetic substrates with Arg or Lys at the P1 position and prefers small side-chain amino acids, such as Ala and Gly, at the P2 position

**Other name(s):** serine protease 14; membrane-type serine protease 1; MT-SP1; prostamin; serine protease TADG-15; tumor-associated differentially-expressed gene 15 protein; ST14; breast cancer 80 kDa protease; epithin; serine endopeptidase SNC19

**Comments:** This trypsin-like integral-membrane serine peptidase has been implicated in breast cancer invasion and metastasis [1631, 1691]. The enzyme can activate hepatocyte growth factor/scattering factor (HGF/SF) by cleavage of the two-chain form at an Arg residue to give active  $\alpha$ - and  $\beta$ -HGF, but It does not activate plasminogen, which shares high homology with HGF [1631]. The enzyme can also activate urokinase plasminogen activator (uPA), which initiates the matrix-degrading peptidase cascade [1631, 1691]. Belongs in peptidase family S1A.

**References:** [1631, 1691]

[EC 3.4.21.109 created 2006]

#### EC 3.4.21.110

**Accepted name:** C5a peptidase

**Reaction:** The primary cleavage site is at His<sup>67</sup>—Lys<sup>68</sup> in human C5a with a minor secondary cleavage site at Ala<sup>58</sup>—Ser<sup>59</sup>

**Other name(s):** streptococcal C5a peptidase; ScpA; ScpB; SCPA

**Comments:** This enzyme is a surface-associated subtilisin-like serine peptidase with very specific substrate specificity. Virulent strains of streptococci, including *Streptococcus pyogenes*, can evade human detection and phagocytosis by destroying the complement chemotaxin C5a. Cleavage of human C5a by this enzyme reduces the ability of C5a to bind receptors on the surface of polymorphonuclear neutrophil leukocytes (PMNLs) and thereby abolishes its chemotactic properties [3167, 52]. Belongs in peptidase family S8A.

**References:** [3167, 254, 461, 52, 2753, 2901]

[EC 3.4.21.110 created 2006]

#### EC 3.4.21.111

**Accepted name:** aqualysin 1

**Reaction:** Exhibits low specificity towards esters of amino acids with small hydrophobic or aromatic residues at the P1 position

**Other name(s):** caldolyysin

**Comments:** This enzyme from the extreme thermophile, *Thermus aquaticus*, is an alkaline serine peptidase. It has three subsites, S1, S2, and S3, in the substrate binding site. The preferred amino acids at the S1 site are Ala and Phe, at the S2 site are Ala and norleucine and at the S3 site are Phe and Ile [2877]. These specificities are similar to those of EC 3.4.21.64 (peptidase K) and EC 3.4.21.62 (subtilisin BPN') [2877]. The enzyme displays broad specificity for cleavage of insulin B-chain and hydrolyses elastin substrates such as succinyl-(Ala)<sub>n</sub>-p-nitroanilide (*n* = 1,2,3) and some peptide esters [1843, 2877]. Belongs in peptidase family S8A.

**References:** [1843, 2876, 2877]

[EC 3.4.21.111 created 2006]

#### EC 3.4.21.112

**Accepted name:** site-1 protease

**Reaction:** Processes precursors containing basic and hydrophobic/aliphatic residues at P4 and P2, respectively, with a relatively relaxed acceptance of amino acids at P1 and P3

**Other name(s):** mammalian subtilisin/kexin isozyme 1; membrane-bound transcription factor site-1 protease; proprotein convertase SKI-1; proprotein convertase SKI-1/S1PPS1; S1P endopeptidase; S1P protease; site-1 peptidase; site-1 protease; SKI-1; SREBP proteinase; SREBP S1 protease; SREBP-1 proteinase; SREBP-2 proteinase; sterol regulatory element-binding protein proteinase; sterol regulatory element-binding protein site 1 protease; sterol-regulated luminal protease; subtilase SKI-1; subtilase SKI-1/S1P; subtilisin/kexin-isozyme 1

**Comments:** Cleaves sterol regulatory element-binding proteins (SREBPs) and thereby initiates a process by which the active fragments of the SREBPs translocate to the nucleus and activate genes controlling the synthesis and uptake of cholesterol and unsaturated fatty acids into the bloodstream [718]. The enzyme also processes pro-brain-derived neurotrophic factor and undergoes autocatalytic activation in the endoplasmic reticulum through sequential cleavages [1647]. The enzyme can also process the unfolded protein response stress factor ATF6 at an Arg-His-Lys-Lys— site [3281, 2606], and the envelope glycoprotein of the highly infectious Lassa virus [1647, 2606] and Crimean Congo hemorrhagic fever virus at Arg-Arg-Lys-Lys— [3067, 2606]. Belongs in peptidase family S8A.

**References:** [718, 422, 2951, 3281, 1647, 167, 3067, 2606]

[EC 3.4.21.112 created 2006]

#### EC 3.4.21.113



**Accepted name:** pestivirus NS3 polyprotein peptidase  
**Reaction:** Leu is conserved at position P1 for all four cleavage sites. Alanine is found at position P1' of the NS4A-NS4B cleavage site, whereas serine is found at position P1' of the NS3-NS4A, NS4B-NS5A and NS5A-NS5B cleavage sites  
**Other name(s):** border disease virus NS3 endopeptidase; BDV NS3 endopeptidase; bovine viral diarrhea virus NS3 endopeptidase; BVDV NS3 endopeptidase; classical swine fever virus NS3 endopeptidase; CSFV NS3 endopeptidase; p80  
**Comments:** The polyprotein of noncytopathogenic pestiviruses is cleaved co- and post-translationally into at least 11 proteins (N<sup>pro</sup>, C, E<sup>ms</sup>, E1, E2, p7, NS2-3, NS4A, NS4B, NS5A, and NS5B) [2893]. The genomes of cytopathogenic pestivirus strains express at least one additional protein, called NS3 (p80) [2893]. This enzyme, which resides in the N-terminal region of NS3 (nonstructural protein 3), is essential for generation of its own C-terminus and for processing of the downstream cleavage sites, leading to the release of the pestivirus nonstructural proteins NS4A, NS4B, NS5A and NS5B [3200, 2893]. Belongs in peptidase family S31.  
**References:** [3200, 2893, 3232, 2894]

[EC 3.4.21.113 created 2006]

#### EC 3.4.21.114

**Accepted name:** equine arterivirus serine peptidase  
**Reaction:** Cleavage of (Glu/Gln)-|(Gly/Ser/Ala) in arterivirus replicase translation products ORF1a and ORF1ab  
**Comments:** In the equine arterivirus (EAV), the replicase gene is translated into open reading frame 1a (ORF1a) and ORF1ab polyproteins. This enzyme is the main viral proteinase and processes five cleavage sites in the ORF1a protein and three in the ORF1b-encoded part of the ORF1ab protein to yield nonstructural proteins (nsp5-nsp12) [162]. It combines the catalytic system of a chymotrypsin-like serine peptidase (His-Asp-Ser catalytic triad) with the substrate specificity of a 3C-like serine peptidase (Glu or Gln) at the P1 position and a small amino-acid residue (Gly, Ser or Ala) at the P1' position [2704]. Cleavage of ORF1ab by this enzyme is essential for viral replication [3046]. Belongs in peptidase family S32.  
**References:** [2704, 3046, 162]

[EC 3.4.21.114 created 2006]

#### EC 3.4.21.115

**Accepted name:** infectious pancreatic necrosis birnavirus Vp4 peptidase  
**Reaction:** Cleaves the (Ser/Thr)-Xaa-Ala-|(Ser/Ala)-Gly motif in the polyprotein NH<sub>2</sub>-pVP2-VP4-VP3-COOH of infectious pancreatic necrosis virus at the pVP2-VP4 and VP4-VP3 junctions  
**Other name(s):** infectious pancreatic necrosis virus protease; IPNV Vp4 protease; IPNV Vp4 peptidase; NS protease; NS-associated protease; Vp4 protease  
**Comments:** Infectious pancreatic necrosis virus (IPNV) is a birnavirus that causes an acute, contagious disease in young salmonid fish [2266]. As with most viruses that infect eukaryotic cells, the proteolytic processing of viral precursor proteins is a crucial step in the life cycle of this virus [2266]. pVP2 is converted into VP2 by cleavage near the carboxy end of pVP2. This cleavage is most likely due to host-cell proteases rather than VP4 [2266, 622]. Differs from most serine peptidases in not having the catalytic triad Ser-His-Asp [2266]. Belongs in peptidase family S50.  
**References:** [1802, 2266, 622]

[EC 3.4.21.115 created 2006]

#### EC 3.4.21.116

**Accepted name:** SpoIVB peptidase  
**Reaction:** Self-cleaves Val<sup>52</sup>-|Asn<sup>53</sup>, Ala<sup>62</sup>-|Phe<sup>63</sup> and Val<sup>74</sup>-|Thr<sup>75</sup> at the N-terminus of SpoIVB  
**Other name(s):** sporulation factor IV B protease



**Comments:** This enzyme plays a central role in a regulatory checkpoint (the  $\sigma^K$  checkpoint), which coordinates gene expression during the later stages of spore formation in *Bacillus subtilis* [3092, 1156]. The enzyme activates proteolytic processing of a sporulation-specific sigma factor, pro- $\sigma^K$ , to its mature and active form,  $\sigma^K$ , by self-cleavage [3092, 1156]. The enzyme is also subject to secondary proteolysis, which presumably inactivates SpoIVB [1156]. The enzyme is also essential for the formation of heat-resistant spores. Belongs in peptidase family S55.

**References:** [3092, 1155, 1156, 635]

[EC 3.4.21.116 created 2006]

#### EC 3.4.21.117

**Accepted name:** stratum corneum chymotryptic enzyme

**Reaction:** Cleavage of proteins with aromatic side chains in the P1 position

**Other name(s):** kallikrein 7; SCCE; KLK7; PRSS6; hK7

**Comments:** This enzyme has wide substrate specificity, being able to degrade heat-denatured bovine casein and the  $\alpha$ -chain of native human fibrinogen. It cleaves the B chain of bovine insulin at Leu<sup>6</sup>—Cya<sup>7</sup>, Tyr<sup>16</sup>—Leu<sup>17</sup>, Phe<sup>25</sup>—Tyr<sup>26</sup> and Tyr<sup>26</sup>—Thr<sup>27</sup> [2687]. It is thought to play a role in the desquamation (skin-shedding) of the outer layer of skin, the stratum corneum, by degrading intercellular cohesive structures [2687, 680]. Belongs in peptidase family S1A.

**References:** [2687, 680, 1046, 3307, 3055]

[EC 3.4.21.117 created 2006]

#### EC 3.4.21.118

**Accepted name:** kallikrein 8

**Reaction:** Cleavage of amide substrates following the basic amino acids Arg or Lys at the P1 position, with a preference for Arg over Lys

**Other name(s):** KLK8; PRSS19; human kallikrein 8; hK8; mK8; ovasin; tumor-associated differentially expressed gene 14; TADG-14; NP; neuropsin

**Comments:** The enzyme is activated by removal of an N-terminal prepropeptide [2633, 1466]. The highest amidolytic activity is observed using Boc-Val-Pro-Arg—7-amido-4-methylcoumarin, which is a substrate of  $\alpha$ -thrombin [2633, 1466]. Substrates lacking basic amino acids in the P1 position are not cleaved [1466]. The enzyme degrades casein, fibronectin, gelatin, collagen type IV, fibrinogen, and high-molecular-mass kininogen [2362] and is associated with diseases such as ovarian cancer and Alzheimer's disease [1466]. Belongs in peptidase family S1A.

**References:** [421, 2633, 2362, 1466]

[EC 3.4.21.118 created 2006]

#### EC 3.4.21.119

**Accepted name:** kallikrein 13

**Reaction:** Hydrolyses mouse Ren2 protein (a species of prorenin present in the submandibular gland) on the carboxy side of the arginine residue at the Lys-Arg— pair in the N-terminus, to yield mature renin

**Other name(s):** KLK13; kallikrein mK13; mGK-13; mK13; mKLK13; prorenin converting enzyme 1; PRECE-1; prorenin-converting enzyme; PRECE; proteinase P

**Comments:** The enzyme is specific for prorenin from the mouse submandibular gland, as prorenin from the mouse kidney (Ren1) and human prorenin are not substrates [2052]. Site-directed mutagenesis studies have shown that the enzyme will also cleave prorenin when Lys-Arg is replaced by Arg-Arg or Gln-Arg but the rate of reaction is much slower when Lys-Lys is used. This enzyme is also able to process pro-interleukin-1 $\beta$  (pro-IL-1 $\beta$ ) in mouse submandibular gland to form IL-1 $\beta$  [3261]. Belongs in peptidase family S1A.

**References:** [2052, 1455, 1439, 3261]

[EC 3.4.21.119 created 2006]

#### EC 3.4.21.120

- Accepted name:** oviductin  
**Reaction:** Preferential cleavage at Gly-Ser-Arg<sup>373</sup>— of glycoprotein gp43 in *Xenopus laevis* coelomic egg envelope to yield gp41  
**Other name(s):** oviductal protease  
**Comments:** The egg envelope of the South African clawed frog (*Xenopus laevis*) is modified during transit of the egg through the pars rectus oviduct, changing the egg envelope from an unfertilizable form to a fertilizable form. This process involves the conversion of glycoprotein gp43 to gp41 (ZPC) by the pars recta protease oviductin. It is thought that the enzymically active protease molecule comprises the N-terminal protease domain coupled to two C-terminal CUB domains, which are related to the mammalian spermadhesin molecules implicated in mediating sperm-envelope interactions [1703]. The enzyme is also found in the Japanese toad (*Bufo japonicus*) [1153]. Belongs in peptidase family S1.  
**References:** [1053, 1703, 1153]

[EC 3.4.21.120 created 2007]

#### EC 3.4.21.121

- Accepted name:** Lys-Lys/Arg-Xaa endopeptidase  
**Reaction:** Cleavage of -Lys-Lys— and -Lys-Arg— bonds.  
**Other name(s):** ASP (*Aeromonas sobria*)-type peptidase; *Aeromonas* extracellular serine protease  
**Comments:** The enzyme is a serine peptidase, which has been shown to cleave prothrombin and prekallikrein. It hydrolyses the complement component C5 releasing complement component C5a.  
**References:** [1489, 2100, 1488, 1241, 2099]

[EC 3.4.21.121 created 2013]

### EC 3.4.22 Cysteine endopeptidases

#### EC 3.4.22.1

- Accepted name:** cathepsin B  
**Reaction:** Hydrolysis of proteins with broad specificity for peptide bonds. Preferentially cleaves -Arg-Arg— bonds in small molecule substrates (thus differing from cathepsin L). In addition to being an endopeptidase, shows peptidyl-dipeptidase activity, liberating C-terminal dipeptides  
**Other name(s):** cathepsin B1 (obsolete); cathepsin II  
**Comments:** An intracellular (lysosomal) enzyme in peptidase family C1 (papain family)  
**References:** [258, 159, 2304, 158, 1464]

[EC 3.4.22.1 created 1972]

#### EC 3.4.22.2

- Accepted name:** papain  
**Reaction:** Hydrolysis of proteins with broad specificity for peptide bonds, but preference for an amino acid bearing a large hydrophobic side chain at the P2 position. Does not accept Val in P1'  
**Other name(s):** papayotin; summetrin; velardon; papaine; Papaya peptidase I  
**Comments:** Type example of peptidase family C1 from latex of the papaya (*Carica papaya*) fruit. Inhibited by compound E-64 and proteins of the cystatin family.  
**References:** [1375, 1884]

[EC 3.4.22.2 created 1961 as EC 3.4.4.10, transferred 1972 to EC 3.4.22.2, modified 1976, modified 2000]

### EC 3.4.22.3

**Accepted name:** ficain

**Reaction:** Similar to that of papain

**Other name(s):** ficin; debricin; higueroxyl delabarre

**Comments:** The major proteolytic component of the latex of fig, *Ficus glabrata*. Cysteine endopeptidases with similar properties are present in other members of the large genus *Ficus*. In peptidase family C1 (papain family).

**References:** [1684, 297]

[EC 3.4.22.3 created 1961 as EC 3.4.4.12, transferred 1972 to EC 3.4.22.3]

[3.4.22.4 Transferred entry. bromelain (stem). Now EC 3.4.22.32 (stem bromelain) and EC 3.4.22.33 (fruit bromelain)]

[EC 3.4.22.4 created 1972, deleted 1992 [EC 3.4.22.5 created 1972, incorporated 1978]]

[3.4.22.5 Transferred entry. bromelain (juice). Now EC 3.4.22.32 (stem bromelain) and EC 3.4.22.33 (fruit bromelain)]

[EC 3.4.22.5 created 1972, deleted 1978]

### EC 3.4.22.6

**Accepted name:** chymopapain

**Reaction:** Similar to that of papain

**Other name(s):** chymopapain A; chymopapain B; chymopapain S

**Comments:** The major endopeptidase of papaya (*Carica papaya*) latex. It has multiple chromatographic forms. In peptidase family C1 (papain family).

**References:** [297, 1293, 339]

[EC 3.4.22.6 created 1961 as EC 3.4.4.11, transferred 1972 to EC 3.4.22.6]

### EC 3.4.22.7

**Accepted name:** asclepain

**Reaction:** Similar to that of papain

**Comments:** From the latex of milkweed, *Asclepias syriaca*. It has multiple forms, and is in peptidase family C1 (papain family)

**References:** [295]

[EC 3.4.22.7 created 1972]

### EC 3.4.22.8

**Accepted name:** clostripain

**Reaction:** Preferential cleavage: Arg—, including Arg—Pro, but not Lys—

**Other name(s):** clostridiopeptidase B; clostridium histolyticum proteinase B;  $\alpha$ -clostridipain; clostridiopeptidase

**Comments:** From the bacterium *Clostridium histolyticum*. It requires  $\text{Ca}^{2+}$  ions and is inhibited by EDTA. Type example of peptidase family C11.

**References:** [1931, 922, 923]

[EC 3.4.22.8 created 1961 as EC 3.4.4.20, transferred 1972 to EC 3.4.22.8]

[3.4.22.9 Transferred entry. yeast proteinase B. Now EC 3.4.21.48, cerevisin]

[EC 3.4.22.9 created 1972, deleted 1981]

### EC 3.4.22.10

**Accepted name:** streptopain

**Reaction:** Preferential cleavage with hydrophobic residues at P2, P1 and P1'

**Other name(s):** *Streptococcus* peptidase A; streptococcal cysteine proteinase; *Streptococcus* protease  
**Comments:** From the bacterium, group A *Streptococcus*. Formed from the proenzyme by limited proteolysis. Type example of peptidase family C10.  
**References:** [693, 1726, 2830, 1730]

[EC 3.4.22.10 created 1961 as EC 3.4.4.18, transferred 1972 to EC 3.4.22.10]

[3.4.22.11 *Transferred entry. insulinase. Now EC 3.4.24.56, insulysin*]

[EC 3.4.22.11 created 1976, deleted 1978 [transferred to EC 3.4.99.45, deleted 1993]]

[3.4.22.12 *Transferred entry.  $\gamma$ -glutamyl hydrolase. Now EC 3.4.19.9,  $\gamma$ -glutamyl hydrolase*]

[EC 3.4.22.12 created 1978, deleted 1992]

[3.4.22.13 *Deleted entry. staphylococcal cysteine proteinase*]

[EC 3.4.22.13 created 1978, modified 1981, deleted 1992]

#### EC 3.4.22.14

**Accepted name:** actinidain  
**Reaction:** Similar to that of papain  
**Other name(s):** actinidin; Actinidia anionic protease; proteinase A<sub>2</sub> of *Actinidia chinensis*  
**Comments:** From the kiwi fruit or Chinese gooseberry (*Actinidia chinensis*). In peptidase family C1 (papain family)  
**References:** [125, 1375, 126]

[EC 3.4.22.14 created 1978]

#### EC 3.4.22.15

**Accepted name:** cathepsin L  
**Reaction:** Similar to that of papain. As compared to cathepsin B, cathepsin L exhibits higher activity towards protein substrates, but has little activity on Z-Arg-Arg-NHMe, and no peptidyl-dipeptidase activity  
**Other name(s):** Aldrichina grahamsi cysteine proteinase  
**Comments:** A lysosomal enzyme in peptidase family C1 (papain family) that is readily inhibited by the diazomethane inhibitor Z-Phe-Phe-CHN<sub>2</sub> or the epoxide inhibitor E-64  
**References:** [159, 158, 1341, 1464]

[EC 3.4.22.15 created 1978 (EC 3.4.99.19 created 1972, incorporated 1981)]

#### EC 3.4.22.16

**Accepted name:** cathepsin H  
**Reaction:** Hydrolysis of proteins, acting as an aminopeptidase (notably, cleaving Arg— bonds) as well as an endopeptidase  
**Other name(s):** cathepsin B3; benzoylarginine-naphthylamide (BANA) hydrolase (obsolete); cathepsin Ba, aleurain; N-benzoylarginine- $\beta$ -naphthylamide hydrolase  
**Comments:** Present in lysosomes of mammalian cells. In peptidase family C1 (papain family)  
**References:** [159, 300, 830]

[EC 3.4.22.16 created 1981, modified 1989]

[3.4.22.17 *Transferred entry. calpain. Now EC 3.4.22.53, calpain-2*]

[EC 3.4.22.17 created 1981 [EC 3.4.24.5 created 1978, part incorporated 1989], deleted 2003]

[3.4.22.18 *Transferred entry. prolyl endopeptidase (thiol-dependent). Now EC 3.4.21.26, prolyl oligopeptidase*]

[EC 3.4.22.18 created 1981, deleted 1992]

[3.4.22.19 *Transferred entry. endo-oligopeptidase. Now EC 3.4.24.15, thimet oligopeptidase*]

[EC 3.4.22.19 created 1989, deleted 1992]

[3.4.22.20 *Deleted entry. dinorphin-converting enzyme*]

[EC 3.4.22.20 created 1989, deleted 1992]

[3.4.22.21 *Transferred entry. yeast cysteine proteinase E. Now EC 3.4.25.1, proteasome endopeptidase complex*]

[EC 3.4.22.21 created 1989, deleted 1992]

[3.4.22.22 *Transferred entry. yeast cysteine proteinase D. Now EC 3.4.24.37, saccharolysin*]

[EC 3.4.22.22 created 1989, deleted 1992]

[3.4.22.23 *Transferred entry. yeast cysteine proteinase F. Now EC 3.4.21.61, kexin*]

[EC 3.4.22.23 created 1989, deleted 1992]

#### EC 3.4.22.24

**Accepted name:** cathepsin T

**Reaction:** Interconversion of the three forms of tyrosine aminotransferase, EC 2.6.1.5

**Comments:** Degrades azocasein and denatured hemoglobin; the only native protein on which it has been shown to act is tyrosine aminotransferase

**References:** [941, 940, 2288]

[EC 3.4.22.24 created 1990]

#### EC 3.4.22.25

**Accepted name:** glycy endopeptidase

**Reaction:** Preferential cleavage: Gly—, in proteins and small molecule substrates

**Other name(s):** papaya peptidase B; papaya proteinase IV; glycine-specific proteinase; chymopapain; Papaya proteinase 4; PPIV; chymopapain M

**Comments:** From the papaya plant, *Carica papaya*. Not inhibited by chicken cystatin, unlike most other homologues of papain, but in peptidase family C1 (papain family)

**References:** [2301, 340, 2430, 342, 341]

[EC 3.4.22.25 created 1992]

#### EC 3.4.22.26

**Accepted name:** cancer procoagulant

**Reaction:** Specific cleavage of Arg—Ile bond in Factor X to form Factor Xa

**Comments:** Apparently produced only by malignant and fetal cells

**References:** [733, 734]

[EC 3.4.22.26 created 1992]

#### EC 3.4.22.27

**Accepted name:** cathepsin S

**Reaction:** Similar to cathepsin L, but with much less activity on Z-Phe-Arg—NHMeC, and more activity on the Z-Val-Val-Arg— compound

**Comments:** A lysosomal cysteine endopeptidase that is unusual amongst such enzymes for its stability to neutral pH. In peptidase family C1 (papain family)

**References:** [3003, 304, 1463]

[EC 3.4.22.27 created 1992]

**EC 3.4.22.28**

**Accepted name:** picornain 3C

**Reaction:** Selective cleavage of Gln—Gly bond in the poliovirus polyprotein. In other picornavirus reactions Glu may be substituted for Gln, and Ser or Thr for Gly

**Other name(s):** picornavirus endopeptidase 3C; poliovirus protease 3C; rhinovirus protease 3C; foot-and-mouth protease 3C; poliovirus proteinase 3C; rhinovirus proteinase 3C; coxsackievirus 3C proteinase; foot-and-mouth-disease virus proteinase 3C; 3C protease; 3C proteinase; cysteine proteinase 3C; hepatitis A virus 3C proteinase; protease 3C; tomato ringspot nepovirus 3C-related protease

**Comments:** From entero-, rhino-, aphto- and cardioviruses. Larger than the homologous virus picornain 2A. Type example of peptidase family C3

**References:** [1278, 179, 1531, 2081]

[EC 3.4.22.28 created 1992]

**EC 3.4.22.29**

**Accepted name:** picornain 2A

**Reaction:** Selective cleavage of Tyr—Gly bond in picornavirus polyprotein

**Other name(s):** picornavirus endopeptidase 2A; poliovirus protease 2A; rhinovirus protease 2A; 2A protease; 2A proteinase; protease 2A; proteinase 2Apro; picornaviral 2A proteinase; Y-G proteinase 2A; poliovirus proteinase 2A; poliovirus protease 2Apro; picornaviral 2A proteinase

**Comments:** From entero-, rhino-, aphto- and cardioviruses. Smaller than the homologous picornain 3C, which is also in peptidase family C3 (picornain 3C family)

**References:** [179, 1508, 1531]

[EC 3.4.22.29 created 1992]

**EC 3.4.22.30**

**Accepted name:** caricain

**Reaction:** Hydrolysis of proteins with broad specificity for peptide bonds, similar to those of papain and chymopapain

**Other name(s):** papaya peptidase A; papaya peptidase II; papaya proteinase ; papaya proteinase III; papaya proteinase 3; proteinase  $\omega$ ; papaya proteinase A; chymopapain S; Pp

**Comments:** From papaya plant, *Carica papaya*. In peptidase family C1 (papain family)

**References:** [2552, 2441, 2302, 296, 3350, 658]

[EC 3.4.22.30 created 1992]

**EC 3.4.22.31**

**Accepted name:** ananain

**Reaction:** Hydrolysis of proteins with broad specificity for peptide bonds. Best reported small molecule substrate Bz-Phe-Val-Arg—NHMeC, but broader specificity than fruit bromelain

**Other name(s):** stem bromelain; fruit bromelain

**Comments:** From stem of pineapple plant, *Ananas comosus*. Differs from stem and fruit bromelains in being inhibited by chicken cystatin. In peptidase family C1 (papain family)

**References:** [2468, 2469]

[EC 3.4.22.31 created 1992]

**EC 3.4.22.32**

**Accepted name:** stem bromelain

**Reaction:** Broad specificity for cleavage of proteins, but strong preference for Z-Arg-Arg-NHMec amongst small molecule substrates  
**Other name(s):** bromelain; pineapple stem bromelain  
**Comments:** The most abundant of the cysteine endopeptidases of the stem of the pineapple plant, *Ananas comosus*. Distinct from the bromelain found in the pineapple fruit (EC 3.4.22.33). Scarcely inhibited by chicken cystatin and also very slowly inactivated by E-64. In peptidase family C1 (papain family).  
**References:** [297, 2468, 2431, 2469]

[EC 3.4.22.32 created 1965 as EC 3.4.4.24, transferred 1972 to EC 3.4.22.4, part transferred 1992 to EC 3.4.22.32]

#### EC 3.4.22.33

**Accepted name:** fruit bromelain  
**Reaction:** Hydrolysis of proteins with broad specificity for peptide bonds. Bz-Phe-Val-Arg-NHMec is a good synthetic substrate, but there is no action on Z-Arg-Arg-NHMec (c.f. stem bromelain)  
**Other name(s):** juice bromelain; ananase; bromelase; bromelin; extranase; juice bromelain; pinase; pineapple enzyme; traumanase; fruit bromelain FA2  
**Comments:** From the pineapple plant, *Ananas comosus*. Scarcely inhibited by chicken cystatin. Another cysteine endopeptidase, with similar action on small molecule substrates, pinguinain, is obtained from the related plant, *Bromelia pinguin*, but pinguinain differs from fruit bromelain in being inhibited by chicken cystatin [2469].  
**References:** [2535, 3241, 2211, 2469]

[EC 3.4.22.33 created 1965 as EC 3.4.4.24, transferred 1972 to EC 3.4.22.4, part transferred 1992 to EC 3.4.22.33]

#### EC 3.4.22.34

**Accepted name:** legumain  
**Reaction:** Hydrolysis of proteins and small molecule substrates at -Asn-Xaa- bonds  
**Other name(s):** asparaginyl endopeptidase; citvac; proteinase B (ambiguous); hemoglobinase (ambiguous); PRSC1 gene product (Homo sapiens); vicilin peptidohydrolase; bean endopeptidase; vicilin peptidohydrolase  
**Comments:** Best known from legume seeds, the trematode *Schistosoma mansoni* and mammalian lysosomes. Not inhibited by compound E-64. Type example of peptidase family C13  
**References:** [1049, 531, 412]

[EC 3.4.22.34 created 1992, modified 2000]

#### EC 3.4.22.35

**Accepted name:** histolysain  
**Reaction:** Hydrolysis of proteins, including basement membrane collagen and azocasein. Preferential cleavage: Arg-Arg-NHMec in small molecule substrates including Z-Arg-Arg-NHMec  
**Other name(s):** histolysin; histolysin; *Entamoeba histolytica* cysteine proteinase; amebapain; *Entamoeba histolytica* cysteine protease; *Entamoeba histolytica* neutral thiol proteinase  
**Comments:** From the protozoan, *Entamoeba histolytica*. In peptidase family C1 (papain family)  
**References:** [1757, 1748]

[EC 3.4.22.35 created 1992]

#### EC 3.4.22.36

**Accepted name:** caspase-1  
**Reaction:** Strict requirement for an Asp residue at position P1 and has a preferred cleavage sequence of Tyr-Val-Ala-Asp-NHMec  
**Other name(s):** interleukin 1 $\beta$ -converting enzyme; protease VII; protease A; interleukin 1 $\beta$  precursor proteinase; interleukin 1 converting enzyme; interleukin 1 $\beta$ -converting endopeptidase; interleukin-1 $\beta$  convertase; interleukin-1 $\beta$  converting enzyme; interleukin-1 $\beta$  precursor proteinase; prointerleukin 1 $\beta$  protease; precursor interleukin-1 $\beta$  converting enzyme; pro-interleukin 1 $\beta$  proteinase; ICE

**Comments:** From mammalian monocytes. This enzyme is part of the family of inflammatory caspases, which also includes caspase-4 (EC 3.4.22.57) and caspase-5 (EC 3.4.22.58) in humans and caspase-11 (EC 3.4.22.64), caspase-12, caspase-13 and caspase-14 in mice. Contains a caspase-recruitment domain (CARD) in its N-terminal prodomain, which plays a role in procaspase activation [1823, 393]. Cleaves pro-interleukin-1 $\beta$  (pro-IL-1 $\beta$ ) to form mature IL-1 $\beta$ , a potent mediator of inflammation. Also activates the proinflammatory cytokine, IL-18, which is also known as interferon- $\gamma$ -inducing factor [1823]. Inhibited by Ac-Tyr-Val-Ala-Asp-CHO. Caspase-11 plays a critical role in the activation of caspase-1 in mice, whereas caspase-4 enhances its activation in humans [393]. Belongs in peptidase family C14.

**References:** [1191, 2919, 2918, 38, 1809, 1823, 393]

[EC 3.4.22.36 created 1993, modified 1997, modified 2007]

#### EC 3.4.22.37

**Accepted name:** gingipain R  
**Reaction:** Hydrolysis of proteins and small molecule substrates, with a preference for Arg in P1  
**Other name(s):** Arg-gingipain; gingipain-1; argingipain; Arg-gingivain-55 proteinase; Arg-gingivain-70 proteinase; Arg-gingivain-75 proteinase; arginine-specific cysteine protease; arginine-specific gingipain; arginine-specific gingivain; RGP-1; RGP  
**Comments:** A secreted endopeptidase from the bacterium *Porphyromonas gingivalis*. Strongly activated by glycine [420], and stabilized by Ca<sup>2+</sup>. Precursor molecule contains a hemagglutinin domain [1465, 2250]. Misleadingly described in some literature as "trypsin-like", being a cysteine peptidase, type example of family C25  
**References:** [420, 1465, 2250]

[EC 3.4.22.37 created 1996]

#### EC 3.4.22.38

**Accepted name:** cathepsin K  
**Reaction:** Broad proteolytic activity. With small-molecule substrates and inhibitors, the major determinant of specificity is P2, which is preferably Leu, Met > Phe, and not Arg  
**Other name(s):** cathepsin O and cathepsin X (both misleading, having been used for other enzymes); cathepsin O<sub>2</sub>  
**Comments:** Prominently expressed in mammalian osteoclasts, and believed to play a role in bone resorption. In peptidase family C1 (papain family)  
**References:** [1248, 268, 301, 3331, 1872]

[EC 3.4.22.38 created 1997]

#### EC 3.4.22.39

**Accepted name:** adenain  
**Reaction:** Cleaves proteins of the adenovirus and its host cell at two consensus sites: -Yaa-Xaa-Gly-Gly-Xaa- and -Yaa-Xaa-Gly-Xaa-Gly- (in which Yaa is Met, Ile or Leu, and Xaa is any amino acid)  
**Comments:** A cysteine endopeptidase from adenoviruses, the type example of peptidase family C5, with a protein fold unlike that known for any other peptidase [617]. Activity is greatly stimulated by the binding to the enzyme of an 11-residue peptide from the adenovirus capsid protein pre-VI at a site separate from the active site [3143]  
**References:** [3143, 617, 3142]

[EC 3.4.22.39 created 2000]

#### EC 3.4.22.40

**Accepted name:** bleomycin hydrolase



**Reaction:** Inactivates bleomycin B2 (a cytotoxic glycometallopeptide) by hydrolysis of a carboxamide bond of  $\beta$ -aminoalanine, but also shows general aminopeptidase activity. The specificity varies somewhat with source, but amino acid arylamides of Met, Leu and Ala are preferred [1]

**Other name(s):** aminopeptidase C (*Lactococcus lactis*) [4]

**Comments:** The molecule is a homohexamer in which the monomers have a papain-like tertiary structure (in peptidase family C1). The active sites are on the walls of a central channel through the molecule, and access of substrate molecules to them is obstructed by this and by the C-terminus of each polypeptide chain [3333]. Bleomycin can scarcely be the natural substrate, and there are reports of limited endopeptidase activity. Known from bacteria as well as eukaryotic organisms. Hydrolase H from chicken muscle has many similarities to bleomycin hydrolase, but hydrolyses Ph-CO-Arg-2-naphthylamine as well as aminopeptidase substrates [13].

**References:** [303, 13, 3333, 1929]

[EC 3.4.22.40 created 2000]

#### EC 3.4.22.41

**Accepted name:** cathepsin F

**Reaction:** The recombinant enzyme cleaves synthetic substrates with Phe and Leu (better than Val) in P2, with high specificity constant ( $k_{cat}/K_m$ ) comparable to that of cathepsin L

**Comments:** Cathepsin F is a lysosomal cysteine endopeptidase of family C1 (papain family), most active at pH 5.9. The enzyme is unstable at neutral pH values and is inhibited by compound E-64. Cathepsin F is expressed in most tissues of human, mouse and rat. Human gene locus: 11q13.1-13.3

**References:** [2525, 2028, 3166, 3104]

[EC 3.4.22.41 created 2000]

#### EC 3.4.22.42

**Accepted name:** cathepsin O

**Reaction:** The recombinant human enzyme hydrolyses synthetic endopeptidase substrates including Z-Phe-Arg-NHMec and Z-Arg-Arg-NHMec

**Comments:** Cathepsin O is a lysosomal cysteine peptidase of family C1 (papain family). The recombinant human enzyme is catalytically active at pH 6.0 and is inhibited by compound E-64. Cathepsin O is ubiquitously expressed in human tissues and the human gene locus is 4q31-32

**References:** [2523, 3057]

[EC 3.4.22.42 created 2000]

#### EC 3.4.22.43

**Accepted name:** cathepsin V

**Reaction:** The recombinant enzyme hydrolyses proteins (serum albumin, collagen) and synthetic substrates (Z-Phe-Arg-NHMec > Z-Leu-Arg-NHMec > Z-Val-Arg-NHMec)

**Other name(s):** Cathepsin L2; cathepsin U

**Comments:** Cathepsin V is a human lysosomal cysteine endopeptidase of family C1 (papain family) that is maximally active at pH 5.7 and unstable at neutral pH. Compound E-64, leupeptin and chicken cystatin are inhibitors. Human cathepsin V shows expression restricted to thymus, testis, corneal epithelium and some colon and breast carcinomas. Human gene locus: 9q22.2

**References:** [302, 14, 2524]

[EC 3.4.22.43 created 2000]

#### EC 3.4.22.44

**Accepted name:** nuclear-inclusion-a endopeptidase

**Reaction:** Hydrolyses glutamyl bonds, and activity is further restricted by preferences for the amino acids in P6 - P1' that vary with the species of potyvirus, e.g. Glu-Xaa-Xaa-Tyr-Xaa-Gln—(Ser or Gly) for the enzyme from tobacco etch virus. The natural substrate is the viral polyprotein, but other proteins and oligopeptides containing the appropriate consensus sequence are also cleaved.

**Other name(s):** potyvirus NIa protease

**Comments:** The potyviruses cause diseases in plants, and inclusion bodies appear in the host cell nuclei; protein a of the inclusion bodies is the endopeptidase. The enzyme finds practical use when encoded in vectors for the artificial expression of recombinant fusion proteins, since it can confer on them the capacity for autolytic cleavage. It is also reported that transgenic plants expressing the enzyme are resistant to viral infection. Type example of peptidase family C4.

**References:** [752, 1443, 2844, 1446]

[EC 3.4.22.44 created 2000]

#### EC 3.4.22.45

**Accepted name:** helper-component proteinase

**Reaction:** Hydrolyses a Gly—Gly bond at its own C-terminus, commonly in the sequence -Tyr-Xaa-Val-Gly—Gly, in the processing of the potyviral polyprotein

**Other name(s):** HC-Pro

**Comments:** Known from many potyviruses. The helper component-proteinase of the tobacco etch virus is a multifunctional protein with several known activities: the N-terminal region is required for aphid transmission and efficient genome amplification, the central region is required for long-distance movement in plants, and the C-terminal domain has cysteine endopeptidase activity. Type example of peptidase family C6.

**References:** [1393, 3058]

[EC 3.4.22.45 created 2001]

#### EC 3.4.22.46

**Accepted name:** L-peptidase

**Reaction:** Autocatalytically cleaves itself from the polyprotein of the foot-and-mouth disease virus by hydrolysis of a Lys—Gly bond, but then cleaves host cell initiation factor eIF-4G at bonds -Gly—Arg- and -Lys—Arg-

**Comments:** Best known from foot-and-mouth disease virus, but occurs in other aphthoviruses and cardioviruses. Destruction of initiation factor eIF-4G has the effect of shutting off host-cell protein synthesis while allowing synthesis of viral proteins to continue. The tertiary structure reveals a distant relationship to papain and, consistent with this, compound E-64 is inhibitory. Type example of peptidase family C28.

**References:** [2276, 997]

[EC 3.4.22.46 created 2001]

#### EC 3.4.22.47

**Accepted name:** gingipain K

**Reaction:** Endopeptidase with strict specificity for lysyl bonds

**Other name(s):** Lys-gingipain; PrtP proteinase

**Comments:** Activity is stimulated by glycine. Known from the bacterium *Porphyromonas gingivalis* and contributes to the pathogenicity of the organism. In peptidase family C25.

**References:** [2284, 522]

[EC 3.4.22.47 created 2003]

#### EC 3.4.22.48

**Accepted name:** staphopain

**Reaction:** Broad endopeptidase action on proteins including elastin, but rather limited hydrolysis of small-molecule substrates. Assays are conveniently made with hemoglobin, casein or Z-Phe-Arg-NHMec as substrate

**Other name(s):** staphylopain

**Comments:** Known from species of Staphylococcus. Type example of peptidase family C47.

**References:** [1157, 2314, 657]

[EC 3.4.22.48 created 2003]

#### EC 3.4.22.49

**Accepted name:** separase

**Reaction:** All bonds known to be hydrolysed by this endopeptidase have arginine in P1 and an acidic residue in P4. P6 is often occupied by an acidic residue or by a hydroxy-amino-acid residue, the phosphorylation of which enhances cleavage

**Other name(s):** separin

**Comments:** In both budding yeast and human cells, cleavage of the cohesin subunit Scc1 by separase is required for sister chromatid separation in mitosis. Budding yeast separase is also known to cleave the Rec8 subunit of a meiotic cohesin complex and the kinetochore protein Slk19. Type example of peptidase family C50.

**References:** [3084]

[EC 3.4.22.49 created 2003]

#### EC 3.4.22.50

**Accepted name:** V-cath endopeptidase

**Reaction:** Endopeptidase of broad specificity, hydrolyzing substrates of both cathepsin L and cathepsin B

**Other name(s):** AcNPV protease; BmNPV protease; NPV protease; baculovirus cathepsin; nucleopolyhedrosis virus protease; viral cathepsin

**Comments:** In peptidase family C1. Contributes to the liquefaction of the tissues of the insect host in the late stages of infection by the baculovirus.

**References:** [2688, 1082]

[EC 3.4.22.50 created 2003]

#### EC 3.4.22.51

**Accepted name:** cruzipain

**Reaction:** Broad endopeptidase specificity similar to that of cathepsin L

**Other name(s):** congopain; cruzain; evansain; trypanopain

**Comments:** In peptidase family C1. Is located in the digestive vacuoles of the parasitic trypanosome and contributes to the nutrition of the organism by digestion of host proteins.

**References:** [383]

[EC 3.4.22.51 created 2003]

#### EC 3.4.22.52

**Accepted name:** calpain-1

**Reaction:** Broad endopeptidase specificity

**Other name(s):**  $\mu$ -calpain; calcium-activated neutral protease I

**Comments:** In peptidase family C2. Requires  $\text{Ca}^{2+}$  at micromolar concentrations for activity. Cytosolic in animal cells. The active enzyme molecule is a heterodimer in which the large subunit contains the peptidase unit, and the small subunit is also a component of EC 3.4.22.53, calpain-2.

**References:** [671]

[EC 3.4.22.52 created 1981 as EC 3.4.22.17, transferred 2003 to EC 3.4.22.52]

#### EC 3.4.22.53

**Accepted name:** calpain-2

**Reaction:** Broad endopeptidase specificity

**Other name(s):** calcium-activated neutral protease II; *m*-calpain; milli-calpain

**Comments:** Type example of peptidase family C2. Requires Ca<sup>2+</sup> at millimolar concentrations for activity. Cytosolic in animal cells. The active enzyme molecule is a heterodimer in which the large subunit contains the peptidase unit, and the small subunit is also a component of EC 3.4.22.52, calpain-1.

**References:** [2783, 671]

[EC 3.4.22.53 created 1981 as EC 3.4.22.17, transferred 2003 to EC 3.4.22.53]

#### EC 3.4.22.54

**Accepted name:** calpain-3

**Reaction:** Broad endopeptidase activity

**Other name(s):** p94; calpain p94; CAPN3; muscle calpain; calpain 3; calcium-activated neutral proteinase 3; muscle-specific calcium-activated neutral protease 3; CANP 3; calpain L3

**Comments:** This Ca<sup>2+</sup>-dependent enzyme is found in skeletal muscle and is genetically linked to limb girdle muscular dystrophy type 2A [2727, 601]. The enzyme is activated by autoproteolytic cleavage of insertion sequence 1 (IS1), which allows substrates and inhibitors gain access to the active site [601]. Substrates include the protein itself [2412, 601] and connectin/titin [2728, 2198]. Belongs in peptidase family C2.

**References:** [2727, 2728, 2412, 601, 2198]

[EC 3.4.22.54 created 2007]

#### EC 3.4.22.55

**Accepted name:** caspase-2

**Reaction:** Strict requirement for an Asp residue at P1, with Asp<sup>316</sup> being essential for proteolytic activity and has a preferred cleavage sequence of Val-Asp-Val-Ala-Asp—

**Other name(s):** ICH-1; NEDD-2; caspase-2L; caspase-2S; neural precursor cell expressed developmentally down-regulated protein 2; CASP-2; NEDD2 protein

**Comments:** Caspase-2 is an initiator caspase, as are caspase-8 (EC 3.4.22.61), caspase-9 (EC 3.4.22.62) and caspase-10 (EC 3.4.22.63) [393]. Contains a caspase-recruitment domain (CARD) in its N-terminal prodomain, which plays a role in procaspase activation [393]. Two forms of caspase-2 with antagonistic effects exist: caspase-2L induces programmed cell death and caspase-2S suppresses cell death [2,3,5]. Caspase-2 is activated by caspase-3 (EC 3.4.22.56), or by a caspase-3-like protease. Activation involves cleavage of the N-terminal prodomain, followed by self-proteolysis between the large and small subunits of pro-caspase-2 and further proteolysis into smaller fragments [1663]. Proteolysis occurs at Asp residues and the preferred substrate for this enzyme is a pentapeptide rather than a tetrapeptide [3334]. Apart from itself, the enzyme can cleave golgin-16, which is present in the Golgi complex and has a cleavage site that is unique for caspase-2 [1796, 3334].  $\alpha$ II-Spectrin, a component of the membrane cytoskeleton, is a substrate of the large isoform of pro-caspase-2 (caspase-2L) but not of the short isoform (caspase-2S). Belongs in peptidase family C14.

**References:** [1550, 3115, 1663, 1796, 3334, 393]

[EC 3.4.22.55 created 2007]

#### EC 3.4.22.56

**Accepted name:** caspase-3

**Reaction:** Strict requirement for an Asp residue at positions P1 and P4. It has a preferred cleavage sequence of Asp-Xaa-Xaa-Asp— with a hydrophobic amino-acid residue at P2 and a hydrophilic amino-acid residue at P3, although Val or Ala are also accepted at this position

**Other name(s):** CPP32; apopain; yama protein

**Comments:** Caspase-3 is an effector/executioner caspase, as are caspase-6 (EC 3.4.22.59) and caspase-7 (EC 3.4.22.60) [393]. These caspases are responsible for the proteolysis of the majority of cellular polypeptides [e.g. poly(ADP-ribose) polymerase (PARP)], which leads to the apoptotic phenotype [2079, 393]. Pro-caspase-3 can be activated by caspase-1 (EC 3.4.22.36), caspase-8 (EC 3.4.22.61), caspase-9 (EC 3.4.22.62) and caspase-10 (EC 3.4.22.63) as well as by the serine protease granzyme B [1533]. Caspase-3 can activate procaspase-2 (EC 3.4.22.55) [1663]. Activation occurs by inter-domain cleavage followed by removal of the N-terminal prodomain [1820]. Although Asp-Glu-(Val/Ile)-Asp is thought to be the preferred cleavage sequence, the enzyme can accommodate different residues at P2 and P3 of the substrate [737]. Like caspase-2, a hydrophobic residue at P5 of caspase-3 leads to more efficient hydrolysis, e.g. (Val/Leu)-Asp-Val-Ala-Asp+ is a better substrate than Asp-Val-Ala-Asp+. This is not the case for caspase-7 [737]. Belongs in peptidase family C14.

**References:** [1533, 1663, 2079, 737, 393, 1820]

[EC 3.4.22.56 created 2007]

#### EC 3.4.22.57

**Accepted name:** caspase-4

**Reaction:** Strict requirement for Asp at the P1 position. It has a preferred cleavage sequence of Tyr-Val-Ala-Asp+ but also cleaves at Asp-Glu-Val-Asp+

**Other name(s):** ICE<sub>rel</sub>II; ICERel-II; Ich-2; transcript X; TX; TX protease; caspase 4; CASP-4

**Comments:** This enzyme is part of the family of inflammatory caspases, which also includes caspase-1 (EC 3.4.22.36) and caspase-5 (EC 3.4.22.58) in humans and caspase-11 (EC 3.4.22.64), caspase-12, caspase-13 and caspase-14 in mice. Contains a caspase-recruitment domain (CARD) in its N-terminal prodomain, which plays a role in procaspase activation [3,5,6]. The enzyme is able to cleave itself and the p30 caspase-1 precursor, but, unlike caspase-1, it is very inefficient at generating mature interleukin-1 $\beta$  (IL-1 $\beta$ ) from pro-IL-1 $\beta$  [744, 741]. Both this enzyme and caspase-5 can cleave pro-caspase-3 to release the small subunit (p12) but not the large subunit (p17) [1369]. The caspase-1 inhibitor Ac-Tyr-Val-Ala-Asp-CHO can also inhibit this enzyme, but more slowly [741]. Belongs in peptidase family C14.

**References:** [744, 1371, 1369, 741, 1823, 393]

[EC 3.4.22.57 created 2007]

#### EC 3.4.22.58

**Accepted name:** caspase-5

**Reaction:** Strict requirement for Asp at the P1 position. It has a preferred cleavage sequence of Tyr-Val-Ala-Asp+ but also cleaves at Asp-Glu-Val-Asp+

**Other name(s):** ICERel-III; Ich-3; ICH-3 protease; transcript Y; TY; CASP-5

**Comments:** This enzyme is part of the family of inflammatory caspases, which also includes caspase-1 (EC 3.4.22.36) and caspase-4 (EC 3.4.22.57) in humans and caspase-11 (EC 3.4.22.64), caspase-12, caspase-13 and caspase-14 in mice. Contains a caspase-recruitment domain (CARD) in its N-terminal prodomain, which plays a role in procaspase activation [3,5,6]. The enzyme is able to cleave itself and the p30 caspase-1 precursor, but is very inefficient at generating mature interleukin-1 $\beta$  (IL-1 $\beta$ ) from pro-IL-1 $\beta$  [743, 741]. Both this enzyme and caspase-4 can cleave pro-caspase-3 to release the small subunit (p12) but not the large subunit (p17) [1696]. Unlike caspase-4, this enzyme can be induced by lipopolysaccharide [1696]. Belongs in peptidase family C14.

**References:** [743, 1369, 1696, 741, 1823, 393]

[EC 3.4.22.58 created 2007]

#### EC 3.4.22.59

**Accepted name:** caspase-6

**Reaction:** Strict requirement for Asp at position P1 and has a preferred cleavage sequence of Val-Glu-His-Asp—

**Other name(s):** CASP-6; apoptotic protease Mch-2; Mch2

**Comments:** Caspase-6 is an effector/executioner caspase, as are caspase-3 (EC 3.4.22.56) and caspase-7 (EC 3.4.22.60) [393]. These caspases are responsible for the proteolysis of the majority of cellular polypeptides [e.g. poly(ADP-ribose) polymerase (PARP)], which leads to the apoptotic phenotype [393]. Caspase-6 can cleave its prodomain to produce mature caspase-6, which directly activates caspase-8 (EC 3.4.22.61) and leads to the release of cytochrome *c* from the mitochondria. The release of cytochrome *c* is an essential component of the intrinsic apoptosis pathway [500]. The enzyme can also cleave and inactivate lamins, the intermediate filament scaffold proteins of the nuclear envelope, leading to nuclear fragmentation in the final phases of apoptosis [2,4,5,6]. Belongs in peptidase family C14.

**References:** [500, 393, 1384, 1629, 1769, 2836]

[EC 3.4.22.59 created 2007]

#### EC 3.4.22.60

**Accepted name:** caspase-7

**Reaction:** Strict requirement for an Asp residue at position P1 and has a preferred cleavage sequence of Asp-Glu-Val-Asp—

**Other name(s):** CASP-7; ICE-like apoptotic protease 3; ICE-LAP3; apoptotic protease Mch-3; Mch3; CMH-1

**Comments:** Caspase-7 is an effector/executioner caspase, as are caspase-3 (EC 3.4.22.56) and caspase-6 (EC 3.4.22.59) [393]. These caspases are responsible for the proteolysis of the majority of cellular polypeptides [e.g. poly(ADP-ribose) polymerase (PARP)], which leads to the apoptotic phenotype [2079]. Although a hydrophobic residue at P5 of caspase-2 (EC 3.4.22.55) and caspase-3 leads to more efficient hydrolysis, the amino-acid residue at this location in caspase-7 has no effect [737]. Caspase-7 is activated by the initiator caspases [caspase-8 (EC 3.4.22.61), caspase-9 (EC 3.4.22.62) and caspase-10 (EC 3.4.22.63)]. Removal of the N-terminal prodomain occurs before cleavage in the linker region between the large and small subunits [582]. Belongs in peptidase family C14.

**References:** [393, 2079, 737, 582]

[EC 3.4.22.60 created 2007]

#### EC 3.4.22.61

**Accepted name:** caspase-8

**Reaction:** Strict requirement for Asp at position P1 and has a preferred cleavage sequence of (Leu/Asp/Val)-Glu-Thr-Asp—(Gly/Ser/Ala)

**Other name(s):** FLICE, FADD-like ICE; MACH; MORT<sub>1</sub>-associated CED-3 homolog; Mch5; mammalian Ced-3 homolog 5; CASP-8; ICE-like apoptotic protease 5; FADD-homologous ICE/CED-3-like protease; apoptotic cysteine protease; apoptotic protease Mch-5; CAP4

**Comments:** Caspase-8 is an initiator caspase, as are caspase-2 (EC 3.4.22.55), caspase-9 (EC 3.4.22.62) and caspase-10 (EC 3.4.22.63) [393]. Caspase-8 is the apical activator of the extrinsic (death receptor) apoptosis pathway, triggered by death receptor ligation [256]. It contains two tandem death effector domains (DEDs) in its N-terminal prodomain, and these play a role in procaspase activation [393]. This enzyme is linked to cell surface death receptors such as Fas [393, 774]. When Fas is aggregated by the Fas ligand, procaspase-8 is recruited to the death receptor where it is activated [393]. The enzyme has a preference for Glu at P3 and prefers small residues, such as Gly, Ser and Ala, at the P1' position. It has very broad P4 specificity, tolerating substrates with Asp, Val or Leu in this position [2,3,4]. Endogenous substrates for caspase-8 include procaspase-3, the pro-apoptotic Bcl-2 family member Bid, RIP, PAK2 and the caspase-8 activity modulator FLIP<sub>1</sub> [2511, 774]. Belongs in peptidase family C14.

**References:** [393, 256, 2014, 2511, 774, 237, 249]

[EC 3.4.22.61 created 2007]

### EC 3.4.22.62

**Accepted name:** caspase-9

**Reaction:** Strict requirement for an Asp residue at position P1 and with a marked preference for His at position P2. It has a preferred cleavage sequence of Leu-Gly-His-Asp—Xaa

**Other name(s):** CASP-9; ICE-like apoptotic protease 6; ICE-LAP6; apoptotic protease Mch-6; apoptotic protease-activating factor 3; APAF-3

**Comments:** Caspase-9 is an initiator caspase, as are caspase -2 (EC 3.4.22.55), caspase-8 (EC 3.4.22.61) and caspase-10 (EC 3.4.22.63) [393]. Caspase-9 contains a caspase-recruitment domain (CARD) in its N-terminal prodomain, which plays a role in procaspase activation [393]. An alternatively spliced version of caspase-9 also exists, caspase-9S, that inhibits apoptosis, similar to the situation found with caspase-2 [393]. Phosphorylation of caspase-9 from some species by Akt, a serine-threonine protein kinase, inhibits caspase activity in vitro and caspase activation in vivo [393]. The activity of caspase-9 is increased dramatically upon association with the apoptosome but the enzyme can be activated without proteolytic cleavage [3284, 250]. Procaspase-3 is the enzyme's physiological substrate [3284]. Belongs in peptidase family C14.

**References:** [393, 3284, 250, 2512]

[EC 3.4.22.62 created 2007]

### EC 3.4.22.63

**Accepted name:** caspase-10

**Reaction:** Strict requirement for Asp at position P1 and has a preferred cleavage sequence of Leu-Gln-Thr-Asp—Gly

**Other name(s):** FLICE2, Mch4; CASP-10; ICE-like apoptotic protease 4; apoptotic protease Mch-4; FAS-associated death domain protein interleukin-1 $\beta$ -converting enzyme 2

**Comments:** Caspase-10 is an initiator caspase, as are caspase-2 (EC 3.4.22.55), caspase-8 (EC 3.4.22.61) and caspase-9 (EC 3.4.22.62) [393]. Like caspase-8, caspase-10 contains two tandem death effector domains (DEDs) in its N-terminal prodomain, and these play a role in procaspase activation [393]. The enzyme has many overlapping substrates in common with caspase-8, such as RIP (the cleavage of which impairs NF- $\kappa$ B survival signalling and starts the cell-death process) and PAK2 (associated with some of the morphological features of apoptosis, such as cell rounding and apoptotic body formation) [774]. Bid, a Bcl2 protein, can be cleaved by caspase-3 (EC 3.4.22.56), caspase-8 and caspase-10 at Lys-Gln-Thr-Asp— to yield the pro-apoptotic p15 fragment. The p15 fragment is N-myristoylated and enhances the release of cytochrome *c* from mitochondria (which, in turn, initiates the intrinsic apoptosis pathway). Bid can be further cleaved by caspase-10 and granzyme B but not by caspase-3 or caspase-8 at Ile-Glu-Thr-Asp— to yield a p13 fragment that is not N-myristoylated [774]. Belongs in peptidase family C14.

**References:** [393, 774, 2631, 249]

[EC 3.4.22.63 created 2007]

### EC 3.4.22.64

**Accepted name:** caspase-11

**Reaction:** Strict requirement for Asp at the P1 position and has a preferred cleavage sequence of (Ile/Leu/Val/Phe)-Gly-His-Asp—

**Other name(s):** CASP-11

**Comments:** This murine enzyme is part of the family of inflammatory caspases, which also includes caspase-1 (EC 3.4.22.36), caspase-4 (EC 3.4.22.57) and caspase-5 (EC 3.4.22.58) in humans and caspase-12, caspase-13 and caspase-14 in mice. Contains a caspase-recruitment domain (CARD) in its N-terminal prodomain, which plays a role in procaspase activation. Like caspase-5, but unlike caspase-4, this enzyme can be induced by lipopolysaccharide [1386]. This enzyme not only activates caspase-1, which is required for the maturation of proinflammatory cytokines such as interleukin-1 $\beta$  (IL-1 $\beta$ ) and IL-18, but it also activates caspase-3 (EC 3.4.22.56), which leads to cellular apoptosis under pathological conditions [1386, 1209]. Belongs in peptidase family C14.

**References:** [1386, 1209, 3118, 703, 393]



[EC 3.4.22.64 created 2007]

#### EC 3.4.22.65

**Accepted name:** peptidase 1 (mite)

**Reaction:** Broad endopeptidase specificity

**Other name(s):** allergen Der f 1; allergen Der p 1; antigen Der p 1; antigen Eur m 1; antigen Pso o 1; major mite fecal allergen Der p 1; Der p 1; Der f 1; Eur m 1; endopeptidase 1 (mite)

**Comments:** This enzyme, derived from the house dust mite, is a major component of the allergic immune response [1366]. The substrate specificity of this enzyme is not altogether clear. It cleaves the low-affinity IgE receptor CD23 at Glu<sup>298</sup>—Ser<sup>299</sup> and Ser<sup>155</sup>—Ser<sup>156</sup> [1879]. It also cleaves the pulmonary structural proteins occludin and claudin at Leu—Leu, Asp—Leu and at Gly—Thr bonds [1879, 1366]. It can also cleave the  $\alpha$  subunit of the interleukin-2 (IL-2) receptor (CD25) [2593]. Using a positional scanning combinatorial library, it was found that the major substrate-specificity determinant is for Ala in the P2 position [1060]. The enzyme shows only a slight preference for basic amino acids in the P1 and P3 positions and a preference for aliphatic amino acids such as Ile, Pro, Val, Leu and norleucine in the P4 position [1060]. Belongs in peptidase family C1A.

**References:** [1879, 1366, 1060, 2593, 2592, 2850]

[EC 3.4.22.65 created 2007]

#### EC 3.4.22.66

**Accepted name:** calicivirin

**Reaction:** Endopeptidase with a preference for cleavage when the P1 position is occupied by Glu— and the P1' position is occupied by Gly—

**Other name(s):** Camberwell virus processing peptidase; Chiba virus processing peptidase; Norwalk virus processing peptidase; Southampton virus processing peptidase; Southampton virus; norovirus virus processing peptidase; calicivirus trypsin-like cysteine protease; calicivirus TCP; calicivirus 3C-like protease; calicivirus endopeptidase; rabbit hemorrhagic disease virus 3C endopeptidase

**Comments:** Viruses that are members of the Norovirus genus (Caliciviridae family) are a major cause of epidemic acute viral gastroenteritis [1715]. The nonstructural proteins of these viruses are produced by proteolytic cleavage of a large precursor polyprotein, performed by a protease that is incorporated into the polyprotein []. Cleavage sites are apparently defined by features based on both sequence and structure since several sites in the polyprotein fulfilling the identified sequence requirements are not cleaved [1899]. The presence of acidic (Asp), basic (Arg), aromatic (Tyr) or aliphatic (Leu) amino acids at the P1' position results in only minor differences in cleavage efficiency, suggesting that steric or conformational constraints may play a role in determining specificity [1899]. Changes to the amino acid at the P2 position do not alter cleavage efficiency [1899, 3199]. Belongs in peptidase family C37.

**References:** [1899, 3199, 40, 1715, 1716]

[EC 3.4.22.66 created 2007]

#### EC 3.4.22.67

**Accepted name:** zingipain

**Reaction:** Preferential cleavage of peptides with a proline residue at the P2 position

**Other name(s):** ginger protease; GP-I; GP-II; ginger protease II (*Zingiber officinale*); zingibain

**Comments:** This enzyme is found in ginger (*Zingiber officinale*) rhizome and is a member of the papain family. GP-II contains two glycosylation sites. The enzyme is inhibited by some divalent metal ions, such as Hg<sup>2+</sup>, Cu<sup>2+</sup>, Cd<sup>2+</sup> and Zn<sup>2+</sup> [2174]. Belongs in peptidase family C1.

**References:** [438, 2174, 439]

[EC 3.4.22.67 created 2007]

#### EC 3.4.22.68



**Accepted name:** Ulp1 peptidase  
**Reaction:** Hydrolysis of the  $\alpha$ -linked peptide bond in the sequence Gly-Gly—Ala-Thr-Tyr at the C-terminal end of the small ubiquitin-like modifier (SUMO) propeptide, Smt3, leading to the mature form of the protein. A second reaction involves the cleavage of an  $\epsilon$ -linked peptide bond between the C-terminal glycine of the mature SUMO and the lysine  $\epsilon$ -amino group of the target protein  
**Other name(s):** Smt3-protein conjugate proteinase; Ubl-specific protease 1; Ulp1; Ulp1 endopeptidase; Ulp1 protease  
**Comments:** The enzyme from *Saccharomyces cerevisiae* can also recognize small ubiquitin-like modifier 1 (SUMO-1) from human as a substrate in both SUMO-processing ( $\alpha$ -linked peptide bonds) and SUMO-deconjugation ( $\epsilon$ -linked peptide bonds) reactions [1,2,3]. Ulp1 has several functions, including an essential role in chromosomal segregation and progression of the cell cycle through the G2/M phase of the cell cycle. Belongs in peptidase family C48.  
**References:** [1689, 1668, 2895, 1669, 1223, 1992]

[EC 3.4.22.68 created 2008, modified 2011]

#### EC 3.4.22.69

**Accepted name:** SARS coronavirus main proteinase  
**Reaction:** TSAVLQ—SGFRK-NH<sub>2</sub> and SGVTFQ—GKFKK the two peptides corresponding to the two self-cleavage sites of the SARS 3C-like proteinase are the two most reactive peptide substrates. The enzyme exhibits a strong preference for substrates containing Gln at P1 position and Leu at P2 position.  
**Other name(s):** 3cLpro; 3C-like protease; coronavirus 3C-like protease; Mpro; SARS 3C-like protease; SARS coronavirus 3CL protease; SARS coronavirus main peptidase; SARS coronavirus main protease; SARS-CoV 3CLpro enzyme; SARS-CoV main protease; SARS-CoV Mpro; severe acute respiratory syndrome coronavirus main protease  
**Comments:** SARS coronavirus main protease is the key enzyme in SARS coronavirus replicase polyprotein processing. In peptidase family C30.  
**References:** [938, 736, 24]

[EC 3.4.22.69 created 2009]

#### EC 3.4.22.70

**Accepted name:** sortase A  
**Reaction:** The enzyme catalyses a cell wall sorting reaction in which a surface protein with a sorting signal containing a LPXTG motif is cleaved between the Thr and Gly residue. The resulting threonine carboxyl end of the protein is covalently attached to a pentaglycine cross-bridge of peptidoglycan.  
**Other name(s):** SrtA; SrtA protein; SrtA sortase  
**Comments:** In peptidase family C60.  
**References:** [2941, 3346, 2351]

[EC 3.4.22.70 created 2009]

#### EC 3.4.22.71

**Accepted name:** sortase B  
**Reaction:** The enzyme catalyses a cell wall sorting reaction in which a surface protein with a sorting signal containing a NPXTN motif is cleaved between the Thr and Asn residue. The resulting threonine carboxyl end of the protein is covalently attached to a pentaglycine cross-bridge of peptidoglycan.  
**Other name(s):** SrtB  
**Comments:** In peptidase family C60.  
**References:** [3347, 213, 470]

[EC 3.4.22.71 created 2009]

## EC 3.4.23 Aspartic endopeptidases

### EC 3.4.23.1

- Accepted name:** pepsin A  
**Reaction:** Preferential cleavage: hydrophobic, preferably aromatic, residues in P1 and P1' positions. Cleaves Phe<sup>1</sup>-Val, Gln<sup>4</sup>-His, Glu<sup>13</sup>-Ala, Ala<sup>14</sup>-Leu, Leu<sup>15</sup>-Tyr, Tyr<sup>16</sup>-Leu, Gly<sup>23</sup>-Phe, Phe<sup>24</sup>-Phe and Phe<sup>25</sup>-Tyr bonds in the B chain of insulin  
**Other name(s):** pepsin; lactated pepsin; pepsin fortior; fundus-pepsin; elixir lactate of pepsin; P I; lactated pepsin elixir; P II; pepsin R; pepsin D  
**Comments:** The predominant endopeptidase in the gastric juice of vertebrates, formed from pepsinogen A by limited proteolysis. Human pepsin A occurs in five molecular forms. Pig pepsin D [1620, 1619] is unphosphorylated pepsin A. Type example of peptidase family A1.  
**References:** [1620, 1619, 793, 1305, 826, 2881, 2300]

[EC 3.4.23.1 created 1961 as EC 3.4.4.1, transferred 1972 to EC 3.4.23.1, modified 1986, modified 1989]

### EC 3.4.23.2

- Accepted name:** pepsin B  
**Reaction:** Degradation of gelatin; little activity on hemoglobin. Specificity on B chain of insulin more restricted than that of pepsin A; does not cleave at Phe<sup>1</sup>-Val, Gln<sup>4</sup>-His or Gly<sup>23</sup>-Phe  
**Other name(s):** parapepsin I; pig gelatinase  
**Comments:** Formed from pig pepsinogen B. In peptidase family A1 (pepsin A family)  
**References:** [2482]

[EC 3.4.23.2 created 1961 as EC 3.4.4.2, transferred 1972 to EC 3.4.23.2, modified 1986]

### EC 3.4.23.3

- Accepted name:** gastricsin  
**Reaction:** More restricted specificity than pepsin A, but shows preferential cleavage at Tyr- bonds. High activity on hemoglobin  
**Other name(s):** pepsin C; pig parapepsin II; parapepsin II  
**Comments:** Formed from progastricsin, apparently in the gastric juice of most vertebrates. In addition to the fundus, progastricsin is also secreted in antrum and proximal duodenum. Seminal plasma contains a zymogen that is immunologically identical with progastricsin [2402]. In peptidase family A1 (pepsin A family).  
**References:** [2482, 2880, 791, 792, 1819, 2402, 1088]

[EC 3.4.23.3 created 1965 as EC 3.4.4.22, transferred 1972 to EC 3.4.23.3, modified 1986]

### EC 3.4.23.4

- Accepted name:** chymosin  
**Reaction:** Broad specificity similar to that of pepsin A. Clots milk by cleavage of a single Ser-Phe<sup>105</sup>-Met-Ala bond in  $\kappa$ -chain of casein  
**Other name(s):** rennin (but this should be avoided since it leads to confusion with renin)  
**Comments:** Neonatal gastric enzyme with high milk clotting and weak general proteolytic activity, formed from prochymosin. Found among mammals with postnatal uptake of immunoglobulins. In peptidase family A1 (pepsin A family)  
**References:** [790, 1063, 3070]

[EC 3.4.23.4 created 1961 as EC 3.4.4.3, transferred 1972 to EC 3.4.23.4, modified 1986]

### EC 3.4.23.5

**Accepted name:** cathepsin D

**Reaction:** Specificity similar to, but narrower than, that of pepsin A. Does not cleave the Gln<sup>4</sup>-His bond in B chain of insulin

**Comments:** Occurs intracellularly, in lysosomes. A zymogen form is known [478]. In peptidase family A1 (pepsin A family).

**References:** [154, 2847, 748, 478]

[EC 3.4.23.5 created 1965 as EC 3.4.4.23, transferred 1972 to EC 3.4.23.5, modified 1986]

[3.4.23.6 *Transferred entry. now EC 3.4.23.30 pycnoporopepsin*]

[EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981, deleted 1992 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978]]

[3.4.23.7 *Transferred entry. Penicillium janthinellum acid proteinase. Now EC 3.4.23.20, penicillopepsin*]

[EC 3.4.23.7 created 1972, modified 1981, deleted 1978 [transferred to EC 3.4.23.6, deleted 1992]]

[3.4.23.8 *Transferred entry. yeast proteinase A. Now EC 3.4.23.25, saccharopepsin*]

[EC 3.4.23.8 created 1972, modified 1981, deleted 1978 [transferred to EC 3.4.23.6, deleted 1992]]

[3.4.23.9 *Transferred entry. Rhizopus acid proteinase. Now EC 3.4.23.21, rhizopuspepsin*]

[EC 3.4.23.9 created 1972, modified 1981, deleted 1978 [transferred to EC 3.4.23.6, deleted 1992]]

[3.4.23.10 *Transferred entry. Endothia acid proteinase. Now EC 3.4.23.22, endothiapepsin*]

[EC 3.4.23.10 created 1972, modified 1981, deleted 1978 [transferred to EC 3.4.23.6, deleted 1992]]

[3.4.23.11 *Deleted entry. thyroid aspartic proteinase*]

[EC 3.4.23.11 created 1978, modified 1981, deleted 1992]

#### EC 3.4.23.12

**Accepted name:** nepenthesin

**Reaction:** Similar to pepsin, but also cleaves on either side of Asp and at Lys—Arg

**Other name(s):** *Nepenthes* aspartic proteinase; *Nepenthes* acid proteinase; nepenthacin; nepenthasin; aspartyl endopeptidase

**Comments:** From the insectivorous plants *Nepenthes* spp. (secretions) and *Drosera peltata* (ground-up leaves). Aspartic endopeptidases are probably present in many other plants, including *Lotus* [2638] and sorghum [879]. In peptidase family A1 (pepsin A family)

**References:** [44, 879, 2638, 43, 2839, 2929]

[EC 3.4.23.12 created 1972 as EC 3.4.99.4, transferred 1978 to EC 3.4.23.12, modified 1981]

[3.4.23.13 *Deleted entry. Lotus aspartic proteinase*]

[EC 3.4.23.13 created 1978, modified 1981, deleted 1992]

[3.4.23.14 *Deleted entry. sorghum aspartic proteinase*]

[EC 3.4.23.14 created 1978, modified 1981, deleted 1992]

#### EC 3.4.23.15

**Accepted name:** renin

**Reaction:** Cleavage of Leu— bond in angiotensinogen to generate angiotensin I

**Other name(s):** angiotensin-forming enzyme; angiotensinogenase

**Comments:** Formed from prorenin in plasma and kidney. In peptidase family A1 (pepsin A family).

**References:** [1247, 2689, 1246, 2655]

[EC 3.4.23.15 created 1961 as EC 3.4.4.15, transferred 1972 to EC 3.4.99.19, transferred 1981 to EC 3.4.23.15]

#### EC 3.4.23.16

**Accepted name:** HIV-1 retropepsin  
**Reaction:** Specific for a P1 residue that is hydrophobic, and P1' variable, but often Pro  
**Other name(s):** human immunodeficiency virus type 1 protease; gag protease; HIV aspartyl protease; HIV proteinase; retroproteinase; HIV-1 protease; HIV-2 protease  
**Comments:** Present in human immunodeficiency virus type 1. Contributes to the maturation of the viral particle, and is a target of antiviral drugs. Active enzyme is a dimer of identical 11-kDa subunits. Similar enzymes occur in other retroviruses [1560]. Type example of peptidase family A2  
**References:** [1560, 667]

[EC 3.4.23.16 created 1992, modified 2000]

#### EC 3.4.23.17

**Accepted name:** *pro*-opiomelanocortin converting enzyme  
**Reaction:** Cleavage at paired basic residues in certain prohormones, either between them, or on the carboxyl side  
**Other name(s):** prohormone converting enzyme; *pro*-opiomelanocortin-converting enzyme; proopiomelanocortin proteinase; PCE  
**Comments:** A 70 kDa membrane-bound enzyme isolated from cattle pituitary secretory vesicle.  
**References:** [1733, 1732, 721]

[EC 3.4.23.17 created 1989 as EC 3.4.99.38, transferred 1992 to EC 3.4.23.17]

#### EC 3.4.23.18

**Accepted name:** aspergillopepsin I  
**Reaction:** Hydrolysis of proteins with broad specificity. Generally favours hydrophobic residues in P1 and P1', but also accepts Lys in P1, which leads to activation of trypsinogen. Does not clot milk  
**Other name(s):** *Aspergillus* acid protease; *Aspergillus* acid proteinase; *Aspergillus* aspartic proteinase; *Aspergillus awamori* acid proteinase; *Aspergillus* carboxyl proteinase; (see also Comments); carboxyl proteinase; *Aspergillus kawachii* aspartic proteinase; *Aspergillus saitoi* acid proteinase; pepsin-type aspartic proteinase; *Aspergillus niger* acid proteinase; sumizyme AP; proctase P; denapsin; denapsin XP 271; proctase  
**Comments:** Found in a variety of *Aspergillus* species (imperfect fungi): *Aspergillus awamori* (awamolin, aspergillopepsin A: [2210]), *A. foetidus* (aspergillopepsin F: [2209]), *A. fumigatus* [2231], *A. kawachii* [3239], *A. niger* (proteinase B, proctase B: [1968, 397]), *A. oryzae* (trypsinogen kinase: [545, 1782]), *A. saitoi* (aspergillopeptidase A: [1782]), and *A. sojae* [2871, 1782]. In peptidase family A1 (pepsin A family). Formerly included in EC 3.4.23.6  
**References:** [1525, 1968, 545, 397, 2871, 2209, 2231, 2210, 3239, 1782]

[EC 3.4.23.18 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992)]

#### EC 3.4.23.19

**Accepted name:** aspergillopepsin II  
**Reaction:** Preferential cleavage in B chain of insulin: Asn<sup>3</sup> + Gln, Gly<sup>13</sup> + Ala, Tyr<sup>26</sup> + Thr  
**Other name(s):** proteinase A; proctase A; *Aspergillus niger* var. *macrosporus* aspartic proteinase  
**Comments:** Isolated from *Aspergillus niger* var. *macrosporus*, distinct from proteinase B (see aspergillopepsin I) in specificity and insensitivity to pepstatin. In peptidase family A4 (scytalidopepsin B family). Formerly included in EC 3.4.23.6  
**References:** [397, 1226]

[EC 3.4.23.19 created 1992 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992)]

#### EC 3.4.23.20

**Accepted name:** penicillopepsin

**Reaction:** Hydrolysis of proteins with broad specificity similar to that of pepsin A, preferring hydrophobic residues at P1 and P1', but also cleaving Gly<sup>20</sup>—Glu in the B chain of insulin. Clots milk, and activates trypsinogen

**Other name(s):** peptidase A; *Penicillium janthinellum* aspartic proteinase; acid protease A; *Penicillium citrinum* acid proteinase; *Penicillium cyclopium* acid proteinase; *Penicillium expansum* acid proteinase; *Penicillium janthinellum* acid proteinase; *Penicillium expansum* aspartic proteinase; *Penicillium* aspartic proteinase; *Penicillium caseicolum* aspartic proteinase; *Penicillium roqueforti* acid proteinase; *Penicillium duponti* aspartic proteinase; *Penicillium citrinum* aspartic proteinase

**Comments:** From the imperfect fungus *Penicillium janthinellum*. In peptidase family A1 (pepsin A family). Closely related enzymes have been isolated from *P. roqueforti* [3321] and *P. duponti* [698].

**References:** [1779, 3321, 698, 1158, 1197]

[EC 3.4.23.20 created 1992 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992)]

#### EC 3.4.23.21

**Accepted name:** rhizopuspepsin

**Reaction:** Hydrolysis of proteins with broad specificity similar to that of pepsin A, preferring hydrophobic residues at P1 and P1'. Clots milk and activates trypsinogen. Does not cleave Gln<sup>4</sup>-His, but does cleave His<sup>10</sup>—Leu and Val<sup>12</sup>—Glu in B chain of insulin

**Other name(s):** *Rhizopus* aspartic proteinase; neurase; *Rhizopus* acid protease; *Rhizopus* acid proteinase

**Comments:** From the zygomycete fungus *Rhizopus chinensis*. A similar endopeptidase is found in *R. niveus* [1574]. In peptidase family A1 (pepsin A family).

**References:** [2987, 1574, 2175, 2796]

[EC 3.4.23.21 created 1992 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992)]

#### EC 3.4.23.22

**Accepted name:** endothiapepsin

**Reaction:** Hydrolysis of proteins with specificity similar to that of pepsin A; prefers hydrophobic residues at P1 and P1', but does not cleave Ala<sup>14</sup>-Leu in the B chain of insulin or Z-Glu-Tyr. Clots milk

**Other name(s):** Endothia aspartic proteinase; Endothia acid proteinase; *Endothia parasitica* acid proteinase; *Endothia parasitica* aspartic proteinase

**Comments:** From the ascomycete *Endothia parasitica*. In peptidase family A1 (pepsin A family).

**References:** [3172, 3192, 143, 486]

[EC 3.4.23.22 created 1992 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992)]

#### EC 3.4.23.23

**Accepted name:** mucorpepsin

**Reaction:** Hydrolysis of proteins, favouring hydrophobic residues at P1 and P1'. Clots milk. Does not accept Lys at P1, and hence does not activate trypsinogen

**Other name(s):** Mucor rennin; Mucor aspartic proteinase; Mucor acid proteinase; Mucor acid protease; *Mucor miehei* aspartic proteinase; *Mucor miehei* aspartic protease; Mucor aspartic proteinase; *Mucor pusillus* empo-  
rase; Fromase 100; *Mucor pusillus* rennin; Fromase 46TL; *Mucor miehei* rennin

**Comments:** Isolated from the zygomycete fungi *Mucor pusillus* and *M. miehei*. The two species variants show 83% sequence identity and are immunologically crossreactive. In peptidase family A1 (pepsin A fam-  
ily). Formerly included in EC 3.4.23.6

**References:** [73, 2214, 2765, 2178, 171]

[EC 3.4.23.23 created 1992 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992)]

#### EC 3.4.23.24

**Accepted name:** candidapepsin

**Reaction:** Preferential cleavage at the carboxyl of hydrophobic amino acids, but fails to cleave Leu<sup>15</sup>-Tyr, Tyr<sup>16</sup>-Leu and Phe<sup>24</sup>-Phe of insulin B chain. Activates trypsinogen, and degrades keratin

**Other name(s):** *Candida albicans* aspartic proteinase; *Candida albicans* carboxyl proteinase; *Candida albicans* secre-  
tory acid proteinase; *Candida olea* acid proteinase; *Candida* aspartic proteinase; *Candida olea* aspar-  
tic proteinase; *Candida albicans* aspartic proteinase

**Comments:** This endopeptidase from the imperfect yeast *Candida albicans* is inhibited by pepstatin, but not by methyl 2-diazoacetamidohexanoate or 1,2-epoxy-3-(*p*-nitrophenoxy)propane. In peptidase family A1 (pepsin A family). Formerly included in EC 3.4.23.6

**References:** [2406, 2475, 2065, 1739]

[EC 3.4.23.24 created 1992 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992)]

#### EC 3.4.23.25

**Accepted name:** saccharopepsin

**Reaction:** Hydrolysis of proteins with broad specificity for peptide bonds. Cleaves -Leu-Leu-Val-Tyr bond in a synthetic substrate. Does not act on esters of Tyr or Arg

**Other name(s):** yeast endopeptidase A; *Saccharomyces* aspartic proteinase; aspartic proteinase yscA; proteinase A; proteinase yscA; yeast proteinase A; *Saccharomyces cerevisiae* aspartic proteinase A; yeast proteinase A; PRA

**Comments:** Located in the vacuole of the baker's yeast (*Saccharomyces cerevisiae*) cell. In peptidase family A1 (pepsin A family).

**References:** [1074, 1894, 47]

[EC 3.4.23.25 created 1992 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992)]

#### EC 3.4.23.26

**Accepted name:** rhodotorulapepsin

**Reaction:** Specificity similar to that of pepsin A. Cleaves Z-Lys-Ala-Ala-Ala and activates trypsinogen

**Other name(s):** Rhodotorula aspartic proteinase; *Cladosporium* acid protease; *Cladosporium* acid proteinase; *Pae-*  
*cilomyces* proteinase; *Cladosporium* aspartic proteinase; *Paecilomyces* proteinase; *Rhodotorula glutini-*  
*nis* aspartic proteinase; *Rhodotorula glutinis* acid proteinase; *Rhodotorula glutinis* aspartic proteinase II; Rhodotorula acid proteinase

**Comments:** From the imperfect yeast *Rhodotorula glutinis*. Somewhat similar enzymes have been isolated from the imperfect yeast-like organism *Cladosporium* sp. [2006, 2129] and the imperfect fungus *Pae-*  
*cilomyces varioti* [2543, 2544].

**References:** [2543, 2544, 1368, 2006, 2130, 2129, 2838, 1782]

[EC 3.4.23.26 created 1992 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992)]

[3.4.23.27 Transferred entry. *physaropepsin*. Now EC 3.4.21.103, *physarolisin*]

[EC 3.4.23.27 created 1992 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992), deleted 2003]

#### EC 3.4.23.28

**Accepted name:** acrocyllindropepsin

**Reaction:** Preference for hydrophobic residues at P1 and P1'. Action on the B chain of insulin is generally similar to that of pepsin A, but it also cleaves Leu<sup>6</sup>—Cys(SO<sub>3</sub>H), Glu<sup>21</sup>—Arg and Asn<sup>3</sup>—Gln, although not Gln<sup>4</sup>-His

**Other name(s):** *Acrocyllindrium* proteinase; *Acrocyllindrium* acid proteinase

**Comments:** From the imperfect fungus *Acrocyllindrium* sp. Has a very low pH optimum on casein of 2.0. In peptidase family A1 (pepsin A family).

**References:** [3007, 1218, 2838]

[EC 3.4.23.28 created 1992 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992)]

#### EC 3.4.23.29

**Accepted name:** polyporopepsin

**Reaction:** Milk clotting activity, broad specificity, but fails to cleave Leu<sup>15</sup>-Tyr or Tyr<sup>16</sup>-Leu of insulin B chain

**Other name(s):** Polyporus aspartic proteinase; *Irpex lacteus* aspartic proteinase; *Irpex lacteus* carboxyl proteinase B

**Comments:** From the basidiomycete *Polyporus tulipiferae* (formerly *Irpex lacteus*). In peptidase family A1 (pepsin A family)

**References:** [1485, 1487]

[EC 3.4.23.29 created 1992]

#### EC 3.4.23.30

**Accepted name:** pycnoporopepsin

**Reaction:** Similar to pepsin A, but narrower, cleaving only three bonds in the B chain of insulin: Ala<sup>14</sup>—Leu, Tyr<sup>16</sup>—Leu, and Phe<sup>24</sup>—Phe

**Other name(s):** proteinase Ia; *Pycnoporus coccineus* aspartic proteinase; *Trametes* acid proteinase

**Comments:** From the basidiomycete *Pycnoporus sanguineus*, formerly known as *P. coccineus* and *Trametes sanguinea*. Formerly included in EC 3.4.23.6

**References:** [2939, 2987, 1219]

[EC 3.4.23.30 created 1992 (EC 3.4.23.6 created 1992 (EC 3.4.23.6 created 1961 as EC 3.4.4.17, transferred 1972 to EC 3.4.23.6, modified 1981 [EC 3.4.23.7, EC 3.4.23.8, EC 3.4.23.9, EC 3.4.23.10, EC 3.4.99.1, EC 3.4.99.15 and EC 3.4.99.25 all created 1972 and incorporated 1978], part incorporated 1992)]

#### EC 3.4.23.31

**Accepted name:** scytalidopepsin A

**Reaction:** Hydrolysis of proteins with specificity similar to that of pepsin A, but also cleaves Cys(SO<sub>3</sub>H)<sup>7</sup>—Gly and Leu<sup>17</sup>—Val in the B chain of insulin

**Other name(s):** Scytalidium aspartic proteinase A; *Scytalidium lignicolum* aspartic proteinase; *Scytalidium lignicolum* aspartic proteinase A-2; *Scytalidium lignicolum* aspartic proteinase A-I; *Scytalidium lignicolum* aspartic proteinase C; *Scytalidium lignicolum* carboxyl proteinase; *Scytalidium lignicolum* acid proteinase



**Comments:** Isolated from the imperfect fungus *Scytalidium lignicolum*. Not inhibited by pepstatin-Ac, methyl 2-diazoacetamidohexanoate or 1,2-epoxy-3-(*p*-nitrophenyl)propane. A related enzyme from the same organism, proteinase C, is also insensitive to these inhibitors and has  $M_r = 406,000$  [2136]

**References:** [2131, 2132, 2136]

[EC 3.4.23.31 created 1992]

#### EC 3.4.23.32

**Accepted name:** scytalidopepsin B

**Reaction:** Hydrolysis of proteins with broad specificity, cleaving Phe<sup>24</sup>-Phe, but not Leu<sup>15</sup>-Tyr and Phe<sup>25</sup>-Tyr in the B chain of insulin

**Other name(s):** Scytalidium aspartic proteinase B; *Ganoderma lucidum* carboxyl proteinase; *Ganoderma lucidum* aspartic proteinase; *Scytalidium lignicolum* aspartic proteinase B; SLB

**Comments:** A second endopeptidase from *Scytalidium lignicolum* (see scytalidopepsin A) that is insensitive to pepstatin and methyl 2-diazoacetamidohexanoate. 1,2-Epoxy-3-(*p*-nitrophenoxy)propane reacts with Glu<sup>53</sup>, which replaces one of the aspartic residues at the active centre. One of the smallest aspartic endopeptidases active as the monomer, with  $M_r$  22,000. Similarly inhibitor-resistant endopeptidases are found in the basidiomycetes *Lentinus edodes* [2902] and *Ganoderma lucidum* [2903], and in *Polyporus tulipiferae* [1486], a second endopeptidase distinct from polyporopepsin, but these are of typical aspartic endopeptidase size,  $M_r$  about 36,000. Type example of peptidase family G1.

**References:** [2902, 1780, 2903, 1486, 2990]

[EC 3.4.23.32 created 1992]

[3.4.23.33 Transferred entry. *xanthomonapepsin*. Now EC 3.4.21.101, *xanthomonalisin*]

[EC 3.4.23.33 created 1992, deleted 2001]

#### EC 3.4.23.34

**Accepted name:** cathepsin E

**Reaction:** Similar to cathepsin D, but slightly broader specificity

**Other name(s):** slow-moving proteinase; erythrocyte membrane aspartic proteinase; SMP; erythrocyte membrane aspartic proteinase; EMAP; non-pepsin proteinase; cathepsin D-like acid proteinase; cathepsin E-like acid proteinase; cathepsin D-type proteinase

**Comments:** Found in stomach, spleen, erythrocyte membrane; not lysosomal. Pro-cathepsin E is an 86 kDa disulfide-linked dimer; activation or reduction produces monomer. In peptidase family A1 (pepsin A family)

**References:** [1599, 3290, 1349, 108]

[EC 3.4.23.34 created 1992]

#### EC 3.4.23.35

**Accepted name:** barrierpepsin

**Reaction:** Selective cleavage of -Leu<sup>6</sup>-Lys- bond in the pheromone  $\alpha$ -mating factor

**Other name(s):** barrier proteinase; Bar proteinase

**Comments:** A secreted endopeptidase known from baker's yeast (*Saccharomyces cerevisiae*). In peptidase family A1 (pepsin A family)

**References:** [1766, 1765]

[EC 3.4.23.35 created 1993]

#### EC 3.4.23.36

**Accepted name:** signal peptidase II



**Reaction:** Release of signal peptides from bacterial membrane prolipoproteins including murein prolipoprotein. Hydrolyses -Xaa-Yaa-Zaa-(S,diacylglyceryl)Cys-, in which Xaa is hydrophobic (preferably Leu), and Yaa (Ala or Ser) and Zaa (Gly or Ala) have small, neutral sidechains

**Other name(s):** premurein-leader peptidase; prolipoprotein signal peptidase; leader peptidase II; premurein leader proteinase; leader peptidase II

**Comments:** An 18-kDa enzyme present in bacterial inner membranes. Inhibited by pepstatin and the antibiotic globomycin. Type example of peptidase family A8.

**References:** [596, 3332, 2519]

[EC 3.4.23.36 created 1984 as EC 3.4.99.35, transferred 1995 to EC 3.4.23.36]

[3.4.23.37 Transferred entry. *pseudomonapepsin*. Now EC 3.4.21.100, *pseudomonalisin*]

[EC 3.4.23.37 created 1995]

#### EC 3.4.23.38

**Accepted name:** plasmepsin I

**Reaction:** Hydrolysis of the -Phe<sup>33</sup>-Leu- bond in the  $\alpha$ -chain of hemoglobin, leading to denaturation of the molecule

**Other name(s):** aspartic hemoglobinase I; PFAPG; malaria aspartic hemoglobinase

**Comments:** Known from the malaria organism, *Plasmodium*. About 37 kDa. In peptidase family A1 (pepsin A family), closest to cathepsin D and renin in structure. Inhibited by pepstatin. Formerly included in EC 3.4.23.6

**References:** [944, 803, 933]

[EC 3.4.23.38 created 1995]

#### EC 3.4.23.39

**Accepted name:** plasmepsin II

**Reaction:** Hydrolysis of the bonds linking certain hydrophobic residues in hemoglobin or globin. Also cleaves the small molecule substrates such as Ala-Leu-Glu-Arg-Thr-Phe-Phe(NO<sub>2</sub>)-Ser-Phe-Pro-Thr [3]

**Other name(s):** aspartic hemoglobinase II; PFAPD

**Comments:** Known from the malaria organism, *Plasmodium*. About 37 kDa. In peptidase family A1 (pepsin A family), and is 73% identical in sequence to plasmepsin I. Inhibited by pepstatin. Formerly included in EC 3.4.23.6

**References:** [532, 933, 1141]

[EC 3.4.23.39 created 1995]

#### EC 3.4.23.40

**Accepted name:** phytepsin

**Reaction:** Prefers hydrophobic residues Phe, Val, Ile, Leu, and Ala at P1 and P1', but also cleaves -Phe-Asp- and -Asp-Asp- bonds in 2S albumin from plant seeds

**Comments:** Known particularly from barley grain, but present in other plants also. In peptidase family A1 (pepsin A family), but structurally distinct in containing an internal region of about 100 amino acids not generally present in the family

**References:** [2478, 1423, 80, 1424]

[EC 3.4.23.40 created 1997]

#### EC 3.4.23.41

**Accepted name:** yapsin 1

**Reaction:** Hydrolyses various precursor proteins with Arg or Lys in P1, and commonly Arg or Lys also in P2. The P3 amino acid is usually non-polar, but otherwise additional basic amino acids are favourable in both non-prime and prime positions

**Other name(s):** yeast aspartic protease 3; Yap3 gene product (*Saccharomyces cerevisiae*)

**Comments:** In peptidase family A1 of pepsin, and weakly inhibited by pepstatin. Can partially substitute for kexin in a deficient strain of yeast. The homologous product of the Mkc7 gene (*Saccharomyces cerevisiae*) has similar catalytic activity and has been termed yapsin 2 [861]

**References:** [382, 861, 2194]

[EC 3.4.23.41 created 2000]

#### EC 3.4.23.42

**Accepted name:** thermopsin

**Reaction:** Similar in specificity to pepsin A preferring bulky hydrophobic amino acids in P1 and P1'

**Comments:** From the thermophilic archaeon *Sulfolobus acidocaldarius*. Maximally active at pH 2 and 90 °C. Weakly inhibited by pepstatin but shows no sequence similarity to pepsin. Type example of peptidase family A5.

**References:** [1695]

[EC 3.4.23.42 created 1992 as EC 3.4.99.43, transferred 2000 to EC 3.4.23.42]

#### EC 3.4.23.43

**Accepted name:** prepilin peptidase

**Reaction:** Typically cleaves a -Gly-Phe- bond to release an N-terminal, basic peptide of 5-8 residues from type IV prepilin, and then N-methylates the new N-terminal amino group, the methyl donor being S-adenosyl-L-methionine

**Comments:** Many species of bacteria carry pili on their cell surfaces. These are virulence determinants in pathogenic strains, and are assembled biosynthetically from type IV prepilin subunits. Before assembly, the prepilin molecules require proteolytic processing, which is done by the prepilin peptidase. Prepilin peptidase and its homologues play a central role not only in type IV pilus biogenesis but also in transport of macromolecules across cell membranes. Although both peptide-bond hydrolysis and N-methylation are catalysed by the same molecule, the methylation can be inhibited without affecting peptidase activity, and it is believed that the enzyme has two separate catalytic sites. Type example of peptidase family A24.

**References:** [1737, 1598]

[EC 3.4.23.43 created 2001]

#### EC 3.4.23.44

**Accepted name:** nodavirus endopeptidase

**Reaction:** Hydrolysis of an asparaginyl bond involved in the maturation of the structural protein of the virus, typically -Asn-Ala- or -Asn-Phe-

**Other name(s):** Black Beetle virus endopeptidase; Flock House virus endopeptidase

**Comments:** A single aspartic residue is critical for activity, and inhibition by EDTA indicates that a metal ion is also important. The enzyme is known from several nodaviruses that are pathogens of insects. Type example of peptidase family A6, and structurally related to the tetravirus endopeptidase in family A21, although in that family, the catalytic residue is thought to be Glu.

**References:** [3345, 1326]

[EC 3.4.23.44 created 2001]

#### EC 3.4.23.45

**Accepted name:** memapsin 1

**Reaction:** Broad endopeptidase specificity. Cleaves Glu-Val-Asn-Leu—Asp-Ala-Glu-Phe in the Swedish variant of Alzheimer's amyloid precursor protein

**Other name(s):**  $\beta$ -secretase;  $\beta$ -site Alzheimer's amyloid precursor protein cleaving enzyme 2 (BACE2); ASP1; Down region aspartic protease

**Comments:** Can cleave  $\beta$ -amyloid precursor protein to form the amyloidogenic  $\beta$ -peptide that is implicated in the pathology of Alzheimer's disease, but is not significantly expressed in human brain. In peptidase family A1, but is atypical in containing a C-terminal membrane-spanning domain.

**References:** [3001]

[EC 3.4.23.45 created 2003]

#### EC 3.4.23.46

**Accepted name:** memapsin 2

**Reaction:** Broad endopeptidase specificity. Cleaves Glu-Val-Asn-Leu—Asp-Ala-Glu-Phe in the Swedish variant of Alzheimer's amyloid precursor protein

**Other name(s):**  $\beta$ -secretase;  $\beta$ -site Alzheimer's amyloid precursor protein cleaving enzyme 1 (BACE1)

**Comments:** Suggested to be the major " $\beta$ -secretase" responsible for the cleavage of the  $\beta$ -amyloid precursor protein to form the amyloidogenic  $\beta$ -peptide that is implicated in the pathology of Alzheimer's disease. In peptidase family A1 but is atypical in containing a C-terminal membrane-spanning domain.

**References:** [3002, 1168]

[EC 3.4.23.46 created 2003]

#### EC 3.4.23.47

**Accepted name:** HIV-2 retropepsin

**Reaction:** Endopeptidase for which the P1 residue is preferably hydrophobic

**Comments:** In peptidase family A2. Responsible for the post-translational processing of the human immunodeficiency virus polyprotein.

**References:** [2957, 419]

[EC 3.4.23.47 created 2003]

#### EC 3.4.23.48

**Accepted name:** plasminogen activator Pla

**Reaction:** Converts human Glu-plasminogen to plasmin by cleaving the Arg<sup>560</sup>—Val peptide bond that is also hydrolysed by the mammalian u-plasminogen activator and t-plasminogen activator. Also cleaves arginyl bonds in other proteins

**Comments:** In peptidase family A26. From the bacterium *Yersinia pestis* that causes plague.

**References:** [1544]

[EC 3.4.23.48 created 2003]

#### EC 3.4.23.49

**Accepted name:** omptin

**Reaction:** Has a virtual requirement for Arg in the P1 position and a slightly less stringent preference for this residue in the P1' position, which can also contain Lys, Gly or Val.

**Other name(s):** protease VII; protease A; gene *ompT* proteins; *ompT* protease; protein a; Pla; OmpT

**Comments:** A product of the *ompT* gene of *Escherichia coli*, and associated with the outer membrane. Omptin shows a preference for cleavage between consecutive basic amino acids, but is capable of cleavage when P1' is a non-basic residue [3049, 1862]. Belongs in peptidase family A26.

**References:** [989, 2794, 1045, 571, 3049, 1528, 1862]

[EC 3.4.23.49 created 1993 as EC 3.4.21.87, transferred 2006 to EC 3.4.23.49]

#### EC 3.4.23.50

**Accepted name:** human endogenous retrovirus K endopeptidase  
**Reaction:** Processing at the authentic HIV-1 PR recognition site and release of the mature p17 matrix and the p24 capsid protein, as a result of the cleavage of the -SQNY-|PIVQ- cleavage site.  
**Other name(s):** human endogenous retrovirus K10 endopeptidase; endogenous retrovirus HERV-K10 putative protease; human endogenous retrovirus K retropepsin; HERV K10 endopeptidase; HERV K10 retropepsin; HERV-K PR; HERV-K protease; HERV-K113 protease; human endogenous retrovirus K113 protease; human retrovirus K10 retropepsin  
**Comments:** In peptidase family A2.  
**References:** [2953]

[EC 3.4.23.50 created 2009]

#### EC 3.4.23.51

**Accepted name:** HycI peptidase  
**Reaction:** This enzyme specifically removes a 32-amino acid peptide from the C-terminus of the precursor of the large subunit of hydrogenase 3 in *Escherichia coli* by cleavage at the C-terminal side of Arg<sup>537</sup>.  
**Other name(s):** HycI; HycE processing protein  
**Comments:** The reaction requires nickel to be bound to the precursor of the large subunit of hydrogenase 3. The endopeptidase uses the metal in the large subunit of [NiFe]-hydrogenases as a recognition motif [2907]. In peptidase family A31.  
**References:** [2907, 3257]

[EC 3.4.23.51 created 2009]

#### EC 3.4.23.52

**Accepted name:** preflagellin peptidase  
**Reaction:** Cleaves the signal peptide of 3 to 12 amino acids from the N-terminal of preflagellin, usually at Arg-Gly-| or Lys-Gly-|, to release flagellin.  
**Other name(s):** FlaK  
**Comments:** An aspartic peptidase from Archaea but not bacteria. In peptidase family A24 (type IV prepilin peptidase family).  
**References:** [139, 2077, 1198]

[EC 3.4.23.52 created 2011]

### EC 3.4.24 Metalloendopeptidases

#### EC 3.4.24.1

**Accepted name:** atrolysin A  
**Reaction:** Cleavage of Asn<sup>3</sup>-|Gln, His<sup>5</sup>-|Leu, His<sup>10</sup>-|Leu, Ala<sup>14</sup>-|Leu and Tyr<sup>16</sup>-|Leu in insulin B chain; removes C-terminal Leu from small peptides  
**Other name(s):** *Crotalus atrox* metalloendopeptidase a; hemorrhagic toxin a; *Crotalus atrox*  $\alpha$ -proteinase; *Crotalus atrox* proteinase; bothropasin  
**Comments:** A hemorrhagic endopeptidase of 68 kDa, one of six hemorrhagic toxins in the venom of western diamondback rattlesnake. The 60 kDa hemorrhagic toxin 1 of *Crotalus ruber ruber* shows identical specificity [1966]. In peptidase family M12 (astacin family). Related metalloendopeptidases from rattlesnake venoms are EC 3.4.24.41 (atrolysin B), EC 3.4.24.42 (atrolysin C), EC 3.4.24.43 (atroxase), EC 3.4.24.44 (atrolysin E), EC 3.4.24.45 (atrolysin F), EC 3.4.24.46 (adamalysin), EC 3.4.24.47 (horilysin), and EC 3.4.24.48 (ruberlysin)  
**References:** [224, 1966, 223, 222]

[EC 3.4.24.1 created 1972, modified 1986]

[3.4.24.2 Deleted entry. *Sepia* proteinase]

[EC 3.4.24.2 created 1972, deleted 1992]

EC 3.4.24.3

**Accepted name:** microbial collagenase

**Reaction:** Digestion of native collagen in the triple helical region at  $\text{---Gly}$  bonds. With synthetic peptides, a preference is shown for Gly at P3 and P1', Pro and Ala at P2 and P2', and hydroxyproline, Ala or Arg at P3'

**Other name(s):** *Clostridium histolyticum* collagenase; clostridiopeptidase A; collagenase A; collagenase I; *Achromobacter iophagus* collagenase; collagenase; aspergillopeptidase C; nucleolysin; azocollase; metallocollagenase; soycollagestin; *Clostridium histolyticum* proteinase A; clostridiopeptidase II; MMP-8; clostridiopeptidase I; collagen peptidase; collagen protease; collagenase MMP-1; metalloproteinase-1; kollaza; matrix metalloproteinase-1; MMP-1; matrix metalloproteinase-8; matrix metalloproteinase-18; interstitial collagenase

**Comments:** Six species of metalloendopeptidase acting on native collagen can be isolated from the medium of *Clostridium histolyticum*. Class I has forms  $\alpha$  (68 kDa),  $\beta$  (115 kDa) and  $\gamma$  (79 kDa); class II has  $\delta$  (100 kDa),  $\epsilon$  (110 kDa) and  $\zeta$  (125 kDa). The two classes are immunologically crossreactive, but have significantly different sequences, and different specificities such that their actions on collagen are complementary. The enzymes also act as peptidyl-tripeptidases. Variants of the enzyme have been purified from *Bacillus cereus* [1783], *Empedobacter collagenolyticum* [1582], *Pseudomonas marinitolus* [1041], and species of *Vibrio*, *Vibrio* B-30 (ATCC 21250) [1890] and *V. alginolyticus* (previously *Achromobacter iophagus*) [1101, 2944]. Also known from *Streptomyces* sp. [702]. The *Vibrio* enzyme is the type example of peptidase family M9.

**References:** [1041, 1890, 1101, 1582, 259, 260, 3132, 2944, 702, 1783]

[EC 3.4.24.3 created 1961 as EC 3.4.4.19, transferred 1972 to EC 3.4.24.3 (EC 3.4.24.8 created 1978, incorporated 1992, EC 3.4.99.5 created 1972, incorporated 1978)]

[3.4.24.4 Transferred entry. now EC 3.4.24.40 serralysin]

[EC 3.4.24.4 created 1972 [EC 3.4.99.13 and EC 3.4.99.22 both created 1972, incorporated 1978], deleted 1992]

[3.4.24.5 Deleted entry. lens neutral proteinase. Now included with EC 3.4.22.53 (calpain-2) and EC 3.4.25.1 (proteasome endopeptidase complex)]

[EC 3.4.24.5 created 1978, deleted 1989]

EC 3.4.24.6

**Accepted name:** leucolysin

**Reaction:** Cleavage of  $\text{Phe}^1\text{---Val}$ ,  $\text{His}^5\text{---Leu}$ ,  $\text{Ala}^{14}\text{---Leu}$ ,  $\text{Gly}^{20}\text{---Glu}$ ,  $\text{Gly}^{23}\text{---Phe}$  and  $\text{Phe}^{24}\text{---Phe}$  bonds in insulin B chain as well as N-blocked dipeptides

**Other name(s):** *Leucostoma* neutral proteinase; *Leucostoma* peptidase A

**Comments:** From the venom of the western cottonmouth moccasin snake (*Agkistrodon piscivorus leucostoma*).

**References:** [3082, 2740]

[EC 3.4.24.6 created 1978]

EC 3.4.24.7

**Accepted name:** interstitial collagenase

**Reaction:** Cleavage of the triple helix of collagen at about three-quarters of the length of the molecule from the N-terminus, at  $\text{Gly}^{775}\text{---Ile}$  in the  $\alpha 1(\text{I})$  chain. Cleaves synthetic substrates and  $\alpha$ -macroglobulins at bonds where P1' is a hydrophobic residue

**Other name(s):** vertebrate collagenase; matrix metalloproteinase 1

**Comments:** The enzyme takes its name from substrates of the interstitial collagen group - types I, II and III, all of which are cleaved in the helical domain. However,  $\alpha$ -macroglobulins are cleaved much more rapidly. The enzyme is widely distributed in vertebrate animals. Type example of peptidase family M10

**References:** [945, 217, 766, 2729]

[EC 3.4.24.7 created 1978]

[3.4.24.8 Transferred entry. *Achromobacter iophagus collagenase*. Now EC 3.4.24.3, microbial collagenase]

[EC 3.4.24.8 created 1978, deleted 1992]

[3.4.24.9 Deleted entry. *Trichophyton schoenleinii collagenase*]

[EC 3.4.24.9 created 1978, deleted 1992]

[3.4.24.10 Deleted entry. *Trichophyton mentagrophytes keratinase*]

[EC 3.4.24.10 created 1972 as EC 3.4.99.12, transferred 1978 to EC 3.4.24.10, deleted 1992]

#### EC 3.4.24.11

**Accepted name:** neprilysin

**Reaction:** Preferential cleavage of polypeptides between hydrophobic residues, particularly with Phe or Tyr at P1'

**Other name(s):** neutral endopeptidase; endopeptidase 24.11; kidney-brush-border neutral peptidase; enkephalinase (misleading); endopeptidase-2; CALLA (common acute lymphoblastic leukemia-associated) antigens; CALLA antigen; endopeptidase; membrane metalloendopeptidase; kidney-brush-border neutral endopeptidase; kidney-brush-border neutral proteinase; endopeptidase-2; CALLA glycoprotein; CALLA; common acute lymphoblastic leukemia antigen; CALLA glycoproteins; common acute lymphoblastic leukemia-associated antigens; neutral metalloendopeptidase; membrane metalloendopeptidase; NEP; neutral endopeptidase 24.11; CD10; neutral endopeptidase; acute lymphoblastic leukemia antigen

**Comments:** A membrane-bound glycoprotein widely distributed in animal tissues. Inhibited by phosphoramidon and thiorphan. Common acute lymphoblastic leukemia antigen (CALLA). Type example of peptidase family M13

**References:** [1836, 1787, 1652, 711]

[EC 3.4.24.11 created 1978, modified 1989]

#### EC 3.4.24.12

**Accepted name:** envelysin

**Reaction:** Hydrolysis of proteins of the fertilization envelope and dimethylcasein

**Other name(s):** sea-urchin-hatching proteinase; hatching enzyme; chorionase; chorion-digesting proteinase; chymotrypsin; sea urchin embryo hatching enzyme

**Comments:** A glycoprotein from various members of the class *Echinoidea*. Extracellular enzyme requiring Ca<sup>2+</sup>. In peptidase family M10 (interstitial collagenase family)

**References:** [160, 1649, 1650, 2108]

[EC 3.4.24.12 created 1978]

#### EC 3.4.24.13

**Accepted name:** IgA-specific metalloendopeptidase

**Reaction:** Cleavage of Pro—Thr bond in the hinge region of the heavy chain of human IgA

**Other name(s):** immunoglobulin A<sub>1</sub> proteinase; IgA protease; IgA1-specific proteinase; IgA1 protease; IgA<sub>1</sub> proteinase

**Comments:** A 190 kDa enzyme found in several pathogenic species of *Streptococcus* such as *sanguis* and *pneumoniae*. Type example of peptidase family M26. There is also an IgA-specific prolyl endopeptidase of the serine-type (see EC 3.4.21.72, IgA-specific serine endopeptidase)

**References:** [1515, 920, 919]

[EC 3.4.24.13 created 1984]

**EC 3.4.24.14**

**Accepted name:** procollagen *N*-endopeptidase

**Reaction:** Cleaves the *N*-propeptide of collagen chain  $\alpha 1(\text{I})$  at Pro—Gln and of  $\alpha 1(\text{II})$  and  $\alpha 2(\text{I})$  at Ala—Gln

**Other name(s):** procollagen *N*-terminal peptidase; procollagen aminopeptidase; aminoprocollagen peptidase; aminoterminal procollagen peptidase; procollagen aminoterminal protease; procollagen *N*-terminal proteinase; type I/II procollagen *N*-proteinase; type III procollagen

**Comments:** Removes the propeptides of type I and II collagens prior to fibril assembly. Does not act on type III collagen. In peptidase family M12 (astacin family)

**References:** [1496, 1160]

[EC 3.4.24.14 created 1984]

**EC 3.4.24.15**

**Accepted name:** thimet oligopeptidase

**Reaction:** Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5-15 residues

**Other name(s):** Pz-peptidase; soluble metalloendopeptidase; endo-oligopeptidase A; tissue-endopeptidase degrading collagenase-synthetic-substrate

**Comments:** Thiol compounds activate at low concentrations. Type example of peptidase family M3.

**References:** [452, 2200, 157, 2279, 2922]

[EC 3.4.24.15 created 1984 (EC 3.4.22.19 created 1989 and EC 3.4.99.31 created 1978 both incorporated 1992)]

**EC 3.4.24.16**

**Accepted name:** neurolysin

**Reaction:** Preferential cleavage in neurotensin: Pro<sup>10</sup>—Tyr

**Other name(s):** neurotensin endopeptidase; endopeptidase 24.16; endo-oligopeptidase B (proline-endopeptidase)

**Comments:** No absolute requirement for a prolyl bond: the enzyme acts on some peptides, such as dynorphin 1-8, that do not contain proline, and does not act on some others that do. In peptidase family M3 (thimet oligopeptidase family)

**References:** [408, 140, 407]

[EC 3.4.24.16 created 1989]

**EC 3.4.24.17**

**Accepted name:** stromelysin 1

**Reaction:** Preferential cleavage where P1', P2' and P3' are hydrophobic residues

**Other name(s):** matrix metalloproteinase 3; proteoglycanase; collagenase activating protein; procollagenase activator; transin; MMP-3; neutral proteoglycanase; stromelysin; collagen-activating protein

**Comments:** An extracellular endopeptidase of vertebrate tissues homologous with interstitial collagenase. Digests proteoglycan, fibronectin, collagen types III, IV, V, IX, and activates procollagenase. In peptidase family M10 (interstitial collagenase family)

**References:** [433, 2181, 623, 700]

[EC 3.4.24.17 created 1990]

**EC 3.4.24.18**

**Accepted name:** meprin A

**Reaction:** Hydrolysis of protein and peptide substrates preferentially on carboxyl side of hydrophobic residues

**Other name(s):** endopeptidase-2; meprin-a; meprin; *N*-benzoyl-L-tyrosyl-*p*-aminobenzoic acid hydrolase; PABA-peptide hydrolase; PPH

**Comments:** A membrane-bound metalloendopeptidase of rat and mouse kidney and intestinal brush borders, and salivary ducts. Differences from neprilysin (EC 3.4.24.11 (astacin family)). Formerly included in EC 3.4.24.11

**References:** [211, 337, 2763, 2764, 147]

[EC 3.4.24.18 created 1992]

#### EC 3.4.24.19

**Accepted name:** procollagen C-endopeptidase

**Reaction:** Cleavage of the C-terminal propeptide at Ala—Asp in type I and II procollagens and at Arg—Asp in type III

**Other name(s):** procollagen C-terminal proteinase; carboxyprocollagen peptidase; procollagen C-terminal peptidase; procollagen C-proteinase; procollagen C-terminal proteinase; procollagen carboxypeptidase; procollagen carboxy-terminal proteinase; procollagen peptidase

**Comments:** A 100 kDa endopeptidase the activity of which is increased by Ca<sup>2+</sup> and by an enhancer glycoprotein. In peptidase family M12 (astacin family)

**References:** [1161, 1426]

[EC 3.4.24.19 created 1992]

#### EC 3.4.24.20

**Accepted name:** peptidyl-Lys metalloendopeptidase

**Reaction:** Preferential cleavage in proteins: -Xaa—Lys- (in which Xaa may be Pro)

**Other name(s):** *Armillaria mellea* neutral proteinase; peptidyllysine metalloproteinase

**Comments:** From the honey fungus *Armillaria mellea*. In peptidase family M35 (deuterolysin family).

**References:** [2483, 1661]

[EC 3.4.24.20 created 1978 as EC 3.4.99.32, transferred 1992 to EC 3.4.24.20 (EC 3.4.99.30 created 1978, incorporated 1992)]

#### EC 3.4.24.21

**Accepted name:** astacin

**Reaction:** Hydrolysis of peptide bonds in substrates containing five or more amino acids, preferentially with Ala in P1', and Pro in P2'

**Other name(s):** *Astacus* proteinase; crayfish small-molecule proteinase

**Comments:** A 22.6 kDa digestive endopeptidase from the cardia of the crayfish *Astacus fluviatilis*. Type example of peptidase family M12.

**References:** [1529, 2924, 2776, 2775]

[EC 3.4.24.21 created 1972 as EC 3.4.99.6, transferred 1992 to EC 3.4.24.21]

#### EC 3.4.24.22

**Accepted name:** stromelysin 2

**Reaction:** Similar to stromelysin 1, but action on collagen types III, IV and V is weak

**Other name(s):** matrix metalloproteinase 10; transin 2; proteoglycanase 2

**Comments:** In peptidase family M10 (interstitial collagenase family). Digests gelatin types I, III, IV, V, fibronectin and proteoglycan

**References:** [282, 1994, 2080]

[EC 3.4.24.22 created 1992]

#### EC 3.4.24.23

**Accepted name:** matrilysin



**Reaction:** Cleavage of Ala<sup>14</sup>—Leu and Tyr<sup>16</sup>—Leu in B chain of insulin. No action on collagen types I, II, IV, V. Cleaves gelatin chain  $\alpha 2(I) \zeta \alpha 1(I)$

**Other name(s):** matrin; uterine metalloendopeptidase; matrix metalloproteinase 7; putative (or punctuated) metalloproteinase-1; matrix metalloproteinase pump 1; MMP 7; PUMP-1 proteinase; PUMP; metalloproteinase pump-1; putative metalloproteinase; MMP

**Comments:** Found in rat uterus; at 19 kDa, the smallest member of peptidase family M10 (interstitial collagenase family). Similar in specificity to stromelysin, but more active on azocoll

**References:** [1994, 3205, 2341, 1938]

[EC 3.4.24.23 created 1992]

#### EC 3.4.24.24

**Accepted name:** gelatinase A

**Reaction:** Cleavage of gelatin type I and collagen types IV, V, VII, X. Cleaves the collagen-like sequence Pro-Gln-Gly—Ile-Ala-Gly-Gln

**Other name(s):** 72-kDa gelatinase; matrix metalloproteinase 2; type IV collagenase (ambiguous); 3/4 collagenase (obsolete); matrix metalloproteinase 5 (obsolete); 72 kDa gelatinase type A; collagenase IV (ambiguous); collagenase type IV (ambiguous); MMP 2; type IV collagen metalloproteinase (ambiguous); type IV collagenase/gelatinase (ambiguous)

**Comments:** A secreted endopeptidase in peptidase family M10 (interstitial collagenase family), but possessing an additional fibronectin-like domain

**References:** [2009, 468, 2182]

[EC 3.4.24.24 created 1992]

#### EC 3.4.24.25

**Accepted name:** vibriolysin

**Reaction:** Preferential cleavage of bonds with bulky hydrophobic groups in P2 and P1'. Phe at P1' is the most favoured residue, which distinguished this enzyme from thermolysin

**Other name(s):** *Aeromonas proteolytica* neutral proteinase; aeromonolysin

**Comments:** Thermostable enzyme from *Vibrio proteolyticus* (formerly *Aeromonas proteolytica*). Specificity related to, but distinct from, those of thermolysin and bacillolysin [1165]. A zinc metallopeptidase in family M4 (thermolysin family). Formerly included in EC 3.4.24.4

**References:** [1165, 3189, 177, 3188, 544]

[EC 3.4.24.25 created 1972 as EC 3.4.24.4, part transferred 1992 to EC 3.4.24.25, modified 1997]

#### EC 3.4.24.26

**Accepted name:** pseudolysin

**Reaction:** Hydrolysis of proteins including elastin, collagen types III and IV, fibronectin and immunoglobulin A, generally with bulky hydrophobic group at P1'. Insulin B chain cleavage pattern identical to that of thermolysin, but specificity differs in other respects

**Other name(s):** *Pseudomonas* elastase; *Pseudomonas aeruginosa* neutral metalloproteinase

**Comments:** In peptidase family M4 (thermolysin family). From the pathogenic bacteria *Pseudomonas aeruginosa* and *Legionella pneumophila*, and causes tissue damage.

**References:** [1971, 2094, 652, 210, 231]

[EC 3.4.24.26 created 1972 as EC 3.4.24.4, part transferred 1992 to EC 3.4.24.26]

#### EC 3.4.24.27

**Accepted name:** thermolysin

**Reaction:** Preferential cleavage: —Leu  $\zeta$  —Phe

**Other name(s):** *Bacillus thermoproteolyticus* neutral proteinase; thermoase; thermoase Y10; TLN

**Comments:** A thermostable extracellular metalloendopeptidase containing four calcium ions. Enzymes that may be species variants of thermolysin are reported from *Micrococcus caseolyticus* [588] and *Aspergillus oryzae* [1970]. Type example of peptidase family M4. Closely related but distinct enzymes are aeromonolysin, pseudolysin, bacillolysin, aureolysin and mycolysin

**References:** [2173, 1973, 1610, 588, 1970, 2923, 1846]

[EC 3.4.24.27 created 1972 as EC 3.4.24.4, part transferred 1992 to EC 3.4.24.27]

#### EC 3.4.24.28

**Accepted name:** bacillolysin

**Reaction:** Similar, but not identical, to that of thermolysin

**Other name(s):** *Bacillus* metalloendopeptidase; *Bacillus subtilis* neutral proteinase; anilozyme P 10; *Bacillus* metalloproteinase; *Bacillus* neutral proteinase; megateriopeptidase

**Comments:** Variants of this enzyme have been found in species of *Bacillus* including *B. subtilis* [1973, 3259], *B. amyloliquefaciens* [3054], *B. megaterium* (megateriopeptidase, [1917]), *B. mesentericus* [2777], *B. cereus* [3,8,9] and *B. stearothermophilus* [2833]. In peptidase family M4 (thermolysin family). Formerly included in EC 3.4.24.4

**References:** [1973, 1917, 749, 1165, 3054, 3259, 2833, 2653, 2249, 2777]

[EC 3.4.24.28 created 1972 as EC 3.4.24.4, part transferred 1992 to EC 3.4.24.28]

#### EC 3.4.24.29

**Accepted name:** aureolysin

**Reaction:** Cleavage of insulin B chain with specificity similar to that of thermolysin, preferring hydrophobic P1' residue. Activates the glutamyl endopeptidase (EC 3.4.21.19) of *Staphylococcus aureus*

**Other name(s):** *Staphylococcus aureus* neutral proteinase; *Staphylococcus aureus* neutral protease

**Comments:** A metalloenzyme from *S. aureus* earlier confused with staphylokinase (a non-enzymic activator of plasminogen).

**References:** [79, 2494, 647, 2315]

[EC 3.4.24.29 created 1972 as EC 3.4.24.4, part transferred 1992 to EC 3.4.24.29]

#### EC 3.4.24.30

**Accepted name:** coccolysin

**Reaction:** Preferential cleavage:  $\text{+Leu, +Phe, +Tyr, +Ala}$

**Other name(s):** *Streptococcus thermophilus* intracellular proteinase; EM 19000

**Comments:** A 30 kDa endopeptidase found intracellularly in *S. thermophilus* [589] and *S. diacetilactis* [590] and in the medium of *S. faecalis* [2699, 1784]. In peptidase family M4 (thermolysin family). Formerly included in EC 3.4.24.4

**References:** [589, 590, 2699, 1784]

[EC 3.4.24.30 created 1972 as EC 3.4.24.4, part transferred 1992 to EC 3.4.24.30]

#### EC 3.4.24.31

**Accepted name:** mycolysin

**Reaction:** Preferential cleavage of bonds with hydrophobic residues in P1'

**Other name(s):** pronase component; *Streptomyces griseus* neutral proteinase; actinase E; SGNPI

**Comments:** From *Streptomyces griseus*, *S. naraensis*, and *S. cacaoi*. Specificity similar to that of thermolysin, but much more sensitive to inhibition by mercaptoacetyl-Phe-Leu. Little structural similarity to other bacterial metalloendopeptidases. Type example of peptidase family M5. Formerly included in EC 3.4.24.4

**References:** [1973, 1146, 246, 395]

[EC 3.4.24.31 created 1972 as EC 3.4.24.4, part transferred 1992 to EC 3.4.24.31]

#### EC 3.4.24.32

**Accepted name:**  $\beta$ -lytic metalloendopeptidase  
**Reaction:** Cleavage of *N*-acetylmuramoyl—Ala, and of the insulin B chain at Gly<sup>23</sup>—Phe<sub>i</sub> Val<sup>18</sup>—Cys  
**Other name(s):** *Myxobacter*  $\beta$ -lytic proteinase; achromopeptidase component;  $\beta$ -lytic metalloproteinase;  $\beta$ -lytic protease; *Myxobacterium sorangium*  $\beta$ -lytic proteinase; *Myxobacter*495  $\beta$ -lytic proteinase  
**Comments:** From *Achromobacter lyticus* and *Lysobacter enzymogenes*. Digests bacterial cell walls. Type example of peptidase family M23.  
**References:** [3171, 3170, 1666]

[EC 3.4.24.32 created 1972 as EC 3.4.24.4, part transferred 1992 to EC 3.4.24.32]

#### EC 3.4.24.33

**Accepted name:** peptidyl-Asp metalloendopeptidase  
**Reaction:** Cleavage of Xaa—Asp, Xaa—Glu and Xaa—cysteic acid bonds  
**Other name(s):** endoproteinase Asp-N; peptidyl-Asp metalloproteinase  
**Comments:** A metalloenzyme isolated from *Pseudomonas fragi*. Useful in protein sequencing applications because of its limited specificity. In peptidase family M72.  
**References:** [2313, 649, 1250]

[EC 3.4.24.33 created 1992]

#### EC 3.4.24.34

**Accepted name:** neutrophil collagenase  
**Reaction:** Cleavage of interstitial collagens in the triple helical domain. Unlike EC 3.4.24.7, interstitial collagenase, this enzyme cleaves type III collagen more slowly than type I  
**Other name(s):** matrix metalloproteinase 8; PMNL collagenase; MMP-8  
**Comments:** Similar to interstitial collagenase in specificity, but the product of a different gene and highly glycosylated. Stored in the specific granules of neutrophil leukocytes. In peptidase family M10 (interstitial collagenase family). Formerly included in EC 3.4.24.7  
**References:** [1072, 1073, 1480]

[EC 3.4.24.34 created 1992]

#### EC 3.4.24.35

**Accepted name:** gelatinase B  
**Reaction:** Cleavage of gelatin types I and V and collagen types IV and V  
**Other name(s):** 92-kDa gelatinase; matrix metalloproteinase 9; type V collagenase; 92-kDa type IV collagenase; macrophage gelatinase; 95 kDa type IV collagenase/gelatinase; collagenase IV (ambiguous); collagenase type IV (ambiguous); gelatinase MMP 9; MMP 9; type IV collagen metalloproteinase (ambiguous)  
**Comments:** Similar to gelatinase A, but possesses a further domain. In peptidase family M10 (interstitial collagenase family)  
**References:** [1136, 3185, 1778]

[EC 3.4.24.35 created 1992]

#### EC 3.4.24.36

**Accepted name:** leishmanolysin  
**Reaction:** Preference for hydrophobic residues at P1 and P1' and basic residues at P2' and P3'. A model non-peptide is cleaved at -Ala-Tyr—Leu-Lys-Lys-

**Other name(s):** promastigote surface endopeptidase; glycoprotein gp63; *Leishmania* metalloproteinase; surface acid proteinase; promastigote surface protease  
**Comments:** A membrane-bound glycoprotein found on the promastigote of various species of *Leishmania* protozoans. Contains consensus sequence for a zinc-binding site; Z-Tyr-Leu-NHOH is a strong inhibitor. The enzyme can activate its proenzyme by cleavage of the Val<sup>100</sup>—Val bond. An acid pH optimum is found with certain protein substrates. Type example of peptidase family M8  
**References:** [343, 271, 405, 272]

[EC 3.4.24.36 created 1992]

#### EC 3.4.24.37

**Accepted name:** saccharolysin  
**Reaction:** Cleavage of Pro—Phe and Ala—Ala bonds  
**Other name(s):** proteinase yscD; yeast cysteine proteinase D (Misleading); *Saccharomyces cerevisiae* proteinase yscD  
**Comments:** An 83 kDa cytoplasmic thiol-dependent metalloendopeptidase from *Saccharomyces cerevisiae*. In peptidase family M3 (thimet oligopeptidase family).  
**References:** [9, 875]

[EC 3.4.24.37 created 1989 as EC 3.4.22.22, transferred 1992 to EC 3.4.24.37]

#### EC 3.4.24.38

**Accepted name:** gametolysin  
**Reaction:** Cleavage of the proline- and hydroxyproline-rich proteins of the *Chlamydomonas* cell wall; also cleaves azocasein, gelatin and Leu-Trp-Met—Arg-Phe-Ala  
**Other name(s):** autolysin, *Chlamydomonas* cell wall degrading protease; lysin; *Chlamydomonas reinhardtii* metalloproteinase; gamete lytic enzyme; gamete autolysin  
**Comments:** A glycoprotein found in the periplasmic space of *Chlamydomonas reinhardtii* gametes in a 62 kDa inactive form; decreased to 60 kDa upon activation. A zinc enzyme, inhibited by phosphoramidon, but also thiol activated. Type example of peptidase family M11  
**References:** [1294, 323, 1837]

[EC 3.4.24.38 created 1992, modified 2000]

#### EC 3.4.24.39

**Accepted name:** deuterolysin  
**Reaction:** Cleavage of the proline- and hydroxyproline-rich proteins of the *Chlamydomonas* cell wall; also cleaves azocasein, gelatin and Leu-Trp-Met—Arg-Phe-Ala  
**Other name(s):** *Penicillium roqueforti* protease II; microbial neutral proteinase II; acid metalloproteinase; neutral proteinase II; *Penicillium roqueforti* metalloproteinase  
**Comments:** Proteolytic activity found in *Penicillium roqueforti* [984], *P. caseicolum* [984], *Aspergillus sojae* [2609] and *A. oryzae* [2033, 3026]. Optimum pH of 5 for digesting various proteins. Strong action on protamine and histones. Insensitive to phosphoramidon. About 20 kDa. A distinct *Aspergillus sojae* endopeptidase is larger and has a neutral pH optimum. Type example of peptidase family M35. Formerly included in EC 3.4.24.4  
**References:** [2033, 983, 2609, 984, 3026]

[EC 3.4.24.39 created 1972 as EC 3.4.24.4, part transferred 1992 to EC 3.4.24.39]

#### EC 3.4.24.40

**Accepted name:** serralysin  
**Reaction:** Preferential cleavage of bonds with hydrophobic residues in P1'  
**Other name(s):** *Pseudomonas aeruginosa* alkaline proteinase; *Escherichia freundii* proteinase; *Serratia marcescens* extracellular proteinase; *Serratia marcescens* metalloproteinase; *Pseudomonas aeruginosa* alk. protease; *Serratia marcescens* metalloprotease

**Comments:** A 50 kDa extracellular endopeptidase from *Pseudomonas aeruginosa* [1,2,6], *Escherichia freundii* [2038], *Serratia marcescens* [4,5,6] and *Erwinia chrysanthemi* [527]. There is broad specificity in cleavage of the insulin B chain, with some species variations. The pH optimum for digesting various proteins is about 9 - 10. In peptidase family M10 (interstitial collagenase family). Formerly included in EC 3.4.24.4

**References:** [1973, 1974, 2038, 565, 627, 2037, 527, 2189]

[EC 3.4.24.40 created 1972 as EC 3.4.24.4, part transferred 1992 to EC 3.4.24.40]

#### EC 3.4.24.41

**Accepted name:** atrolysin B

**Reaction:** Cleavage of His<sup>5</sup>—Leu, His<sup>10</sup>—Leu, Ala<sup>14</sup>—Leu, Tyr<sup>16</sup>—Leu and Gly<sup>23</sup>—Phe of insulin B chain; identical to the cleavage of insulin B chain by atrolysin C. Also cleaves —Ser bonds in glucagon

**Other name(s):** *Crotalus atrox* metalloendopeptidase b; hemorrhagic toxin b; Ht-b

**Comments:** From the venom of the western diamondback rattlesnake (*Crotalus atrox*). In peptidase family M12 (astacin family)

**References:** [224, 223]

[EC 3.4.24.41 created 1992]

#### EC 3.4.24.42

**Accepted name:** atrolysin C

**Reaction:** Cleavage of His<sup>5</sup>—Leu, His<sup>10</sup>—Leu, Ala<sup>14</sup>—Leu, Tyr<sup>16</sup>—Leu and Gly<sup>23</sup>—Phe bonds in B chain of insulin. With small molecule substrates prefers hydrophobic residue at P2' and small residue such as Ala, Gly at P1

**Other name(s):** *Crotalus atrox* metalloendopeptidase c; hemorrhagic toxin c and d

**Comments:** A 24 kDa hemorrhagic endopeptidase from the venom of the western diamondback rattlesnake (*Crotalus atrox*) that digests type IV collagen, and exists as two forms, c and d. Phosphoramidon inhibits in the 0.1 mM range. In peptidase family M12 (astacin family). Hemorrhagic toxin-2 of *C. ruber ruber* has the same  $M_r$  and specificity and is a homologue [1966, 2864].

**References:** [224, 799, 221, 1966, 2618, 2864]

[EC 3.4.24.42 created 1992]

#### EC 3.4.24.43

**Accepted name:** atroxase

**Reaction:** Cleavage of His<sup>5</sup>—Leu, Ser<sup>9</sup>—His, His<sup>10</sup>—Leu, Ala<sup>14</sup>—Leu and Tyr<sup>16</sup>—Leu of insulin B chain

**Comments:** A nonhemorrhagic endopeptidase from the venom of the western diamondback rattlesnake (*Crotalus atrox*) that cleaves fibrinogen. In peptidase family M12 (astacin family)

**References:** [3193]

[EC 3.4.24.43 created 1992]

#### EC 3.4.24.44

**Accepted name:** atrolysin E

**Reaction:** Cleavage of Asn<sup>3</sup>—Gln, Ser<sup>9</sup>—His and Ala<sup>14</sup>—Leu bonds in insulin B chain and Tyr<sup>14</sup>—Gln and Thr<sup>8</sup>—Ser in A chain. Cleaves type IV collagen at Ala<sup>73</sup>—Gln in  $\alpha$ 1(IV) and at Gly<sup>7</sup>—Leu in  $\alpha$ 2(IV)

**Other name(s):** *Crotalus atrox* metalloendopeptidase e; hemorrhagic toxin e

**Comments:** A 25.7 kDa hemorrhagic endopeptidase from the venom of the western diamondback rattlesnake (*Crotalus atrox*) that digests basement membrane components, including the triple helix of type IV collagen. Such action is believed to contribute to the hemorrhagic property by weakening capillary walls. In peptidase family M12 (astacin family)

**References:** [224, 220, 136]

[EC 3.4.24.44 created 1992]

#### EC 3.4.24.45

**Accepted name:** atrolysin F

**Reaction:** Cleavage of Val<sup>2</sup>-Asn, Gln<sup>4</sup>-His, Leu<sup>6</sup>-Cys, His<sup>10</sup>-Leu, Ala<sup>14</sup>-Leu and Tyr<sup>16</sup>-Leu bonds in insulin B chain

**Other name(s):** *Crotalus atrox* metalloendopeptidase; hemorrhagic toxin f; *Crotalus atrox* metalloendopeptidase f

**Comments:** A 64 kDa hemorrhagic endopeptidase from the venom of the western diamondback rattlesnake (*Crotalus atrox*) that digests the  $\gamma$  chain of fibrinogen. Immunologically distinct from EC 3.4.24.1, atrolysin A.

**References:** [2085]

[EC 3.4.24.45 created 1992]

#### EC 3.4.24.46

**Accepted name:** adamalysin

**Reaction:** Cleavage of Phe<sup>1</sup>-Val, His<sup>5</sup>-Leu, His<sup>10</sup>-Leu, Ala<sup>14</sup>-Leu, Leu<sup>15</sup>-Tyr, and Tyr<sup>16</sup>-Leu of insulin B chain

**Other name(s):** *Crotalus adamanteus* metalloendopeptidase; proteinase I and II; *Crotalus adamanteus* venom proteinase II; adamalysin II

**Comments:** From the venom of the eastern diamondback rattlesnake (*Crotalus adamanteus*). Two isoenzymes of approx. 24 kDa that inactivate  $\alpha_1$ -proteinase inhibitor by a single cleavage. In peptidase family M12 (astacin family)

**References:** [1565]

[EC 3.4.24.46 created 1992]

#### EC 3.4.24.47

**Accepted name:** horrilysin

**Reaction:** Cleavage of only the single bond Ala<sup>14</sup>-Leu in the insulin B chain, Ser<sup>12</sup>-Leu in the A chain, and Ile-Gly, Pro-Ala, and Ser-Trp in melittin

**Other name(s):** *Crotalus horridus* metalloendopeptidase; hemorrhagic proteinase IV; *Crotalus horridus horridus* venom hemorrhagic proteinase

**Comments:** A 56 kDa hemorrhagic endopeptidase from the venom of the timber rattlesnake (*Crotalus horridus horridus*) that cleaves basement membrane, hide powder and fibrinogen.

**References:** [455, 456]

[EC 3.4.24.47 created 1992]

#### EC 3.4.24.48

**Accepted name:** ruberlysin

**Reaction:** Cleavage of His<sup>10</sup>-Leu, Ala<sup>14</sup>-Leu, Tyr<sup>16</sup>-Leu and Gly<sup>23</sup>-Phe bonds in the B chain of insulin; His-Pro, Pro-Phe, and Trp-Ser of angiotensin I; and Gly-Phe of Met enkephalin

**Other name(s):** *Crotalus ruber* metalloendopeptidase II; hemorrhagic toxin II

**Comments:** A 25 kDa hemorrhagic endopeptidase from the venom of the red rattlesnake (*Crotalus ruber ruber*) that cleaves fibrinogen. In peptidase family M12 (astacin family)

**References:** [1966, 2864]

[EC 3.4.24.48 created 1992]

#### EC 3.4.24.49

**Accepted name:** bothropasin  
**Reaction:** Cleavage of His<sup>5</sup>—Leu, His<sup>10</sup>—Leu, Ala<sup>14</sup>—Leu, Tyr<sup>16</sup>—Leu and Phe<sup>24</sup>—Phe in insulin B chain  
**Other name(s):** *Bothrops jararaca* venom metalloproteinase  
**Comments:** Caseinolytic endopeptidase of jararaca snake (*Bothrops jararaca*) venom; 48 kDa. In peptidase family M12  
**References:** [1797]

[EC 3.4.24.49 created 1992]

#### EC 3.4.24.50

**Accepted name:** bothrolysin  
**Reaction:** Cleavage of Gln<sup>4</sup>—His, Ser<sup>9</sup>—His and Ala<sup>14</sup>—Leu of insulin B chain and Pro—Phe of angiotensin I  
**Other name(s):** Bothrops metalloendopeptidase J; J protease  
**Comments:** A 22.5 kDa endopeptidase from the venom of the jararaca snake (*Bothrops jararaca*), insensitive to phosphoramidon at 0.5 mM. In peptidase family M12 (astacin family)  
**References:** [2884]

[EC 3.4.24.50 created 1992]

#### EC 3.4.24.51

**Accepted name:** ophiolysin  
**Reaction:** Cleavage of Asn<sup>3</sup>—Gln, Gln<sup>4</sup>—His, His<sup>10</sup>—Leu, Ala<sup>14</sup>—Leu, and Tyr<sup>16</sup>—Leu in insulin B chain  
**Other name(s):** Ophiophagus metalloendopeptidase  
**Comments:** A 70 kDa endopeptidase from the venom of the king cobra (*Ophiophagus hannah*)  
**References:** [3246]

[EC 3.4.24.51 created 1992]

#### EC 3.4.24.52

**Accepted name:** trimereylisin I  
**Reaction:** Cleavage of only two bonds His<sup>10</sup>—Leu and Ala<sup>14</sup>—Leu in the insulin B chain  
**Other name(s):** Trimeresurus metalloendopeptidase I; hemorrhagic proteinase HR1A; hemorrhagic metalloproteinase HR1A; metalloproteinase HR1A  
**Comments:** A 60 kDa hemorrhagic endopeptidase of pI 4.4 from the venom of the habu snake (*Trimeresurus flavoviridis*). In peptidase family M12 (astacin family)  
**References:** [2197, 3247, 2863]

[EC 3.4.24.52 created 1992]

#### EC 3.4.24.53

**Accepted name:** trimereylisin II  
**Reaction:** Cleavage of Asn<sup>3</sup>—Gln, His<sup>10</sup>—Leu and Ala<sup>14</sup>—Leu in the insulin B chain, and the bond Z-Gly-Pro—Leu-Gly-Pro in a small molecule substrate of microbial collagenase  
**Other name(s):** Trimeresurus metalloendopeptidase II; proteinase H<sub>2</sub>; H<sub>2</sub>-proteinase  
**Comments:** A 24 kDa nonhemorrhagic endopeptidase from the venom of the habu snake (*Trimeresurus flavoviridis*). In peptidase family M12 (astacin family)  
**References:** [2845, 2861]

[EC 3.4.24.53 created 1992]



#### EC 3.4.24.54

- Accepted name:** mucrolysin  
**Reaction:** Cleavage of Ser<sup>9</sup>—His, His<sup>10</sup>—Leu, Ala<sup>14</sup>—Leu, Leu<sup>15</sup>—Tyr and Tyr<sup>16</sup>—Leu bonds in insulin B chain  
**Other name(s):** Trimeresurus metalloendopeptidase A; mucrotoxin A  
**Comments:** A 94 kDa hemorrhagic and fibrinogenolytic endopeptidase from the Chinese habu snake (*Trimeresurus mucrosquamatus*) venom. In peptidase family M12 (astacin family)  
**References:** [2793, 1467]

[EC 3.4.24.54 created 1992]

#### EC 3.4.24.55

- Accepted name:** pitrilysin  
**Reaction:** Preferential cleavage of -Tyr<sup>16</sup>—Leu- and -Phe<sup>25</sup>—Tyr-bonds of oxidized insulin B chain. Also acts on other substrates of less than 7 kDa such as insulin and glucagon  
**Other name(s):** *Escherichia coli* protease III; protease Pi; proteinase Pi; PTR; *Escherichia coli* metalloproteinase Pi  
**Comments:** From the periplasmic space of *Escherichia coli*. Inhibited by EDTA and 1,10-phenanthroline; not thiol-dependent. Type example of peptidase family M16  
**References:** [768, 20, 183, 618, 49]

[EC 3.4.24.55 created 1992 as EC 3.4.99.44, transferred 1993 to EC 3.4.24.55 (EC 3.4.99.45 created 1992, incorporated 1993)]

#### EC 3.4.24.56

- Accepted name:** insulysin  
**Reaction:** Degradation of insulin, glucagon and other polypeptides. No action on proteins  
**Other name(s):** insulinase; insulin-degrading enzyme; insulin protease; insulin proteinase; insulin-degrading neutral proteinase; insulin-specific protease; insulin-glucagon protease; metalloinsulinase; IDE  
**Comments:** A 110 kDa cytosolic enzyme, known from mammals and the fruit fly, *Drosophila melanogaster*. Inhibited by bacitracin, chelating agents EDTA and 1,10-phenanthroline, and by thiol-blocking reagents such as *N*-ethylmaleimide, but not by phosphoramidon. In peptidase family M16 (pitrilysin family).  
**References:** [660, 21, 661, 1562, 618]

[EC 3.4.24.56 created 1972 as EC 3.4.99.10, transferred 1976 EC 3.4.22.11, transferred 1978 to EC 3.4.99.45, transferred 1993 to to EC 3.4.24.56 (EC 3.4.99.46 created 1992, incorporated 2000)]

#### EC 3.4.24.57

- Accepted name:** *O*-sialoglycoprotein endopeptidase  
**Reaction:** Hydrolysis of *O*-sialoglycoproteins; cleaves -Arg<sup>31</sup>—Asp- bond in glycoporphin A. Does not cleave unglycosylated proteins, desialylated glycoproteins or glycoproteins that are only *N*-glycosylated  
**Other name(s):** glycoprotease; glycoporphin A proteinase; glycoproteinase; sialoglycoprotease; sialoglycoproteinase  
**Comments:** An enzyme secreted by the bacterium *Pasteurella haemolytica*. Inhibited by EDTA (100 mM) and 1,10-phenanthroline. Type example of peptidase family M22  
**References:** [3, 4, 2808]

[EC 3.4.24.57 created 1993]

#### EC 3.4.24.58

- Accepted name:** russellysin  
**Reaction:** Specifically activates several components of the blood clotting system, including coagulation factor X, coagulation factor IX and protein C by cleavage of -Arg— bonds. Has no action on insulin B chain  
**Other name(s):** Russell's viper venom factor X activator, RVV-X; blood-coagulation factor X activating enzyme; metalloproteinase RVV-x; *Vipera russelli* proteinase; Russell's viper blood coagulation factor X activator; RVV-V



**Comments:** This enzyme from the venom of Russell's viper (*Vipera russelli*) of 79 kDa comprises a heavy (59 kDa) and a heterogeneous light (18-21 kDa) chain. Contains Ca<sup>2+</sup> and Zn<sup>2+</sup>. The heavy chain contains the zinc-binding endopeptidase domain and a disintegrin. In peptidase family M12 (astacin family)

**References:** [864, 1702, 2862]

[EC 3.4.24.58 created 1993]

#### EC 3.4.24.59

**Accepted name:** mitochondrial intermediate peptidase

**Reaction:** Release of an N-terminal octapeptide as second stage of processing of some proteins imported into the mitochondrion

**Other name(s):** mitochondrial intermediate precursor-processing proteinase; MIP

**Comments:** A homologue of thimet oligopeptidase. Natural substrates are precursor proteins that have already been processed by mitochondrial processing peptidase. In peptidase family M3 (thimet oligopeptidase family)

**References:** [1256, 1257]

[EC 3.4.24.59 created 1993]

#### EC 3.4.24.60

**Accepted name:** dactylysin

**Reaction:** Hydrolysis of peptides of at least six residues, with bulky hydrophobic residues in the P1' position. Shows a preference for hydrophobic doublets such as -Phe-|Phe- and -Phe-|Leu- in somatostatin-(1-14)-peptide and dynorphin A-(1-6)-peptide, respectively

**Other name(s):** peptide hormone inactivating endopeptidase; PHIE

**Comments:** An endopeptidase of 100 kDa secreted from the skin of the amphibian, *Xenopus laevis* (Dactylète du Cap). Resembles neprilysin in insensitivity to 1 μM captopril, but differs from it in being insensitive to thiorphan (1 μM) and unable to digest [Met<sup>5</sup>]enkephalin, [Leu<sup>5</sup>]enkephalin, oxytocin, and substance P-(7-11)-peptide. A similar endopeptidase is found in human neuroblastoma cells [577]

**References:** [378, 577, 1343]

[EC 3.4.24.60 created 1995]

#### EC 3.4.24.61

**Accepted name:** nardilysin

**Reaction:** Hydrolysis of polypeptides, preferably at -Xaa-|Arg-Lys-, and less commonly at -Arg-|Arg-Xaa-, in which Xaa is not Arg or Lys

**Other name(s):** *N*-arginine dibasic convertase; NRD-convertase

**Comments:** Enzyme of 133 kDa from rat brain and testis. A homologue of pitrilysin containing the His-Phe-Leu-Glu-His zinc-binding sequence, and a highly acidic stretch of 71 residues. Unusually for a metalloendopeptidase, inhibited by bestatin, amastatin and *N*-ethylmaleimide. In peptidase family M16 (pitrilysin family)

**References:** [953, 932, 425, 2280]

[EC 3.4.24.61 created 1995]

#### EC 3.4.24.62

**Accepted name:** magnolysin

**Reaction:** Hydrolysis of polypeptides with Arg or Lys in P1 and P2, e.g. to hydrolyse *pro*-oxytocin at -Lys-Arg-|Ala-Val-. The specificity further depends on the organization of a β-turn-α-helix of nine or more residues containing the paired basic amino acids near the centre [3]

**Other name(s):** bovine neurosecretory granule protease cleaving *pro*-oxytocin/neurophysin; *pro*-oxytocin/neurophysin convertase; prooxyphysin proteinase; *pro*-oxytocin convertase

**Comments:** An endopeptidase of 58 kDa known from bovine pituitary neurosecretory granules and bovine and human corpus luteum [2293, 998]. Inhibited by EDTA [457]  
**References:** [457, 503, 276, 2293, 998]

[EC 3.4.24.62 created 1995]

#### EC 3.4.24.63

**Accepted name:** meprin B  
**Reaction:** Hydrolysis of proteins, including azocasein, and peptides. Hydrolysis of -His<sup>5</sup>-Leu-, -Leu<sup>6</sup>-Cys-, -Ala<sup>14</sup>-Leu- and -Cys<sup>19</sup>-Gly- bonds in insulin B chain  
**Other name(s):** meprin-b  
**Comments:** A brush border membrane-bound metalloendopeptidase known from the intestine of all mouse strains that have been tested, and the kidney of certain inbred strains. A tetramer of meprin  $\beta$  subunits (in contrast to meprin A, which contains both  $\alpha$  and  $\beta$  subunits). Occurs in the kidney as a proenzyme that can be activated by trypsin. Meprin B is inhibited by both EDTA and 1,10-phenanthroline, but not by phosphoramidon, captopril or thiorphan. In peptidase family M12 (astacin family)  
**References:** [1522, 957, 1325, 3210]

[EC 3.4.24.63 created 1995]

#### EC 3.4.24.64

**Accepted name:** mitochondrial processing peptidase  
**Reaction:** Release of N-terminal targeting peptides from precursor proteins imported into the mitochondrion, typically with Arg in position P2  
**Other name(s):** processing enhancing peptidase (for one of two subunits); mitochondrial protein precursor-processing proteinase; matrix peptidase; matrix processing peptidase; matrix processing proteinase; mitochondrial protein precursor-processing proteinase; MPP  
**Comments:** Known from the mitochondrial matrix of fungi and mammals. Formed from two subunits, both homologous with pitrilysin [2382], and the products of the *MAS1* and *MAS2* genes in yeast. In peptidase family M16 (pitrilysin family).  
**References:** [1319, 3202, 2382, 1365, 320]

[EC 3.4.24.64 created 1989/90 as EC 3.4.99.41, transferred 1995 to EC 3.4.24.64]

#### EC 3.4.24.65

**Accepted name:** macrophage elastase  
**Reaction:** Hydrolysis of soluble and insoluble elastin [1]. Specific cleavages are also produced at -Ala<sup>14</sup>-Leu- and -Tyr<sup>16</sup>-Leu- in the B chain of insulin [2]  
**Other name(s):** metalloelastase; human macrophage metalloelastase (HME)  
**Comments:** This enzyme is synthesized as a proenzyme of 53 kDa that is converted to an active form of 22 kDa. cDNA sequences have been obtained for the mouse [2623] and human [2624] enzymes. In peptidase family M10 (interstitial collagenase family)  
**References:** [135, 1430, 2623, 2624]

[EC 3.4.24.65 created 1995]

#### EC 3.4.24.66

**Accepted name:** choriolysin L  
**Reaction:** Hydrolysis of the inner layer of fish egg envelope. Also hydrolysis of casein and small molecule substrates such as succinyl-Leu-Leu-Val-Tyr-7-(4-methyl)coumarylamide  
**Other name(s):** teleost hatching enzyme (component); low choriolytic enzyme (LCE)  
**Comments:** Known from the teleost fish *Oryzias latipes* (medaka). Efficient dissolution of the egg membrane requires concerted action with choriolysin H. A 24 kDa peptidase family M12 (astacin family)

**References:** [3272, 3273, 3275, 3277]

[EC 3.4.24.66 created 1995]

#### EC 3.4.24.67

**Accepted name:** choriolysin H

**Reaction:** Hydrolysis of the inner layer of fish egg envelope. Also hydrolysis of casein and small molecule substrates such as succinyl-Leu-Leu-Val-Tyr-7-(4-methyl)coumarylamide

**Other name(s):** teleost hatching enzyme (component); high choriolytic enzyme (HCE)

**Comments:** Known from the teleost fish *Oryzias latipes* (medaka). Efficient dissolution of the egg membrane requires concerted action with choriolysin L. A 25.5 kDa peptidase in family M12 (astacin family)

**References:** [3243, 3274, 3276, 3277, 1625]

[EC 3.4.24.67 created 1995]

#### EC 3.4.24.68

**Accepted name:** tentoxilysin

**Reaction:** Hydrolysis of -Gln<sup>76</sup>-Phe- bond in synaptobrevin (also known as neuronal vesicle-associated membrane protein, VAMP)

**Other name(s):** tetanus neurotoxin

**Comments:** Zinc enzyme produced by *Clostridium tetani*. Proenzyme of 150 kDa is processed to disulfide-linked subunits of 100 and 50 kDa, the latter being responsible for the endopeptidase activity. Weakly inhibited by captopril, and phosphoramidon. The clostridial neurotoxins disable the neuroexocytosis apparatus, and have been described as the most toxic substances known. Tentoxilysin acts at the spinal inhibitory interneurons, blocking the release of various neurotransmitters to produce spastic paralysis. Type example of peptidase family M27 (tentoxilysin family)

**References:** [832, 2557, 2561, 1954, 2559]

[EC 3.4.24.68 created 1995]

#### EC 3.4.24.69

**Accepted name:** bontoxilysin

**Reaction:** Limited hydrolysis of proteins of the neuroexocytosis apparatus, synaptobrevin (also known as neuronal vesicle-associated membrane protein, VAMP), synaptosome-associated protein of 25 kDa (SNAP25) or syntaxin. No detected action on small molecule substrates

**Other name(s):** botulinum neurotoxin; BoNT

**Comments:** This zinc enzyme, produced by *Clostridium botulinum*, occurs as forms A-G that differ in specificity of action on the proteins of the neuroexocytosis apparatus [2560, 2562, 2563, 2558, 1954]. The 150-kDa proenzymes of bontoxilysin are processed to disulfide-linked subunits of 100 and 50 kDa, the latter being responsible for the endopeptidase activities. Weakly inhibited by captopril, and phosphoramidon. Toxicity is due to action at the neuromuscular junctions that blocks release of acetylcholine, causing flaccid paralysis, in contrast to the spastic paralysis caused by tentoxilysin. In peptidase family M27 (tentoxilysin family)

**References:** [2560, 2562, 2563, 2558, 1954, 2559]

[EC 3.4.24.69 created 1995]

#### EC 3.4.24.70

**Accepted name:** oligopeptidase A

**Reaction:** Hydrolysis of oligopeptides, with broad specificity. Gly or Ala commonly occur as P1 or P1' residues, but more distant residues are also important, as is shown by the fact that Z-Gly-Pro-Gly-Gly-Pro-Ala is cleaved, but not Z-(Gly)<sub>5</sub> [4]

**Other name(s):** 68000-M signalpeptide hydrolase

**Comments:** Known from *Escherichia coli* and *Salmonella typhimurium*. A zinc metallopeptidase, in peptidase family M3 (thimet oligopeptidase family), but differs from thimet oligopeptidase in lack of thiol-activation

**References:** [2118, 476, 475, 474]

[EC 3.4.24.70 created 1996]

#### EC 3.4.24.71

**Accepted name:** endothelin-converting enzyme 1

**Reaction:** Hydrolysis of the -Trp<sup>21</sup>-Val- bond in big endothelin to form endothelin 1

**Other name(s):** endothelin-converting enzyme; ECE-1

**Comments:** A phosphoramidon-sensitive metalloendopeptidase in peptidase family M13 (neprilysin family). An integral membrane protein predominantly of endothelial cells, which generates the potent vasoconstrictor endothelin 1 from its inactive precursor

**References:** [2840, 2632, 3231]

[EC 3.4.24.71 created 1996]

#### EC 3.4.24.72

**Accepted name:** fibrolase

**Reaction:** Hydrolysis of -Ala<sup>14</sup>-Leu- in insulin B chain and -Lys<sup>413</sup>-Leu- in A $\alpha$ -chain of fibrinogen

**Other name(s):** fibrinolytic proteinase; *Agkistrodon contortrix contortrix* metalloproteinase; *Agkistrodon contortrix contortrix* venom metalloproteinase

**Comments:** A 23-kDa, non-hemorrhagic enzyme from the venom of the southern copperhead snake (*Agkistrodon contortrix contortrix*). In peptidase family M12 (astacin family)

**References:** [12, 996, 2372, 1731, 2411]

[EC 3.4.24.72 created 1996]

#### EC 3.4.24.73

**Accepted name:** jararhagin

**Reaction:** Hydrolysis of -His<sup>10</sup>-Leu-, -Ala<sup>14</sup>-Leu-, -Tyr<sup>16</sup>-Leu- and -Phe<sup>24</sup>-Phe- bonds in insulin B chain

**Other name(s):** HF2-proteinase; JF1

**Comments:** Hemorrhagic endopeptidase from the venom of the jararaca snake (*Bothrops jararaca*). The 52-kDa enzyme contains a disintegrin domain [2224]. In peptidase family M12 (astacin family)

**References:** [1798, 90, 2224]

[EC 3.4.24.73 created 1996]

#### EC 3.4.24.74

**Accepted name:** fragilysin

**Reaction:** Broad proteolytic specificity, bonds hydrolysed including -Gly-Leu-, -Met-Leu-, -Phe-Leu-, -Cys-Leu-, Leu-Gly

**Other name(s):** *Bacteroides fragilis* (entero)toxin

**Comments:** Thought to be a cause of diarrhoea in animals and humans. Hydrolyses extracellular matrix proteins, and disrupts tight junctions of intestinal epithelial cells. Also degrades intracellular, cytoskeletal proteins actin, myosin and others. In peptidase family M10 (interstitial collagenase family)

**References:** [1951, 2124, 632, 1516, 1476]

[EC 3.4.24.74 created 1997]

#### EC 3.4.24.75

**Accepted name:** lysostaphin  
**Reaction:** Hydrolysis of the -Gly—Gly- bond in the pentaglycine inter-peptide link joining staphylococcal cell wall peptidoglycans  
**Other name(s):** glycyl-glycine endopeptidase  
**Comments:** A zinc-dependent, 25-kDa endopeptidase from *Staphylococcus simulans*. Lyses cells of *S. aureus*, in particular, by its action on the cross-bridges of the cell wall. Type example of peptidase family M23.  
**References:** [2392, 109, 2920]

[EC 3.4.24.75 created 1997]

#### EC 3.4.24.76

**Accepted name:** flavastacin  
**Reaction:** Hydrolyses polypeptides on the amino-side of Asp in -Xaa—Asp-. Acts very slowly on -Xaa—Glu  
**Comments:** A zinc metalloendopeptidase in peptidase family M12 (astacin family), secreted by the bacterium *Flavobacterium meningosepticum*. The specificity is similar to that of EC 3.4.24.33, peptidyl-Asp metalloendopeptidase from *Pseudomonas fragi* but the two are reported to be structurally dissimilar  
**References:** [2889]

[EC 3.4.24.76 created 2000]

#### EC 3.4.24.77

**Accepted name:** snapalysin  
**Reaction:** Hydrolyses proteins with a preference for Tyr or Phe in the P1' position. Has no action on amino-acid *p*-nitroanilides  
**Other name(s):** small neutral protease; SnpA gene product (*Streptomyces lividans*)  
**Comments:** Type example of peptidase family M7.  
**References:** [1570, 336, 1569]

[EC 3.4.24.77 created 2001]

#### EC 3.4.24.78

**Accepted name:** gpr endopeptidase  
**Reaction:** Endopeptidase action with P4 Glu or Asp, P1 preferably Glu > Asp, P1' hydrophobic and P2' Ala  
**Other name(s):** germination proteinase  
**Comments:** Initiates the degradation of small, acid-soluble proteins during spore germination in *Bacillus megaterium*. Type example of peptidase family M63.  
**References:** [2309]

[EC 3.4.24.78 created 2003]

#### EC 3.4.24.79

**Accepted name:** pappalysin-1  
**Reaction:** Cleavage of the Met<sup>135</sup>—Lys bond in insulin-like growth factor binding protein (IGFBP)-4, and the Ser<sup>143</sup>—Lys bond in IGFBP-5  
**Other name(s):** insulin-like growth factor binding protein-4 protease; pregnancy-associated plasma protein-A  
**Comments:** A 400-kDa disulfide-linked dimer. Circulates in human pregnancy mainly as a complex with the pro-form of eosinophil major basic protein, which acts as an inhibitor of the peptidase. The rate of hydrolysis of IGFBP-4 is increased about 20-fold by the presence of insulin-like growth factor (IGF), whereas that of IGFBP-5 is decreased about two-fold. In peptidase family M43.  
**References:** [1611, 409]

[EC 3.4.24.79 created 2003]

#### EC 3.4.24.80

**Accepted name:** membrane-type matrix metalloproteinase-1

**Reaction:** Endopeptidase activity. Activates progelatinase A by cleavage of the propeptide at Asn<sup>37</sup>—Leu. Other bonds hydrolysed include Gly<sup>35</sup>—Ile in the propeptide of collagenase 3, and Asn<sup>341</sup>—Phe, Asp<sup>441</sup>—Leu and Gln<sup>354</sup>—Thr in the aggrecan interglobular domain

**Other name(s):** matrix metalloproteinase 14

**Comments:** In peptidase family M10, but, unlike most members of the family, is membrane-anchored. Believed to play an important role in the activation of progelatinase A at cell surfaces.

**References:** [1277]

[EC 3.4.24.80 created 2003]

#### EC 3.4.24.81

**Accepted name:** ADAM10 endopeptidase

**Reaction:** Endopeptidase of broad specificity

**Other name(s):** Kuzbanian protein; myelin-associated disintegrin metalloproteinase

**Comments:** In peptidase family M12. Partially responsible for the "α-secretase" activity in brain that degrades the potentially harmful β-amyloid peptide. Work with ADAM10-deficient mice supports a role in Notch signalling.

**References:** [1010]

[EC 3.4.24.81 created 2003]

#### EC 3.4.24.82

**Accepted name:** ADAMTS-4 endopeptidase

**Reaction:** Glutamyl endopeptidase; bonds cleaved include -Thr-Glu-Gly-Glu<sup>373</sup>—Ala-Arg-Gly-Ser- in the interglobular domain of mammalian aggrecan

**Other name(s):** aggrecanase-1

**Comments:** In peptidase family M12. Thought to be biologically significant for the degradation of the aggrecan component of cartilage matrix.

**References:** [3165]

[EC 3.4.24.82 created 2003]

#### EC 3.4.24.83

**Accepted name:** anthrax lethal factor endopeptidase

**Reaction:** Preferred amino acids around the cleavage site can be denoted BBBBxHx—H, in which B denotes Arg or Lys, H denotes a hydrophobic amino acid, and x is any amino acid. The only known protein substrates are mitogen-activated protein (MAP) kinase kinases

**Other name(s):** lethal toxin

**Comments:** From the bacterium *Bacillus anthracis* that causes anthrax. One of three proteins that are collectively termed anthrax toxin. Cleaves several MAP kinase kinases near their N-termini, preventing them from phosphorylating the downstream mitogen-activated protein kinases. In peptidase family M34.

**References:** [2232]

[EC 3.4.24.83 created 2003]

#### EC 3.4.24.84

**Accepted name:** Ste24 endopeptidase

**Reaction:** The peptide bond hydrolysed can be designated -C—aaX in which C is an *S*-isoprenylated cysteine residue, a is usually aliphatic and X is the C-terminal residue of the substrate protein, and may be any of several amino acids

**Comments:** Type example of peptidase family M48. One of two enzymes that can catalyse this processing step for mating a-factor in yeast. Subsequently, the *S*-isoprenylated cysteine residue that forms the new C-terminus is methyl-esterified and forms a hydrophobic membrane-anchor.

**References:** [2865]

[EC 3.4.24.84 created 2003]

#### EC 3.4.24.85

**Accepted name:** S2P endopeptidase

**Reaction:** Cleaves several transcription factors that are type-2 transmembrane proteins within membrane-spanning domains. Known substrates include sterol regulatory element-binding protein (SREBP)-1, SREBP-2 and forms of the transcriptional activator ATF6. SREBP-2 is cleaved at the site DRSRILL<sup>483</sup>—CVLTLCLCSFNPLTSLLQWGGA, in which the membrane-spanning segment is underlined. The residues NP (bold), 11 residues distal to the site of cleavage in the membrane-spanning domain, are important for cleavage by S2P endopeptidase. Replacement of either of these residues does not prevent cleavage, but there is no cleavage if both of these residues are replaced.

**Comments:** Type example of peptidase family M50. The transcription factors SREBP-1 and -2 are synthesized as precursor proteins that are attached to the membranes of the endoplasmic reticulum and two cleavages are needed to release the active factor so that it can move to the nucleus. This enzyme cleaves the second of these, and is thus the "site 2 protease", S2P.

**References:** [310]

[EC 3.4.24.85 created 2003]

#### EC 3.4.24.86

**Accepted name:** ADAM 17 endopeptidase

**Reaction:** Narrow endopeptidase specificity. Cleaves Pro-Leu-Ala-Gln-Ala—Val-Arg-Ser-Ser-Ser in the membrane-bound, 26-kDa form of tumour necrosis factor  $\alpha$  (TNF $\alpha$ ). Similarly cleaves other membrane-anchored, cell-surface proteins to "shed" the extracellular domains

**Other name(s):** tumor necrosis factor  $\alpha$ -converting enzyme; TACE

**Comments:** In peptidase family M12. *In vivo*, the cleavage of tumour necrosis factor  $\alpha$  precursor releases the soluble, 17-kDa TNF $\alpha$ , which induces inflammation.

**References:** [228]

[EC 3.4.24.86 created 2003]

#### EC 3.4.24.87

**Accepted name:** ADAMTS13 endopeptidase

**Reaction:** The enzyme cleaves the von Willebrand factor at bond Tyr<sup>842</sup>—Met<sup>843</sup> within the A2 domain

**Other name(s):** ADAMTS VWF cleaving metalloprotease; ADAMTS-13; ADAMTS13; vWF-cleaving protease; VWF-CP; vWF-degrading protease; Upshaw factor; von Willebrand factor cleaving protease; ADAMTS13 peptidase

**Comments:** In peptidase family M12.

**References:** [841, 634]

[EC 3.4.24.87 created 2009]

[3.4.24.88 *Transferred entry. desampylase. Transferred to EC 3.4.19.15 desampylase*]

[EC 3.4.24.88 created 2015, deleted 2016]

#### EC 3.4.24.89

**Accepted name:** Pro-Pro endopeptidase

**Reaction:** The enzyme catalyses the hydrolytic cleavage of peptide bonds between two proline residues



**Other name(s):** metalloprotease CD2830  
**Comments:** This metalloprotease, which is secreted by the bacterium *Peptoclostridium difficile*, contains zinc.  
**References:** [351, 1116, 1115]

[EC 3.4.24.89 created 2015]

## EC 3.4.25 Threonine endopeptidases

### EC 3.4.25.1

**Accepted name:** proteasome endopeptidase complex  
**Reaction:** Cleavage of peptide bonds with very broad specificity  
**Other name(s):** ingensin; macropain; multicatalytic endopeptidase complex; prosome; multicatalytic proteinase (complex); MCP; proteasome; large multicatalytic protease; multicatalytic proteinase; proteasome organelle; alkaline protease; 26S protease; tricorn proteinase; tricorn protease  
**Comments:** A 20-S protein composed of 28 subunits arranged in four rings of seven. The outer rings are composed of  $\alpha$  subunits, but the  $\beta$  subunits forming the inner rings are responsible for peptidase activity. In eukaryotic organisms there are up to seven different types of  $\beta$  subunits, three of which may carry the N-terminal threonine residues that are the nucleophiles in catalysis, and show different specificities. The molecule is barrel-shaped, and the active sites are on the inner surfaces. Terminal apertures restrict access of substrates to the active sites. There is evidence that catalytic subunits are replaced by others under some conditions so as to alter the specificity of proteolysis, perhaps optimizing it for the formation of antigenic peptides. A complex of the 20-S proteasome endopeptidase complex with a 19-S regulatory unit is the 26-S proteasome that degrades ubiquitin-protein conjugates. Type example of peptidase family T1.  
**References:** [2600, 499, 991, 605]

[EC 3.4.25.1 created 1978 as EC 3.4.24.5, part transferred 1989 to EC 3.4.22.21, transferred 1992 to EC 3.4.99.46, transferred 2000 to EC 3.4.25.1]

### EC 3.4.25.2

**Accepted name:** HslU—HslV peptidase  
**Reaction:** ATP-dependent cleavage of peptide bonds with broad specificity.  
**Other name(s):** HslUV; HslV-HslU; HslV peptidase; ATP-dependent HslV-HslU proteinase; caseinolytic protease X; caseinolytic proteinase X; ClpXP ATP-dependent protease; ClpXP protease; ClpXP serine proteinase; *Escherichia coli* ClpXP serine proteinase; HslUV protease; HslUV proteinase; HslVU protease; HslVU proteinase; protease HslVU; proteinase HslUV  
**Comments:** The HslU subunit of the HslU—HslV complex functions as an ATP dependent ‘unfoldase’. The binding of ATP and its subsequent hydrolysis by HslU are essential for unfolding of protein substrates subsequently hydrolysed by HslV [3291]. HslU recognizes the N-terminal part of its protein substrates and unfolds these before they are guided to HslV for hydrolysis [333]. In peptidase family T1.  
**References:** [3112, 2092, 2364, 3292, 3291, 1382, 333]

[EC 3.4.25.2 created 2009, modified 2010]

## EC 3.4.99 Endopeptidases of unknown catalytic mechanism (sub-subclass is currently empty)

[3.4.99.1 Transferred entry. *acrocylicum* proteinase. Now EC 3.4.23.28, *acrocylicdropepsin*]

[EC 3.4.99.1 created 1972, deleted 1978 [transferred to EC 3.4.23.6, deleted 1992]]

[3.4.99.2 Deleted entry. *agavain*]

[EC 3.4.99.2 created 1972, deleted 1992]



- [3.4.99.3 Deleted entry. *angiotensinase*]  
[EC 3.4.99.3 created 1972, deleted 1992]
- [3.4.99.4 Transferred entry. *aspartylendopeptidase*. Now EC 3.4.23.12, *nepenthesin*]  
[EC 3.4.99.4 created 1972, deleted 1978]
- [3.4.99.5 Transferred entry. *Clostridium histolyticum collagenase 2*. Now EC 3.4.24.3, *microbial collagenase*]  
[EC 3.4.99.5 created 1972, deleted 1978]
- [3.4.99.6 Transferred entry. *crayfish low-molecular-weight proteinase*. Now EC 3.4.24.21, *astacin*]  
[EC 3.4.99.6 created 1972, deleted 1992]
- [3.4.99.7 Deleted entry. *euphorbain*]  
[EC 3.4.99.7 created 1972, deleted 1989]
- [3.4.99.8 Deleted entry. *Gliocladium proteinase*]  
[EC 3.4.99.8 created 1972, deleted 1992]
- [3.4.99.9 Deleted entry. *hurain*. Now considered EC 3.4.21.25, *cucumisin*]  
[EC 3.4.99.9 created 1972, deleted 1992]
- [3.4.99.10 Transferred entry. *insulinase*. Now EC 3.4.24.56, *insulysin*]  
[EC 3.4.99.10 created 1972, transferred 1976 to EC 3.4.22.11, transferred 1978 to EC 3.4.99.45, transferred 1993 to EC 3.4.24.56]
- [3.4.99.11 Deleted entry. *Streptomyces alkalophilic keratinase*]  
[EC 3.4.99.11 created 1965 as EC 3.4.4.25, transferred 1972 to EC 3.4.99.11, deleted 1992]
- [3.4.99.12 Deleted entry. *Trichophyton mentagrophytes keratinase*]  
[EC 3.4.99.12 created 1972, deleted 1978 [transferred to EC 3.4.24.10, deleted 1992]]
- [3.4.99.13 Transferred entry.  $\beta$ -lytic proteinase (*Mycobacterium sporangium*). Now EC 3.4.24.32,  $\beta$ -lytic metalloendopeptidase]  
[EC 3.4.99.13 created 1972, deleted 1978 [transferred to EC 3.4.24.4, deleted 1992]]
- [3.4.99.14 Deleted entry. *mexicanain*]  
[EC 3.4.99.14 created 1972, deleted 1992]
- [3.4.99.15 Deleted entry. *Paecilomyces proteinase*]  
[EC 3.4.99.15 created 1972, deleted 1978 [transferred to EC 3.4.23.6, deleted 1992]]
- [3.4.99.16 Deleted entry. *Penicillium notatum extracellular proteinase*]  
[EC 3.4.99.16 created 1972, deleted 1992]
- [3.4.99.17 Deleted entry. *peptidoglycan endopeptidase*]  
[EC 3.4.99.17 created 1972, deleted 1992]
- [3.4.99.18 Deleted entry. *pinguinain*]  
[EC 3.4.99.18 created 1972, deleted 1992]
- [3.4.99.19 Transferred entry. *renin*. Now EC 3.4.23.15, *renin*]  
[EC 3.4.99.19 created 1972, deleted 1981]

- [3.4.99.20 Deleted entry. *Scopulariopsis proteinase*]  
[EC 3.4.99.20 created 1972, deleted 1992]
- [3.4.99.21 Deleted entry. *solanain*. Now considered EC 3.4.21.25, *cucumisin*]  
[EC 3.4.99.21 created 1972, deleted 1992]
- [3.4.99.22 Transferred entry. *staphylokinase*. Now EC 3.4.24.29, *aureolysin*]  
[EC 3.4.99.22 created 1972, modified 1976, deleted 1978 [transferred to EC 3.4.24.4, deleted 1992]]
- [3.4.99.23 Deleted entry. *tabernamontanain*. Now considered EC 3.4.21.25, *cucumisin*]  
[EC 3.4.99.23 created 1972, deleted 1992]
- [3.4.99.24 Deleted entry. *Tenebrio*  $\alpha$ -*proteinase*]  
[EC 3.4.99.24 created 1972, deleted 1978 [transferred to EC 3.4.21.18, deleted 1992]]
- [3.4.99.25 Transferred entry. *trametes acid proteinase*. Now EC 3.4.23.21, *rhizopuspepsin*]  
[EC 3.4.99.25 created 1972, deleted 1978 [transferred to EC 3.4.23.6, deleted 1992]]
- [3.4.99.26 Transferred entry. *urokinase*. Now EC 3.4.21.68, *t-plasminogen activator*]  
[EC 3.4.99.26 created 1972, deleted 1978 [transferred to EC 3.4.21.31, deleted 1992]]
- [3.4.99.27 Deleted entry. *Echis carinatus prothrombin-activating proteinase*]  
[EC 3.4.99.27 created 1978, deleted 1992]
- [3.4.99.28 Transferred entry. *Oxyuranus scutellatus prothrombin-activating proteinase*. Now EC 3.4.21.60, *scutellarin*]  
[EC 3.4.99.28 created 1978, deleted 1992]
- [3.4.99.29 Deleted entry. *Myxobacter AL-1 proteinase I*]  
[EC 3.4.99.29 created 1978, deleted 1992]
- [3.4.99.30 Transferred entry. *Myxobacter AL-1 proteinase II*. Now EC 3.4.24.20, *peptidyl-Lys metalloendopeptidase*]  
[EC 3.4.99.30 created 1978, deleted 1992]
- [3.4.99.31 Transferred entry. *tissue endopeptidase degrading collagenase synthetic substrate*. Now EC 3.4.24.15, *thimet oligopeptidase*]  
[EC 3.4.99.31 created 1978, deleted 1992]
- [3.4.99.32 Transferred entry. *Armillaria mellea neutral proteinase*. Now EC 3.4.24.20, *peptidyl-Lys metalloendopeptidase*]  
[EC 3.4.99.32 created 1978, deleted 1992]
- [3.4.99.33 Deleted entry. *cathepsin R*]  
[EC 3.4.99.33 created 1981, deleted 1984 [transferred to EC 3.4.21.52, deleted 1992]]
- [3.4.99.34 Deleted entry. *mytilidase*]  
[EC 3.4.99.34 created 1981, deleted 1992]
- [3.4.99.35 Transferred entry. *premurein-leader peptidase*. Now EC 3.4.23.36, *signal peptidase II*]  
[EC 3.4.99.35 created 1984, deleted 1995]
- [3.4.99.36 Transferred entry. *leader peptidase*. Now EC 3.4.21.89, *signal peptidase I*]  
[EC 3.4.99.36 created 1984, deleted 1995]

- [3.4.99.37 Deleted entry. *RecA* peptidase]  
[EC 3.4.99.37 created 1989, deleted 1992]
- [3.4.99.38 Transferred entry. *pro-opiomelanotropin-converting proteinase*. Now EC 3.4.23.17, *pro-opiomelanocortin converting enzyme*]  
[EC 3.4.99.38 created 1989, deleted 1992]
- [3.4.99.39 Deleted entry. *pseudomurein endopeptidase*]  
[EC 3.4.99.39 created 1989, deleted 1992]
- [3.4.99.40 Deleted entry. *Pro-gonadoliberin proteinase*]  
[EC 3.4.99.40 created 1989, deleted 1992]
- [3.4.99.41 Transferred entry. *mitochondrial processing peptidase*. Now EC 3.4.24.64, *mitochondrial processing peptidase*]  
[EC 3.4.99.41 created 1989/90, deleted 1995]
- [3.4.99.42 Deleted entry. *leucyllysine endopeptidase*]  
[EC 3.4.99.42 created 1990, deleted 1992]
- [3.4.99.43 Transferred entry. *thermopsin*. Now EC 3.4.23.42, *thermopsin*]  
[EC 3.4.99.43 created 1992, deleted 2000]
- [3.4.99.44 Transferred entry. *pitrilysin*. Now EC 3.4.24.55, *pitrilysin*]  
[EC 3.4.99.44 created 1992, deleted 1993]
- [3.4.99.45 Transferred entry. *insulinase*. Now EC 3.4.24.56, *insulysin*]  
[EC 3.4.99.45 created 1992, deleted 1993]
- [3.4.99.46 Transferred entry. *multicatalytic endopeptidase complex*. Now EC 3.4.25.1, *proteasome endopeptidase complex*]  
[EC 3.4.99.46 created 1992, deleted 2000]

## EC 3.5 Acting on carbon-nitrogen bonds, other than peptide bonds

This subclass contains those enzymes that hydrolyse amides, amidines and other C-N bonds. Sub-subclasses are based on the substrate: linear amides (EC 3.5.1), cyclic amides (EC 3.5.2), linear amidines (EC 3.5.3), cyclic amidines (EC 3.5.4), nitriles (EC 3.5.5) and other compounds (EC 3.5.99).

### EC 3.5.1 In linear amides

#### EC 3.5.1.1

**Accepted name:** asparaginase  
**Reaction:** L-asparagine + H<sub>2</sub>O = L-aspartate + NH<sub>3</sub>  
**Other name(s):** asparaginase II; L-asparaginase; colaspase; elspar; leunase; crasnitin; α-asparaginase  
**Systematic name:** L-asparagine amidohydrolase  
**References:** [1031, 1154, 2797]

[EC 3.5.1.1 created 1961]

#### EC 3.5.1.2

**Accepted name:** glutaminase  
**Reaction:** L-glutamine + H<sub>2</sub>O = L-glutamate + NH<sub>3</sub>  
**Other name(s):** glutaminase I; L-glutaminase; glutamine aminohydrolase  
**Systematic name:** L-glutamine amidohydrolase  
**References:** [1553, 2435]

[EC 3.5.1.2 created 1961]

#### EC 3.5.1.3

**Accepted name:** ω-amidase  
**Reaction:** a monoamide of a dicarboxylate + H<sub>2</sub>O = a dicarboxylate + NH<sub>3</sub>  
**Other name(s):** α-keto acid-ω-amidase  
**Systematic name:** ω-amidodicarboxylate amidohydrolase  
**Comments:** Acts on glutaramate, succinamate and their 2-oxo derivatives.  
**References:** [1881, 1882]

[EC 3.5.1.3 created 1961]

#### EC 3.5.1.4

**Accepted name:** amidase  
**Reaction:** a monocarboxylic acid amide + H<sub>2</sub>O = a monocarboxylate + NH<sub>3</sub>  
**Other name(s):** acylamidase; acylase (misleading); amidohydrolase (ambiguous); deaminase (ambiguous); fatty acylamidase; *N*-acetylaminohydrolase (ambiguous)  
**Systematic name:** acylamide amidohydrolase  
**References:** [280, 281]

[EC 3.5.1.4 created 1961, modified 2011]

#### EC 3.5.1.5

**Accepted name:** urease  
**Reaction:** urea + H<sub>2</sub>O = CO<sub>2</sub> + 2 NH<sub>3</sub>  
**Systematic name:** urea amidohydrolase  
**Comments:** A nickel protein.  
**References:** [620, 2799, 3053]

[EC 3.5.1.5 created 1961]

#### EC 3.5.1.6

**Accepted name:** β-ureidopropionase  
**Reaction:** 3-ureidopropoate + H<sub>2</sub>O = β-alanine + CO<sub>2</sub> + NH<sub>3</sub>  
**Other name(s):** *N*-carbamoyl-β-alanine amidohydrolase  
**Systematic name:** 3-ureidopropoate amidohydrolase  
**Comments:** The animal enzyme also acts on β-ureidoisobutyrate.  
**References:** [361, 368, 2961]

[EC 3.5.1.6 created 1961]

#### EC 3.5.1.7

**Accepted name:** ureidosuccinase  
**Reaction:** *N*-carbamoyl-L-aspartate + H<sub>2</sub>O = L-aspartate + CO<sub>2</sub> + NH<sub>3</sub>  
**Systematic name:** *N*-carbamoyl-L-aspartate amidohydrolase  
**References:** [1683]

[EC 3.5.1.7 created 1961]

#### EC 3.5.1.8

**Accepted name:** formylaspartate deformylase  
**Reaction:** *N*-formyl-L-aspartate + H<sub>2</sub>O = formate + L-aspartate  
**Other name(s):** formylaspartic formylase (formylase I, formylase II)  
**Systematic name:** *N*-formyl-L-aspartate amidohydrolase  
**References:** [2163]

[EC 3.5.1.8 created 1961]

#### EC 3.5.1.9

**Accepted name:** arylformamidase  
**Reaction:** *N*-formyl-L-kynurenine + H<sub>2</sub>O = formate + L-kynurenine  
**Other name(s):** kynurenine formamidase; formylase; formylkynureninase; formylkynurenine formamidase; formamidase I; formamidase II  
**Systematic name:** aryl-formylamine amidohydrolase  
**Comments:** Also acts on other aromatic formylamines.  
**References:** [1086, 1298, 1878]

[EC 3.5.1.9 created 1961]

#### EC 3.5.1.10

**Accepted name:** formyltetrahydrofolate deformylase  
**Reaction:** 10-formyltetrahydrofolate + H<sub>2</sub>O = formate + tetrahydrofolate  
**Systematic name:** 10-formyltetrahydrofolate amidohydrolase  
**References:** [1206]

[EC 3.5.1.10 created 1961]

#### EC 3.5.1.11

**Accepted name:** penicillin amidase  
**Reaction:** penicillin + H<sub>2</sub>O = a carboxylate + 6-aminopenicillanate  
**Other name(s):** penicillin acylase; benzylpenicillin acylase; novozym 217; semacylase; α-acylamino-β-lactam acylhydrolase; ampicillin acylase  
**Systematic name:** penicillin amidohydrolase  
**References:** [2500]

[EC 3.5.1.11 created 1961]

#### EC 3.5.1.12

**Accepted name:** biotinidase  
**Reaction:** biotin amide + H<sub>2</sub>O = biotin + NH<sub>3</sub>  
**Other name(s):** amidohydrolase biotinidase  
**Systematic name:** biotin-amide amidohydrolase  
**Comments:** Also acts on biotin esters.  
**References:** [1479, 2912]

[EC 3.5.1.12 created 1961]

#### EC 3.5.1.13

**Accepted name:** aryl-acylamidase

**Reaction:** an anilide + H<sub>2</sub>O = a carboxylate + aniline  
**Other name(s):** AAA-1; AAA-2; brain acetylcholinesterase (is associated with AAA-2); pseudocholinesterase (associated with arylacylamidase)  
**Systematic name:** aryl-acylamide amidohydrolase  
**Comments:** Also acts on 4-substituted anilides.  
**References:** [2090]

[EC 3.5.1.13 created 1965]

#### EC 3.5.1.14

**Accepted name:** *N*-acyl-aliphatic-L-amino acid amidohydrolase  
**Reaction:** (1) an *N*-acyl-aliphatic-L-amino acid + H<sub>2</sub>O = an aliphatic L-amino acid + a carboxylate  
(2) an *N*-acetyl-L-cysteine-*S*-conjugate + H<sub>2</sub>O = an L-cysteine-*S*-conjugate + acetate  
**Other name(s):** aminoacylase 1; aminoacylase I; dehydropeptidase II; histozyme; hippuricase; benzamidase; acylase I; hippurase; amido acid deacylase; L-aminoacylase; acylase; aminoacylase; L-amino-acid acylase; α-*N*-acylaminoacid hydrolase; long acyl amidoacylase; short acyl amidoacylase; ACY1 (gene name); *N*-acyl-L-amino-acid amidohydrolase  
**Systematic name:** *N*-acyl-aliphatic-L-amino acid amidohydrolase (carboxylate-forming)  
**Comments:** Contains Zn<sup>2+</sup>. The enzyme is found in animals and is involved in the hydrolysis of *N*-acylated or *N*-acetylated amino acids (except L-aspartate). It acts on mercapturic acids (*S*-conjugates of *N*-acetyl-L-cysteine) and neutral aliphatic *N*-acyl-α-amino acids. Some bacterial aminoacylases demonstrate substrate specificity of both EC 3.5.1.14 and EC 3.5.1.114. *cf.* EC 3.5.1.15, aspartoacylase and EC 3.5.1.114, *N*-acyl-aromatic-L-amino acid amidohydrolase.  
**References:** [219, 796, 1117, 1097, 2228, 3024, 1701]

[EC 3.5.1.14 created 1965, modified 2013]

#### EC 3.5.1.15

**Accepted name:** aspartoacylase  
**Reaction:** *N*-acyl-L-aspartate + H<sub>2</sub>O = a carboxylate + L-aspartate  
**Other name(s):** aminoacylase II; *N*-acetylaspartate amidohydrolase; acetyl-aspartic deaminase; acylase II  
**Systematic name:** *N*-acyl-L-aspartate amidohydrolase  
**References:** [218, 219]

[EC 3.5.1.15 created 1965]

#### EC 3.5.1.16

**Accepted name:** acetylornithine deacetylase  
**Reaction:** *N*<sup>2</sup>-acetyl-L-ornithine + H<sub>2</sub>O = acetate + L-ornithine  
**Other name(s):** acetylornithinase; *N*-acetylornithinase; 2-*N*-acetyl-L-ornithine amidohydrolase  
**Systematic name:** *N*<sup>2</sup>-acetyl-L-ornithine amidohydrolase  
**Comments:** Also hydrolyses *N*-acetylmethionine.  
**References:** [3072, 3073]

[EC 3.5.1.16 created 1965]

#### EC 3.5.1.17

**Accepted name:** acyl-lysine deacylase  
**Reaction:** *N*<sup>6</sup>-acyl-L-lysine + H<sub>2</sub>O = a carboxylate + L-lysine  
**Other name(s):** ε-lysine acylase; 6-*N*-acyl-L-lysine amidohydrolase  
**Systematic name:** *N*<sup>6</sup>-acyl-L-lysine amidohydrolase  
**References:** [2223]

[EC 3.5.1.17 created 1965]

**EC 3.5.1.18**

**Accepted name:** succinyl-diaminopimelate desuccinylase  
**Reaction:** *N*-succinyl-LL-2,6-diaminoheptanedioate + H<sub>2</sub>O = succinate + LL-2,6-diaminoheptanedioate  
**Other name(s):** *N*-succinyl-L- $\alpha$ , $\epsilon$ -diaminopimelic acid deacylase  
**Systematic name:** *N*-succinyl-LL-2,6-diaminoheptanedioate amidohydrolase  
**References:** [1459]

[EC 3.5.1.18 created 1965]

**EC 3.5.1.19**

**Accepted name:** nicotinamidase  
**Reaction:** nicotinamide + H<sub>2</sub>O = nicotinate + NH<sub>3</sub>  
**Other name(s):** nicotinamide deaminase; nicotinamide amidase; YNDase  
**Systematic name:** nicotinamide amidohydrolase  
**References:** [2268, 2532]

[EC 3.5.1.19 created 1972]

**EC 3.5.1.20**

**Accepted name:** citrullinase  
**Reaction:** L-citrulline + H<sub>2</sub>O = L-ornithine + CO<sub>2</sub> + NH<sub>3</sub>  
**Other name(s):** citrulline ureidase; citrulline hydrolase; L-citrulline 5-*N*-carbamoyldihydrolase  
**Systematic name:** L-citrulline *N*<sup>5</sup>-carbamoyldihydrolase  
**References:** [1140]

[EC 3.5.1.20 created 1972]

**EC 3.5.1.21**

**Accepted name:** *N*-acetyl- $\beta$ -alanine deacetylase  
**Reaction:** *N*-acetyl- $\beta$ -alanine + H<sub>2</sub>O = acetate +  $\beta$ -alanine  
**Systematic name:** *N*-acetyl- $\beta$ -alanine amidohydrolase  
**References:** [842]

[EC 3.5.1.21 created 1972]

**EC 3.5.1.22**

**Accepted name:** pantothenase  
**Reaction:** (*R*)-pantothenate + H<sub>2</sub>O = (*R*)-pantoate +  $\beta$ -alanine  
**Other name(s):** pantothenate hydrolase; pantothenate amidohydrolase  
**Systematic name:** (*R*)-pantothenate amidohydrolase  
**References:** [2122]

[EC 3.5.1.22 created 1972]

**EC 3.5.1.23**

**Accepted name:** ceramidase  
**Reaction:** a ceramide + H<sub>2</sub>O = a carboxylate + sphingosine  
**Other name(s):** acylsphingosine deacylase; glycosphingolipid ceramide deacylase  
**Systematic name:** *N*-acylsphingosine amidohydrolase  
**References:** [2088, 3279]

[EC 3.5.1.23 created 1972, modified 1990]

#### EC 3.5.1.24

**Accepted name:** choloylglycine hydrolase  
**Reaction:** glycocholate + H<sub>2</sub>O = cholate + glycine  
**Other name(s):** glycocholase; bile salt hydrolase; choloyltaurine hydrolase; 3 $\alpha$ ,7 $\alpha$ ,12 $\alpha$ -trihydroxy-5 $\beta$ -cholan-24-oylglycine amidohydrolase  
**Systematic name:** glycocholate amidohydrolase  
**Comments:** Also acts on the 3 $\alpha$ ,12 $\alpha$ -dihydroxy-derivative, and on choloyl-aurine.  
**References:** [2030, 2762]

[EC 3.5.1.24 created 1972]

#### EC 3.5.1.25

**Accepted name:** *N*-acetylglucosamine-6-phosphate deacetylase  
**Reaction:** *N*-acetyl-D-glucosamine 6-phosphate + H<sub>2</sub>O = D-glucosamine 6-phosphate + acetate  
**Other name(s):** acetylglucosamine phosphate deacetylase; acetylaminodeoxyglucosephosphate acetylhydrolase; 2-acetamido-2-deoxy-D-glucose-6-phosphate amidohydrolase  
**Systematic name:** *N*-acetyl-D-glucosamine-6-phosphate amidohydrolase  
**References:** [3174, 3251]

[EC 3.5.1.25 created 1972 (EC 3.5.1.80 created 1999, incorporated 2002)]

#### EC 3.5.1.26

**Accepted name:** *N*<sup>4</sup>-( $\beta$ -*N*-acetylglucosaminyl)-L-asparaginase  
**Reaction:** *N*<sup>4</sup>-( $\beta$ -*N*-acetyl-D-glucosaminyl)-L-asparagine + H<sub>2</sub>O = *N*-acetyl- $\beta$ -D-glucosaminylamine + L-aspartate  
**Other name(s):** aspartylglucosylamine deaspartylase; aspartylglucosylaminase; aspartylglucosaminidase; aspartylglycosylamine amidohydrolase; *N*-aspartyl- $\beta$ -glucosaminidase; glucosylamidase;  $\beta$ -aspartylglucosylamine amidohydrolase; 4-*N*-( $\beta$ -*N*-acetyl-D-glucosaminyl)-L-asparagine amidohydrolase  
**Systematic name:** *N*<sup>4</sup>-( $\beta$ -*N*-acetyl-D-glucosaminyl)-L-asparagine amidohydrolase  
**Comments:** Acts only on asparagine-oligosaccharides containing one amino acid, i.e., the asparagine has free  $\alpha$ -amino and  $\alpha$ -carboxyl groups [*cf.* EC 3.5.1.52, peptide-*N*<sup>4</sup>-(*N*-acetyl- $\beta$ -glucosaminyl)asparagine amidase]  
**References:** [1497, 1774, 2887]

[EC 3.5.1.26 created 1972 (EC 3.5.1.37 created 1972, incorporated 1976)]

[3.5.1.27 Deleted entry. *N*-formylmethionylaminoacyl-tRNA deformylase. The activity is covered by EC 3.5.1.88, peptide deformylase]

[EC 3.5.1.27 created 1972, deleted 2014]

#### EC 3.5.1.28

**Accepted name:** *N*-acetylmuramoyl-L-alanine amidase  
**Reaction:** Hydrolyses the link between *N*-acetylmuramoyl residues and L-amino acid residues in certain cell-wall glycopeptides  
**Other name(s):** acetylmuramyl-L-alanine amidase; *N*-acetylmuramyl-L-alanine amidase; *N*-acylmuramyl-L-alanine amidase; acetylmuramoyl-alanine amidase; *N*-acetylmuramic acid L-alanine amidase; acetylmuramyl-alanine amidase; *N*-acetylmuramylalanine amidase; murein hydrolase; *N*-acetylmuramoyl-L-alanine amidase type I; *N*-acetylmuramoyl-L-alanine amidase type II  
**Systematic name:** peptidoglycan amidohydrolase  
**References:** [908, 1127, 1126, 3126]



[EC 3.5.1.28 created 1972 (EC 3.4.19.10 created 1992, incorporated 1997)]

#### EC 3.5.1.29

**Accepted name:** 2-(acetamidomethylene)succinate hydrolase  
**Reaction:** 2-(acetamidomethylene)succinate + 2 H<sub>2</sub>O = acetate + succinate semialdehyde + NH<sub>3</sub> + CO<sub>2</sub>  
**Other name(s):** α-(*N*-acetylaminomethylene)succinic acid hydrolase  
**Systematic name:** 2-(acetamidomethylene)succinate amidohydrolase (deaminating, decarboxylating)  
**Comments:** Involved in the degradation of pyridoxin in *Pseudomonas*.  
**References:** [1212, 2123]

[EC 3.5.1.29 created 1972]

#### EC 3.5.1.30

**Accepted name:** 5-aminopentanamidase  
**Reaction:** 5-aminopentanamide + H<sub>2</sub>O = 5-aminopentanoate + NH<sub>3</sub>  
**Other name(s):** 5-aminovaleramidase; 5-aminonorvaleramidase  
**Systematic name:** 5-aminopentanamide amidohydrolase  
**Comments:** The enzyme from *Pseudomonas putida* also acts on 4-aminobutanamide and, more slowly, on 6-aminohexanamide.  
**References:** [2404, 2855]

[EC 3.5.1.30 created 1972, modified 1976]

#### EC 3.5.1.31

**Accepted name:** formylmethionine deformylase  
**Reaction:** *N*-formyl-L-methionine + H<sub>2</sub>O = formate + L-methionine  
**Systematic name:** *N*-formyl-L-methionine amidohydrolase  
**References:** [75]

[EC 3.5.1.31 created 1972]

#### EC 3.5.1.32

**Accepted name:** hippurate hydrolase  
**Reaction:** hippurate + H<sub>2</sub>O = benzoate + glycine  
**Systematic name:** *N*-benzoylamino-acid amidohydrolase  
**Comments:** Acts on various *N*-benzoylamino acids.  
**References:** [2452, 2453]

[EC 3.5.1.32 created 1972]

#### EC 3.5.1.33

**Accepted name:** *N*-acetylglucosamine deacetylase  
**Reaction:** *N*-acetyl-D-glucosamine + H<sub>2</sub>O = D-glucosamine + acetate  
**Other name(s):** acetylaminodeoxyglucose acetylhydrolase; *N*-acetyl-D-glucosaminyl *N*-deacetylase  
**Systematic name:** *N*-acetyl-D-glucosamine amidohydrolase  
**References:** [2459]

[EC 3.5.1.33 created 1972]

[3.5.1.34 Deleted entry. acetylhistidine deacetylase. Identical with EC 3.4.13.5, Xaa-methyl-His dipeptidase]

[EC 3.5.1.34 created 1972, deleted 1981]

### EC 3.5.1.35

**Accepted name:** D-glutaminase  
**Reaction:** D-glutamine + H<sub>2</sub>O = D-glutamate + NH<sub>3</sub>  
**Systematic name:** D-glutamine amidohydrolase  
**References:** [631]

[EC 3.5.1.35 created 1972]

### EC 3.5.1.36

**Accepted name:** N-methyl-2-oxoglutaramate hydrolase  
**Reaction:** N-methyl-2-oxoglutaramate + H<sub>2</sub>O = 2-oxoglutarate + methylamine  
**Other name(s):** 5-hydroxy-N-methylpyroglutamate synthase  
**Systematic name:** N-methyl-2-oxoglutaramate methylamidohydrolase  
**Comments:** In the reverse reaction, the product cyclizes non-enzymically to 2-hydroxy-N-methyl-5-oxo-L-proline.  
**References:** [1132, 1133]

[EC 3.5.1.36 created 1972]

[3.5.1.37 Deleted entry. 4-L-aspartylglycosylamine amidohydrolase. Identical with EC 3.5.1.26 N<sup>4</sup>-(β-N-acetylglucosaminyl)-L-asparaginase]

[EC 3.5.1.37 created 1972, deleted 1976]

### EC 3.5.1.38

**Accepted name:** glutamin-(asparagin-)ase  
**Reaction:** (1) L-glutamine + H<sub>2</sub>O = L-glutamate + NH<sub>3</sub>  
(2) L-asparagine + H<sub>2</sub>O = L-aspartate + NH<sub>3</sub>  
**Other name(s):** glutaminase-asparaginase; *ansB* (gene name); L-asparagine/L-glutamine amidohydrolase; L-ASNase/L-GLNase  
**Systematic name:** L-glutamine(L-asparagine) amidohydrolase  
**Comments:** The enzyme from the bacterium *Achromobacter* hydrolyses L-asparagine at 0.8 of the rate of L-glutamine; the D-isomers are also hydrolysed, but more slowly. *cf.* EC 3.5.1.2, glutaminase and EC 3.5.1.1, asparaginase.  
**References:** [2436, 2872, 1750, 2205]

[EC 3.5.1.38 created 1976]

### EC 3.5.1.39

**Accepted name:** alkylamidase  
**Reaction:** N-methylhexanamide + H<sub>2</sub>O = hexanoate + methylamine  
**Systematic name:** N-methylhexanamide amidohydrolase  
**Comments:** The enzyme hydrolyses N-monosubstituted and N,N-disubstituted amides, and there is some activity towards primary amides. It has little or no activity towards short-chain substrates.  
**References:** [416]

[EC 3.5.1.39 created 1976]

### EC 3.5.1.40

**Accepted name:** acylagmatine amidase  
**Reaction:** benzoylagmatine + H<sub>2</sub>O = benzoate + agmatine  
**Other name(s):** acylagmatine amidohydrolase; acylagmatine deacylase  
**Systematic name:** benzoylagmatine amidohydrolase  
**Comments:** Also acts on acetylglutamine, propanoylagmatine and bleomycin B2  
**References:** [3009]

[EC 3.5.1.40 created 1976]

**EC 3.5.1.41**

**Accepted name:** chitin deacetylase  
**Reaction:** chitin + H<sub>2</sub>O = chitosan + acetate  
**Systematic name:** chitin amidohydrolase  
**Comments:** Hydrolyses the *N*-acetamido groups of *N*-acetyl-D-glucosamine residues in chitin.  
**References:** [69]

[EC 3.5.1.41 created 1976]

**EC 3.5.1.42**

**Accepted name:** nicotinamide-nucleotide amidase  
**Reaction:** β-nicotinamide D-ribonucleotide + H<sub>2</sub>O = β-nicotinate D-ribonucleotide + NH<sub>3</sub>  
**Other name(s):** NMN deamidase; nicotinamide mononucleotide deamidase; nicotinamide mononucleotide amidohydrolase  
**Systematic name:** nicotinamide-D-ribonucleotide amidohydrolase  
**Comments:** Also acts more slowly on β-nicotinamide D-ribonucleoside.  
**References:** [1235]

[EC 3.5.1.42 created 1976]

**EC 3.5.1.43**

**Accepted name:** peptidyl-glutaminase  
**Reaction:** α-*N*-peptidyl-L-glutamine + H<sub>2</sub>O = α-*N*-peptidyl-L-glutamate + NH<sub>3</sub>  
**Other name(s):** peptidoglutaminase I; peptideglutaminase; peptidoglutaminase  
**Systematic name:** peptidyl-L-glutamine amidohydrolase  
**Comments:** Specific for the hydrolysis of the γ-amide of glutamine substituted at the α-amino group, *e.g.*, glycyl-L-glutamine, *N*-acetyl-L-glutamine and L-leucylglycyl-L-glutamine.  
**References:** [1440]

[EC 3.5.1.43 created 1976]

**EC 3.5.1.44**

**Accepted name:** protein-glutamine glutaminase  
**Reaction:** protein L-glutamine + H<sub>2</sub>O = protein L-glutamate + NH<sub>3</sub>  
**Other name(s):** peptidoglutaminase II; glutaminyl-peptide glutaminase; destabilase; peptidylglutaminase II  
**Systematic name:** protein-L-glutamine amidohydrolase  
**Comments:** Specific for the hydrolysis of the γ-amide of glutamine substituted at the carboxyl position or both the α-amino and carboxyl positions, *e.g.*, L-glutaminylglycine and L-phenylalanyl-L-glutaminylglycine.  
**References:** [1440]

[EC 3.5.1.44 created 1976, modified 1983]

[3.5.1.45 Deleted entry. urease (ATP-hydrolysing). Now listed only as EC 6.3.4.6 urea carboxylase]

[EC 3.5.1.45 created 1978, deleted 1986]

**EC 3.5.1.46**

**Accepted name:** 6-aminohexanoate-oligomer exohydrolase  
**Reaction:** (1) [N-(6-aminohexanoyl)]<sub>n</sub> + H<sub>2</sub>O = [N-(6-aminohexanoyl)]<sub>n-1</sub> + 6-aminohexanoate  
(2) N-(6-aminohexanoyl)-6-aminohexanoate + H<sub>2</sub>O = 2 6-aminohexanoate

**Other name(s):** 6-aminohexanoate-dimer hydrolase; *nylB* (gene name); 6-aminohexanoic acid oligomer hydrolase (ambiguous); *N*-(6-aminohexanoyl)-6-aminohexanoate amidohydrolase; nylon-6 hydrolase (ambiguous)

**Systematic name:** *N*-(6-aminohexanoyl)-6-aminohexanoate exoamidohydrolase

**Comments:** The enzyme is involved in degradation of nylon-6 oligomers. It degrades linear oligomers of 6-aminohexanoate with a degree of polymerization of 2–20 by exo-type cleavage, removing residues sequentially from the N-terminus. Activity decreases with the increase of the polymerization number of the oligomer. *cf.* EC 3.5.1.117, 6-aminohexanoate-oligomer endohydrolase and EC 3.5.2.12, 6-aminohexanoate-cyclic-dimer hydrolase.

**References:** [1461]

[EC 3.5.1.46 created 1983, modified 2014]

#### EC 3.5.1.47

**Accepted name:** *N*-acetyldiaminopimelate deacetylase

**Reaction:** *N*-acetyl-LL-2,6-diaminoheptanedioate + H<sub>2</sub>O = acetate + LL-2,6-diaminoheptanedioate

**Other name(s):** *N*-acetyl-L-diaminopimelic acid deacylase; *N*-acetyl-LL-diaminopimelate deacylase; 6-*N*-acetyl-LL-2,6-diaminoheptanedioate amidohydrolase

**Systematic name:** *N*<sup>6</sup>-acetyl-LL-2,6-diaminoheptanedioate amidohydrolase

**References:** [166, 2509, 2803]

[EC 3.5.1.47 created 1984 (EC 3.1.1.62 created 1989, incorporated 1992)]

#### EC 3.5.1.48

**Accepted name:** acetylspermidine deacetylase

**Reaction:** *N*<sup>8</sup>-acetylspermidine + H<sub>2</sub>O = acetate + spermidine

**Other name(s):** *N*<sup>8</sup>-monoacetylspermidine deacetylase; *N*<sup>8</sup>-acetylspermidine deacetylase; *N*-acetylspermidine deacetylase; *N*<sup>1</sup>-acetylspermidine amidohydrolase (incorrect); 8-*N*-acetylspermidine amidohydrolase

**Systematic name:** *N*<sup>8</sup>-acetylspermidine amidohydrolase

**Comments:** It was initially thought that *N*<sup>1</sup>-acetylspermidine was the substrate for this deacetylase reaction [1680] but this has since been disproved by Marchant et al. [1806].

**References:** [1680, 242, 1806]

[EC 3.5.1.48 created 1984, modified 2005]

#### EC 3.5.1.49

**Accepted name:** formamidase

**Reaction:** formamide + H<sub>2</sub>O = formate + NH<sub>3</sub>

**Systematic name:** formamide amidohydrolase

**Comments:** Also acts, more slowly, on acetamide, propanamide and butanamide.

**References:** [460, 821]

[EC 3.5.1.49 created 1984]

#### EC 3.5.1.50

**Accepted name:** pentanamidase

**Reaction:** pentanamide + H<sub>2</sub>O = pentanoate + NH<sub>3</sub>

**Other name(s):** valeramidase

**Systematic name:** pentanamide amidohydrolase

**Comments:** Also acts, more slowly, on other short-chain aliphatic amides. Different from EC 3.5.1.49 formamidase.

**References:** [821]

[EC 3.5.1.50 created 1984]

#### EC 3.5.1.51

**Accepted name:** 4-acetamidobutyryl-CoA deacetylase  
**Reaction:** 4-acetamidobutanoyl-CoA + H<sub>2</sub>O = acetate + 4-aminobutanoyl-CoA  
**Other name(s):** aminobutyryl-CoA thiolesterase; deacetylase-thiolesterase  
**Systematic name:** 4-acetamidobutanoyl-CoA amidohydrolase  
**Comments:** The enzyme also hydrolyses 4-aminobutanoyl-CoA to aminobutanoate and coenzyme A.  
**References:** [2168]

[EC 3.5.1.51 created 1984]

#### EC 3.5.1.52

**Accepted name:** peptide-*N*<sup>4</sup>-(*N*-acetyl-β-glucosaminyl)asparagine amidase  
**Reaction:** Hydrolysis of an *N*<sup>4</sup>-(acetyl-β-D-glucosaminyl)asparagine residue in which the glucosamine residue may be further glycosylated, to yield a (substituted) *N*-acetyl-β-D-glucosaminylamine and a peptide containing an aspartate residue  
**Other name(s):** glycopeptide *N*-glycosidase; glycopeptidase; *N*-oligosaccharide glycopeptidase; *N*-glycanase; Jack-bean glycopeptidase; PNGase A; PNGase F  
**Systematic name:** *N*-linked-glycopeptide-(*N*-acetyl-β-D-glucosaminyl)-*L*-asparagine amidohydrolase  
**Comments:** Does not act on (GlcNAc)Asn, because it requires the presence of more than two amino-acid residues in the substrate [*cf.* EC 3.5.1.26, *N*<sup>4</sup>-(β-*N*-acetylglucosaminyl)-*L*-asparaginase]. The plant enzyme was previously erroneously listed as EC 3.2.2.18.  
**References:** [2295, 2841, 2843, 2886]

[EC 3.5.1.52 created 1984, modified 1989 (EC 3.2.2.18 created 1984, incorporated 1989)]

#### EC 3.5.1.53

**Accepted name:** *N*-carbamoylputrescine amidase  
**Reaction:** *N*-carbamoylputrescine + H<sub>2</sub>O = putrescine + CO<sub>2</sub> + NH<sub>3</sub>  
**Other name(s):** carbamoylputrescine hydrolase; NCP  
**Systematic name:** *N*-carbamoylputrescine amidohydrolase  
**References:** [3253]

[EC 3.5.1.53 created 1986]

#### EC 3.5.1.54

**Accepted name:** allophanate hydrolase  
**Reaction:** urea-1-carboxylate + H<sub>2</sub>O = 2 CO<sub>2</sub> + 2 NH<sub>3</sub>  
**Other name(s):** allophanate lyase; AtzF; TrzF  
**Systematic name:** urea-1-carboxylate amidohydrolase  
**Comments:** Along with EC 3.5.2.15 (cyanuric acid amidohydrolase) and EC 3.5.1.84 (biuret amidohydrolase), this enzyme forms part of the cyanuric-acid metabolism pathway, which degrades *s*-triazide herbicides, such as atrazine [2-chloro-4-(ethylamino)-6-(isopropylamino)-1,3,5-triazine], in bacteria. The yeast enzyme (but not that from green algae) also catalyses the reaction of EC 6.3.4.6, urea carboxylase, thus bringing about the hydrolysis of urea to CO<sub>2</sub> and NH<sub>3</sub> in the presence of ATP and bicarbonate. The enzyme from *Pseudomonas* sp. strain ADP has a narrow substrate specificity, being unable to use the structurally analogous compounds urea, hydroxyurea or methylcarbamate as substrate [2621].  
**References:** [1781, 2456, 2800, 1377, 423, 2621, 2619]

[EC 3.5.1.54 created 1986, modified 2008]

#### EC 3.5.1.55

**Accepted name:** long-chain-fatty-acyl-glutamate deacylase  
**Reaction:** *N*-long-chain-fatty-acyl-L-glutamate + H<sub>2</sub>O = a long-chain carboxylate + L-glutamate  
**Other name(s):** long-chain aminoacylase; long-chain-fatty-acyl-glutamate deacylase; long-chain acylglutamate amidase; *N*-acyl-D-glutamate deacylase  
**Systematic name:** *N*-long-chain-fatty-acyl-L-glutamate amidohydrolase  
**Comments:** Does not act on acyl derivatives of other amino acids. Optimum chain length of acyl residue is 12 to 16.  
**References:** [857]

[EC 3.5.1.55 created 1986]

#### EC 3.5.1.56

**Accepted name:** *N,N*-dimethylformamidase  
**Reaction:** *N,N*-dimethylformamide + H<sub>2</sub>O = dimethylamine + formate  
**Other name(s):** dimethylformamidase; DMFase  
**Systematic name:** *N,N*-dimethylformamide amidohydrolase  
**Comments:** An iron protein. Also acts on *N*-ethylformamide and *N*-methylformamide and, more slowly, on *N,N*-diethylformamide, *N,N*-dimethylacetamide and unsubstituted acyl amides.  
**References:** [2554]

[EC 3.5.1.56 created 1989]

#### EC 3.5.1.57

**Accepted name:** tryptophanamidase  
**Reaction:** L-tryptophanamide + H<sub>2</sub>O = L-tryptophan + NH<sub>3</sub>  
**Other name(s):** tryptophan aminopeptidase; L-tryptophan aminopeptidase  
**Systematic name:** L-tryptophanamide amidohydrolase  
**Comments:** Requires Mn<sup>2+</sup>. Also acts on *N*-ethylformamide and L-tyrosinamide, and on some tryptophan dipeptides.  
**References:** [1283]

[EC 3.5.1.57 created 1989]

#### EC 3.5.1.58

**Accepted name:** *N*-benzyloxycarbonylglycine hydrolase  
**Reaction:** *N*-benzyloxycarbonylglycine + H<sub>2</sub>O = benzyl alcohol + CO<sub>2</sub> + glycine  
**Other name(s):** benzyloxycarbonylglycine hydrolase; *N*<sup>α</sup>-carbobenzoxyamino acid amidohydrolase; *N*<sup>α</sup>-benzyloxycarbonyl amino acid urethane hydrolase; *N*<sup>α</sup>-benzyloxycarbonyl amino acid urethane hydrolase I  
**Systematic name:** *N*-benzyloxycarbonylglycine urethanehydrolase  
**Comments:** Also acts, more slowly, on *N*-benzyloxycarbonylalanine, but not on the corresponding derivatives of other amino acids or on *N*-benzyloxycarbonylpeptides. Requires Co<sup>2+</sup> or Zn<sup>2+</sup>. *cf.* EC 3.5.1.64, *N*<sup>α</sup>-benzyloxycarbonylleucine hydrolase.  
**References:** [2005]

[EC 3.5.1.58 created 1989]

#### EC 3.5.1.59

**Accepted name:** *N*-carbamoylsarcosine amidase  
**Reaction:** *N*-carbamoylsarcosine + H<sub>2</sub>O = sarcosine + CO<sub>2</sub> + NH<sub>3</sub>  
**Other name(s):** carbamoylsarcosine amidase  
**Systematic name:** *N*-carbamoylsarcosine amidohydrolase  
**References:** [568]

[EC 3.5.1.59 created 1989]

**EC 3.5.1.60**

**Accepted name:** *N*-(long-chain-acyl)ethanolamine deacylase  
**Reaction:** *N*-(long-chain-acyl)ethanolamine + H<sub>2</sub>O = a long-chain carboxylate + ethanolamine  
**Other name(s):** *N*-acylethanolamine amidohydrolase; acylethanolamine amidase  
**Systematic name:** *N*-(long-chain-acyl)ethanolamine amidohydrolase  
**Comments:** Does not act on *N*-acylsphingosine or *N,O*-diacylethanolamine.  
**References:** [2569]

[EC 3.5.1.60 created 1989]

**EC 3.5.1.61**

**Accepted name:** mimosinase  
**Reaction:** (*S*)-2-amino-3-(3-hydroxy-4-oxo-4*H*-pyridin-1-yl)propanoate + H<sub>2</sub>O = 3-hydroxy-4*H*-pyrid-4-one + L-serine  
**Systematic name:** mimosine amidohydrolase  
**Comments:** An enzyme from *Leucaena leucocephala* leaf, which also contains the toxic amino acid, mimosine.  
**References:** [2882]

[EC 3.5.1.61 created 1989]

**EC 3.5.1.62**

**Accepted name:** acetylputrescine deacetylase  
**Reaction:** *N*-acetylputrescine + H<sub>2</sub>O = acetate + putrescine  
**Systematic name:** *N*-acetylputrescine acetylhydrolase  
**Comments:** The enzyme from *Micrococcus luteus* also acts on *N*<sup>8</sup>-acetylspermidine and acetylcadaverine, but more slowly.  
**References:** [2813]

[EC 3.5.1.62 created 1989]

**EC 3.5.1.63**

**Accepted name:** 4-acetamidobutyrate deacetylase  
**Reaction:** 4-acetamidobutanoate + H<sub>2</sub>O = acetate + 4-aminobutanoate  
**Systematic name:** 4-acetamidobutanoate amidohydrolase  
**Comments:** Also acts on *N*-acetyl-β-alanine and 5-acetamidopentanoate.  
**References:** [1090]

[EC 3.5.1.63 created 1989]

**EC 3.5.1.64**

**Accepted name:** *N*<sup>α</sup>-benzyloxycarbonylleucine hydrolase  
**Reaction:** *N*<sup>α</sup>-benzyloxycarbonyl-L-leucine + H<sub>2</sub>O = benzyl alcohol + CO<sub>2</sub> + L-leucine  
**Other name(s):** benzyloxycarbonylleucine hydrolase; *N*<sup>α</sup>-benzyloxycarbonyl amino acid urethane hydrolase IV; α-*N*-benzyloxycarbonyl-L-leucine urethanehydrolase  
**Systematic name:** *N*<sup>α</sup>-benzyloxycarbonyl-L-leucine urethanehydrolase  
**Comments:** Also acts on *N*<sup>α</sup>-*t*-butoxycarbonyl-L-leucine, and, more slowly, on the corresponding derivatives of L-aspartate, L-methionine, L-glutamate and L-alanine. *cf.* EC 3.5.1.58 *N*-benzyloxycarbonylglycine hydrolase.  
**References:** [1838]

[EC 3.5.1.64 created 1989]

**EC 3.5.1.65**

**Accepted name:** theanine hydrolase  
**Reaction:**  $N^5$ -ethyl-L-glutamine + H<sub>2</sub>O = L-glutamate + ethylamine  
**Other name(s):** L-theanine amidohydrolase; 5-*N*-ethyl-L-glutamine amidohydrolase  
**Systematic name:**  $N^5$ -ethyl-L-glutamine amidohydrolase  
**Comments:** Also acts on other *N*-alkyl-L-glutamines.  
**References:** [2991]

[EC 3.5.1.65 created 1989]

**EC 3.5.1.66**

**Accepted name:** 2-(hydroxymethyl)-3-(acetamidomethylene)succinate hydrolase  
**Reaction:** 2-(hydroxymethyl)-3-(acetamidomethylene)succinate + 2 H<sub>2</sub>O = acetate + 2-(hydroxymethyl)-4-oxobutanoate + NH<sub>3</sub> + CO<sub>2</sub>  
**Other name(s):** compound B hydrolase;  $\alpha$ -hydroxymethyl- $\alpha'$ -(*N*-acetylamino)methylene)succinic acid hydrolase  
**Systematic name:** 2-(hydroxymethyl)-3-(acetamidomethylene)succinate amidohydrolase (deaminating, decarboxylating)  
**Comments:** Involved in the degradation of pyridoxin by *Pseudomonas* and *Arthrobacter*.  
**References:** [1212]

[EC 3.5.1.66 created 1989]

**EC 3.5.1.67**

**Accepted name:** 4-methyleneglutaminase  
**Reaction:** 4-methylene-L-glutamine + H<sub>2</sub>O = 4-methylene-L-glutamate + NH<sub>3</sub>  
**Other name(s):** 4-methyleneglutamine deamidase; 4-methyleneglutamine amidohydrolase  
**Systematic name:** 4-methylene-L-glutamine amidohydrolase  
**References:** [1217]

[EC 3.5.1.67 created 1989]

**EC 3.5.1.68**

**Accepted name:** *N*-formylglutamate deformylase  
**Reaction:** *N*-formyl-L-glutamate + H<sub>2</sub>O = formate + L-glutamate  
**Other name(s):**  $\beta$ -citryl-L-glutamate hydrolase; formylglutamate deformylase; *N*-formylglutamate hydrolase;  $\beta$ -citrylglutamate amidase;  $\beta$ -citryl-L-glutamate amidohydrolase;  $\beta$ -citryl-L-glutamate amidase;  $\beta$ -citryl-L-glutamate-hydrolyzing enzyme  
**Systematic name:** *N*-formyl-L-glutamate amidohydrolase  
**Comments:** The animal enzyme also acts on  $\beta$ -citryl-L-glutamate and  $\beta$ -citryl-L-glutamine.  
**References:** [1199, 1937]

[EC 3.5.1.68 created 1989]

**EC 3.5.1.69**

**Accepted name:** glycosphingolipid deacylase  
**Reaction:** Hydrolysis of gangliosides and neutral glycosphingolipids, releasing fatty acids to form the lyso-derivatives  
**Other name(s):** glycosphingolipid ceramide deacylase  
**Systematic name:** glycosphingolipid amidohydrolase  
**Comments:** Does not act on sphingolipids such as ceramide. Not identical with EC 3.5.1.23 ceramidase.  
**References:** [1143]



[EC 3.5.1.69 created 1990]

#### EC 3.5.1.70

**Accepted name:** aculeacin-A deacylase  
**Reaction:** Hydrolysis of the amide bond in aculeacin A and related neutral lipopeptide antibiotics, releasing the long-chain fatty acid side-chain  
**Other name(s):** aculeacin A acylase  
**Systematic name:** aculeacin-A amidohydrolase  
**References:** [2859]

[EC 3.5.1.70 created 1992]

#### EC 3.5.1.71

**Accepted name:** *N*-feruloylglycine deacylase  
**Reaction:** *N*-feruloylglycine + H<sub>2</sub>O = ferulate + glycine  
**Other name(s):** *N*-feruloylglycine hydrolase  
**Systematic name:** *N*-feruloylglycine amidohydrolase  
**Comments:** Hydrolyses a range of L-amino acids from the cinnamoyl and substituted cinnamoyl series. Not identical with EC 3.5.1.14 aminoacylase.  
**References:** [1817, 1816]

[EC 3.5.1.71 created 1992]

#### EC 3.5.1.72

**Accepted name:** D-benzoylarginine-4-nitroanilide amidase  
**Reaction:** *N*-benzoyl-D-arginine-4-nitroanilide + H<sub>2</sub>O = *N*-benzoyl-D-arginine + 4-nitroaniline  
**Other name(s):** benzoyl-D-arginine arylamidase; D-BAPA-ase  
**Systematic name:** *N*-benzoyl-D-arginine-4-nitroanilide amidohydrolase  
**References:** [939]

[EC 3.5.1.72 created 1992]

#### EC 3.5.1.73

**Accepted name:** carnitinamidase  
**Reaction:** L-carnitinamide + H<sub>2</sub>O = L-carnitine + NH<sub>3</sub>  
**Other name(s):** L-carnitinamidase; carnitine amidase; L-carnitine amidase  
**Systematic name:** L-carnitinamide amidohydrolase  
**Comments:** Does not act on D-carnitinamide.  
**References:** [2051]

[EC 3.5.1.73 created 1992]

#### EC 3.5.1.74

**Accepted name:** chenodeoxycholoyltaurine hydrolase  
**Reaction:** chenodeoxycholoyltaurine + H<sub>2</sub>O = chenodeoxycholate + taurine  
**Systematic name:** chenodeoxycholoyltaurine amidohydrolase  
**Comments:** Some other taurine conjugates are hydrolysed, but not glycine conjugates of bile acids.  
**References:** [1406]

[EC 3.5.1.74 created 1992]

#### EC 3.5.1.75

**Accepted name:** urethanase  
**Reaction:** urethane + H<sub>2</sub>O = ethanol + CO<sub>2</sub> + NH<sub>3</sub>  
**Other name(s):** urethane hydrolase  
**Systematic name:** urethane amidohydrolase (decarboxylating)  
**References:** [1484]

[EC 3.5.1.75 created 1992]

#### EC 3.5.1.76

**Accepted name:** arylalkyl acylamidase  
**Reaction:** *N*-acetylarylalkylamine + H<sub>2</sub>O = arylalkylamine + acetate  
**Other name(s):** aralkyl acylamidase  
**Systematic name:** *N*-acetylarylalkylamine amidohydrolase  
**Comments:** Identified in *Pseudomonas putida*. Strict specificity for *N*-acetyl arylalkylamines, including *N*-acetyl-2-phenylethylamine, *N*-acetyl-3-phenylpropylamine, *N*-acetyldopamine, *N*-acetyl-serotonin and melatonin. It also accepts arylalkyl acetates but not acetanilide derivatives, which are common substrates of EC 3.5.1.13, aryl acylamidase.  
**References:** [2634]

[EC 3.5.1.76 created 1999]

#### EC 3.5.1.77

**Accepted name:** *N*-carbamoyl-D-amino-acid hydrolase  
**Reaction:** an *N*-carbamoyl-D-amino acid + H<sub>2</sub>O = a D-amino acid + NH<sub>3</sub> + CO<sub>2</sub>  
**Other name(s):** D-*N*-carbamoylase; *N*-carbamoylase (ambiguous); *N*-carbamoyl-D-amino acid hydrolase  
**Systematic name:** *N*-carbamoyl-D-amino-acid amidohydrolase  
**Comments:** This enzyme, along with EC 3.5.1.87 (*N*-carbamoyl-L-amino-acid hydrolase), EC 5.1.99.5 (hydantoin racemase) and hydantoinase, forms part of the reaction cascade known as the "hydantoinase process", which allows the total conversion of D,L-5-monosubstituted hydantoins into optically pure D- or L-amino acids [41]. It has strict stereospecificity for *N*-carbamoyl-D-amino acids and does not act upon the corresponding L-amino acids or on the *N*-formyl amino acids, *N*-carbamoyl-sarcosine, -citrulline, -allantoin and -ureidopropanoate, which are substrates for other amidohydrolases.  
**References:** [2146, 41]

[EC 3.5.1.77 created 1999, modified 2008]

#### EC 3.5.1.78

**Accepted name:** glutathionylspermidine amidase  
**Reaction:** glutathionylspermidine + H<sub>2</sub>O = glutathione + spermidine  
**Other name(s):** glutathionylspermidine amidohydrolase (spermidine-forming)  
**Systematic name:** γ-L-glutamyl-L-cysteinyl-glycine:spermidine amidase  
**Comments:** Spermidine is numbered so that atom N-1 is in the amino group of the aminopropyl part of the molecule. The enzyme from *Escherichia coli* is bifunctional and also catalyses the glutathionylspermidine synthase (EC 6.3.1.8) reaction, resulting in a net hydrolysis of ATP.  
**References:** [257]

[EC 3.5.1.78 created 1999]

#### EC 3.5.1.79

**Accepted name:** phthalyl amidase  
**Reaction:** a phthalylamide + H<sub>2</sub>O = phthalic acid + a substituted amine  
**Systematic name:** phthalyl-amide amidohydrolase

**Comments:** In the entry, "phthalyl" is used to mean "2-carboxybenzoyl". The enzyme from *Xanthobacter agilis* hydrolyses phthalylated amino acids, peptides,  $\beta$ -lactams, aromatic and aliphatic amines. The substituent on nitrogen may be an alkyl group, but may also be complex, giving an amino acid or peptide derivative. Substitutions on the phthalyl ring include 6-F, 6-NH<sub>2</sub>, 3-OH, and a nitrogen in the aromatic ring *ortho* to the carboxy group attached to the amine. No cofactors are required

**References:** [291, 230, 495, 290]

[EC 3.5.1.79 created 1999]

[3.5.1.80 Deleted entry. *N*-acetylgalactosamine-6-phosphate deacetylase. Identical to EC 3.5.1.25, *N*-acetylglucosamine-6-phosphate deacetylase]

[EC 3.5.1.80 created 1999, deleted 2002]

#### EC 3.5.1.81

**Accepted name:** *N*-acyl-D-amino-acid deacylase  
**Reaction:** *N*-acyl-D-amino acid + H<sub>2</sub>O = a carboxylate + D-amino acid  
**Systematic name:** *N*-acyl-D-amino acid amidohydrolase  
**Comments:** The enzyme from *Alcaligenes denitrificans* subsp. *xylosoxydans* and *Alcaligenes xylosoxydans* subsp. *xylosoxydans* has wide specificity; hydrolyses *N*-acyl derivative of neutral D-amino acids. Used in separating D- and L-amino acids. Requires zinc.  
**References:** [3088, 3087]

[EC 3.5.1.81 created 1999]

#### EC 3.5.1.82

**Accepted name:** *N*-acyl-D-glutamate deacylase  
**Reaction:** *N*-acyl-D-glutamate + H<sub>2</sub>O = a carboxylate + D-glutamate  
**Systematic name:** *N*-acyl-D-glutamate amidohydrolase  
**Comments:** The enzyme from *Alcaligenes xylosoxydans* subsp. *xylosoxydans* and *Pseudomonas* sp. is specific for *N*-acyl-D-glutamate. Requires zinc.  
**References:** [3086, 3089, 3090]

[EC 3.5.1.82 created 1999]

#### EC 3.5.1.83

**Accepted name:** *N*-acyl-D-aspartate deacylase  
**Reaction:** *N*-acyl-D-aspartate + H<sub>2</sub>O = a carboxylate + D-aspartate  
**Systematic name:** *N*-acyl-D-aspartate amidohydrolase  
**Comments:** The enzyme from *Alcaligenes xylosoxydans* subsp. *xylosoxydans* is specific for *N*-acyl-D-aspartate. Requires zinc.  
**References:** [1967, 3091]

[EC 3.5.1.83 created 1999]

#### EC 3.5.1.84

**Accepted name:** biuret amidohydrolase  
**Reaction:** biuret + H<sub>2</sub>O = urea-1-carboxylate + NH<sub>3</sub>  
**Systematic name:** biuret amidohydrolase

**Comments:** Along with EC 3.5.2.15 (cyanuric acid amidohydrolase) and EC 3.5.1.54 (allophanate hydrolase), this enzyme forms part of the cyanuric-acid metabolism pathway, which degrades *s*-triazide herbicides, such as atrazine [2-chloro-4-(ethylamino)-6-(isopropylamino)-1,3,5-triazine], in bacteria. Urea-1-carboxylate rather than urea (as was thought previously) is the 2-nitrogen intermediate in cyanuric-acid metabolism in bacteria [423]. The product, urea-1-carboxylate, can spontaneously decarboxylate under acidic conditions to form urea but, under physiological conditions, it can be converted into CO<sub>2</sub> and ammonia by the action of EC 3.5.1.54 [423].

**References:** [484, 423, 2621]

[EC 3.5.1.84 created 2000, modified 2008]

#### EC 3.5.1.85

**Accepted name:** (*S*)-*N*-acetyl-1-phenylethylamine hydrolase  
**Reaction:** *N*-acetylphenylethylamine + H<sub>2</sub>O = phenylethylamine + acetate  
**Systematic name:** (*S*)-*N*-acetylphenylethylamine:H<sub>2</sub>O hydrolase  
**Comments:** Inhibited by phenylmethanesulfonyl fluoride. Some related acetylated compounds are hydrolysed with variable enantiomeric selectivities.  
**References:** [317]

[EC 3.5.1.85 created 2000, modified 2002]

#### EC 3.5.1.86

**Accepted name:** mandelamide amidase  
**Reaction:** (*R*)-mandelamide + H<sub>2</sub>O = (*R*)-mandelate + NH<sub>3</sub>  
**Other name(s):** *Pseudomonas* mandelamide hydrolase  
**Systematic name:** mandelamide hydrolase  
**References:** [3249]

[EC 3.5.1.86 created 2000]

#### EC 3.5.1.87

**Accepted name:** *N*-carbamoyl-L-amino-acid hydrolase  
**Reaction:** an *N*-carbamoyl-L-2-amino acid (a 2-ureido carboxylate) + H<sub>2</sub>O = an L-2-amino acid + NH<sub>3</sub> + CO<sub>2</sub>  
**Other name(s):** *N*-carbamyl L-amino acid amidohydrolase; *N*-carbamoyl-L-amino acid amidohydrolase; L-*N*-carbamoylase; *N*-carbamoylase (ambiguous)  
**Systematic name:** *N*-carbamoyl-L-amino-acid amidohydrolase  
**Comments:** This enzyme, along with EC 3.5.1.77 (*N*-carbamoyl-D-amino-acid hydrolase), EC 5.1.99.5 (hydantoin racemase) and hydantoinase, forms part of the reaction cascade known as the "hydantoinase process", which allows the total conversion of D,L-5-monosubstituted hydantoins into optically pure D- or L-amino acids [41]. The enzyme from *Alcaligenes xylosoxidans* has broad specificity for carbamoyl-L-amino acids, although it is inactive on the carbamoyl derivatives of glutamate, aspartate, arginine, tyrosine or tryptophan. The enzyme from *Sinorhizobium meliloti* requires a divalent cation for activity and can hydrolyse *N*-carbamoyl-L-tryptophan as well as *N*-carbamoyl L-amino acids with aliphatic substituents [1822]. The enzyme is inactive on derivatives of D-amino acids. In addition to *N*-carbamoyl L-amino acids, the enzyme can also hydrolyse formyl and acetyl derivatives to varying degrees [2145, 1822].  
**References:** [2145, 1822, 41]

[EC 3.5.1.87 created 2001, modified 2008]

#### EC 3.5.1.88

**Accepted name:** peptide deformylase  
**Reaction:** formyl-L-methionyl peptide + H<sub>2</sub>O = formate + methionyl peptide

**Other name(s):** *N*-formylmethionylaminoacyl-tRNA deformylase  
**Systematic name:** formyl-L-methionyl peptide amidohydrolase  
**Comments:** Requires Fe(II). Also requires at least a dipeptide for an efficient rate of reaction. N-terminal L-methionine is a prerequisite for activity but the enzyme has broad specificity at other positions. Differs in substrate specificity from EC 3.5.1.31 (formylmethionine deformylase).  
**References:** [15, 1855, 391, 182, 181, 2361, 986, 2360, 1200, 2356, 916, 2252]

[EC 3.5.1.88 created 2001]

#### EC 3.5.1.89

**Accepted name:** *N*-acetylglucosaminylphosphatidylinositol deacetylase  
**Reaction:** 6-(*N*-acetyl- $\alpha$ -D-glucosaminy)-1-phosphatidyl-1D-*myo*-inositol + H<sub>2</sub>O = 6-( $\alpha$ -D-glucosaminy)-1-phosphatidyl-1D-*myo*-inositol + acetate  
**Other name(s):** *N*-acetyl-D-glucosaminylphosphatidylinositol acetylhydrolase; *N*-acetylglucosaminylphosphatidylinositol de-*N*-acetylase; GlcNAc-PI de-*N*-acetylase; GlcNAc-PI deacetylase; acetylglucosaminylphosphatidylinositol deacetylase  
**Systematic name:** 6-(*N*-acetyl- $\alpha$ -D-glucosaminy)-1-phosphatidyl-1D-*myo*-inositol acetylhydrolase  
**Comments:** Involved in the second step of glycosylphosphatidylinositol (GPI) anchor formation in all eukaryotes. The enzyme appears to be composed of a single subunit (PIG-L in mammalian cells and GPI12 in yeast). In some species, the long-chain *sn*-1-acyl group of the phosphatidyl group is replaced by a long-chain alkyl or alk-1-enyl group.  
**References:** [626, 2045, 3133, 2702]

[EC 3.5.1.89 created 1992 as EC 3.1.1.69, transferred 2002 to EC 3.5.1.89, modified 2002]

#### EC 3.5.1.90

**Accepted name:** adenosylcobinamide hydrolase  
**Reaction:** adenosylcobinamide + H<sub>2</sub>O = adenosylcobyric acid + (*R*)-1-aminopropan-2-ol  
**Other name(s):** CbiZ; AdoCbi amidohydrolase  
**Systematic name:** adenosylcobinamide amidohydrolase  
**Comments:** Involved in the salvage pathway of cobinamide in archaea. *Archaea* convert adenosylcobinamide (AdoCbi) into adenosylcobinamide phosphate (AdoCbi-P) in two steps. First, the amidohydrolase activity of CbiZ cleaves off the aminopropanol moiety of AdoCbi yielding adenosylcobyric acid (AdoCby); second, AdoCby is converted into AdoCbi-P by the action of EC 6.3.1.10, adenosylcobinamide-phosphate synthase (CbiB).  
**References:** [3213]

[EC 3.5.1.90 created 2004]

#### EC 3.5.1.91

**Accepted name:** *N*-substituted formamide deformylase  
**Reaction:** *N*-benzylformamide + H<sub>2</sub>O = formate + benzylamine  
**Other name(s):** NfdA  
**Systematic name:** *N*-benzylformamide amidohydrolase  
**Comments:** Zinc is a cofactor. While *N*-benzylformamide is the best substrate, the enzyme from *Arthrobacter pascens* can also act on the *N*-substituted formamides *N*-butylformamide, *N*-allylformamide, *N*-[2-(cyclohex-1-enyl)ethyl]formamide and *N*-(1-phenylethyl)formamide, but much more slowly. Amides of other acids do not act as substrates.  
**References:** [856]

[EC 3.5.1.91 created 2005]

#### EC 3.5.1.92

**Accepted name:** pantetheine hydrolase  
**Reaction:** (*R*)-pantetheine + H<sub>2</sub>O = (*R*)-pantothenate + 2-aminoethanethiol  
**Other name(s):** pantetheinase; vanin; vanin-1  
**Systematic name:** (*R*)-pantetheine amidohydrolase  
**Comments:** The enzyme hydrolyses only one of the amide bonds of pantetheine. The substrate analogues phosphopantetheine and CoA are not substrates. The enzyme recycles pantothenate (vitamin B<sub>5</sub>) and produces 2-aminoethanethiol (cysteamine), a potent anti-oxidant [2287].  
**References:** [669, 670, 1805, 103, 2287, 1818, 2220]

[EC 3.5.1.92 created 2006]

#### EC 3.5.1.93

**Accepted name:** glutaryl-7-aminocephalosporanic-acid acylase  
**Reaction:** (*7R*)-7-(4-carboxybutanamido)cephalosporanate + H<sub>2</sub>O = (*7R*)-7-aminocephalosporanate + glutarate  
**Other name(s):** 7β-(4-carboxybutanamido)cephalosporanic acid acylase; cephalosporin C acylase; glutaryl-7-ACA acylase; CA; GCA; GA; cephalosporin acylase; glutaryl-7-aminocephalosporanic acid acylase; GL-7-ACA acylase  
**Systematic name:** (*7R*)-7-(4-carboxybutanamido)cephalosporanate amidohydrolase  
**Comments:** Forms 7-aminocephalosporanic acid, a key intermediate in the synthesis of cephem antibiotics. It reacts only weakly with cephalosporin C.  
**References:** [1262, 1462, 1955, 1581, 1456, 1204, 1449]

[EC 3.5.1.93 created 2005]

#### EC 3.5.1.94

**Accepted name:** γ-glutamyl-γ-aminobutyrate hydrolase  
**Reaction:** 4-(γ-L-glutamylamino)butanoate + H<sub>2</sub>O = 4-aminobutanoate + L-glutamate  
**Other name(s):** γ-glutamyl-GABA hydrolase; PuuD; YcjL; 4-(γ-glutamylamino)butanoate amidohydrolase; 4-(L-γ-glutamylamino)butanoate amidohydrolase  
**Systematic name:** 4-(γ-L-glutamylamino)butanoate amidohydrolase  
**Comments:** Forms part of a putrescine-utilizing pathway in *Escherichia coli*, in which it has been hypothesized that putrescine is first glutamylated to form γ-glutamylputrescine, which is oxidized to 4-(γ-glutamylamino)butanal and then to 4-(γ-glutamylamino)butanoate. The enzyme can also catalyse the reactions of EC 3.5.1.35 (D-glutaminase) and EC 3.5.1.65 (theanine hydrolase).  
**References:** [1566]

[EC 3.5.1.94 created 2006, modified 2011]

#### EC 3.5.1.95

**Accepted name:** *N*-malonylurea hydrolase  
**Reaction:** 3-oxo-3-ureidopropanoate + H<sub>2</sub>O = malonate + urea  
**Other name(s):** ureidomalonase  
**Systematic name:** 3-oxo-3-ureidopropanoate amidohydrolase (urea- and malonate-forming)  
**Comments:** Forms part of the oxidative pyrimidine-degrading pathway in some microorganisms, along with EC 1.17.99.4 (uracil/thymine dehydrogenase) and EC 3.5.2.1 (barbiturase).  
**References:** [2725, 2724]

[EC 3.5.1.95 created 2006]

#### EC 3.5.1.96

**Accepted name:** succinylglutamate desuccinylase  
**Reaction:** *N*-succinyl-L-glutamate + H<sub>2</sub>O = succinate + L-glutamate  
**Other name(s):** *N*<sup>2</sup>-succinylglutamate desuccinylase; SGDS; AstE

**Systematic name:** *N*-succinyl-L-glutamate amidohydrolase  
**Comments:** Requires Co<sup>2+</sup> for maximal activity [3139]. *N*<sup>2</sup>-Acetylglutamate is not a substrate. This is the final enzyme in the arginine succinyltransferase (AST) pathway for the catabolism of arginine [3139]. This pathway converts the carbon skeleton of arginine into glutamate, with the concomitant production of ammonia and conversion of succinyl-CoA into succinate and CoA. The five enzymes involved in this pathway are EC 2.3.1.109 (arginine *N*-succinyltransferase), EC 3.5.3.23 (*N*-succinylarginine dihydrolase), EC 2.6.1.11 (acetylornithine transaminase), EC 1.2.1.71 (succinylglutamate-semialdehyde dehydrogenase) and EC 3.5.1.96 (succinylglutamate desuccinylase).  
**References:** [3139, 517, 518, 1276, 2575]

[EC 3.5.1.96 created 2006]

#### EC 3.5.1.97

**Accepted name:** acyl-homoserine-lactone acylase  
**Reaction:** an *N*-acyl-L-homoserine lactone + H<sub>2</sub>O = L-homoserine lactone + a carboxylate  
**Other name(s):** acyl-homoserine lactone acylase; AHL-acylase; AiiD; *N*-acyl-homoserine lactone acylase; PA2385 protein; quorum-quenching AHL acylase; quorum-quenching enzyme; QuiP  
**Systematic name:** *N*-acyl-L-homoserine-lactone amidohydrolase  
**Comments:** Acyl-homoserine lactones (AHLs) are produced by a number of bacterial species and are used by them to regulate the expression of virulence genes in a process known as quorum-sensing. Each bacterial cell has a basal level of AHL and, once the population density reaches a critical level, it triggers AHL-signalling which, in turn, initiates the expression of particular virulence genes. Plants or animals capable of degrading AHLs would have a therapeutic advantage in avoiding bacterial infection as they could prevent AHL-signalling and the expression of virulence genes in quorum-sensing bacteria. This quorum-quenching enzyme removes the fatty-acid side chain from the homoserine lactone ring of AHL-dependent quorum-sensing signal molecules. It has broad specificity for AHLs with side chains ranging in length from 11 to 14 carbons. Substituents at the 3'-position, as found in *N*-(3-oxododecanoyl)-L-homoserine lactone, do not affect this activity.  
**References:** [1697, 2674]

[EC 3.5.1.97 created 2007]

#### EC 3.5.1.98

**Accepted name:** histone deacetylase  
**Reaction:** Hydrolysis of an *N*<sup>6</sup>-acetyl-lysine residue of a histone to yield a deacetylated histone  
**Other name(s):** HDAC  
**Systematic name:** histone amidohydrolase  
**Comments:** A class of enzymes that remove acetyl groups from *N*<sup>6</sup>-acetyl-lysine residues on a histone. The reaction of this enzyme is opposite to that of EC 2.3.1.48, histone acetyltransferase. Histone deacetylases (HDACs) can be organized into three classes, HDAC1, HDAC2 and HDAC3, depending on sequence similarity and domain organization. Histone acetylation plays an important role in regulation of gene expression. In eukaryotes, HDACs play a key role in the regulation of transcription and cell proliferation [2722]. May be identical to EC 3.5.1.17, acyl-lysine deacetylase.  
**References:** [1535, 572, 2217, 2722, 770, 2274, 560]

[EC 3.5.1.98 created 2008]

#### EC 3.5.1.99

**Accepted name:** fatty acid amide hydrolase  
**Reaction:** (1) anandamide + H<sub>2</sub>O = arachidonic acid + ethanolamine  
(2) oleamide + H<sub>2</sub>O = oleic acid + NH<sub>3</sub>  
**Other name(s):** FAAH; oleamide hydrolase; anandamide amidohydrolase  
**Systematic name:** fatty acylamide amidohydrolase



**Comments:** Integral membrane protein, the enzyme is responsible for the catabolism of neuromodulatory fatty acid amides, including anandamide and oleamide, occurs in mammalia.

**References:** [253, 2245, 2244]

[EC 3.5.1.99 created 2009]

#### EC 3.5.1.100

**Accepted name:** (*R*)-amidase

**Reaction:** (1) (*R*)-piperazine-2-carboxamide + H<sub>2</sub>O = (*R*)-piperazine-2-carboxylate + NH<sub>3</sub>  
(2) β-alaninamide + H<sub>2</sub>O = β-alanine + NH<sub>3</sub>

**Other name(s):** R-stereospecific amidase; R-amidase

**Systematic name:** (*R*)-piperazine-2-carboxamide amidohydrolase

**Comments:** In addition (*R*)-piperidine-3-carboxamide is hydrolysed to (*R*)-piperidine-3-carboxylic acid and NH<sub>3</sub>, and (*R*)-*N*-*tert*-butylpiperazine-2-carboxamide is hydrolysed to (*R*)-piperazine-2-carboxylic acid and *tert*-butylamine with lower activity. The enzyme does not act on the other amide substrates which are hydrolysed by EC 3.5.1.4 (amidase).

**References:** [1502]

[EC 3.5.1.100 created 2009, modified 2011]

#### EC 3.5.1.101

**Accepted name:** L-proline amide hydrolase

**Reaction:** (1) (*S*)-piperidine-2-carboxamide + H<sub>2</sub>O = (*S*)-piperidine-2-carboxylate + NH<sub>3</sub>  
(2) L-prolinamide + H<sub>2</sub>O = L-proline + NH<sub>3</sub>

**Other name(s):** *S*-stereoselective piperazine-2-*tert*-butylcarboxamide hydrolase; LaaA; L-amino acid amidase

**Systematic name:** (*S*)-piperidine-2-carboxamide amidohydrolase

**References:** [1503]

[EC 3.5.1.101 created 2009]

#### EC 3.5.1.102

**Accepted name:** 2-amino-5-formylamino-6-ribosylaminopyrimidin-4(3*H*)-one 5'-monophosphate deformylase

**Reaction:** 2-amino-5-formylamino-6-(5-phospho-D-ribosylamino)pyrimidin-4(3*H*)-one + H<sub>2</sub>O = 2,5-diamino-6-(5-phospho-D-ribosylamino)pyrimidin-4(3*H*)-one + formate

**Other name(s):** ArfB

**Systematic name:** 2-amino-5-formylamino-6-(5-phospho-D-ribosylamino)pyrimidin-4(3*H*)-one amidohydrolase

**Comments:** The enzyme catalyses the second step in archaeal riboflavin and 7,8-didemethyl-8-hydroxy-5-deazariboflavin biosynthesis. The first step is catalysed by EC 3.5.4.29 (GTP cyclohydrolase IIa). The bacterial enzyme, EC 3.5.4.25 (GTP cyclohydrolase II) catalyses both reactions.

**References:** [988]

[EC 3.5.1.102 created 2010, modified 2011]

#### EC 3.5.1.103

**Accepted name:** *N*-acetyl-1-*D*-*myo*-inositol-2-amino-2-deoxy-α-*D*-glucopyranoside deacetylase

**Reaction:** 1-*O*-(2-acetamido-2-deoxy-α-*D*-glucopyranosyl)-1*D*-*myo*-inositol + H<sub>2</sub>O = 1-*O*-(2-amino-2-deoxy-α-*D*-glucopyranosyl)-1*D*-*myo*-inositol + acetate

**Other name(s):** MshB

**Systematic name:** 1-(2-acetamido-2-deoxy-α-*D*-glucopyranosyl)-1*D*-*myo*-inositol acetylhydrolase

**Comments:** This enzyme is considered the key enzyme and rate limiting step in the mycothiol biosynthesis pathway [2380]. In addition to acetylase activity, the enzyme possesses weak activity of EC 3.5.1.115, mycothiol *S*-conjugate amidase, and shares sequence similarity with that enzyme [2076]. The enzyme requires a divalent transition metal ion for activity, believed to be Zn<sup>2+</sup> [1854].

**References:** [2380, 2076, 1854]



[EC 3.5.1.103 created 2010]

#### EC 3.5.1.104

- Accepted name:** peptidoglycan-*N*-acetylglucosamine deacetylase  
**Reaction:** peptidoglycan-*N*-acetyl-D-glucosamine + H<sub>2</sub>O = peptidoglycan-D-glucosamine + acetate  
**Other name(s):** HP310; PgdA; SpPgdA; BC1960; peptidoglycan deacetylase; *N*-acetylglucosamine deacetylase; peptidoglycan GlcNAc deacetylase; peptidoglycan *N*-acetylglucosamine deacetylase; PG *N*-deacetylase  
**Systematic name:** peptidoglycan-*N*-acetylglucosamine amidohydrolase  
**Comments:** Modification of peptidoglycan by *N*-deacetylation is an important factor in virulence of *Helicobacter pylori*, *Listeria monocytogenes* and *Streptococcus suis* [3108, 2312, 778]. The enzyme from *Streptococcus pneumoniae* is a metalloenzyme using a His-His-Asp zinc-binding triad with a nearby aspartic acid and histidine acting as the catalytic base and acid, respectively [233].  
**References:** [2328, 2970, 233, 3108, 2312, 778]

[EC 3.5.1.104 created 2010]

#### EC 3.5.1.105

- Accepted name:** chitin disaccharide deacetylase  
**Reaction:** *N,N'*-diacetylchitobiose + H<sub>2</sub>O = *N*-acetyl-β-D-glucosaminy1-(1→4)-D-glucosamine + acetate  
**Other name(s):** chitobiose amidohydrolase; COD; chitin oligosaccharide deacetylase; chitin oligosaccharide amidohydrolase; 2-(acetylamino)-4-*O*-[2-(acetylamino)-2-deoxy-β-D-glucopyranosyl]-2-deoxy-β-D-glucopyranose acetylhydrolase  
**Systematic name:** *N,N'*-diacetylchitobiose acetylhydrolase  
**Comments:** Chitin oligosaccharide deacetylase is a key enzyme in the chitin catabolic cascade of chitinolytic *Vibrio* strains. Besides being a nutrient, the heterodisaccharide product 4-*O*-(*N*-acetyl-β-D-glucosaminy1)-D-glucosamine is a unique inducer of chitinase production in *Vibrio parahaemolyticus* [1148]. In contrast to EC 3.5.1.41 (chitin deacetylase) this enzyme is specific for the chitin disaccharide [1352, 2160]. It also deacetylates the chitin trisaccharide with lower efficiency [2160]. No activity with higher polymers of GlcNAc [1352, 2160].  
**References:** [1352, 1148, 2160, 2159]

[EC 3.5.1.105 created 2010]

#### EC 3.5.1.106

- Accepted name:** *N*-formylmaleamate deformylase  
**Reaction:** *N*-formylmaleamic acid + H<sub>2</sub>O = maleamate + formate  
**Other name(s):** NicD  
**Systematic name:** *N*-formylmaleamic acid amidohydrolase  
**Comments:** The reaction is involved in the aerobic catabolism of nicotinic acid.  
**References:** [1322]

[EC 3.5.1.106 created 2010]

#### EC 3.5.1.107

- Accepted name:** maleamate amidohydrolase  
**Reaction:** maleamate + H<sub>2</sub>O = maleate + NH<sub>3</sub>  
**Other name(s):** NicF  
**Systematic name:** maleamate amidohydrolase  
**Comments:** The reaction is involved in the aerobic catabolism of nicotinic acid.  
**References:** [1322]

[EC 3.5.1.107 created 2010]

### EC 3.5.1.108

- Accepted name:** UDP-3-*O*-acyl-*N*-acetylglucosamine deacetylase  
**Reaction:** UDP-3-*O*-[(3*R*)-3-hydroxymyristoyl]-*N*-acetyl- $\alpha$ -D-glucosamine + H<sub>2</sub>O = UDP-3-*O*-[(3*R*)-3-hydroxymyristoyl]- $\alpha$ -D-glucosamine + acetate  
**Other name(s):** LpxC protein; LpxC enzyme; LpxC deacetylase; deacetylase LpxC; UDP-3-*O*-acyl-GlcNAc deacetylase; UDP-3-*O*-((*R*)-3-hydroxymyristoyl)-*N*-acetylglucosamine deacetylase; UDP-(3-*O*-acyl)-*N*-acetylglucosamine deacetylase; UDP-3-*O*-(*R*-3-hydroxymyristoyl)-*N*-acetylglucosamine deacetylase; UDP-(3-*O*-(*R*-3-hydroxymyristoyl))-*N*-acetylglucosamine deacetylase; UDP-3-*O*-[(3*R*)-3-hydroxymyristoyl]-*N*-acetylglucosamine amidohydrolase  
**Systematic name:** UDP-3-*O*-[(3*R*)-3-hydroxymyristoyl]-*N*-acetyl- $\alpha$ -D-glucosamine amidohydrolase  
**Comments:** A zinc protein. The enzyme catalyses a committed step in the biosynthesis of lipid A.  
**References:** [1131, 1288, 1214, 3122, 3176, 1943]

[EC 3.5.1.108 created 2010]

### EC 3.5.1.109

- Accepted name:** sphingomyelin deacylase  
**Reaction:** (1) an *N*-acyl-sphingosylphosphorylcholine + H<sub>2</sub>O = a fatty acid + sphingosylphosphorylcholine  
(2) a D-glucosyl-*N*-acylsphingosine + H<sub>2</sub>O = a fatty acid + D-glucosyl-sphingosine  
**Other name(s):** SM deacylase; GcSM deacylase; glucosylceramide sphingomyelin deacylase; sphingomyelin glucosylceramide deacylase; SM glucosylceramide GCer deacylase; SM-GCer deacylase; SMGCer deacylase  
**Systematic name:** *N*-acyl-sphingosylphosphorylcholine amidohydrolase  
**Comments:** The enzyme is involved in the sphingolipid metabolism in the epidermis.  
**References:** [1048, 1139, 1258]

[EC 3.5.1.109 created 2011]

### EC 3.5.1.110

- Accepted name:** peroxyureidoacrylate/ureidoacrylate amidohydrolase  
**Reaction:** (1) (*Z*)-3-ureidoacrylate peracid + H<sub>2</sub>O = (*Z*)-3-peroxyaminoacrylate + CO<sub>2</sub> + NH<sub>3</sub> (overall reaction)  
(1a) (*Z*)-3-ureidoacrylate peracid + H<sub>2</sub>O = (*Z*)-3-peroxyaminoacrylate + carbamate  
(1b) carbamate = CO<sub>2</sub> + NH<sub>3</sub> (spontaneous)  
(2) (*Z*)-2-methylureidoacrylate peracid + H<sub>2</sub>O = (*Z*)-2-methylperoxyaminoacrylate + CO<sub>2</sub> + NH<sub>3</sub> (overall reaction)  
(2a) (*Z*)-2-methylureidoacrylate peracid + H<sub>2</sub>O = (*Z*)-2-methylperoxyaminoacrylate + carbamate  
(2b) carbamate = CO<sub>2</sub> + NH<sub>3</sub> (spontaneous)  
**Other name(s):** RutB  
**Systematic name:** (*Z*)-3-ureidoacrylate peracid amidohydrolase  
**Comments:** The enzyme also shows activity towards ureidoacrylate. Part of the Rut pyrimidine catabolic pathway.  
**References:** [1452]

[EC 3.5.1.110 created 2012]

### EC 3.5.1.111

- Accepted name:** 2-oxoglutaramate amidase  
**Reaction:** 2-oxoglutaramate + H<sub>2</sub>O = 2-oxoglutarate + NH<sub>3</sub>  
**Other name(s):**  $\omega$ -amidase (ambiguous)  
**Systematic name:** 5-amino-2,5-dioxopentanoate amidohydrolase  
**Comments:** The enzyme, which is highly specific for its substrate, participates in the nicotine degradation pathway of several Gram-positive bacteria.  
**References:** [462]

[EC 3.5.1.111 created 2012]

#### EC 3.5.1.112

- Accepted name:** 2'-*N*-acetylparomamine deacetylase  
**Reaction:** 2'-*N*-acetylparomamine + H<sub>2</sub>O = paromamine + acetate  
**Other name(s):** *btrD* (gene name); *neoL* (gene name); *kanN* (gene name)  
**Systematic name:** 2'-*N*-acetylparomamine hydrolase (acetate-forming)  
**Comments:** Involved in the biosynthetic pathways of several clinically important aminocyclitol antibiotics, including kanamycin, butirosin, neomycin and ribostamycin. The enzyme from the bacterium *Streptomyces fradiae* can also accept 2'''-acetyl-6'''-hydroxyneomycin C as substrate, *cf.* EC 3.5.1.113, 2'''-acetyl-6'''-hydroxyneomycin C deacetylase [3288].  
**References:** [2967, 3288]

[EC 3.5.1.112 created 2012]

#### EC 3.5.1.113

- Accepted name:** 2'''-acetyl-6'''-hydroxyneomycin C deacetylase  
**Reaction:** 2'''-acetyl-6'''-deamino-6'''-hydroxyneomycin C + H<sub>2</sub>O = 6'''-deamino-6'''-hydroxyneomycin C + acetate  
**Other name(s):** *neoL* (gene name)  
**Systematic name:** 2'''-acetyl-6'''-hydroxyneomycin C hydrolase (acetate-forming)  
**Comments:** Involved in the biosynthetic pathway of aminoglycoside antibiotics of the neomycin family. The enzyme from the bacterium *Streptomyces fradiae* also catalyses EC 3.5.1.112, 2'-*N*-acetylparomamine deacetylase.  
**References:** [3288]

[EC 3.5.1.113 created 2012]

#### EC 3.5.1.114

- Accepted name:** *N*-acyl-aromatic-L-amino acid amidohydrolase  
**Reaction:** (1) an *N*-acyl-aromatic-L-amino acid + H<sub>2</sub>O = an aromatic-L-amino acid + a carboxylate  
(2) an *N*-acetyl-L-cysteine-*S*-conjugate + H<sub>2</sub>O = an L-cysteine-*S*-conjugate + acetate  
**Other name(s):** aminoacylase 3; aminoacylase III; ACY3 (gene name)  
**Systematic name:** *N*-acyl-aromatic-L-amino acid amidohydrolase (carboxylate-forming)  
**Comments:** This enzyme is found in animals and is involved in the hydrolysis of *N*-acylated or *N*-acetylated amino acids (except L-aspartate). It preferentially deacetylates *N*<sup>α</sup>-acetylated aromatic amino acids and mercapturic acids (*S*-conjugates of *N*-acetyl-L-cysteine) that are usually not deacetylated by EC 3.5.1.14, *N*-acyl-aliphatic-L-amino acid amidohydrolase. The enzyme is significantly activated by Co<sup>2+</sup> and Ni<sup>2+</sup> [2975]. Some bacterial aminoacylases demonstrate substrate specificity for both EC 3.5.1.14 and EC 3.5.1.114. *cf.* EC 3.5.1.14, *N*-acyl-aliphatic-L-amino acid amidohydrolase and EC 3.5.1.15, aspartoacylase.  
**References:** [2334, 2072, 2975, 1196, 2974]

[EC 3.5.1.114 created 2013]

#### EC 3.5.1.115

- Accepted name:** mycothiol *S*-conjugate amidase  
**Reaction:** a mycothiol *S*-conjugate + H<sub>2</sub>O = an *N*-acetyl L-cysteine-*S*-conjugate + 1-*O*-(2-amino-2-deoxy-α-D-glucopyranosyl)-1D-*myo*-inositol  
**Other name(s):** MCA  
**Systematic name:** mycothiol *S*-conjugate 1D-*myo*-inositol 2-amino-2-deoxy-α-D-glucopyranosyl-hydrolase  
**Comments:** The enzyme that is found in actinomycetes is involved in the detoxification of oxidizing agents and electrophilic antibiotics. The enzyme has low activity with 1-*O*-(2-acetamido-2-deoxy-α-D-glucopyranosyl)-1D-*myo*-inositol as substrate (*cf.* EC 3.5.1.103, *N*-acetyl-1-D-*myo*-inositol-2-amino-2-deoxy-α-D-glucopyranoside deacetylase) [2759].  
**References:** [2075, 2759]

[EC 3.5.1.115 created 2013]

#### EC 3.5.1.116

**Accepted name:** ureidoglycolate amidohydrolase  
**Reaction:** (*S*)-ureidoglycolate + H<sub>2</sub>O = glyoxylate + 2 NH<sub>3</sub> + CO<sub>2</sub>  
**Other name(s):** ureidoglycolate hydrolase; UAH (gene name)  
**Systematic name:** (*S*)-ureidoglycolate amidohydrolase (decarboxylating)  
**Comments:** This plant enzyme is involved in the degradation of ureidoglycolate, an intermediate of purine degradation. Not to be confused with EC 4.3.2.3, ureidoglycolate lyase, which releases urea rather than ammonia.  
**References:** [3197, 3161, 3162]

[EC 3.5.1.116 created 1992 as EC 3.5.3.19, transferred 2014 to EC 3.5.1.116]

#### EC 3.5.1.117

**Accepted name:** 6-aminohexanoate-oligomer endohydrolase  
**Reaction:** [*N*-(6-aminohexanoyl)]<sub>n</sub> + H<sub>2</sub>O = [*N*-(6-aminohexanoyl)]<sub>n-x</sub> + [*N*-(6-aminohexanoyl)]<sub>x</sub>  
**Other name(s):** endo-type 6-aminohexanoate oligomer hydrolase; Ahx endo-type-oligomer hydrolase; 6-aminohexanoate oligomer hydrolase; Ahx-oligomer hydrolase; nylon hydrolase; nylon-oligomer hydrolase; NylC; nylon-6 hydrolase (ambiguous)  
**Systematic name:** 6-aminohexanoate oligomer endoamidohydrolase  
**Comments:** The enzyme is involved in degradation of nylon-6 oligomers. It degrades linear or cyclic oligomers of poly(6-aminohexanoate) with a degree of polymerization greater than three ( $n > 3$ ) by endo-type cleavage, to oligomers of a length of two or more ( $2 \leq x < n$ ). It shows negligible activity with *N*-(6-aminohexanoyl)-6-aminohexanoate (*cf.* EC 3.5.1.46, 6-aminohexanoate-oligomer exo hydrolase) or with 1,8-diazacyclotetradecane-2,9-dione (*cf.* EC 3.5.2.12, 6-aminohexanoate-cyclic-dimer hydrolase).  
**References:** [1358, 3271, 2067]

[EC 3.5.1.117 created 2014]

#### EC 3.5.1.118

**Accepted name:**  $\gamma$ -glutamyl hercynylcysteine *S*-oxide hydrolase  
**Reaction:**  $\gamma$ -L-glutamyl-*S*-(hercyn-2-yl)-L-cysteine *S*-oxide + H<sub>2</sub>O = *S*-(hercyn-2-yl)-L-cysteine *S*-oxide + L-glutamate  
**Other name(s):** EgtC  
**Systematic name:**  $\gamma$ -glutamyl-*S*-(hercyn-2-yl)cysteine *S*-oxide amidohydrolase  
**Comments:** The enzyme is part of the biosynthesis pathway of ergothioneine in mycobacteria.  
**References:** [2599]

[EC 3.5.1.118 created 2015]

#### EC 3.5.1.119

**Accepted name:** Pup amidohydrolase  
**Reaction:** [prokaryotic ubiquitin-like protein]-L-glutamine + H<sub>2</sub>O = [prokaryotic ubiquitin-like protein]-L-glutamate + NH<sub>3</sub>  
**Other name(s):** *dop* (gene name); Pup deamidase; depupylase/deamidase; DPUP; depupylase  
**Systematic name:** [prokaryotic ubiquitin-like protein]-L-glutamine amidohydrolase

**Comments:** The enzyme has been characterized from the bacterium *Mycobacterium tuberculosis*. It catalyses the hydrolysis of the amido group of the C-terminal glutamine of prokaryotic ubiquitin-like protein (Pup), thus activating it for ligation to target proteins, a process catalysed by EC 6.3.1.19, prokaryotic ubiquitin-like protein ligase. The reaction requires ATP as cofactor but not its hydrolysis. The enzyme also catalyses the hydrolytic cleavage of the bond formed by the ligase, between an ε-amino group of a lysine residue of the target protein and the γ-carboxylate of the C-terminal glutamate of the prokaryotic ubiquitin-like protein.

**References:** [2782, 331, 2781]

[EC 3.5.1.119 created 2015]

[3.5.1.120 *Transferred entry. 2-aminomuconate deaminase (2-hydroxymuconate-forming). Now EC 3.5.99.11, 2-aminomuconate deaminase (2-hydroxymuconate-forming) ]*

[EC 3.5.1.120 created 2016, deleted 2017]

#### EC 3.5.1.121

**Accepted name:** protein N-terminal asparagine amidohydrolase  
**Reaction:** N-terminal L-asparaginyl-[protein] + H<sub>2</sub>O = N-terminal L-aspartyl-[protein] + NH<sub>3</sub>  
**Other name(s):** NTAN1 (gene name)  
**Systematic name:** protein N-terminal asparagine amidohydrolase  
**Comments:** This enzyme participates in the eukaryotic ubiquitin-dependent Arg/N-end rule pathway of protein degradation, promoting the turnover of intracellular proteins that initiate with Met-Asn. Following the acetylation and removal of the initiator methionine, the exposed N-terminal asparagine is deaminated, resulting in its conversion to L-aspartate. The latter serves as a substrate for EC 2.3.2.8, arginyltransferase, making the protein susceptible to arginylation, polyubiquitination and degradation as specified by the N-end rule.

**References:** [2770, 979, 367]

[EC 3.5.1.121 created 2016]

#### EC 3.5.1.122

**Accepted name:** protein N-terminal glutamine amidohydrolase  
**Reaction:** N-terminal L-glutamyl-[protein] + H<sub>2</sub>O = N-terminal L-glutamyl-[protein] + NH<sub>3</sub>  
**Other name(s):** NTAQ1 (gene name)  
**Systematic name:** protein N-terminal glutamine amidohydrolase  
**Comments:** This enzyme participates in the eukaryotic ubiquitin-dependent Arg/N-end rule pathway of protein degradation, promoting the turnover of intracellular proteins that initiate with Met-Gln. Following the acetylation and removal of the initiator methionine, the exposed N-terminal glutamine is deaminated, resulting in its conversion to L-glutamate. The latter serves as a substrate for EC 2.3.2.8, arginyltransferase, making the protein susceptible to arginylation, polyubiquitination and degradation as specified by the N-end rule.

**References:** [3110]

[EC 3.5.1.122 created 2016]

#### EC 3.5.1.123

**Accepted name:** γ-glutamylanilide hydrolase  
**Reaction:** N<sup>5</sup>-phenyl-L-glutamine + H<sub>2</sub>O = L-glutamate + aniline  
**Other name(s):** *atdA2* (gene name)  
**Systematic name:** N<sup>5</sup>-phenyl-L-glutamine amidohydrolase  
**Comments:** The enzyme, characterized from the bacterium *Acinetobacter* sp. YAA, catalyses the opposite reaction from that catalysed by EC 6.3.1.18, γ-glutamylanilide synthase, which is part of an aniline degradation pathway. Its purpose is likely to maintain a low concentration of N<sup>5</sup>-phenyl-L-glutamine, which is potentially toxic.

**References:** [2858]

[EC 3.5.1.123 created 2016]

#### EC 3.5.1.124

**Accepted name:** protein deglycase  
**Reaction:** (1) an  $N^{\omega}$ -(1-hydroxy-2-oxopropyl)-[protein]-L-arginine + H<sub>2</sub>O = a [protein]-L-arginine + lactate  
(2) an  $N^{\epsilon}$ -(1-hydroxy-2-oxopropyl)-[protein]-L-lysine + H<sub>2</sub>O = a [protein]-L-lysine + lactate  
(3) an  $S$ -(1-hydroxy-2-oxopropyl)-[protein]-L-cysteine + H<sub>2</sub>O = a [protein]-L-cysteine + lactate  
**Other name(s):** PARK7 (gene name); DJ-1 protein; *yhbO* (gene name); *yajL* (gene name); glyoxylase III (incorrect)  
**Systematic name:** a [protein]-L-amino acid-1-hydroxypropan-2-one hydrolase [(*R*)-lactate-forming]  
**Comments:** The enzyme, previously thought to be a glyoxalase, acts on glycated L-arginine, L-lysine, and L-cysteine residues within proteins that have been attacked and modified by glyoxal or 2-oxopropanal. The attack forms hemithioacetal in the case of cysteines and aminocarbinols in the case of arginines and lysines. The enzyme repairs the amino acids, releasing glycolate or lactate (70-80% (*S*)-lactate and 20-30% (*R*)-lactate), depending on whether the attacking agent was glyoxal or 2-oxopropanal, respectively [2421, 1908].  
**References:** [1928, 2785, 2421, 1908, 2]

[EC 3.5.1.124 created 2016]

#### EC 3.5.1.125

**Accepted name:**  $N^2$ -acetyl-L-2,4-diaminobutanoate deacetylase  
**Reaction:** (2*S*)-2-acetamido-4-aminobutanoate + H<sub>2</sub>O = L-2,4-diaminobutanoate + acetate  
**Other name(s):** *doeB* (gene name)  
**Systematic name:** (2*S*)-2-acetamido-4-aminobutanoate amidohydrolase  
**Comments:** The enzyme, found in bacteria, has no activity with (2*S*)-4-acetamido-2-aminobutanoate (*cf.* EC 3.5.4.44, ectoine hydrolase).  
**References:** [2597]

[EC 3.5.1.125 created 2017]

#### EC 3.5.1.126

**Accepted name:** oxamate amidohydrolase  
**Reaction:** oxamate + H<sub>2</sub>O = oxalate + NH<sub>3</sub>  
**Other name(s):** HpxW  
**Systematic name:** oxamate amidohydrolase  
**Comments:** The enzyme has been characterized from the bacterium *Klebsiella pneumoniae*.  
**References:** [1137]

[EC 3.5.1.126 created 2017]

#### EC 3.5.1.127

**Accepted name:** jasmonoyl-L-amino acid hydrolase  
**Reaction:** a jasmonoyl-L-amino acid + H<sub>2</sub>O = jasmonate + an L-amino acid  
**Other name(s):** IAR3 (gene name); ILL4 (gene name); ILL6 (gene name)  
**Systematic name:** jasmonoyl-L amino acid amidohydrolase  
**Comments:** This entry includes a family of enzymes that recycle jasmonoyl-amino acid conjugates back to jasmonates. The enzymes from *Arabidopsis thaliana* have been shown to also act on 12-hydroxyjasmonoyl-L-isoleucine, generating tuberonic acid.  
**References:** [3179]

[EC 3.5.1.127 created 2017]

### EC 3.5.1.128

- Accepted name:** deaminated glutathione amidase  
**Reaction:**  $N$ -(4-oxoglutaryl)-L-cysteinylglycine + H<sub>2</sub>O = 2-oxoglutarate + L-cysteinylglycine  
**Other name(s):** dGSH deaminase; NIT1 (gene name)  
**Systematic name:**  $N$ -(4-oxoglutaryl)-L-cysteinylglycine amidohydrolase  
**Comments:** The enzyme, present in animals, fungi and bacteria, is involved in clearing cells of the toxic compound deaminated glutathione, which can be produced as an unwanted side product by several transaminases.  
**References:** [2257]

[EC 3.5.1.128 created 2018]

### EC 3.5.1.129

- Accepted name:**  $N^5$ -(cytidine 5'-diphosphoramidyl)-L-glutamine hydrolase  
**Reaction:**  $N^5$ -(cytidine 5'-diphosphoramidyl)-L-glutamine + H<sub>2</sub>O = cytidine 5'-diphosphoramidate + L-glutamate  
**Other name(s):**  $N^5$ -(cytidine 5'-diphosphoramidyl)-L-glutamine deacylase  
**Systematic name:**  $N^5$ -(cytidine 5'-diphosphoramidyl)-L-glutamine amidohydrolase  
**Comments:** The enzyme, characterized from the bacterium *Campylobacter jejuni*, is involved in formation of a unique *O*-methyl phosphoramidate modification on specific sugar residues within the bacterium's capsular polysaccharides.  
**References:** [2896]

[EC 3.5.1.129 created 2018]

## EC 3.5.2 In cyclic amides

### EC 3.5.2.1

- Accepted name:** barbiturase  
**Reaction:** barbiturate + H<sub>2</sub>O = 3-oxo-3-ureidopropanoate  
**Systematic name:** barbiturate amidohydrolase (3-oxo-3-ureidopropanoate-forming)  
**Comments:** Contains zinc and is specific for barbiturate as substrate [2724]. Forms part of the oxidative pyrimidine-degrading pathway in some microorganisms, along with EC 1.17.99.4 (uracil/thymine dehydrogenase) and EC 3.5.1.95 (*N*-malonylurea hydrolase). It was previously thought that the end-products of the reaction were malonate and urea but this has since been disproved [2725]. May be involved in the regulation of pyrimidine metabolism, along with EC 2.4.2.9, uracil phosphoribosyltransferase.  
**References:** [1083, 2725, 2724]

[EC 3.5.2.1 created 1961, modified 2006]

### EC 3.5.2.2

- Accepted name:** dihydropyrimidinase  
**Reaction:** 5,6-dihydrouracil + H<sub>2</sub>O = 3-ureidopropanoate  
**Other name(s):** hydantoinase; hydroxypyrimidine hydrase; hydantoin peptidase; pyrimidine hydrase; D-hydantoinase  
**Systematic name:** 5,6-dihydropyrimidine amidohydrolase  
**Comments:** Also acts on dihydrothymine and hydantoin.  
**References:** [305, 673]

[EC 3.5.2.2 created 1961]

### EC 3.5.2.3



**Accepted name:** dihydroorotase  
**Reaction:** (S)-dihydroorotate + H<sub>2</sub>O = N-carbamoyl-L-aspartate  
**Other name(s):** carbamoylaspartic dehydrase; dihydroorotate hydrolase  
**Systematic name:** (S)-dihydroorotate amidohydrolase  
**References:** [485, 1682]

[EC 3.5.2.3 created 1961]

#### EC 3.5.2.4

**Accepted name:** carboxymethylhydantoinase  
**Reaction:** L-5-carboxymethylhydantoin + H<sub>2</sub>O = N-carbamoyl-L-aspartate  
**Other name(s):** hydantoin hydrolase  
**Systematic name:** L-5-carboxymethylhydantoin amidohydrolase  
**References:** [1682]

[EC 3.5.2.4 created 1961]

#### EC 3.5.2.5

**Accepted name:** allantoinase  
**Reaction:** (S)-allantoin + H<sub>2</sub>O = allantoate  
**Systematic name:** (S)-allantoin amidohydrolase  
**References:** [786]

[EC 3.5.2.5 created 1961]

#### EC 3.5.2.6

**Accepted name:** β-lactamase  
**Reaction:** a β-lactam + H<sub>2</sub>O = a substituted β-amino acid  
**Other name(s):** penicillinase; cephalosporinase; neutrapen; penicillin β-lactamase; exopenicillinase; ampicillinase; penicillin amido-β-lactamhydrolase; penicillinase I, II; β-lactamase I-III; β-lactamase A, B, C; β-lactamase AME I; cephalosporin-β-lactamase  
**Systematic name:** β-lactam hydrolase  
**Comments:** A group of enzymes of varying specificity hydrolysing β-lactams; some act more rapidly on penicillins, some more rapidly on cephalosporins. The latter were formerly listed as EC 3.5.2.8, cephalosporinase.  
**References:** [454, 1112, 1578, 2306, 2307, 2462]

[EC 3.5.2.6 created 1961, modified 1981 (EC 3.5.2.8 created 1972, incorporated 1978)]

#### EC 3.5.2.7

**Accepted name:** imidazolonepropionase  
**Reaction:** (S)-3-(5-oxo-4,5-dihydro-3H-imidazol-4-yl)propanoate + H<sub>2</sub>O = N-formimidoyl-L-glutamate + H<sup>+</sup>  
**Other name(s):** 4(5)-imidazolone-5(4)-propionic acid hydrolase; imidazolone propionic acid hydrolase  
**Systematic name:** 3-(5-oxo-4,5-dihydro-3H-imidazol-4-yl)propanoate amidohydrolase  
**References:** [2373, 2705]

[EC 3.5.2.7 created 1965, modified 2001]

[3.5.2.8 Deleted entry. cephalosporinase. Now included with EC 3.5.2.6 β-lactamase]

[EC 3.5.2.8 created 1972, deleted 1978]

#### EC 3.5.2.9



**Accepted name:** 5-oxoprolinase (ATP-hydrolysing)  
**Reaction:** ATP + 5-oxo-L-proline + 2 H<sub>2</sub>O = ADP + phosphate + L-glutamate  
**Other name(s):** pyroglutamase (ATP-hydrolysing); oxoprolinase; pyroglutamase; 5-oxoprolinase; pyroglutamate hydrolase; pyroglutamic hydrolase; L-pyroglutamate hydrolase; 5-oxo-L-prolinase; pyroglutamase  
**Systematic name:** 5-oxo-L-proline amidohydrolase (ATP-hydrolysing)  
**References:** [3043]

[EC 3.5.2.9 created 1976]

#### EC 3.5.2.10

**Accepted name:** creatininase  
**Reaction:** creatinine + H<sub>2</sub>O = creatine  
**Other name(s):** creatinine hydrolase  
**Systematic name:** creatinine amidohydrolase  
**References:** [2989]

[EC 3.5.2.10 created 1978]

#### EC 3.5.2.11

**Accepted name:** L-lysine-lactamase  
**Reaction:** (S)-2-aminohexano-6-lactam + H<sub>2</sub>O = L-lysine  
**Other name(s):** L- $\alpha$ -aminocaprolactam hydrolase; L-lysinamidase; L-lysine-1,6-lactam lactamhydrolase  
**Systematic name:** (S)-2-aminohexano-6-lactam lactamhydrolase  
**Comments:** Also hydrolyses L-lysinamide.  
**References:** [860, 2651]

[EC 3.5.2.11 created 1981, modified 1989]

#### EC 3.5.2.12

**Accepted name:** 6-aminohexanoate-cyclic-dimer hydrolase  
**Reaction:** 1,8-diazacyclotetradecane-2,9-dione + H<sub>2</sub>O = *N*-(6-aminohexanoyl)-6-aminohexanoate  
**Systematic name:** 1,8-diazacyclotetradecane-2,9-dione lactamhydrolase  
**Comments:** The cyclic dimer of 6-aminohexanoate is converted to the linear dimer.  
**References:** [1460]

[EC 3.5.2.12 created 1983]

#### EC 3.5.2.13

**Accepted name:** 2,5-dioxopiperazine hydrolase  
**Reaction:** 2,5-dioxopiperazine + H<sub>2</sub>O = glycylglycine  
**Other name(s):** cyclo(Gly-Gly) hydrolase; cyclo(glycylglycine) hydrolase  
**Systematic name:** 2,5-dioxopiperazine amidohydrolase  
**Comments:** Highly specific; does not hydrolyse other dioxopiperazines, glycylglycine, proteins or barbiturates.  
**References:** [2818]

[EC 3.5.2.13 created 1989]

#### EC 3.5.2.14

**Accepted name:** *N*-methylhydantoinase (ATP-hydrolysing)  
**Reaction:** ATP + *N*-methylhydantoin + 2 H<sub>2</sub>O = ADP + phosphate + *N*-carbamoylsarcosine  
**Other name(s):** *N*-methylhydantoin amidohydrolase; methylhydantoin amidase; *N*-methylhydantoin hydrolase; *N*-methylhydantoinase; *N*-methylimidazolidine-2,4-dione amidohydrolase (ATP-hydrolysing)

**Systematic name:** *N*-methylhydantoin amidohydrolase (ATP-hydrolysing)

**References:** [1451]

[EC 3.5.2.14 created 1989]

#### EC 3.5.2.15

**Accepted name:** cyanuric acid amidohydrolase

**Reaction:** cyanuric acid + H<sub>2</sub>O = biuret + CO<sub>2</sub>

**Other name(s):** AtzD

**Systematic name:** cyanuric acid amidohydrolase

**Comments:** Along with EC 3.5.1.54 (allophanate hydrolase) and EC 3.5.1.84 (biuret amidohydrolase), this enzyme forms part of the cyanuric-acid metabolism pathway, which degrades *s*-triazide herbicides, such as atrazine [2-chloro-4-(ethylamino)-6-(isopropylamino)-1,3,5-triazine], in bacteria. This is a key enzyme in the pathway, catalysing the ring cleavage of cyanuric acid. The enzyme is specific for cyanuric acid as substrate as neither the structurally related compounds ammeline (2,4-diamino-6-hydroxy-*s*-triazine) and ammelide (2-amino-4,6-dihydroxy-*s*-triazine) nor a number of pyrimidine compounds, such as uracil and cytosine, can act as substrates [1389].

**References:** [676, 675, 1389, 824]

[EC 3.5.2.15 created 2000, modified 2008]

#### EC 3.5.2.16

**Accepted name:** maleimide hydrolase

**Reaction:** maleimide + H<sub>2</sub>O = maleamic acid

**Other name(s):** imidase; cyclic imide hydrolase; cyclic-imide amidohydrolase (decyclizing) [misprint]; cyclic-imide amidohydrolase (decyclizing)

**Systematic name:** cyclic-imide amidohydrolase (ring-opening)

**Comments:** Succinimide and glutarimide, and sulfur-containing cyclic imides, such as rhodanine, can also act as substrates for the enzyme from *Blastobacter* sp. A17p-4. The reverse, cyclization, reaction is also catalysed, but much more slowly. It has lower activity towards cyclic ureides, which are the substrates of EC 3.5.2.2, dihydropyrimidinase.

**References:** [2147]

[EC 3.5.2.16 created 2001]

#### EC 3.5.2.17

**Accepted name:** hydroxyisourate hydrolase

**Reaction:** 5-hydroxyisourate + H<sub>2</sub>O = 5-hydroxy-2-oxo-4-ureido-2,5-dihydro-1*H*-imidazole-5-carboxylate

**Other name(s):** HIUHase; 5-hydroxyisourate hydrolase

**Systematic name:** 5-hydroxyisourate amidohydrolase

**Comments:** The reaction is the first stage in the conversion of 5-hydroxyisourate into *S*-allantoin. This reaction will also occur spontaneously but more slowly.

**References:** [2387, 2386, 2531]

[EC 3.5.2.17 created 2004]

#### EC 3.5.2.18

**Accepted name:** enamidase

**Reaction:** 6-oxo-1,4,5,6-tetrahydronicotinate + 2 H<sub>2</sub>O = 2-formylglutarate + NH<sub>3</sub>

**Systematic name:** 6-oxo-1,4,5,6-tetrahydronicotinate amidohydrolase

**Comments:** Contains iron and  $Zn^{2+}$ . Forms part of the nicotinate-fermentation catabolism pathway in *Eubacterium barkeri*. Other enzymes involved in this pathway are EC 1.17.1.5 (nicotinate dehydrogenase), EC 1.3.7.1 (6-hydroxynicotinate reductase), EC 1.1.1.291 (2-hydroxymethylglutarate dehydrogenase), EC 5.4.99.4 (2-methyleneglutarate mutase), EC 5.3.3.6 (methylitaconate  $\Delta$ -isomerase), EC 4.2.1.85 (dimethylmaleate hydratase) and EC 4.1.3.32 (2,3-dimethylmalate lyase).

**References:** [34]

[EC 3.5.2.18 created 2006]

#### EC 3.5.2.19

**Accepted name:** streptothricin hydrolase  
**Reaction:** streptothricin-F +  $H_2O$  = streptothricin-F acid  
**Other name(s):** *sttH* (gene name)  
**Systematic name:** streptothricin-F hydrolase  
**Comments:** The enzyme also catalyses the hydrolysis of streptothricin-D to streptothricin-D acid [1826]. The enzyme is responsible for streptothricin resistance in *Streptomyces albulus* and *Streptomyces noursei* [1826, 1033].  
**References:** [1826, 1033]

[EC 3.5.2.19 created 2011]

#### EC 3.5.2.20

**Accepted name:** isatin hydrolase  
**Reaction:** isatin +  $H_2O$  = isatinic acid  
**Systematic name:** isatin amidohydrolase  
**Comments:** Requires  $Mn^{2+}$ . This enzyme, found in several bacterial species, is involved in the degradation of indole-3-acetic acid.  
**References:** [2713, 225]

[EC 3.5.2.20 created 2014]

### EC 3.5.3 In linear amidines

#### EC 3.5.3.1

**Accepted name:** arginase  
**Reaction:** L-arginine +  $H_2O$  = L-ornithine + urea  
**Other name(s):** arginine amidinase; canavanase; L-arginase; arginine transamidinase  
**Systematic name:** L-arginine amidinohydrolase  
**Comments:** Also hydrolyses  $\alpha$ -N-substituted L-arginines and canavanine.  
**References:** [114, 345, 665, 971, 972]

[EC 3.5.3.1 created 1961]

#### EC 3.5.3.2

**Accepted name:** guanidinoacetase  
**Reaction:** guanidinoacetate +  $H_2O$  = glycine + urea  
**Other name(s):** glycoyaminate hydrolase  
**Systematic name:** guanidinoacetate amidinohydrolase  
**Comments:** Requires  $Mn^{2+}$ .  
**References:** [2444, 3297]

[EC 3.5.3.2 created 1961]

### EC 3.5.3.3

**Accepted name:** creatinase  
**Reaction:** creatine + H<sub>2</sub>O = sarcosine + urea  
**Systematic name:** creatine amidinohydrolase  
**References:** [2444, 3304]

[EC 3.5.3.3 created 1961]

### EC 3.5.3.4

**Accepted name:** allantoicase  
**Reaction:** allantoate + H<sub>2</sub>O = (*S*)-ureidoglycolate + urea  
**Systematic name:** allantoate amidinohydrolase  
**Comments:** Also hydrolyses (*R*)-ureidoglycolate to glyoxylate and urea.  
**References:** [786, 2963, 3038, 2484]

[EC 3.5.3.4 created 1961]

### EC 3.5.3.5

**Accepted name:** formimidoylaspartate deiminase  
**Reaction:** *N*-formimidoyl-L-aspartate + H<sub>2</sub>O = *N*-formyl-L-aspartate + NH<sub>3</sub>  
**Other name(s):** formiminoaspartate deiminase  
**Systematic name:** *N*-formimidoyl-L-aspartate iminohydrolase  
**References:** [1087]

[EC 3.5.3.5 created 1961, modified 2000]

### EC 3.5.3.6

**Accepted name:** arginine deiminase  
**Reaction:** L-arginine + H<sub>2</sub>O = L-citrulline + NH<sub>3</sub>  
**Other name(s):** arginine dihydrolase; citrulline iminase; L-arginine deiminase  
**Systematic name:** L-arginine iminohydrolase  
**Comments:** Also acts on canavanine.  
**References:** [2152, 2269, 2377]

[EC 3.5.3.6 created 1961]

### EC 3.5.3.7

**Accepted name:** guanidinobutyrase  
**Reaction:** 4-guanidinobutanoate + H<sub>2</sub>O = 4-aminobutanoate + urea  
**Other name(s):** γ-guanidinobutyrase; 4-guanidinobutyrate amidinobutyrase; γ-guanidinobutyrate amidinohydrolase; G-Base; GBH; guanidinobutyrate ureahydrolase  
**Systematic name:** 4-guanidinobutanoate amidinohydrolase  
**Comments:** Requires Mn<sup>2+</sup>. Also acts, very slowly, on 5-guanidinopentanoate and 6-guanidinohexanoate.  
**References:** [1959, 2911, 3294, 3295]

[EC 3.5.3.7 created 1972]

### EC 3.5.3.8

**Accepted name:** formimidoylglutamase  
**Reaction:** *N*-formimidoyl-L-glutamate + H<sub>2</sub>O = L-glutamate + formamide  
**Other name(s):** formiminoglutamase; *N*-formiminoglutamate hydrolase; *N*-formimino-L-glutamate formiminohydro-  
lase

**Systematic name:** *N*-formimidoyl-L-glutamate formimidoylhydrolase

**References:** [1372, 1752]

[EC 3.5.3.8 created 1972, modified 2000, modified 2001]

#### EC 3.5.3.9

**Accepted name:** allantoate deiminase

**Reaction:** allantoate + H<sub>2</sub>O = (*S*)-ureidoglycine + NH<sub>3</sub> + CO<sub>2</sub>

**Other name(s):** allantoate amidohydrolase

**Systematic name:** allantoate amidinohydrolase (decarboxylating)

**Comments:** This enzyme is part of the ureide pathway, which permits certain organisms to recycle the nitrogen in purine compounds. This enzyme, which liberates ammonia from allantoate, is present in plants and bacteria. In plants it is localized in the endoplasmic reticulum. Requires manganese.

**References:** [3074, 2613]

[EC 3.5.3.9 created 1972, modified 2010]

#### EC 3.5.3.10

**Accepted name:** D-arginase

**Reaction:** D-arginine + H<sub>2</sub>O = D-ornithine + urea

**Systematic name:** D-arginine amidinohydrolase

**References:** [2022]

[EC 3.5.3.10 created 1972]

#### EC 3.5.3.11

**Accepted name:** agmatinase

**Reaction:** agmatine + H<sub>2</sub>O = putrescine + urea

**Other name(s):** agmatine ureohydrolase; SpeB

**Systematic name:** agmatine amidinohydrolase

**References:** [1151, 3065]

[EC 3.5.3.11 created 1972]

#### EC 3.5.3.12

**Accepted name:** agmatine deiminase

**Reaction:** agmatine + H<sub>2</sub>O = *N*-carbamoylputrescine + NH<sub>3</sub>

**Other name(s):** agmatine amidinohydrolase

**Systematic name:** agmatine iminohydrolase

**Comments:** The plant enzyme also catalyses the reactions of EC 2.1.3.3 (ornithine carbamoyltransferase), EC 2.1.3.6 (putrescine carbamoyltransferase) and EC 2.7.2.2 (carbamate kinase), thus functioning as a putrescine synthase, converting agmatine and ornithine into putrescine and citrulline, respectively.

**References:** [2701, 2751]

[EC 3.5.3.12 created 1972]

#### EC 3.5.3.13

**Accepted name:** formimidoylglutamate deiminase

**Reaction:** *N*-formimidoyl-L-glutamate + H<sub>2</sub>O = *N*-formyl-L-glutamate + NH<sub>3</sub>

**Other name(s):** formiminoglutamate deiminase; formiminoglutamic iminohydrolase

**Systematic name:** *N*-formimidoyl-L-glutamate iminohydrolase

**References:** [3178]

[EC 3.5.3.13 created 1975, modified 2000]

#### EC 3.5.3.14

**Accepted name:** amidinoaspartase  
**Reaction:**  $N$ -amidino-L-aspartate + H<sub>2</sub>O = L-aspartate + urea  
**Other name(s):** amidinoaspartic amidinohydrolase  
**Systematic name:**  $N$ -amidino-L-aspartate amidinohydrolase  
**Comments:** Also acts slowly on  $N$ -amidino-L-glutamate.  
**References:** [1919]

[EC 3.5.3.14 created 1976]

#### EC 3.5.3.15

**Accepted name:** protein-arginine deiminase  
**Reaction:** protein L-arginine + H<sub>2</sub>O = protein L-citrulline + NH<sub>3</sub>  
**Other name(s):** peptidylarginine deiminase; PAD  
**Systematic name:** protein-L-arginine iminohydrolase  
**Comments:** Also acts on  $N$ -acyl-L-arginine and, more slowly, on L-arginine esters.  
**References:** [847]

[EC 3.5.3.15 created 1983]

#### EC 3.5.3.16

**Accepted name:** methylguanidinase  
**Reaction:** methylguanidine + H<sub>2</sub>O = methylamine + urea  
**Other name(s):** methylguanidine hydrolase  
**Systematic name:** methylguanidine amidinohydrolase  
**Comments:** Acts on some other alkylguanidines, but very slowly.  
**References:** [2039]

[EC 3.5.3.16 created 1984]

#### EC 3.5.3.17

**Accepted name:** guanidinopropionase  
**Reaction:** 3-guanidinopropanoate + H<sub>2</sub>O =  $\beta$ -alanine + urea  
**Other name(s):** GPase; GPH  
**Systematic name:** 3-guanidinopropanoate amidinopropionase  
**Comments:** Requires Mn<sup>2+</sup>. Also acts, more slowly, on taurocyamine and 4-guanidinobutanoate.  
**References:** [3296]

[EC 3.5.3.17 created 1989]

#### EC 3.5.3.18

**Accepted name:** dimethylargininase  
**Reaction:**  $N^{\omega},N^{\omega'}$ -dimethyl-L-arginine + H<sub>2</sub>O = dimethylamine + L-citrulline  
**Other name(s):** dimethylarginine dimethylaminohydrolase;  $N^G,N^G$ -dimethylarginine dimethylaminohydrolase;  $N^G,N^G$ -dimethyl-L-arginine dimethylamidohydrolase;  $\omega,\omega'$ -di- $N$ -methyl-L-arginine dimethylamidohydrolase;  $N^{\omega},N^{\omega'}$ -methyl-L-arginine dimethylamidohydrolase (incorrect)  
**Systematic name:**  $N^{\omega},N^{\omega'}$ -dimethyl-L-arginine dimethylamidohydrolase  
**Comments:** Also acts on  $N^{\omega}$ -methyl-L-arginine.  
**References:** [2150]

[EC 3.5.3.18 created 1992]

[3.5.3.19 Transferred entry. ureidoglycolate hydrolase. Now EC 3.5.1.116, ureidoglycolate amidohydrolase]

[EC 3.5.3.19 created 1992, deleted 2014]

#### EC 3.5.3.20

**Accepted name:** diguanidinobutanase  
**Reaction:** 1,4-diguanidinobutane + H<sub>2</sub>O = agmatine + urea  
**Systematic name:** 1,4-diguanidinobutane amidohydrolase  
**Comments:** Other diguanidinoalkanes with 3 to 10 methylene groups can also act, but more slowly.  
**References:** [3293]

[EC 3.5.3.20 created 1992]

#### EC 3.5.3.21

**Accepted name:** methylenediurea deaminase  
**Reaction:** methylenediurea + 2 H<sub>2</sub>O = *N*-(hydroxymethyl)urea + 2 NH<sub>3</sub> + CO<sub>2</sub> (overall reaction)  
(1a) methylenediurea + H<sub>2</sub>O = *N*-(carboxyaminoethyl)urea + NH<sub>3</sub>  
(1b) *N*-(carboxyaminoethyl)urea = *N*-(aminomethyl)urea + CO<sub>2</sub> (spontaneous)  
(1c) *N*-(aminomethyl)urea + H<sub>2</sub>O = *N*-(hydroxymethyl)urea + NH<sub>3</sub> (spontaneous)  
**Other name(s):** methylenediurease  
**Systematic name:** methylenediurea aminohydrolase  
**Comments:** Methylenediurea is hydrolysed and decarboxylated to give an aminated methylurea, which then spontaneously hydrolyses to hydroxymethylurea. The enzyme from *Ochrobactrum anthropi* also hydrolyses dimethylenetriurea and trimethylenetetraurea as well as ureidoglycolate, which is hydrolysed to urea and glyoxylate, and allantoate, which is hydrolysed to ureidoglycolate, ammonia and carbon dioxide.  
**References:** [1297]

[EC 3.5.3.21 created 1999]

#### EC 3.5.3.22

**Accepted name:** proclavamate amidohydrolase  
**Reaction:** amidinoproclavamate + H<sub>2</sub>O = proclavamate + urea  
**Other name(s):** PAH; proclavamate amidino hydrolase  
**Systematic name:** amidinoproclavamate amidohydrolase  
**Comments:** Forms part of the pathway for the biosynthesis of the β-lactamase inhibitor clavulanate in *Streptomyces clavuligerus*. It carries out an intermediary reaction between the first reaction of EC 1.14.11.21, clavamate synthase, and the second and third reactions of that enzyme. Requires Mn<sup>2+</sup>.  
**References:** [2510, 3335, 2955, 3222]

[EC 3.5.3.22 created 2003]

#### EC 3.5.3.23

**Accepted name:** *N*-succinylarginine dihydrolase  
**Reaction:** *N*<sup>2</sup>-succinyl-L-arginine + 2 H<sub>2</sub>O = *N*<sup>2</sup>-succinyl-L-ornithine + 2 NH<sub>3</sub> + CO<sub>2</sub>  
**Other name(s):** *N*<sup>2</sup>-succinylarginine dihydrolase; arginine succinylhydrolase; SADH; AruB; AstB; 2-*N*-succinyl-L-arginine iminohydrolase (decarboxylating)  
**Systematic name:** *N*<sup>2</sup>-succinyl-L-arginine iminohydrolase (decarboxylating)

**Comments:** Arginine, *N*<sup>2</sup>-acetylarginine and *N*<sup>2</sup>-glutamylarginine do not act as substrates [3139]. This is the second enzyme in the arginine succinyltransferase (AST) pathway for the catabolism of arginine [2575]. This pathway converts the carbon skeleton of arginine into glutamate, with the concomitant production of ammonia and conversion of succinyl-CoA into succinate and CoA. The five enzymes involved in this pathway are EC 2.3.1.109 (arginine *N*-succinyltransferase), EC 3.5.3.23 (*N*-succinylarginine dihydrolase), EC 2.6.1.81 (succinylornithine transaminase), EC 1.2.1.71 (succinylglutamate semialdehyde dehydrogenase) and EC 3.5.1.96 (succinylglutamate desuccinylase).

**References:** [2575, 2927, 3139, 517, 1276]

[EC 3.5.3.23 created 2006]

#### EC 3.5.3.24

**Accepted name:** *N*<sup>1</sup>-aminopropylagmatine ureohydrolase  
**Reaction:** *N*<sup>1</sup>-aminopropylagmatine + H<sub>2</sub>O = spermidine + urea  
**Systematic name:** *N*<sup>1</sup>-aminopropylagmatine amidinohydrolase  
**Comments:** The enzyme, which has been characterized from the hyperthermophilic archaeon *Pyrococcus kodakarensis* and the thermophilic Gram-negative bacterium *Thermus thermophilus*, is involved in the biosynthesis of spermidine.  
**References:** [2165, 1975]

[EC 3.5.3.24 created 2013]

#### EC 3.5.3.25

**Accepted name:** *N*<sup>ω</sup>-hydroxy-L-arginine amidinohydrolase  
**Reaction:** *N*<sup>ω</sup>-hydroxy-L-arginine + H<sub>2</sub>O = L-ornithine + hydroxyurea  
**Other name(s):** *dcsB* (gene name)  
**Systematic name:** *N*<sup>ω</sup>-hydroxy-L-arginine amidinohydrolase  
**Comments:** The enzyme participates in the biosynthetic pathway of D-cycloserine, an antibiotic substance produced by several *Streptomyces* species.  
**References:** [1547, 1548]

[EC 3.5.3.25 created 2013]

#### EC 3.5.3.26

**Accepted name:** (*S*)-ureidoglycine aminohydrolase  
**Reaction:** (*S*)-2-ureidoglycine + H<sub>2</sub>O = (*S*)-ureidoglycolate + NH<sub>3</sub>  
**Other name(s):** UGlyAH; UGHY; *ylbA* (gene name)  
**Systematic name:** (*S*)-ureidoglycine aminohydrolase  
**Comments:** Binds Mn<sup>2+</sup>. This enzyme, found in plants and bacteria, is part of the ureide pathway, which enables the recycling of the nitrogen in purine compounds. In plants it is localized in the endoplasmic reticulum.  
**References:** [2613, 3162, 2637]

[EC 3.5.3.26 created 2013]

### EC 3.5.4 In cyclic amidines

#### EC 3.5.4.1

**Accepted name:** cytosine deaminase  
**Reaction:** cytosine + H<sub>2</sub>O = uracil + NH<sub>3</sub>  
**Other name(s):** isocytosine deaminase  
**Systematic name:** cytosine aminohydrolase



**Comments:** Also acts on 5-methylcytosine.

**References:** [465, 1532]

[EC 3.5.4.1 created 1961]

#### EC 3.5.4.2

**Accepted name:** adenine deaminase

**Reaction:** adenine + H<sub>2</sub>O = hypoxanthine + NH<sub>3</sub>

**Other name(s):** adenase; adenine aminase; ADase

**Systematic name:** adenine aminohydrolase

**References:** [243, 1124]

[EC 3.5.4.2 created 1961]

#### EC 3.5.4.3

**Accepted name:** guanine deaminase

**Reaction:** guanine + H<sub>2</sub>O = xanthine + NH<sub>3</sub>

**Other name(s):** guanase; guanine aminase; GAH

**Systematic name:** guanine aminohydrolase

**References:** [1152, 1361, 2348]

[EC 3.5.4.3 created 1961]

#### EC 3.5.4.4

**Accepted name:** adenosine deaminase

**Reaction:** adenosine + H<sub>2</sub>O = inosine + NH<sub>3</sub>

**Other name(s):** deoxyadenosine deaminase

**Systematic name:** adenosine aminohydrolase

**References:** [1388, 2320]

[EC 3.5.4.4 created 1961]

#### EC 3.5.4.5

**Accepted name:** cytidine deaminase

**Reaction:** (1) cytidine + H<sub>2</sub>O = uridine + NH<sub>3</sub>

(2) 2'-deoxycytidine + H<sub>2</sub>O = 2'-deoxyuridine + NH<sub>3</sub>

**Other name(s):** cytosine nucleoside deaminase; (deoxy)cytidine deaminase; *cdd* (gene name); CDA (gene name)

**Systematic name:** cytidine/2'-deoxycytidine aminohydrolase

**Comments:** Contains zinc. Catalyses the deamination of cytidine and 2'-deoxycytidine with similar efficiencies. The enzyme, which is widely distributed among organisms, is involved in salvage of both exogenous and endogenous cytidine and 2'-deoxycytidine for UMP synthesis.

**References:** [2434, 3120, 2714, 1591, 3068]

[EC 3.5.4.5 created 1961, modified 2013]

#### EC 3.5.4.6

**Accepted name:** AMP deaminase

**Reaction:** AMP + H<sub>2</sub>O = IMP + NH<sub>3</sub>

**Other name(s):** adenylic acid deaminase; AMP aminase; adenylic deaminase; adenylate deaminase; 5-AMP deaminase; adenosine 5-monophosphate deaminase; 5-adenylate deaminase; adenyl deaminase; 5-adenylic acid deaminase; adenosine monophosphate deaminase; adenylate aminohydrolase; adenylate desaminase; adenosine 5-phosphate aminohydrolase; 5-adenylate deaminase

**Systematic name:** AMP aminohydrolase  
**Comments:** cf. EC 3.5.4.17 adenosine-phosphate deaminase.  
**References:** [1361, 1634, 1635, 1636, 1886, 3000, 3152]

[EC 3.5.4.6 created 1961]

#### EC 3.5.4.7

**Accepted name:** ADP deaminase  
**Reaction:** ADP + H<sub>2</sub>O = IDP + NH<sub>3</sub>  
**Other name(s):** adenosine diphosphate deaminase; adenosinepyrophosphate deaminase  
**Systematic name:** ADP aminohydrolase  
**References:** [594]

[EC 3.5.4.7 created 1961]

#### EC 3.5.4.8

**Accepted name:** aminoimidazolase  
**Reaction:** 4-aminoimidazole + H<sub>2</sub>O = imidazol-4-one + NH<sub>3</sub>  
**Other name(s):** 4-aminoimidazole hydrolase; 4-aminoimidazole deaminase  
**Systematic name:** 4-aminoimidazole aminohydrolase  
**Comments:** Requires Fe<sup>2+</sup>. This enzyme forms part of the xanthine-degradation pathway in some bacteria. The product of the reaction, imidazol-4-one, can be converted non-enzymically into formiminoglycine. An enzyme has been identified in *Clostridium cylindrosporium* that can perform this hydrolysis reaction [811, 3075].  
**References:** [2349, 811, 3075, 578]

[EC 3.5.4.8 created 1961]

#### EC 3.5.4.9

**Accepted name:** methenyltetrahydrofolate cyclohydrolase  
**Reaction:** 5,10-methenyltetrahydrofolate + H<sub>2</sub>O = 10-formyltetrahydrofolate  
**Other name(s):** Citrovorum factor cyclodehydrase; cyclohydrolase; formyl-methenyl-methylenetetrahydrofolate synthetase (combined); 5,10-methenyltetrahydrofolate 5-hydrolase (decyclizing)  
**Systematic name:** 5,10-methenyltetrahydrofolate 5-hydrolase (ring-opening)  
**Comments:** In eukaryotes, the enzyme occurs as a trifunctional enzyme that also has methylenetetrahydrofolate dehydrogenase (NADP<sup>+</sup>) (EC 1.5.1.5) and formate—tetrahydrofolate ligase (EC 6.3.4.3) activity. In some prokaryotes, it occurs as a bifunctional enzyme that also has dehydrogenase (EC 1.5.1.5) activity or formimidoyltetrahydrofolate cyclodeaminase (EC 4.3.1.4) activity.  
**References:** [2350, 2827]

[EC 3.5.4.9 created 1961]

#### EC 3.5.4.10

**Accepted name:** IMP cyclohydrolase  
**Reaction:** IMP + H<sub>2</sub>O = 5-formamido-1-(5-phospho-D-ribose)imidazole-4-carboxamide  
**Other name(s):** inosinicase; inosinate cyclohydrolase; IMP 1,2-hydrolase (decyclizing)  
**Systematic name:** IMP 1,2-hydrolase (ring-opening)  
**References:** [780]

[EC 3.5.4.10 created 1961, modified 2000]

#### EC 3.5.4.11

**Accepted name:** pterin deaminase  
**Reaction:** a 2-amino-4-hydroxypteridine + H<sub>2</sub>O = a 2,4-dihydroxypteridine + NH<sub>3</sub>  
**Other name(s):** acrasinase  
**Systematic name:** 2-amino-4-hydroxypteridine aminohydrolase  
**Comments:** The animal enzyme is specific for pterin, isoxanthopterin and tetrahydropterin.  
**References:** [1654, 2405]

[EC 3.5.4.11 created 1965]

#### EC 3.5.4.12

**Accepted name:** dCMP deaminase  
**Reaction:** dCMP + H<sub>2</sub>O = dUMP + NH<sub>3</sub>  
**Other name(s):** deoxycytidylate deaminase; deoxy-CMP-deaminase; deoxycytidylate aminohydrolase; deoxycytidine monophosphate deaminase; deoxycytidine-5'-phosphate deaminase; deoxycytidine-5'-monophosphate aminohydrolase  
**Systematic name:** dCMP aminohydrolase  
**Comments:** Also acts on some 5-substituted dCMPs.  
**References:** [2550, 2551, 2612]

[EC 3.5.4.12 created 1965]

#### EC 3.5.4.13

**Accepted name:** dCTP deaminase  
**Reaction:** dCTP + H<sub>2</sub>O = dUTP + NH<sub>3</sub>  
**Other name(s):** deoxycytidine triphosphate deaminase; 5-methyl-dCTP deaminase  
**Systematic name:** dCTP aminohydrolase  
**References:** [2934]

[EC 3.5.4.13 created 1972]

[3.5.4.14 *Transferred entry. deoxycytidine deaminase. Now included in EC 3.5.4.5, (deoxy)cytidine deaminase*]

[EC 3.5.4.14 created 1972, transferred 2013 to EC 3.5.4.5., deleted 2013]

#### EC 3.5.4.15

**Accepted name:** guanosine deaminase  
**Reaction:** guanosine + H<sub>2</sub>O = xanthosine + NH<sub>3</sub>  
**Other name(s):** guanosine aminase  
**Systematic name:** guanosine aminohydrolase  
**References:** [1264]

[EC 3.5.4.15 created 1972]

#### EC 3.5.4.16

**Accepted name:** GTP cyclohydrolase I  
**Reaction:** GTP + H<sub>2</sub>O = formate + 7,8-dihydroneopterin 3'-triphosphate  
**Other name(s):** GTP cyclohydrolase; guanosine triphosphate cyclohydrolase; guanosine triphosphate 8-deformylase; dihydroneopterin triphosphate synthase; GTP 8-formylhydrolase  
**Systematic name:** GTP 7,8-8,9-dihydrolase  
**Comments:** The reaction involves hydrolysis of two C-N bonds and isomerization of the pentose unit; the recyclization may be non-enzymic. This enzyme is involved in the de novo synthesis of tetrahydrobiopterin from GTP, with the other enzymes involved being EC 1.1.1.153 (sepiapterin reductase) and EC 4.2.3.12 (6-pyruvoyltetrahydropterin synthase) [2805].  
**References:** [326, 3208, 2805]

[EC 3.5.4.16 created 1972]

#### EC 3.5.4.17

- Accepted name:** adenosine-phosphate deaminase  
**Reaction:** (1)  $\text{AMP} + \text{H}_2\text{O} = \text{IMP} + \text{NH}_3$   
(2)  $\text{ADP} + \text{H}_2\text{O} = \text{IDP} + \text{NH}_3$   
(3)  $\text{ATP} + \text{H}_2\text{O} = \text{ITP} + \text{NH}_3$   
**Other name(s):** adenylate deaminase; adenine nucleotide deaminase; adenosine (phosphate) deaminase  
**Systematic name:** adenosine-phosphate aminohydrolase  
**Comments:** Acts on AMP, ADP, ATP,  $\text{NAD}^+$  and adenosine, in decreasing order of activity. The bacterial enzyme can also accept the deoxy derivatives. *cf.* EC 3.5.4.6, AMP deaminase.  
**References:** [2784, 3278]

[EC 3.5.4.17 created 1972, modified 1980, modified 2014]

#### EC 3.5.4.18

- Accepted name:** ATP deaminase  
**Reaction:**  $\text{ATP} + \text{H}_2\text{O} = \text{ITP} + \text{NH}_3$   
**Other name(s):** adenosine triphosphate deaminase  
**Systematic name:** ATP aminohydrolase  
**References:** [450]

[EC 3.5.4.18 created 1972]

#### EC 3.5.4.19

- Accepted name:** phosphoribosyl-AMP cyclohydrolase  
**Reaction:**  $1\text{-(5-phospho-}\beta\text{-D-ribose)}\text{-AMP} + \text{H}_2\text{O} = 1\text{-(5-phospho-}\beta\text{-D-ribose)}\text{-5-}[(5\text{-phospho-}\beta\text{-D-ribose)}\text{amino)methylideneamino]imidazole-4-carboxamide}$   
**Other name(s):** PRAMP-cyclohydrolase; phosphoribosyladenosine monophosphate cyclohydrolase; 1-(5-phospho-D-ribose)-AMP 1,6-hydrolase  
**Systematic name:** 1-(5-phospho- $\beta$ -D-ribose)-AMP 1,6-hydrolase  
**Comments:** The *Neurospora crassa* enzyme also catalyses the reactions of EC 1.1.1.23 (histidinol dehydrogenase) and EC 3.6.1.31 (phosphoribosyl-ATP diphosphatase).  
**References:** [1925]

[EC 3.5.4.19 created 1972, modified 1976, modified 1981, modified 2000]

#### EC 3.5.4.20

- Accepted name:** pyrithiamine deaminase  
**Reaction:**  $1\text{-(4-amino-2-methylpyrimid-5-ylmethyl)}\text{-3-(2-hydroxyethyl)}\text{-2-methylpyridinium} + \text{H}_2\text{O} = 1\text{-(4-hydroxy-2-methylpyrimid-5-ylmethyl)}\text{-3-(2-hydroxyethyl)}\text{-2-methylpyridinium} + \text{NH}_3$   
**Other name(s):** 1-(4-amino-2-methylpyrimid-5-ylmethyl)-3-( $\beta$ -hydroxyethyl)-2-methylpyridinium-bromide aminohydrolase  
**Systematic name:** 1-(4-amino-2-methylpyrimid-5-ylmethyl)-3-(2-hydroxyethyl)-2-methylpyridinium aminohydrolase  
**References:** [2673]

[EC 3.5.4.20 created 1972, modified 2014]

#### EC 3.5.4.21

- Accepted name:** creatinine deaminase  
**Reaction:**  $\text{creatinine} + \text{H}_2\text{O} = \text{N-methylhydantoin} + \text{NH}_3$   
**Other name(s):** creatinine hydrolase; creatinine desiminase

**Systematic name:** creatinine iminohydrolase  
**References:** [2823]

[EC 3.5.4.21 created 1972]

#### EC 3.5.4.22

**Accepted name:** 1-pyrroline-4-hydroxy-2-carboxylate deaminase  
**Reaction:** 1-pyrroline-4-hydroxy-2-carboxylate + H<sub>2</sub>O = 2,5-dioxopentanoate + NH<sub>3</sub>  
**Other name(s):** HPC deaminase; 1-pyrroline-4-hydroxy-2-carboxylate aminohydrolase (decyclizing)  
**Systematic name:** 1-pyrroline-4-hydroxy-2-carboxylate aminohydrolase (ring-opening)  
**References:** [2671, 2672]

[EC 3.5.4.22 created 1976]

#### EC 3.5.4.23

**Accepted name:** blasticidin-S deaminase  
**Reaction:** blasticidin S + H<sub>2</sub>O = deaminohydroxyblasticidin S + NH<sub>3</sub>  
**Systematic name:** blasticidin-S aminohydrolase  
**Comments:** Catalyses the deamination of the cytosine moiety of the antibiotics blasticidin S, cytomycin and acetylblasticidin S.  
**References:** [3245]

[EC 3.5.4.23 created 1976]

#### EC 3.5.4.24

**Accepted name:** sepiapterin deaminase  
**Reaction:** sepiapterin + H<sub>2</sub>O = xanthopterin-B2 + NH<sub>3</sub>  
**Systematic name:** sepiapterin aminohydrolase  
**Comments:** Also acts on isosepiapterin, but more slowly.  
**References:** [2992]

[EC 3.5.4.24 created 1976]

#### EC 3.5.4.25

**Accepted name:** GTP cyclohydrolase II  
**Reaction:** GTP + 3 H<sub>2</sub>O = formate + 2,5-diamino-6-hydroxy-4-(5-phospho-D-ribosylamino)pyrimidine + diphosphate  
**Other name(s):** guanosine triphosphate cyclohydrolase II; GTP-8-formylhydrolase  
**Systematic name:** GTP 7,8-8,9-dihydrolase (diphosphate-forming)  
**Comments:** Two C-N bonds are hydrolysed, releasing formate, with simultaneous removal of the terminal diphosphate.  
**References:** [797]

[EC 3.5.4.25 created 1984, modified 2011]

#### EC 3.5.4.26

**Accepted name:** diaminohydroxyphosphoribosylaminopyrimidine deaminase  
**Reaction:** 2,5-diamino-6-hydroxy-4-(5-phospho-D-ribosylamino)pyrimidine + H<sub>2</sub>O = 5-amino-6-(5-phospho-D-ribosylamino)uracil + NH<sub>3</sub>  
**Systematic name:** 2,5-diamino-6-hydroxy-4-(5-phospho-D-ribosylamino)pyrimidine 2-aminohydrolase  
**Comments:** The substrate is the product of EC 3.5.4.25 GTP cyclohydrolase II.  
**References:** [332]

[EC 3.5.4.26 created 1984, modified 2011]

#### EC 3.5.4.27

- Accepted name:** methenyltetrahydromethanopterin cyclohydrolase  
**Reaction:** 5,10-methenyl-5,6,7,8-tetrahydromethanopterin + H<sub>2</sub>O = 5-formyl-5,6,7,8-tetrahydromethanopterin  
**Other name(s):** 5,10-methenyltetrahydromethanopterin cyclohydrolase; N<sup>5</sup>,N<sup>10</sup>-methenyltetrahydromethanopterin cyclohydrolase; methenyl-H<sub>4</sub>MPT cyclohydrolase; 5,10-methenyltetrahydromethanopterin 10-hydrolase (decyclizing)  
**Systematic name:** 5,10-methenyltetrahydromethanopterin 10-hydrolase (ring-opening)  
**Comments:** Methanopterin is a pterin analogue. The enzyme is involved in the formation of methane from CO<sub>2</sub> in *Methanobacterium thermoautotrophicum*.  
**References:** [640]

[EC 3.5.4.27 created 1989]

#### EC 3.5.4.28

- Accepted name:** S-adenosylhomocysteine deaminase  
**Reaction:** S-adenosyl-L-homocysteine + H<sub>2</sub>O = S-inosyl-L-homocysteine + NH<sub>3</sub>  
**Other name(s):** adenosylhomocysteine deaminase  
**Systematic name:** S-adenosyl-L-homocysteine aminohydrolase  
**References:** [3351]

[EC 3.5.4.28 created 1992]

#### EC 3.5.4.29

- Accepted name:** GTP cyclohydrolase IIa  
**Reaction:** GTP + 3 H<sub>2</sub>O = 2-amino-5-formylamino-6-(5-phospho-D-ribosylamino)pyrimidin-4(3H)-one + 2 phosphate  
**Systematic name:** GTP 8,9-hydrolase (phosphate-forming)  
**Comments:** Requires Mg<sup>2+</sup>. This enzyme catalyses the hydrolysis of the imidazole ring of guanosine 5'-triphosphate, N<sup>7</sup>-methylguanosine 5'-triphosphate or inosine 5'-triphosphate. Xanthosine 5'-triphosphate and ATP are not substrates. It also catalyses the hydrolysis of diphosphate to form two equivalents of phosphate. Unlike GTP cyclohydrolase II (EC 3.5.4.25), this enzyme does not release formate, but does hydrolyse the diphosphate from GTP to phosphate.  
**References:** [963]

[EC 3.5.4.29 created 2003, modified 2011]

#### EC 3.5.4.30

- Accepted name:** dCTP deaminase (dUMP-forming)  
**Reaction:** dCTP + 2 H<sub>2</sub>O = dUMP + diphosphate + NH<sub>3</sub>  
**Systematic name:** dCTP aminohydrolase (dUMP-forming)  
**Comments:** Requires Mg<sup>2+</sup>. Is highly specific for dCTP as substrate as dCMP, CTP, CDP, CMP, cytosine or deoxycytosine are not deaminated. While most bacteria require two enzymes to form dUMP from dCTP (EC 3.5.4.13, dCTP deaminase and EC 3.6.1.23, dUTP diphosphatase), the archaeon *Methanocaldococcus jannaschii* uses a single enzyme to carry out both functions. This enzyme can also act as a dUTP diphosphatase, but more slowly.  
**References:** [1664]

[EC 3.5.4.30 created 2003]

#### EC 3.5.4.31

**Accepted name:** S-methyl-5'-thioadenosine deaminase  
**Reaction:** S-methyl-5'-thioadenosine + H<sub>2</sub>O = S-methyl-5'-thioinosine + NH<sub>3</sub>  
**Other name(s):** MTA deaminase; 5-methylthioadenosine deaminase  
**Systematic name:** S-methyl-5'-thioadenosine amidohydrolase  
**Comments:** The enzyme from *Thermotoga maritima* also functions as S-adenosylhomocysteine deaminase (EC 3.5.4.28) and has some activity against adenosine. Adenosine 5'-phosphate and S-adenosyl-L-methionine (SAM) are not substrates.  
**References:** [1130]

[EC 3.5.4.31 created 2011]

#### EC 3.5.4.32

**Accepted name:** 8-oxoguanine deaminase  
**Reaction:** 8-oxoguanine + H<sub>2</sub>O = urate + NH<sub>3</sub>  
**Other name(s):** 8-OGD  
**Systematic name:** 8-oxoguanine aminohydrolase  
**Comments:** Zn<sup>2+</sup> is bound in the active site. 8-Oxoguanine is formed via the oxidation of guanine within DNA by reactive oxygen species. If uncorrected, this modification leads to the incorporation of 8-oxoG:A mismatches and eventually to G:C to T:A transversions.  
**References:** [1027]

[EC 3.5.4.32 created 2012]

#### EC 3.5.4.33

**Accepted name:** tRNA(adenine<sup>34</sup>) deaminase  
**Reaction:** adenine<sup>34</sup> in tRNA + H<sub>2</sub>O = hypoxanthine<sup>34</sup> in tRNA + NH<sub>3</sub>  
**Other name(s):** tRNA:A34 deaminase; *tadA* protein; ADAT2-ADAT3 complex; TADA; tRNA adenosine deaminase arginine; AtTadA; *tadA/ecADAT2*; tRNA A:34 deaminase  
**Systematic name:** tRNA(adenine<sup>34</sup>) aminohydrolase  
**Comments:** The enzyme is involved in editing of tRNA. The active site contains Zn<sup>2+</sup> [2737].  
**References:** [2737, 575, 1564, 3207, 1633, 2357]

[EC 3.5.4.33 created 2013]

#### EC 3.5.4.34

**Accepted name:** tRNA<sup>Ala</sup>(adenine<sup>37</sup>) deaminase  
**Reaction:** adenine<sup>37</sup> in tRNA<sup>Ala</sup> + H<sub>2</sub>O = hypoxanthine<sup>37</sup> in tRNA<sup>Ala</sup> + NH<sub>3</sub>  
**Other name(s):** ADAT1; Tad1p  
**Systematic name:** tRNA<sup>Ala</sup>(adenine<sup>37</sup>) aminohydrolase  
**Comments:** The enzyme deaminates adenosine<sup>37</sup> to inosine in eukaryotic tRNA<sup>Ala</sup> [1762]. tRNA editing is strictly dependent on Mg<sup>2+</sup> [892].  
**References:** [1762, 892, 1410]

[EC 3.5.4.34 created 2013]

#### EC 3.5.4.35

**Accepted name:** tRNA(cytosine<sup>8</sup>) deaminase  
**Reaction:** cytosine<sup>8</sup> in tRNA + H<sub>2</sub>O = uracil<sup>8</sup> in tRNA + NH<sub>3</sub>  
**Other name(s):** CDAT8  
**Systematic name:** tRNA(cytosine<sup>8</sup>) aminohydrolase  
**Comments:** The enzyme from *Methanopyrus kandleri* specifically catalyses the deamination of cytosine at position 8 of tRNA in 30 different tRNAs. This cytosine-to-uracil editing guarantees the proper folding and functionality of the tRNAs.

**References:** [2370]

[EC 3.5.4.35 created 2013]

#### EC 3.5.4.36

**Accepted name:** mRNA(cytosine<sup>666</sup>) deaminase  
**Reaction:** cytosine<sup>666</sup> in apolipoprotein B mRNA + H<sub>2</sub>O = uracil<sup>666</sup> in apolipoprotein B mRNA + NH<sub>3</sub>  
**Other name(s):** APOBEC-1 (catalytic component of an RNA-editing complex); APOBEC1 (catalytic subunit); apolipoprotein B mRNA-editing enzyme 1 (catalytic component of an RNA-editing complex); *apoB* mRNA-editing enzyme catalytic polypeptide 1 (catalytic component of an RNA-editing complex); *apoB* mRNA editing complex; apolipoprotein B mRNA editing enzyme; REPR  
**Systematic name:** mRNA(cytosine<sup>666</sup>) aminohydrolase  
**Comments:** The apolipoprotein B mRNA editing enzyme complex catalyses the editing of apolipoprotein B mRNA at cytidine<sup>666</sup> to uridine, thereby transforming the codon for glutamine-2153 to a termination codon. Editing results in translation of a truncated apolipoprotein B isoform (*apoB*-48) with distinct functions in lipid transport. The catalytic component (APOBEC-1) contains zinc at the active site [145].  
**References:** [427, 846, 145, 426]

[EC 3.5.4.36 created 2013]

#### EC 3.5.4.37

**Accepted name:** double-stranded RNA adenine deaminase  
**Reaction:** adenine in double-stranded RNA + H<sub>2</sub>O = hypoxanthine in double-stranded RNA + NH<sub>3</sub>  
**Other name(s):** ADAR; double-stranded RNA adenosine deaminase; dsRAD; dsRNA adenosine deaminase; DRADA1; double-stranded RNA-specific adenosine deaminase  
**Systematic name:** double-stranded RNA adenine aminohydrolase  
**Comments:** This eukaryotic enzyme is involved in RNA editing. It destabilizes double-stranded RNA through conversion of adenosine to inosine. Inositol hexakisphosphate is required for activity [1763].  
**References:** [1187, 2127, 3211, 1763]

[EC 3.5.4.37 created 2013]

#### EC 3.5.4.38

**Accepted name:** single-stranded DNA cytosine deaminase  
**Reaction:** cytosine in single-stranded DNA + H<sub>2</sub>O = uracil in single-stranded DNA + NH<sub>3</sub>  
**Other name(s):** AID; activation-induced deaminase; AICDA (gene name); activation-induced cytidine deaminase  
**Systematic name:** single-stranded DNA cytosine aminohydrolase  
**Comments:** The enzyme exclusively catalyses deamination of cytosine in single-stranded DNA. It preferentially deaminates five-nucleotide bubbles. The optimal target consists of a single-stranded NWRCN motif (W = A or T, R = A or G) [1601]. The enzyme initiates antibody diversification processes by deaminating immunoglobulin sequences.  
**References:** [2709, 1601, 278, 1600, 3062]

[EC 3.5.4.38 created 2013]

#### EC 3.5.4.39

**Accepted name:** GTP cyclohydrolase IV  
**Reaction:** GTP + H<sub>2</sub>O = 7,8-dihydroneopterin 2',3'-cyclic phosphate + formate + diphosphate  
**Other name(s):** MptA; GTP cyclohydrolase MptA  
**Systematic name:** GTP 7,8-8,9-dihydrolase (cyclizing, formate-releasing, diphosphate-releasing)  
**Comments:** Requires Fe<sup>2+</sup>. A zinc protein. The enzyme is involved in methanopterin biosynthesis in methanogenic archaea. *cf.* GTP cyclohydrolase I (EC 3.5.4.16), GTP cyclohydrolase II (EC 3.5.4.25) and GTP cyclohydrolase IIa (EC 3.5.4.29).



**References:** [987]

[EC 3.5.4.39 created 2013]

#### EC 3.5.4.40

**Accepted name:** aminodeoxyfutalosine deaminase  
**Reaction:** 6-amino-6-deoxyfutalosine + H<sub>2</sub>O = futalosine + NH<sub>3</sub>  
**Other name(s):** AFL deaminase; aminofutalosine deaminase; *mqnX* (gene name)  
**Systematic name:** 6-amino-6-deoxyfutalosine deaminase  
**Comments:** The enzyme, found in several bacterial species, is part of the futalosine pathway for menaquinone biosynthesis.  
**References:** [65, 935]

[EC 3.5.4.40 created 2014]

#### EC 3.5.4.41

**Accepted name:** 5'-deoxyadenosine deaminase  
**Reaction:** 5'-deoxyadenosine + H<sub>2</sub>O = 5'-deoxyinosine + NH<sub>3</sub>  
**Other name(s):** MJ1541 (gene name); DadD  
**Systematic name:** 5'-deoxyadenosine aminohydrolase  
**Comments:** The enzyme from the archaeon *Methanocaldococcus jannaschii* is involved in the recycling of 5'-deoxyadenosine.  
**References:** [1912]

[EC 3.5.4.41 created 2014]

#### EC 3.5.4.42

**Accepted name:** *N*-isopropylammelide isopropylaminohydrolase  
**Reaction:** *N*-isopropylammelide + H<sub>2</sub>O = cyanuric acid + isopropylamine  
**Other name(s):** *atzC* (gene name)  
**Systematic name:** *N*-isopropylammelide isopropylaminohydrolase  
**Comments:** Requires Zn<sup>2+</sup>. This bacterial enzyme is involved in degradation of the herbicide atrazine. It can hydrolyse other *N*-substituted amino dihydroxy-*s*-triazine molecules, and prefers substrates with linear *N*-alkyl groups to those with branched alkyl groups.  
**References:** [2490, 2620, 130]

[EC 3.5.4.42 created 2000 as EC 3.5.99.4, transferred 2016 to EC 3.5.4.42]

#### EC 3.5.4.43

**Accepted name:** hydroxydechloroatrazine ethylaminohydrolase  
**Reaction:** hydroxyatrazine + H<sub>2</sub>O = *N*-isopropylammelide + ethylamine  
**Other name(s):** *atzB* (gene name); 2,4-dihydroxy-6-(isopropylamino)-1,3,5-triazine ethylaminohydrolase  
**Systematic name:** hydroxyatrazine ethylaminohydrolase  
**Comments:** Contains Zn<sup>2+</sup>. This bacterial enzyme is involved in degradation of the herbicide atrazine. The enzyme has a broad substrate range, and requires a monohydroxylated *s*-triazine ring with a minimum of one primary or secondary amine substituent and either a chloride or amine leaving group. It catalyses both deamination and dechlorination reactions.  
**References:** [269, 2601]

[EC 3.5.4.43 created 2000 as EC 3.5.99.3, transferred 2016 to EC 3.5.4.43]

#### EC 3.5.4.44

**Accepted name:** ectoine hydrolase  
**Reaction:** ectoine + H<sub>2</sub>O = (2*S*)-2-acetamido-4-aminobutanoate  
**Other name(s):** *doeA* (gene name)  
**Systematic name:** ectoine aminohydrolase  
**Comments:** The enzyme, found in some halophilic bacteria, is involved in the degradation of the compatible solute ectoine. The enzyme, which belongs to peptidase family M24, only acts in the direction of ectoine hydrolysis. It also produces smaller amounts of (2*S*)-4-acetamido-2-aminobutanoate, which is recycled back to ectoine by EC 4.2.1.108, ectoine synthase.  
**References:** [2597]

[EC 3.5.4.44 created 2017]

#### EC 3.5.4.45

**Accepted name:** melamine deaminase  
**Reaction:** (1) melamine + H<sub>2</sub>O = ammeline + NH<sub>3</sub>  
(2) ammeline + H<sub>2</sub>O = ammelide + NH<sub>3</sub>  
**Other name(s):** *triA* (gene name)  
**Systematic name:** melamine aminohydrolase  
**Comments:** The enzyme, isolated from the bacterium *Acidovorax citrulli*, performs the deamination of melamine 15-fold faster than the deamination of ammeline. It also has activity with 2-chloro-4,6-diamino-*s*-triazine, but has no activity toward halo-substituted triazine ring compounds such as atrazine (*cf.* EC 3.8.1.8, atrazine chlorohydrolase).  
**References:** [2602]

[EC 3.5.4.45 created 2017]

#### EC 3.5.4.46

**Accepted name:** cAMP deaminase  
**Reaction:** 3',5'-cyclic AMP + H<sub>2</sub>O = 3',5'-cyclic IMP + NH<sub>3</sub>  
**Other name(s):** cyclic adenylylase; CadD  
**Systematic name:** 3',5'-cyclic AMP aminohydrolase  
**Comments:** Requires Zn<sup>2+</sup>. The enzyme, isolated from the bacterium *Leptospira interrogans*, is specific for cAMP.  
**References:** [934]

[EC 3.5.4.46 created 2017]

### EC 3.5.5 In nitriles

#### EC 3.5.5.1

**Accepted name:** nitrilase  
**Reaction:** a nitrile + 2 H<sub>2</sub>O = a carboxylate + NH<sub>3</sub>  
**Other name(s):** acetonitrilase; benzonitrilase  
**Systematic name:** nitrile aminohydrolase  
**Comments:** Acts on a wide range of aromatic nitriles including (indol-3-yl)acetonitrile, and also on some aliphatic nitriles, and on the corresponding acid amides. *cf.* EC 4.2.1.84 nitrile hydratase.  
**References:** [1058, 2910, 2220]

[EC 3.5.5.1 created 1965, modified 1989]

#### EC 3.5.5.2

**Accepted name:** ricinine nitrilase

**Reaction:** ricinine + 2 H<sub>2</sub>O = 3-carboxy-4-methoxy-*N*-methyl-2-pyridone + NH<sub>3</sub>  
**Systematic name:** ricinine aminohydrolase  
**References:** [2442, 1171, 2220]

[EC 3.5.5.2 created 1972]

[3.5.5.3 *Transferred entry. cyanate hydrolase. Now EC 4.2.1.104, cyanate hydratase*]

[EC 3.5.5.3 created 1972, deleted 1990]

#### EC 3.5.5.4

**Accepted name:** cyanoalanine nitrilase  
**Reaction:** 3-cyano-L-alanine + 2 H<sub>2</sub>O = L-aspartate + NH<sub>3</sub> (overall reaction)  
(1a) 3-cyano-L-alanine + H<sub>2</sub>O = L-asparagine  
(1b) L-asparagine + H<sub>2</sub>O = L-aspartate + NH<sub>3</sub>  
**Other name(s):** β-cyanoalanine nitrilase  
**Systematic name:** 3-cyano-L-alanine aminohydrolase  
**Comments:** L-Asparagine is formed as an intermediate. *cf.* EC 4.2.1.65, 3-cyanoalanine hydratase and EC 3.5.1.1, asparaginase.  
**References:** [3256]

[EC 3.5.5.4 created 1986]

#### EC 3.5.5.5

**Accepted name:** arylacetone nitrilase  
**Reaction:** 4-chlorophenylacetone nitrile + 2 H<sub>2</sub>O = 4-chlorophenylacetate + NH<sub>3</sub>  
**Systematic name:** arylacetone nitrile aminohydrolase  
**Comments:** Requires thiol compounds. Also hydrolyses other 4-substituted phenylacetone nitriles, thien-2-ylacetone nitrile, tolylacetone nitriles, and, more slowly, benzyl cyanide.  
**References:** [1848, 2024]

[EC 3.5.5.5 created 1992]

#### EC 3.5.5.6

**Accepted name:** bromoxynil nitrilase  
**Reaction:** 3,5-dibromo-4-hydroxybenzoyl nitrile + 2 H<sub>2</sub>O = 3,5-dibromo-4-hydroxybenzoate + NH<sub>3</sub>  
**Systematic name:** 3,5-dibromo-4-hydroxybenzoyl nitrile aminohydrolase  
**Comments:** Involved in the bacterial degradation of the herbicide bromoxynil. Highly specific.  
**References:** [2754]

[EC 3.5.5.6 created 1992]

#### EC 3.5.5.7

**Accepted name:** aliphatic nitrilase  
**Reaction:** R-CN + 2 H<sub>2</sub>O = R-COOH + NH<sub>3</sub>  
**Systematic name:** aliphatic nitrile aminohydrolase  
**Comments:** Preferentially hydrolyses aliphatic nitriles, some of which are apparently not substrates for other known nitrilases (EC 3.5.5.1). Substrates include crotononitrile, acrylonitrile and glutaronitrile.  
**References:** [1491, 2220]

[EC 3.5.5.7 created 1999]

#### EC 3.5.5.8

**Accepted name:** thiocyanate hydrolase  
**Reaction:** thiocyanate + 2 H<sub>2</sub>O = carbonyl sulfide + NH<sub>3</sub> + HO<sup>-</sup>  
**Systematic name:** thiocyanate aminohydrolase  
**Comments:** The enzyme from *Thiobacillus thioparus* catalyses the first step in the degradation of thiocyanate.  
**References:** [1394, 1395]

[EC 3.5.5.8 created 2000]

## EC 3.5.99 In other compounds

### EC 3.5.99.1

**Accepted name:** riboflavinase  
**Reaction:** riboflavin + H<sub>2</sub>O = ribitol + lumichrome  
**Systematic name:** riboflavin hydrolase  
**References:** [3254]

[EC 3.5.99.1 created 1961]

### EC 3.5.99.2

**Accepted name:** aminopyrimidine aminohydrolase  
**Reaction:** (1) 4-amino-5-aminomethyl-2-methylpyrimidine + H<sub>2</sub>O = 4-amino-5-hydroxymethyl-2-methylpyrimidine + NH<sub>3</sub>  
(2) thiamine + H<sub>2</sub>O = 4-amino-5-hydroxymethyl-2-methylpyrimidine + 5-(2-hydroxyethyl)-4-methylthiazole  
**Other name(s):** thiaminase; thiaminase II; *tenA* (gene name)  
**Systematic name:** 4-amino-5-aminomethyl-2-methylpyrimidine aminohydrolase  
**Comments:** Previously known as thiaminase II, this enzyme is involved in the regeneration of the thiamine pyrimidine from degraded products, rather than in thiamine degradation, and participates in thiamine salvage pathways.  
**References:** [849, 1231, 2940, 191, 1317, 1318, 809]

[EC 3.5.99.2 created 1961, modified 2011]

[3.5.99.3 *Transferred entry. hydroxydechloroatrazine ethylaminohydrolase. Now EC 3.5.4.43, hydroxydechloroatrazine ethylaminohydrolase*]

[EC 3.5.99.3 created 2000, deleted 2016]

[3.5.99.4 *Transferred entry. N-isopropylammelide isopropylaminohydrolase. Now EC 3.5.4.42, N-isopropylammelide isopropylaminohydrolase*]

[EC 3.5.99.4 created 2000, deleted 2016]

### EC 3.5.99.5

**Accepted name:** 2-aminomuconate deaminase  
**Reaction:** 2-aminomuconate + H<sub>2</sub>O = (3E)-2-oxohex-3-enedioate + NH<sub>3</sub>  
**Other name(s):** *amnD* (gene name); *nbaF* (gene name)  
**Systematic name:** 2-aminomuconate aminohydrolase  
**Comments:** 2-Aminomuconate is an intermediate in the bacterial biodegradation of nitrobenzene. The enzyme has been isolated from several species, including *Pseudomonas pseudocaligenes* JS45, *Pseudomonas fluorescens* KU-7, *Pseudomonas* sp. AP3 and *Burkholderia cenocepacia* J2315. The reaction is spontaneous in acid conditions.  
**References:** [1092, 1093, 2857, 2004]

[EC 3.5.99.5 created 2000, modified 2012]

### EC 3.5.99.6

- Accepted name:** glucosamine-6-phosphate deaminase  
**Reaction:**  $\alpha$ -D-glucosamine 6-phosphate + H<sub>2</sub>O = D-fructose 6-phosphate + NH<sub>3</sub>  
**Other name(s):** glucosaminophosphate isomerase; glucosamine-6-phosphate isomerase; phosphoglucosaminomerase; glucosamine phosphate deaminase; aminodeoxyglucosephosphate isomerase; phosphoglucosamine isomerase; 2-amino-2-deoxy-D-glucose-6-phosphate aminohydrolase (ketol isomerizing)  
**Systematic name:** 2-amino-2-deoxy- $\alpha$ -D-glucose-6-phosphate aminohydrolase (ketol isomerizing)  
**Comments:** The enzyme uses ring opening and isomerization of the aldose-ketose type to convert the -CH(-NH<sub>2</sub>)-CH=O group of glucosamine 6-phosphate into -C(=NH)-CH<sub>2</sub>-OH, forming 2-deoxy-2-imino-D-arabino-hexitol, which then hydrolyses to yield fructose 6-phosphate and ammonia. *N*-Acetyl-D-glucosamine 6-phosphate, which is not broken down, activates the enzyme.  
**References:** [3209, 469, 2246, 1717]

[EC 3.5.99.6 created 1961 as EC 5.3.1.10, transferred 2000 to EC 3.5.99.6]

### EC 3.5.99.7

- Accepted name:** 1-aminocyclopropane-1-carboxylate deaminase  
**Reaction:** 1-aminocyclopropane-1-carboxylate + H<sub>2</sub>O = 2-oxobutanoate + NH<sub>3</sub> (overall reaction)  
(1a) 1-aminocyclopropane-1-carboxylate = 2-aminobut-2-enoate  
(1b) 2-aminobut-2-enoate = 2-iminobutanoate (spontaneous)  
(1c) 2-iminobutanoate + H<sub>2</sub>O = 2-oxobutanoate + NH<sub>3</sub> (spontaneous)  
**Other name(s):** 1-aminocyclopropane-1-carboxylate endolyase (deaminating); ACC deaminase; 1-aminocyclopropane carboxylic acid deaminase  
**Systematic name:** 1-aminocyclopropane-1-carboxylate aminohydrolase (isomerizing)  
**Comments:** A pyridoxal 5'-phosphate enzyme. The enzyme, found in certain soil bacteria and fungi, catalyses the ring opening of 1-aminocyclopropane-1-carboxylate, the immediate precursor to ethylene, an important plant hormone that regulates fruit ripening and other processes. The enzyme releases an unstable enamine product that tautomerizes to an imine form, which undergoes a hydrolytic deamination. The latter reaction, which can occur spontaneously, can also be catalysed by EC 3.5.99.10, 2-iminobutanoate/2-iminopropanoate deaminase. The enzyme has been used to make fruit ripening dependent on externally added ethylene, as it removes the substrate for endogenous ethylene formation.  
**References:** [1169, 3262, 2908]

[EC 3.5.99.7 created 1981 as EC 4.1.99.4, transferred 2002 to EC 3.5.99.7, modified 2014]

### EC 3.5.99.8

- Accepted name:** 5-nitroanthranilic acid aminohydrolase  
**Reaction:** 5-nitroanthranilate + H<sub>2</sub>O = 5-nitrosalicylate + NH<sub>3</sub>  
**Other name(s):** *naaA* (gene name); 5NAA deaminase  
**Systematic name:** 5-nitroanthranilate amidohydrolase  
**Comments:** The enzyme catalyses the initial step in biodegradation of 5-nitroanthranilic acid by *Bradyrhizobium* sp. strain JS329.  
**References:** [2338]

[EC 3.5.99.8 created 2011]

### EC 3.5.99.9

- Accepted name:** 2-nitroimidazole nitrohydrolase  
**Reaction:** 2-nitroimidazole + H<sub>2</sub>O = imidazol-2-one + nitrite  
**Other name(s):** NnhA; 2NI nitrohydrolase; 2NI denitrase

**Systematic name:** 2-nitroimidazole nitrohydrolase  
**Comments:** The enzyme catalyses the initial step in the biodegradation of 2-nitroimidazole by the soil bacterium *Mycobacterium* sp. JS330  
**References:** [2339]

[EC 3.5.99.9 created 2012]

#### EC 3.5.99.10

**Accepted name:** 2-iminobutanoate/2-iminopropanoate deaminase  
**Reaction:** (1) 2-iminobutanoate + H<sub>2</sub>O = 2-oxobutanoate + NH<sub>3</sub>  
(2) 2-iminopropanoate + H<sub>2</sub>O = pyruvate + NH<sub>3</sub>  
**Other name(s):** *yjgF* (gene name); *ridA* (gene name); enamine/imine deaminase (ambiguous)  
**Systematic name:** 2-iminobutanoate aminohydrolase  
**Comments:** This enzyme, which has been found in all species and tissues examined, catalyses the hydrolytic deamination of imine intermediates formed by several types of pyridoxal-5'-phosphate-dependent dehydratases, such as EC 4.3.1.19, threonine ammonia-lyase and EC 4.3.1.17, L-serine ammonia-lyase. The reactions, which can occur spontaneously, are accelerated to minimize the cellular damage that could be caused by these reactive intermediates.  
**References:** [1595]

[EC 3.5.99.10 created 2014]

#### EC 3.5.99.11

**Accepted name:** 2-aminomuconate deaminase (2-hydroxymuconate-forming)  
**Reaction:** 2-aminomuconate + H<sub>2</sub>O = (2Z,4E)-2-hydroxyhexa-2,4-dienedioate + NH<sub>3</sub>  
**Other name(s):** *cnbZ* (gene name)  
**Systematic name:** 2-aminomuconate aminohydrolase [(2Z,4E)-2-hydroxyhexa-2,4-dienedioate-forming]  
**Comments:** The enzyme, characterized from the bacterium *Comamonas testosteroni* CNB-1, converts 2-aminomuconate to 2-hydroxyhexa-2,4-dienedioate, unlike the enzymes from *Pseudomonas*, which produce (3E)-2-oxohex-3-enedioate (see EC 3.5.99.5, 2-aminomuconate deaminase). The enzyme also acts on 2-amino-5-chloromuconate.  
**References:** [1723]

[EC 3.5.99.11 created 2016 as EC 3.5.1.120, transferred 2017 to EC 3.5.99.11]

## EC 3.6 Acting on acid anhydrides

To this subclass belong mainly the enzymes acting on diphosphate bonds in compounds such as nucleoside di- and tri-phosphates (EC 3.6.1), on sulfonyl-containing anhydrides such as adenylylsulfate (EC 3.6.2) and on acid anhydrides; catalysing transmembrane movement of substances (EC 3.6.3).

### EC 3.6.1 In phosphorus-containing anhydrides

#### EC 3.6.1.1

**Accepted name:** inorganic diphosphatase  
**Reaction:** diphosphate + H<sub>2</sub>O = 2 phosphate  
**Systematic name:** diphosphate phosphohydrolase  
**Comments:** Specificity varies with the source and with the activating metal ion. The enzyme from some sources may be identical with EC 3.1.3.1 (alkaline phosphatase) or EC 3.1.3.9 (glucose-6-phosphatase). *cf.* EC 7.1.3.1, H<sup>+</sup>-exporting diphosphatase.  
**References:** [120, 1556, 2355]

[EC 3.6.1.1 created 1961, modified 2000, modified 2018]

#### EC 3.6.1.2

**Accepted name:** trimetaphosphatase  
**Reaction:** trimetaphosphate + H<sub>2</sub>O = triphosphate  
**Other name(s):** inorganic trimetaphosphatase  
**Systematic name:** trimetaphosphate hydrolase  
**References:** [1513, 1898]

[EC 3.6.1.2 created 1961]

#### EC 3.6.1.3

**Accepted name:** adenosinetriphosphatase  
**Reaction:** ATP + H<sub>2</sub>O = ADP + phosphate  
**Other name(s):** adenylpyrophosphatase; ATP monophosphatase; triphosphatase; ATPase (ambiguous); SV40 T-antigen; adenosine 5'-triphosphatase; ATP hydrolase; complex V (mitochondrial electron transport); (Ca<sup>2+</sup> + Mg<sup>2+</sup>)-ATPase; HCO<sub>3</sub><sup>-</sup>-ATPase; adenosine triphosphatase  
**Systematic name:** ATP phosphohydrolase  
**Comments:** Many enzymes previously listed under this number are now listed separately under EC 3.6.3 and EC 3.6.4.  
**References:** [888, 1437, 1821, 2101, 2427, 2925]

[EC 3.6.1.3 created 1961 (EC 3.6.1.4 created 1961, incorporated 1965)]

[3.6.1.4 Deleted entry. *adenosinetriphosphatase (Mg-activated)*. Now included with EC 3.6.1.3 *adenosinetriphosphatase*]

[EC 3.6.1.4 created 1961, deleted 1965]

#### EC 3.6.1.5

**Accepted name:** apyrase  
**Reaction:** a nucleoside 5'-triphosphate + 2 H<sub>2</sub>O = a nucleoside 5'-phosphate + 2 phosphate (overall reaction)  
(1a) a nucleoside 5'-triphosphate + H<sub>2</sub>O = a nucleoside 5'-diphosphate + phosphate  
(1b) a nucleoside 5'-diphosphate + H<sub>2</sub>O = a nucleoside 5'-phosphate + phosphate  
**Other name(s):** ATP-diphosphatase; adenosine diphosphatase; ADPase; ATP diphosphohydrolase [ambiguous]  
**Systematic name:** nucleoside triphosphate phosphohydrolase (nucleoside monophosphate-forming)  
**Comments:** Apyrases are active against both di- and triphosphate nucleotides (NDPs and NTPs) and hydrolyse NTPs to nucleotide monophosphates (NMPs) in two distinct successive phosphate-releasing steps, with NDPs as intermediates. They differ from ATPases, which specifically hydrolyse ATP, by hydrolysing both ATP and ADP. The eukaryotic enzymes requires Ca<sup>2+</sup>, but Mg<sup>2+</sup> can substitute. Most of the ecto-ATPases that occur on the cell surface and hydrolyse extracellular nucleotides belong to this enzyme family.  
**References:** [1536, 1681, 418, 444, 3119, 872, 3237]

[EC 3.6.1.5 created 1961, modified 1976, modified 2000, modified 2013]

#### EC 3.6.1.6

**Accepted name:** nucleoside diphosphate phosphatase  
**Reaction:** a nucleoside diphosphate + H<sub>2</sub>O = a nucleoside phosphate + phosphate  
**Other name(s):** nucleoside-diphosphatase; thiaminpyrophosphatase; UDPase; inosine diphosphatase; adenosine diphosphatase; IDPase; ADPase; adenosinepyrophosphatase; guanosine diphosphatase; guanosine 5'-diphosphatase; inosine 5'-diphosphatase; uridine diphosphatase; uridine 5'-diphosphatase; type B nucleoside diphosphatase; GDPase; CDPase; nucleoside 5'-diphosphatase; type L nucleoside diphosphatase; NDPase; nucleoside diphosphate phosphohydrolase  
**Systematic name:** nucleoside-diphosphate phosphohydrolase

**Comments:** The enzyme, which appears to be limited to metazoa, acts on multiple nucleoside diphosphates as well as on D-ribose 5-diphosphate. Specificity depends on species and isoform.

**References:** [913, 1176, 3283, 731, 3005]

[EC 3.6.1.6 created 1961]

#### EC 3.6.1.7

**Accepted name:** acylphosphatase

**Reaction:** an acylphosphate + H<sub>2</sub>O = a carboxylate + phosphate

**Other name(s):** acetylphosphatase; 1,3-diphosphoglycerate phosphatase; acetic phosphatase; Ho 1-3; GP 1-3

**Systematic name:** acylphosphate phosphohydrolase

**References:** [2359, 2367, 2368, 2641]

[EC 3.6.1.7 created 1961]

#### EC 3.6.1.8

**Accepted name:** ATP diphosphatase

**Reaction:** ATP + H<sub>2</sub>O = AMP + diphosphate

**Other name(s):** ATPase (ambiguous); ATP pyrophosphatase; adenosine triphosphate pyrophosphatase; ATP diphosphohydrolase (ambiguous)

**Systematic name:** ATP diphosphohydrolase (diphosphate-forming)

**Comments:** Also acts on ITP, GTP, CTP and UTP.

**References:** [1123, 1329]

[EC 3.6.1.8 created 1961]

#### EC 3.6.1.9

**Accepted name:** nucleotide diphosphatase

**Reaction:** a nucleoside triphosphate + H<sub>2</sub>O = a nucleotide + diphosphate

**Other name(s):** ENPP1 (gene name); nucleotide pyrophosphatase; nucleotide-sugar pyrophosphatase; nucleoside-triphosphate diphosphatase

**Systematic name:** nucleoside-triphosphate diphosphohydrolase

**Comments:** The enzyme preferentially hydrolyses ATP, but can also hydrolyse other nucleoside 5' triphosphates such as GTP, CTP, TTP and UTP to their corresponding monophosphates. *In vitro* the enzyme also acts as a nucleotidohydrolase on ADP, NAD<sup>+</sup>, NADP<sup>+</sup>, FAD, and CoA.

**References:** [424, 1418, 1597, 3325]

[EC 3.6.1.9 created 1961 (EC 3.6.1.19 created 1972, incorporated 2016), modified 2016]

#### EC 3.6.1.10

**Accepted name:** endopolyphosphatase

**Reaction:** polyphosphate + *n* H<sub>2</sub>O = (*n*+1) oligophosphate

**Other name(s):** polyphosphate depolymerase; metaphosphatase; polyphosphatase; polymetaphosphatase

**Systematic name:** polyphosphate polyphosphohydrolase

**Comments:** The product contains 4 or 5 phosphate residues.

**References:** [1793, 1844]

[EC 3.6.1.10 created 1961]

#### EC 3.6.1.11

**Accepted name:** exopolyphosphatase

**Reaction:** (polyphosphate)<sub>*n*</sub> + H<sub>2</sub>O = (polyphosphate)<sub>*n*-1</sub> + phosphate



**Other name(s):** metaphosphatase; acid phosphoanhydride phosphohydrolase; Gra-Pase  
**Systematic name:** polyphosphate phosphohydrolase  
**References:** [993, 1536, 1793]

[EC 3.6.1.11 created 1965]

#### EC 3.6.1.12

**Accepted name:** dCTP diphosphatase  
**Reaction:**  $\text{dCTP} + \text{H}_2\text{O} = \text{dCMP} + \text{diphosphate}$   
**Other name(s):** DCTPP1 (gene name); deoxycytidine-triphosphatase; dCTPase; dCTP pyrophosphatase; deoxycytidine triphosphatase; deoxy-CTPase  
**Systematic name:** dCTP nucleotidohydrolase  
**Comments:** The mammalian enzyme also displays weak activity against dTTP and dATP, but none against dGTP. Activity is highest with analogs including 5-iodo-dCTP and 5-methyl-dCTP.  
**References:** [3343, 1982, 3217, 2109, 2410]

[EC 3.6.1.12 created 1965]

#### EC 3.6.1.13

**Accepted name:** ADP-ribose diphosphatase  
**Reaction:**  $\text{ADP-D-ribose} + \text{H}_2\text{O} = \text{AMP} + \text{D-ribose 5-phosphate}$   
**Other name(s):** ADPribose pyrophosphatase; adenosine diphosphoribose pyrophosphatase; ADPR-PPase; ADP-ribose ribophosphohydrolase  
**Systematic name:** ADP-D-ribose ribophosphohydrolase  
**References:** [630]

[EC 3.6.1.13 created 1965]

#### EC 3.6.1.14

**Accepted name:** adenosine-tetraphosphatase  
**Reaction:**  $\text{adenosine 5'-tetraphosphate} + \text{H}_2\text{O} = \text{ATP} + \text{phosphate}$   
**Systematic name:** adenosine-tetraphosphate phosphohydrolase  
**Comments:** Also acts on inosine tetraphosphate and tripolyphosphate but shows little or no activity with other nucleotides or polyphosphates.  
**References:** [2692]

[EC 3.6.1.14 created 1972]

#### EC 3.6.1.15

**Accepted name:** nucleoside-triphosphate phosphatase  
**Reaction:**  $\text{a nucleoside triphosphate} + \text{H}_2\text{O} = \text{a nucleoside diphosphate} + \text{phosphate}$   
**Other name(s):** nucleoside-triphosphatase; nucleoside triphosphate phosphohydrolase; nucleoside-5-triphosphate phosphohydrolase; nucleoside 5-triphosphatase; unspecific diphosphate phosphohydrolase  
**Systematic name:** nucleoside-triphosphate phosphohydrolase  
**Comments:** The enzyme is found in eukaryotes and thermophilic bacteria, but appears to be absent from mesophilic bacteria. Also hydrolyses nucleoside diphosphates, thiamine diphosphate and FAD. The enzyme from the plant *Pisum sativum* (garden pea) is regulated by calmodulin [1195].  
**References:** [292, 1660, 1840, 2942, 1195, 1477, 2290]

[EC 3.6.1.15 created 1972]

#### EC 3.6.1.16

**Accepted name:** CDP-glycerol diphosphatase  
**Reaction:** CDP-glycerol + H<sub>2</sub>O = CMP + *sn*-glycerol 3-phosphate  
**Other name(s):** CDP-glycerol pyrophosphatase; cytidine diphosphoglycerol pyrophosphatase  
**Systematic name:** CDP-glycerol phosphoglycerohydrolase  
**References:** [926]

[EC 3.6.1.16 created 1972]

#### EC 3.6.1.17

**Accepted name:** bis(5'-nucleosyl)-tetraphosphatase (asymmetrical)  
**Reaction:** P<sup>1</sup>,P<sup>4</sup>-bis(5'-guanosyl) tetraphosphate + H<sub>2</sub>O = GTP + GMP  
**Other name(s):** bis(5'-guanosyl)-tetraphosphatase; bis(5'-adenosyl)-tetraphosphatase; diguanosinetetraphosphatase (asymmetrical); dinucleosidetetraphosphatase (asymmetrical); diadenosine P<sup>1</sup>,P<sup>4</sup>-tetraphosphatase; dinucleoside tetraphosphatase; 1-*P*,4-*P*-bis(5'-nucleosyl)-tetraphosphate nucleotidohydrolase  
**Systematic name:** P<sup>1</sup>,P<sup>4</sup>-bis(5'-nucleosyl)-tetraphosphate nucleotidohydrolase  
**Comments:** Also acts on bis(5'-xanthosyl)-tetraphosphate and, more slowly, on bis(5'-adenosyl)-tetraphosphate and bis(5'-uridylyl)-tetraphosphate [*cf.* EC 3.6.1.41 bis(5'-nucleosyl)-tetraphosphatase (symmetrical)]  
**References:** [1302, 3029, 3128]

[EC 3.6.1.17 created 1972, modified 1976, modified 1986]

#### EC 3.6.1.18

**Accepted name:** FAD diphosphatase  
**Reaction:** FAD + H<sub>2</sub>O = AMP + FMN  
**Other name(s):** FAD pyrophosphatase; riboflavin adenine dinucleotide pyrophosphatase; flavin adenine dinucleotide pyrophosphatase; riboflavine adenine dinucleotide pyrophosphatase; flavine adenine dinucleotide pyrophosphatase  
**Systematic name:** FAD nucleotidohydrolase  
**Comments:** The plant enzyme also hydrolyses NAD<sup>+</sup> and NADH; the animal enzyme hydrolyses NAD<sup>+</sup> and CoA at about half of the rate of hydrolysis of FAD. May be identical with EC 3.6.1.9 nucleotide diphosphatase.  
**References:** [2379, 2636]

[EC 3.6.1.18 created 1972]

[3.6.1.19 *Transferred entry. nucleoside-triphosphate diphosphatase. Now EC 3.6.1.9, nucleotide diphosphatase*]

[EC 3.6.1.19 created 1972, deleted 2016]

#### EC 3.6.1.20

**Accepted name:** 5'-acylphosphoadenosine hydrolase  
**Reaction:** 5'-acylphosphoadenosine + H<sub>2</sub>O = AMP + a carboxylate  
**Other name(s):** 5-phosphoadenosine hydrolase  
**Systematic name:** 5'-acylphosphoadenosine acylhydrolase  
**Comments:** Also acts on inosine and uridine compounds.  
**References:** [1414]

[EC 3.6.1.20 created 1972]

#### EC 3.6.1.21

**Accepted name:** ADP-sugar diphosphatase  
**Reaction:** ADP-sugar + H<sub>2</sub>O = AMP + α-D-aldose 1-phosphate  
**Other name(s):** ADP-sugar pyrophosphatase; adenosine diphosphosugar pyrophosphatase  
**Systematic name:** ADP-sugar sugarphosphohydrolase

**Comments:** Has a specificity that is distinct from that of UDP-sugar diphosphatase (EC 3.6.1.45).

**References:** [2451]

[EC 3.6.1.21 created 1972, modified 1999]

#### EC 3.6.1.22

**Accepted name:** NAD<sup>+</sup> diphosphatase

**Reaction:** NAD(H) + H<sub>2</sub>O = AMP + NMN(H)

**Other name(s):** NPY1 (gene name); *nudC* (gene name); NUDT7 (gene name); nicotinamide adenine dinucleotide pyrophosphatase; NADP pyrophosphatase; NADH pyrophosphatase; NAD<sup>+</sup> phosphohydrolase

**Systematic name:** NAD(H) phosphohydrolase

**Comments:** This enzyme, described from plants, animals, and bacteria, can act on both reduced and oxidized forms of its substrate, although enzymes from different organisms have different preferences. Also acts on other dinucleotides, including NADP(H), FAD(H<sub>2</sub>), and the thionicotinamide analogues of NAD<sup>+</sup> and NADP<sup>+</sup>.

**References:** [1512, 1291, 2821, 1551, 51, 2042, 815, 3236, 1304]

[EC 3.6.1.22 created 1972]

#### EC 3.6.1.23

**Accepted name:** dUTP diphosphatase

**Reaction:** dUTP + H<sub>2</sub>O = dUMP + diphosphate

**Other name(s):** DUT (gene name); deoxyuridine-triphosphatase; dUTPase; dUTP pyrophosphatase; desoxyuridine 5'-triphosphate nucleotidohydrolase; desoxyuridine 5'-triphosphatase

**Systematic name:** dUTP nucleotidohydrolase

**Comments:** The enzyme catalyses the Mg<sup>2+</sup>-dependent hydrolysis of dUTP to dUMP, providing the substrate for EC 2.1.1.45, thymidylate synthase, leading to production of thymidine nucleotides. By reducing the effective ratio of dUTP to TTP, the enzyme also reduces the possibility of dUTP incorporation into DNA.

**References:** [974, 205, 980, 2645, 924, 384, 1588, 123, 3052]

[EC 3.6.1.23 created 1972]

#### EC 3.6.1.24

**Accepted name:** nucleoside phosphoacylhydrolase

**Reaction:** Hydrolyses mixed phospho-anhydride bonds

**Systematic name:** nucleoside-5'-phosphoacylate acylhydrolase

**Comments:** Attacks ribonucleoside 5'-nitrophenylphosphates, but is inactive against phosphodiesteres.

**References:** [2736]

[EC 3.6.1.24 created 1972]

#### EC 3.6.1.25

**Accepted name:** triphosphatase

**Reaction:** triphosphate + H<sub>2</sub>O = diphosphate + phosphate

**Other name(s):** inorganic triphosphatase

**Systematic name:** triphosphate phosphohydrolase

**References:** [1545, 3010]

[EC 3.6.1.25 created 1976]

#### EC 3.6.1.26

**Accepted name:** CDP-diacylglycerol diphosphatase  
**Reaction:** CDP-diacylglycerol + H<sub>2</sub>O = CMP + phosphatidate  
**Other name(s):** cytidine diphosphodiacylglycerol pyrophosphatase; CDP diacylglycerol hydrolase  
**Systematic name:** CDP-diacylglycerol phosphatidylhydrolase  
**References:** [2354]

[EC 3.6.1.26 created 1976]

#### EC 3.6.1.27

**Accepted name:** undecaprenyl-diphosphate phosphatase  
**Reaction:** *ditrans*,*octacis*-undecaprenyl diphosphate + H<sub>2</sub>O = *ditrans*,*octacis*-undecaprenyl phosphate + phosphate  
**Other name(s):** C<sub>55</sub>-isoprenyl diphosphatase; C<sub>55</sub>-isoprenyl pyrophosphatase; isoprenyl pyrophosphatase (ambiguous); undecaprenyl pyrophosphate phosphatase; undecaprenyl pyrophosphate pyrophosphatase; UPP phosphatase; Und-PP pyrophosphatase; UppP (ambiguous); BacA; undecaprenyl-diphosphate phosphohydrolase; undecaprenyl-diphosphatase  
**Systematic name:** *ditrans*,*octacis*-undecaprenyl-diphosphate phosphohydrolase  
**Comments:** Isolated from the bacteria *Micrococcus lysodeikticus* [950], *Escherichia coli* [2,3,5,6] and *Bacillus subtilis* [198]. The product of the reaction, *ditrans*,*octacis*-undecaprenyl phosphate, is essential for cell wall polysaccharide biosynthesis in these strains.  
**References:** [950, 900, 901, 198, 2892, 2952]

[EC 3.6.1.27 created 1978, modified 2002, modified 2012]

#### EC 3.6.1.28

**Accepted name:** thiamine-triphosphatase  
**Reaction:** thiamine triphosphate + H<sub>2</sub>O = thiamine diphosphate + phosphate  
**Systematic name:** thiamine-triphosphate phosphohydrolase  
**References:** [1071]

[EC 3.6.1.28 created 1978]

#### EC 3.6.1.29

**Accepted name:** bis(5'-adenosyl)-triphosphatase  
**Reaction:** P<sup>1</sup>,P<sup>3</sup>-bis(5'-adenosyl) triphosphate + H<sub>2</sub>O = ADP + AMP  
**Other name(s):** dinucleosidetriphosphatase; diadenosine 5,5-P<sup>1</sup>,P<sup>3</sup>-triphosphatase; 1-P,3-P-bis(5'-adenosyl)-triphosphate adenylohydrolase  
**Systematic name:** P<sup>1</sup>,P<sup>3</sup>-bis(5'-adenosyl)-triphosphate adenylohydrolase  
**References:** [1302, 2659]

[EC 3.6.1.29 created 1978]

[3.6.1.30 Deleted entry. *m*<sup>7</sup>G(5')pppN diphosphatase. Now covered by EC 3.6.1.59 [*m*<sup>7</sup>GpppX diphosphatase] and EC 3.6.1.62 [*m*<sup>7</sup>GpppN-mRNA hydrolase].]

[EC 3.6.1.30 created 1978, deleted 2012]

#### EC 3.6.1.31

**Accepted name:** phosphoribosyl-ATP diphosphatase  
**Reaction:** 1-(5-phospho-β-D-ribose)-ATP + H<sub>2</sub>O = 1-(5-phospho-β-D-ribose)-AMP + diphosphate  
**Other name(s):** phosphoribosyl-ATP pyrophosphatase; phosphoribosyladenosine triphosphate pyrophosphatase; 1-(5-phosphoribosyl)-ATP diphosphohydrolase  
**Systematic name:** 1-(5-phospho-β-D-ribose)-ATP diphosphohydrolase

**Comments:** The *Neurospora crassa* enzyme also catalyses the reactions of EC 1.1.1.23 (histidinol dehydrogenase) and EC 3.5.4.19 (phosphoribosyl-AMP cyclohydrolase).

**References:** [2697]

[EC 3.6.1.31 created 1981]

[3.6.1.32 *Transferred entry. myosin ATPase. Now EC 3.6.4.1, myosin ATPase*]

[EC 3.6.1.32 created 1984, deleted 2000]

[3.6.1.33 *Transferred entry. dynein ATPase. Now EC 3.6.4.2, dynein ATPase*]

[EC 3.6.1.33 created 1984, deleted 2000]

[3.6.1.34 *Transferred entry. H<sup>+</sup>-transporting ATP synthase. Now EC 3.6.3.14, H<sup>+</sup>-transporting two-sector ATPase*]

[EC 3.6.1.34 created 1984, deleted 2000]

[3.6.1.35 *Transferred entry. H<sup>+</sup>-transporting ATPase. Now EC 3.6.3.6, H<sup>+</sup>-exporting ATPase*]

[EC 3.6.1.35 created 1984, deleted 2000]

[3.6.1.36 *Transferred entry. H<sup>+</sup>/K<sup>+</sup> exchanging ATPase. Now EC 3.6.3.10, H<sup>+</sup>/K<sup>+</sup>-exchanging ATPase*]

[EC 3.6.1.36 created 1984, deleted 2000]

[3.6.1.37 *Transferred entry. Na<sup>+</sup>/K<sup>+</sup> exchanging ATPase. Now EC 3.6.3.9, Na<sup>+</sup>/K<sup>+</sup>-exchanging ATPase*]

[EC 3.6.1.37 created 1984, deleted 2000]

[3.6.1.38 *Transferred entry. Ca<sup>2+</sup>-transporting ATPase. Now EC 3.6.3.8, Ca<sup>2+</sup>-transporting ATPase*]

[EC 3.6.1.38 created 1984, deleted 2000]

#### EC 3.6.1.39

**Accepted name:** thymidine-triphosphatase  
**Reaction:** dTTP + H<sub>2</sub>O = dTDP + phosphate  
**Other name(s):** thymidine triphosphate nucleotidohydrolase; dTTPase; deoxythymidine-5'-triphosphatase  
**Systematic name:** dTTP nucleotidohydrolase  
**Comments:** Also acts, more slowly, on dUTP and UTP.  
**References:** [528]

[EC 3.6.1.39 created 1984]

#### EC 3.6.1.40

**Accepted name:** guanosine-5'-triphosphate,3'-diphosphate phosphatase  
**Reaction:** guanosine 5'-triphosphate 3'-diphosphate + H<sub>2</sub>O = guanosine 3',5'-bis(diphosphate) + phosphate  
**Other name(s):** pppGpp 5'-phosphohydrolase; guanosine 5'-triphosphate-3'-diphosphate 5'-phosphohydrolase; guanosine pentaphosphatase; guanosine pentaphosphate phosphatase; guanosine 5'-triphosphate 3'-diphosphate 5'-phosphatase; guanosine pentaphosphate phosphohydrolase  
**Systematic name:** guanosine-5'-triphosphate-3'-diphosphate 5'-phosphohydrolase  
**Comments:** Also hydrolyses other guanosine 5'-triphosphate derivatives with at least one unsubstituted phosphate group on the 3'-position, but not GTP, ATP or adenosine 5'-triphosphate 3'-diphosphate.  
**References:** [1047]

[EC 3.6.1.40 created 1986, modified 2010]

#### EC 3.6.1.41

**Accepted name:** bis(5'-nucleosyl)-tetraphosphatase (symmetrical)

**Reaction:**  $P^1, P^4$ -bis(5'-adenosyl) tetraphosphate + H<sub>2</sub>O = 2 ADP  
**Other name(s):** diadenosinetetraphosphatase (symmetrical); dinucleosidetetraphosphatase (symmetrical); symmetrical diadenosine tetraphosphate hydrolase; adenosine tetraphosphate phosphodiesterase; Ap4A hydrolase; bis(5'-adenosyl) tetraphosphatase; diadenosine tetraphosphate hydrolase; diadenosine polyphosphate hydrolase; diadenosine 5',5'''-P<sup>1</sup>,P<sup>4</sup>-tetraphosphatase; diadenosinetetraphosphatase (symmetrical); 1-*P*,4-*P*-bis(5'-nucleosyl)-tetraphosphate nucleosidebisphosphohydrolase  
**Systematic name:**  $P^1, P^4$ -bis(5'-nucleosyl)-tetraphosphate nucleosidebisphosphohydrolase  
**Comments:** Also acts on bis(5'-guanosyl) tetraphosphate and bis(5'-adenosyl) pentaphosphate and, more slowly, on some other polyphosphates, forming a nucleoside bisphosphate as one product in all cases [*cf.* EC 3.6.1.17 bis(5'-nucleosyl)-tetraphosphatase (asymmetrical)].  
**References:** [148, 1002]

[EC 3.6.1.41 created 1986]

#### EC 3.6.1.42

**Accepted name:** guanosine-diphosphatase  
**Reaction:** GDP + H<sub>2</sub>O = GMP + phosphate  
**Other name(s):** GDPase  
**Systematic name:** GDP phosphohydrolase  
**Comments:** Also acts on UDP but not on other nucleoside diphosphates and triphosphates.  
**References:** [2388]

[EC 3.6.1.42 created 1989]

#### EC 3.6.1.43

**Accepted name:** dolichyldiphosphatase  
**Reaction:** dolichyl diphosphate + H<sub>2</sub>O = dolichyl phosphate + phosphate  
**Other name(s):** dolichol diphosphatase; dolichyl pyrophosphatase; dolichyl pyrophosphate phosphatase; dolichyl diphosphate phosphohydrolase; Dol-*P-P* phosphohydrolase  
**Systematic name:** dolichyl-diphosphate phosphohydrolase  
**References:** [2058]

[EC 3.6.1.43 created 1989]

#### EC 3.6.1.44

**Accepted name:** oligosaccharide-diphosphodolichol diphosphatase  
**Reaction:** oligosaccharide-diphosphodolichol + H<sub>2</sub>O = oligosaccharide phosphate + dolichyl phosphate  
**Other name(s):** oligosaccharide-diphosphodolichol pyrophosphatase  
**Systematic name:** oligosaccharide-diphosphodolichol phosphodolichohydrolase  
**References:** [186]

[EC 3.6.1.44 created 1992]

#### EC 3.6.1.45

**Accepted name:** UDP-sugar diphosphatase  
**Reaction:** UDP-sugar + H<sub>2</sub>O = UMP + α-D-aldose 1-phosphate  
**Other name(s):** nucleosidediphosphate-sugar pyrophosphatase; nucleosidediphosphate-sugar diphosphatase; UDP-sugar hydrolase; UDP-sugar pyrophosphatase  
**Systematic name:** UDP-sugar sugarphosphohydrolase  
**Comments:** A divalent cation is required for activity. UDP-sugar is the best substrate, although other nucleoside-sugar diphosphates are used as substrates with similar *K<sub>m</sub>* values but much lower maximum velocities. Thus, this enzyme has a specificity distinct from that of ADP-sugar diphosphatase (EC 3.6.1.21). Some but not all enzymes of this class also appear to have 5'-nucleotidase (see EC 3.1.3.5) activity.  
**References:** [881, 927]

[EC 3.6.1.45 created 1999]

[3.6.1.46 *Transferred entry. heterotrimeric G-protein GTPase. Now EC 3.6.5.1, heterotrimeric G-protein GTPase*]

[EC 3.6.1.46 created 2000, deleted 2003]

[3.6.1.47 *Transferred entry. small monomeric GTPase. Now EC 3.6.5.2, small monomeric GTPase*]

[EC 3.6.1.47 created 2000, deleted 2003]

[3.6.1.48 *Transferred entry. protein-synthesizing GTPase. Now EC 3.6.5.3, protein-synthesizing GTPase*]

[EC 3.6.1.48 created 2000, deleted 200]

[3.6.1.49 *Transferred entry. signal-recognition-particle GTPase. Now EC 3.6.5.4, signal-recognition-particle GTPase*]

[EC 3.6.1.49 created 2000, deleted 2003]

[3.6.1.50 *Transferred entry. dynamin GTPase. Now EC 3.6.5.5, dynamin GTPase*]

[EC 3.6.1.50 created 2000, deleted 2003]

[3.6.1.51 *Transferred entry. tubulin GTPase. Now EC 3.6.5.6, tubulin GTPase*]

[EC 3.6.1.51 created 2000, deleted 2003]

#### EC 3.6.1.52

**Accepted name:** diphosphoinositol-polyphosphate diphosphatase  
**Reaction:** diphospho-*myo*-inositol polyphosphate + H<sub>2</sub>O = *myo*-inositol polyphosphate + phosphate  
**Other name(s):** diphosphoinositol-polyphosphate phosphohydrolase; DIPP  
**Systematic name:** diphospho-*myo*-inositol-polyphosphate diphosphohydrolase  
**Comments:** This enzyme hydrolyses the diphosphate bond, leaving a phospho group where a diphospho group had been. It can also act on bis(adenosine) diphosphate.  
**References:** [2491, 352]

[EC 3.6.1.52 created 2002]

#### EC 3.6.1.53

**Accepted name:** Mn<sup>2+</sup>-dependent ADP-ribose/CDP-alcohol diphosphatase  
**Reaction:** (1) CDP-choline + H<sub>2</sub>O = CMP + phosphocholine  
(2) ADP-D-ribose + H<sub>2</sub>O = AMP + D-ribose 5-phosphate  
**Other name(s):** Mn<sup>2+</sup>-dependent ADP-ribose/CDP-alcohol pyrophosphatase; ADPRibase-Mn  
**Systematic name:** CDP-choline phosphohydrolase  
**Comments:** Requires Mn<sup>2+</sup>. Unlike EC 3.6.1.13, ADP-ribose diphosphatase, it cannot utilize Mg<sup>2+</sup>. ADP-D-ribose, CDP-choline, CDP-ethanolamine and ADP are substrates for this enzyme but ADP-D-glucose, UDP-D-glucose, CDP-D-glucose, CDP, CMP and AMP are not hydrolysed [362]. The mammalian enzyme hydrolyses cyclic ADP-ribose to 1-(5-phospho-β-D-ribose)-AMP with 100-fold lower efficiency than ADP-D-ribose [363]. In rat, the enzyme is found predominantly in thymus and spleen.  
**References:** [364, 362, 363, 2448]

[EC 3.6.1.53 created 2008]

#### EC 3.6.1.54

**Accepted name:** UDP-2,3-diacylglucosamine diphosphatase  
**Reaction:** UDP-2-*N*,3-*O*-bis[(3*R*)-3-hydroxytetradecanoyl]-α-D-glucosamine + H<sub>2</sub>O = 2-*N*,3-*O*-bis[(3*R*)-3-hydroxytetradecanoyl]-α-D-glucosaminyl 1-phosphate + UMP  
**Other name(s):** UDP-2,3-diacylglucosamine hydrolase; UDP-2,3-diacylglucosamine pyrophosphatase; *ybbF* (gene name); *lpxH* (gene name); UDP-2,3-bis[(3*R*)-3-hydroxymyristoyl]-α-D-glucosamine 2,3-bis[(3*R*)-3-hydroxymyristoyl]-β-D-glucosaminyl 1-phosphate phosphohydrolase (incorrect)

**Systematic name:** UDP-2-*N*,3-*O*-bis[(3*R*)-3-hydroxytetradecanoyl]- $\alpha$ -D-glucosamine 2-*N*,3-*O*-bis[(3*R*)-3-hydroxytetradecanoyl]- $\alpha$ -D-glucosaminyl 1-phosphate phosphohydrolase  
**Comments:** The enzyme catalyses a step in the biosynthesis of lipid A.  
**References:** [111, 110]

[EC 3.6.1.54 created 2010]

#### EC 3.6.1.55

**Accepted name:** 8-oxo-dGTP diphosphatase  
**Reaction:** 8-oxo-dGTP + H<sub>2</sub>O = 8-oxo-dGMP + diphosphate  
**Other name(s):** MutT; 7,8-dihydro-8-oxoguanine triphosphatase; 8-oxo-dGTPase; 7,8-dihydro-8-oxo-dGTP pyrophosphohydrolase  
**Systematic name:** 8-oxo-dGTP diphosphohydrolase  
**Comments:** This enzyme hydrolyses the phosphoanhydride bond between the  $\alpha$  and  $\beta$  phosphate of 8-oxoguanine-containing nucleoside di- and triphosphates thereby preventing misincorporation of the oxidized purine nucleoside triphosphates into DNA. It does not hydrolyse 2-hydroxy-dATP (*cf.* EC 3.6.1.56, 2-hydroxy-dATP diphosphatase) [3289]. Requires Mg<sup>2+</sup>.  
**References:** [1267, 3305, 2047, 3289]

[EC 3.6.1.55 created 2011]

#### EC 3.6.1.56

**Accepted name:** 2-hydroxy-dATP diphosphatase  
**Reaction:** 2-hydroxy-dATP + H<sub>2</sub>O = 2-hydroxy-dAMP + diphosphate  
**Other name(s):** NUDT1; MTH1; MTH<sub>2</sub>; oxidized purine nucleoside triphosphatase; (2'-deoxy) ribonucleoside 5'-triphosphate pyrophosphohydrolase  
**Systematic name:** 2-hydroxy-dATP diphosphohydrolase  
**Comments:** The enzyme hydrolyses oxidized purine nucleoside triphosphates such as 2-hydroxy-dATP, thereby preventing their misincorporation into DNA. It can also recognize 8-oxo-dGTP and 8-oxo-dATP, but with lower efficiency (*cf.* EC 3.6.1.55, 8-oxo-dGTP diphosphatase) [838].  
**References:** [2506, 1360, 838, 2503, 839]

[EC 3.6.1.56 created 2011]

#### EC 3.6.1.57

**Accepted name:** UDP-2,4-diacetamido-2,4,6-trideoxy- $\beta$ -L-altropyranose hydrolase  
**Reaction:** UDP-2,4-diacetamido-2,4,6-trideoxy- $\beta$ -L-altropyranose + H<sub>2</sub>O = 2,4-diacetamido-2,4,6-trideoxy- $\beta$ -L-altropyranose + UDP  
**Other name(s):** PseG; UDP-6-deoxy-AltdiNAc hydrolase; Cj1312; UDP-2,4-bis(acetamido)-2,4,6-trideoxy- $\beta$ -L-altropyranose hydrolase  
**Systematic name:** UDP-2,4-diacetamido-2,4,6-trideoxy- $\beta$ -L-altropyranose hydrolase  
**Comments:** The enzyme is involved in biosynthesis of pseudaminic acid.  
**References:** [1719, 2581]

[EC 3.6.1.57 created 2011]

#### EC 3.6.1.58

**Accepted name:** 8-oxo-dGDP phosphatase  
**Reaction:** 8-oxo-dGDP + H<sub>2</sub>O = 8-oxo-dGMP + phosphate  
**Other name(s):** NUDT5; MTH3 (gene name); NUDT18  
**Systematic name:** 8-oxo-dGDP phosphohydrolase



**Comments:** The enzyme catalyses the hydrolysis of both 8-oxo-dGDP and 8-oxo-GDP thereby preventing translational errors caused by oxidative damage. The preferred *in vivo* substrate is not known. The enzyme does not degrade 8-oxo-dGTP and 8-oxo-GTP to the monophosphates (*cf.* EC 3.6.1.55, 8-oxo-dGTP diphosphatase) [1259, 1260]. Ribonucleotide diphosphates and deoxyribonucleotide diphosphates are hydrolysed with broad specificity. The bifunctional enzyme NUDT5 also hydrolyses ADP-ribose to AMP and D-ribose 5-phosphate (*cf.* EC 3.6.1.13, ADP-ribose diphosphatase) [1268]. The human enzyme NUDT18 also hydrolyses 8-oxo-dADP and 2-hydroxy-dADP, the latter at a slower rate [2835].

**References:** [1259, 1260, 1373, 1268, 3323, 2835]

[EC 3.6.1.58 created 2012]

#### EC 3.6.1.59

**Accepted name:** 5'-(*N*<sup>7</sup>-methyl 5'-triphosphoguanosine)-[mRNA] diphosphatase

**Reaction:** a 5'-(*N*<sup>7</sup>-methyl 5'-triphosphoguanosine)-[mRNA] + H<sub>2</sub>O = *N*<sup>7</sup>-methylguanosine 5'-phosphate + a 5'-diphospho-[mRNA]

**Other name(s):** DcpS; m<sup>7</sup>GpppX pyrophosphatase; m<sup>7</sup>GpppN m<sup>7</sup>GMP phosphohydrolase; m<sup>7</sup>GpppX diphosphatase; m<sup>7</sup>G5'ppp5'N m<sup>7</sup>GMP phosphohydrolase

**Systematic name:** 5'-(*N*<sup>7</sup>-methyl 5'-triphosphoguanosine)-[mRNA] *N*<sup>7</sup>-methylguanosine 5'-phosphate phosphohydrolase

**Comments:** The enzyme removes (decaps) the *N*<sup>7</sup>-methylguanosine 5-phosphate cap from an mRNA degraded to a maximal length of 10 nucleotides [1720, 464]. Decapping is an important process in the control of eukaryotic mRNA degradation. The enzyme functions to clear the cell of cap structure following decay of the RNA body [1725]. The nematode enzyme can also decap triply methylated substrates, 5'-(*N*<sup>2</sup>,*N*<sup>2</sup>,*N*<sup>7</sup>-trimethyl 5'-triphosphoguanosine)-[mRNA] [3045].

**References:** [1795, 1725, 1720, 3045, 414, 464, 3227]

[EC 3.6.1.59 created 2012, modified 2013]

#### EC 3.6.1.60

**Accepted name:** diadenosine hexaphosphate hydrolase (AMP-forming)

**Reaction:** (1) *P*<sup>1</sup>,*P*<sup>6</sup>-bis(5'-adenosyl)hexaphosphate + H<sub>2</sub>O = adenosine 5'-pentaphosphate + AMP  
(2) *P*<sup>1</sup>,*P*<sup>5</sup>-bis(5'-adenosyl)pentaphosphate + H<sub>2</sub>O = adenosine 5'-tetraphosphate + AMP

**Other name(s):** hAps1; NUDT11 (gene name); hAps2; NUDT10 (gene name)

**Systematic name:** *P*<sup>1</sup>,*P*<sup>6</sup>-bis(5'-adenosyl)hexaphosphate nucleotidohydrolase (AMP-forming)

**Comments:** A divalent cation is essential for activity. Mn<sup>2+</sup> (2–6 mM) is most effective. The enzyme controls intracellular levels of *P*<sup>1</sup>,*P*<sup>5</sup>-bis(5'-adenosyl)pentaphosphate and *P*<sup>1</sup>,*P*<sup>6</sup>-bis(5'-adenosyl)hexaphosphate. Weak activity with *P*<sup>1</sup>,*P*<sup>4</sup>-bis(5'-adenosyl)tetraphosphate. Marked preference for adenine over guanine nucleotides.

**References:** [1651, 2492]

[EC 3.6.1.60 created 2012]

#### EC 3.6.1.61

**Accepted name:** diadenosine hexaphosphate hydrolase (ATP-forming)

**Reaction:** (1) *P*<sup>1</sup>,*P*<sup>6</sup>-bis(5'-adenosyl)hexaphosphate + H<sub>2</sub>O = 2 ATP  
(2) *P*<sup>1</sup>,*P*<sup>5</sup>-bis(5'-adenosyl)pentaphosphate + H<sub>2</sub>O = ATP + ADP  
(3) *P*<sup>1</sup>,*P*<sup>4</sup>-bis(5'-adenosyl)tetraphosphate + H<sub>2</sub>O = ATP + AMP

**Other name(s):** Ndx1

**Systematic name:** *P*<sup>1</sup>,*P*<sup>6</sup>-bis(5'-adenosyl)hexaphosphate nucleotidohydrolase (ATP-forming)

**Comments:** The enzyme requires the presence of the divalent cations (Mn<sup>2+</sup>, Mg<sup>2+</sup>, Zn<sup>2+</sup>, and Co<sup>2+</sup>). It hydrolyses *P*<sup>1</sup>,*P*<sup>4</sup>-bis(5'-guanosyl) tetraphosphate very slowly [*cf.* EC 3.6.1.17, bis(5-nucleosyl)-tetraphosphatase (asymmetrical)].

**References:** [1281]

[EC 3.6.1.61 created 2012]

#### EC 3.6.1.62

- Accepted name:** 5'-(*N*<sup>7</sup>-methylguanosine 5'-triphospho)-[mRNA] hydrolase  
**Reaction:** a 5'-(*N*<sup>7</sup>-methylguanosine 5'-triphospho)-[mRNA] + H<sub>2</sub>O = *N*<sup>7</sup>-methylguanosine 5'-diphosphate + a 5'-phospho-[mRNA]  
**Other name(s):** Dcp2; NUDT16; D10 protein; D9 protein; D10 decapping enzyme; decapping enzyme; m<sup>7</sup>GpppN-mRNA hydrolase; m<sup>7</sup>GpppN-mRNA m<sup>7</sup>GDP phosphohydrolase  
**Systematic name:** 5'-(*N*<sup>7</sup>-methylguanosine 5'-triphospho)-[mRNA] *N*<sup>7</sup>-methylguanosine-5'-diphosphate phosphohydrolase  
**Comments:** Decapping of mRNA is a critical step in eukaryotic mRNA turnover. The enzyme is unable to cleave a free cap structure (m<sup>7</sup>GpppG) [3044]. The enzyme from *Vaccinia* virus is synergistically activated in the presence of Mg<sup>2+</sup> and Mn<sup>2+</sup> [2732].  
**References:** [3233, 1744, 3044, 2239, 2732, 2238, 2720]

[EC 3.6.1.62 created 2012, modified 2013]

#### EC 3.6.1.63

- Accepted name:** α-D-ribose 1-methylphosphonate 5-triphosphate diphosphatase  
**Reaction:** α-D-ribose 1-methylphosphonate 5-triphosphate + H<sub>2</sub>O = α-D-ribose 1-methylphosphonate 5-phosphate + diphosphate  
**Other name(s):** *phnM* (gene name)  
**Systematic name:** α-D-ribose-1-methylphosphonate-5-triphosphate diphosphohydrolase  
**Comments:** Isolated from the bacterium *Escherichia coli*.  
**References:** [1370]

[EC 3.6.1.63 created 2012]

#### EC 3.6.1.64

- Accepted name:** inosine diphosphate phosphatase  
**Reaction:** (1) IDP + H<sub>2</sub>O = IMP + phosphate  
(2) dIDP + H<sub>2</sub>O = dIMP + phosphate  
**Other name(s):** (deoxy)inosine diphosphatase; NUDT16  
**Systematic name:** inosine diphosphate phosphatase  
**Comments:** The human enzyme also hydrolyses GDP and dGDP, and to a lesser extent ITP, dITP and XTP.  
**References:** [1284]

[EC 3.6.1.64 created 2013]

#### EC 3.6.1.65

- Accepted name:** (d)CTP diphosphatase  
**Reaction:** (1) CTP + H<sub>2</sub>O = CMP + diphosphate  
(2) dCTP + H<sub>2</sub>O = dCMP + diphosphate  
**Other name(s):** (d)CTP pyrophosphohydrolase; (d)CTP diphosphohydrolase; *nudG* (gene name)  
**Systematic name:** (deoxy)cytidine 5'-triphosphate diphosphohydrolase  
**Comments:** The enzyme, characterized from the bacterium *Escherichia coli*, is specific for the pyrimidine nucleotides CTP and dCTP. It also acts on 5-methyl-dCTP, 5-hydroxy-dCTP and 8-hydroxy-dGTP.  
**References:** [2155, 840, 1374, 1224]

[EC 3.6.1.65 created 2013]

#### EC 3.6.1.66

**Accepted name:** XTP/dITP diphosphatase  
**Reaction:** (1)  $XTP + H_2O = XMP + \text{diphosphate}$   
(2)  $dITP + H_2O = dIMP + \text{diphosphate}$   
(2)  $ITP + H_2O = IMP + \text{diphosphate}$   
**Other name(s):** hypoxanthine/xanthine dNTP pyrophosphatase; *rdgB* (gene name)  
**Systematic name:** XTP/dITP diphosphohydrolase (diphosphate-forming)  
**Comments:** The enzymes from the bacterium *Escherichia coli* and the archaea *Methanococcus jannaschii* and *Archaeoglobus fulgidus* are highly specific for XTP, dITP and ITP. The activity is dependent on divalent cations,  $Mg^{2+}$  is preferred.  
**References:** [1213, 448, 449, 2537]

[EC 3.6.1.66 created 2013]

#### EC 3.6.1.67

**Accepted name:** dihydroneopterin triphosphate diphosphatase  
**Reaction:**  $7,8\text{-dihydroneopterin } 3'\text{-triphosphate} + H_2O = 7,8\text{-dihydroneopterin } 3'\text{-phosphate} + \text{diphosphate}$   
**Other name(s):** *folQ* (gene name); *nudB* (gene name); NUDT1 (gene name); dihydroneopterin triphosphate pyrophosphohydrolase  
**Systematic name:** 7,8-dihydroneopterin 3'-triphosphate diphosphohydrolase  
**Comments:** The enzyme participates in a folate biosynthesis pathway, which is found in bacteria, fungi, and plants. Requires  $Mg^{2+}$ .  
**References:** [2816, 2156, 1472, 867]

[EC 3.6.1.67 created 2014]

#### EC 3.6.1.68

**Accepted name:** geranyl diphosphate phosphohydrolase  
**Reaction:**  $\text{geranyl diphosphate} + H_2O = \text{geranyl phosphate} + \text{phosphate}$   
**Other name(s):** NUDX1 (gene name)  
**Systematic name:** geranyl-diphosphate phosphohydrolase  
**Comments:** The enzyme, characterized from roses, is involved in a cytosolic pathway for the biosynthesis of free monoterpene alcohols that contribute to fragrance. *In vitro* the enzyme also acts on (2*E*,6*E*)-farnesyl diphosphate.  
**References:** [1771]

[EC 3.6.1.68 created 2015 as EC 3.1.3.98, transferred 2016 to EC 3.6.1.68]

### EC 3.6.2 In sulfonyl-containing anhydrides

#### EC 3.6.2.1

**Accepted name:** adenylylsulfatase  
**Reaction:**  $\text{adenylyl sulfate} + H_2O = \text{AMP} + \text{sulfate}$   
**Other name(s):** adenosine 5-phosphosulfate sulfohydrolase; adenylylsulfate sulfohydrolase  
**Systematic name:** adenylyl-sulfate sulfohydrolase  
**References:** [122]

[EC 3.6.2.1 created 1972]

#### EC 3.6.2.2

**Accepted name:** phosphoadenylylsulfatase  
**Reaction:**  $3'\text{-phosphoadenylyl sulfate} + H_2O = \text{adenosine } 3',5'\text{-bisphosphate} + \text{sulfate}$   
**Other name(s):** 3-phosphoadenylyl sulfatase; 3-phosphoadenosine 5-phosphosulfate sulfatase; PAPS sulfatase; 3'-phosphoadenylylsulfate sulfohydrolase

**Systematic name:** 3'-phosphoadenylyl-sulfate sulfohydrolase  
**Comments:** Requires Mn<sup>2+</sup>.  
**References:** [128]

[EC 3.6.2.2 created 1972]

### EC 3.6.3 Acting on acid anhydrides to catalyse transmembrane movement of substances

Several types of ATP phosphohydrolase are listed here. Entries EC 3.6.3.1 to EC 3.6.3.12 and EC 3.6.3.53 are enzymes undergoing covalent phosphorylation of an aspartate residue during the transport cycle; entries EC 3.6.3.14 and EC 3.6.3.15 refer to enzymes of complicated membrane and non-membrane location that can also serve in ATP synthesis; entry EC 3.6.3.16 is a multisubunit enzyme that is involved in arsenite transport only; entries EC 3.6.3.17 to EC 3.6.3.50 are two-domain enzymes of the ABC family; entries EC 3.6.3.51 and EC 3.6.3.52 are parts of a complex protein-transporting machinery in mitochondria and chloroplasts.

[3.6.3.1 *Transferred entry. phospholipid-translocating ATPase. Now EC 7.6.2.1, P-type phospholipid transporter*]

[EC 3.6.3.1 created 2000 (EC 3.6.3.13 created 2000, incorporated 2001), deleted 2018]

[3.6.3.2 *Transferred entry. Mg<sup>2+</sup>-importing ATPase. Now EC 7.2.2.14, P-type Mg<sup>2+</sup> transporter*]

[EC 3.6.3.2 created 2000, modified 2001, deleted 2018]

#### EC 3.6.3.3

**Accepted name:** Cd<sup>2+</sup>-exporting ATPase  
**Reaction:** ATP + H<sub>2</sub>O + Cd<sup>2+</sup><sub>in</sub> = ADP + phosphate + Cd<sup>2+</sup><sub>out</sub>  
**Systematic name:** ATP phosphohydrolase (Cd<sup>2+</sup>-exporting)  
**Comments:** A P-type ATPase that undergoes covalent phosphorylation during the transport cycle. This enzyme occurs in protozoa, fungi and plants.  
**References:** [2660, 2968]

[EC 3.6.3.3 created 2000]

[3.6.3.4 *Transferred entry. Cu<sup>2+</sup>-exporting ATPase. Now EC 7.2.2.9, Cu<sup>2+</sup>-exporting ATPase*]

[EC 3.6.3.4 created 2000, modified 2013, deleted 2018]

[3.6.3.5 *Transferred entry. Zn<sup>2+</sup>-exporting ATPase. Now EC 7.2.2.12, Zn<sup>2+</sup>-exporting ATPase*]

[EC 3.6.3.5 created 2000, modified 2001, modified 2006, deleted 2018]

[3.6.3.6 *Transferred entry. H<sup>+</sup>-exporting ATPase. Now EC 7.1.2.1, P-type H<sup>+</sup>-exporting transporter*]

[EC 3.6.3.6 created 1984 as EC 3.6.1.35, transferred 2000 to EC 3.6.3.6, deleted 2018]

[3.6.3.7 *Transferred entry. Na<sup>+</sup>-exporting ATPase. Now EC 7.2.2.3, P-type Na<sup>+</sup> transporter*]

[EC 3.6.3.7 created 2000, modified 2001, transferred 2018 to EC 7.2.2.3, deleted 2018]

[3.6.3.8 *Transferred entry. Ca<sup>2+</sup>-transporting ATPase. Now EC 7.2.2.10, Ca<sup>2+</sup>-transporting ATPase*]

[EC 3.6.3.8 created 1984 as EC 3.6.1.38, transferred 2000 to EC 3.6.3.8, modified 2001, modified 2011, deleted 2018]

[3.6.3.9 *Transferred entry. Na<sup>+</sup>/K<sup>+</sup>-exchanging ATPase. Now EC 7.2.2.13, Na<sup>+</sup>/K<sup>+</sup>-exchanging ATPase*]

[EC 3.6.3.9 created 1984 as EC 3.6.1.37, transferred 2000 to EC 3.6.3.9, modified 2001, deleted 2018]

[3.6.3.10 *Transferred entry. H<sup>+</sup>/K<sup>+</sup>-exchanging ATPase. Now EC 7.2.2.19, H<sup>+</sup>/K<sup>+</sup>-exchanging ATPase*]

[EC 3.6.3.10 created 1984 as EC 3.6.1.36, transferred 2000 to EC 3.6.3.10, deleted 2018]

### EC 3.6.3.11

**Accepted name:** Cl<sup>-</sup>-transporting ATPase  
**Reaction:** ATP + H<sub>2</sub>O + Cl<sup>-</sup><sub>out</sub> = ADP + phosphate + Cl<sup>-</sup><sub>in</sub>  
**Other name(s):** Cl<sup>-</sup>-translocating ATPase; Cl<sup>-</sup>-motive ATPase  
**Systematic name:** ATP phosphohydrolase (Cl<sup>-</sup>-importing)  
**Comments:** A P-type ATPase that undergoes covalent phosphorylation during the transport cycle. An animal and plant enzyme involved in the import of chloride anions.  
**References:** [2158, 893, 1245]

[EC 3.6.3.11 created 2000]

[3.6.3.12 *Transferred entry. K<sup>+</sup>-transporting ATPase. Now EC 7.2.2.6, K<sup>+</sup>-transporting ATPase]*

[EC 3.6.3.12 created 2000, deleted 2018]

[3.6.3.13 *Deleted entry. aminophospholipid-transporting ATPase. Identical to EC 3.6.3.1, phospholipid-translocating ATPase]*

[EC 3.6.3.13 created 2000, deleted 2001]

[3.6.3.14 *Transferred entry. H<sup>+</sup>-transporting two-sector ATPase. Now EC 7.1.2.2, H<sup>+</sup>-transporting two-sector ATPase]*

[EC 3.6.3.14 created 1984 as EC 3.6.1.34, transferred 2000 to EC 3.6.3.14, deleted 2018]

[3.6.3.15 *Transferred entry. Na<sup>+</sup>-transporting two-sector ATPase. Now EC 7.2.2.1, Na<sup>+</sup>-transporting two-sector ATPase]*

[EC 3.6.3.15 created 2000, transferred 2018 to EC 7.2.2.1, deleted 2018]

### EC 3.6.3.16

**Accepted name:** arsenite-transporting ATPase  
**Reaction:** ATP + H<sub>2</sub>O + arsenite<sub>in</sub> = ADP + phosphate + arsenite<sub>out</sub>  
**Systematic name:** ATP phosphohydrolase (arsenite-exporting)  
**Comments:** A multisubunit non-phosphorylated ATPase that is involved in the transport of ions. A bacterial enzyme that usually contains two subunits where one (with 12 membrane-spanning segments) forms the 'channel' part and the other (occurring in pairs peripherally to the membrane) contains the ATP-binding site. Exports arsenite and antimonite anions.  
**References:** [2661, 2460, 316, 3338]

[EC 3.6.3.16 created 2000]

### EC 3.6.3.17

**Accepted name:** monosaccharide-transporting ATPase  
**Reaction:** ATP + H<sub>2</sub>O + monosaccharide<sub>out</sub> = ADP + phosphate + monosaccharide<sub>in</sub>  
**Systematic name:** ATP phosphohydrolase (monosaccharide-importing)  
**Comments:** ABC-type (ATP-binding cassette-type) ATPase, characterized by the presence of two similar ATP-binding domains. Does not undergo phosphorylation during the transport process. Family of bacterial enzymes importing ribose, xylose, arabinose, galactose and methylgalactoside.  
**References:** [1138, 1541, 1419, 2495, 2721, 978]

[EC 3.6.3.17 created 2000]

[3.6.3.18 *Transferred entry. oligosaccharide-transporting ATPase. Now EC 7.5.2.2, ABC-type oligosaccharide transporter]*

[EC 3.6.3.18 created 2000, deleted 2018]

[3.6.3.19 *Transferred entry. maltose-transporting ATPase. Now EC 7.5.2.1, ABC-type maltose transporter]*

[EC 3.6.3.19 created 2000, deleted 2018]

- [3.6.3.20 *Transferred entry. glycerol-3-phosphate-transporting ATPase. Now EC 7.6.2.10, glycerol-3-phosphate-transporting ATPase*]  
[EC 3.6.3.20 created 2000, deleted 2018]
- [3.6.3.21 *Transferred entry. polar-amino-acid-transporting ATPase. Now EC 7.4.2.1, ABC-type polar-amino-acid transporter*]  
[EC 3.6.3.21 created 2000, deleted 2018]
- [3.6.3.22 *Transferred entry. nonpolar-amino-acid-transporting ATPase. Now EC 7.4.2.2, ABC-type nonpolar-amino-acid transporter*]  
[EC 3.6.3.22 created 2000, deleted 2018]
- [3.6.3.23 *Transferred entry. oligopeptide-transporting ATPase. Now EC 7.4.2.6, oligopeptide-transporting ATPase*]  
[EC 3.6.3.23 created 2000, deleted 2018]
- [3.6.3.24 *Transferred entry. nickel-transporting ATPase. Now EC 7.2.2.11, nickel-transporting ATPase*]  
[EC 3.6.3.24 created 2000, deleted 2018]
- [3.6.3.25 *Transferred entry. sulfate-transporting ATPase. Now EC 7.3.2.3, sulfate-transporting ATPase*]  
[EC 3.6.3.25 created 2000, deleted 2018]
- [3.6.3.26 *Transferred entry. nitrate-transporting ATPase. Now EC 7.3.2.4, nitrate-transporting ATPase*]  
[EC 3.6.3.26 created 2000, deleted 2018]
- [3.6.3.27 *Transferred entry. phosphate-transporting ATPase. Now EC 7.3.2.1, ABC-type phosphate transporter*]  
[EC 3.6.3.27 created 2000, deleted 2018]
- [3.6.3.28 *Transferred entry. phosphonate-transporting ATPase. Now EC 7.3.2.2, ABC-type phosphonate transporter*]  
[EC 3.6.3.28 created 2000, deleted 2018]
- [3.6.3.29 *Transferred entry. molybdate-transporting ATPase. Now EC 7.3.2.5, molybdate-transporting ATPase*]  
[EC 3.6.3.29 created 2000, deleted 2018]
- [3.6.3.30 *Transferred entry. Fe<sup>3+</sup>-transporting ATPase. Now EC 7.2.2.7, Fe<sup>3+</sup>-transporting ATPase*]  
[EC 3.6.3.30 created 2000, deleted 2018]
- [3.6.3.31 *Transferred entry. polyamine-transporting ATPase. Now EC 7.6.2.11, polyamine-transporting ATPase*]  
[EC 3.6.3.31 created 2000, deleted 2018]
- [3.6.3.32 *Transferred entry. quaternary-amine-transporting ATPase. Now EC 7.6.2.9, quaternary-amine-transporting ATPase*]  
[EC 3.6.3.32 created 2000, deleted 2018]
- [3.6.3.33 *Transferred entry. vitamin B<sub>12</sub>-transporting ATPase. Now EC 7.6.2.8, vitamin B<sub>12</sub>-transporting ATPase*]  
[EC 3.6.3.33 created 2000, deleted 2018]
- [3.6.3.34 *Transferred entry. iron-chelate-transporting ATPase; now recognized to be at least 3 separate enzymes EC 7.2.2.16, iron(III) hydroxamate ABC transporter, EC 7.2.2.17, ferric enterobactin ABC transporter, and EC 7.2.2.18, ferric citrate ABC transporter*]  
[EC 3.6.3.34 created 2000, deleted 2018]
- [3.6.3.35 *Transferred entry. manganese-transporting ATPase. Now EC 7.2.2.5, manganese-transporting ATPase*]

- [EC 3.6.3.35 created 2000, deleted 2018]
- [3.6.3.36 *Transferred entry. taurine-transporting ATPase. Now EC 7.6.2.7, taurine-transporting ATPase*]  
[EC 3.6.3.36 created 2000, deleted 2018]
- [3.6.3.37 *Transferred entry. guanine-transporting ATPase. Now EC 7.6.2.6, guanine-transporting ATPase*]  
[EC 3.6.3.37 created 2000, deleted 2018]
- [3.6.3.38 *Transferred entry. capsular-polysaccharide-transporting ATPase. Now EC 7.6.2.2, ABC-type capsular-polysaccharide transporter*]  
[EC 3.6.3.38 created 2000, deleted 2018]
- [3.6.3.39 *Transferred entry. lipopolysaccharide-transporting ATPase. Now EC 7.5.2.5, lipopolysaccharide-transporting ATPase*]  
[EC 3.6.3.39 created 2000, deleted 2018]
- [3.6.3.40 *Transferred entry. teichoic-acid-transporting ATPase. Now EC 7.5.2.4, teichoic-acid-transporting ATPase*]  
[EC 3.6.3.40 created 2000, deleted 2018]
- [3.6.3.41 *Transferred entry. heme-transporting ATPase. Now EC 7.6.2.5, heme-transporting ATPase*]  
[EC 3.6.3.41 created 2000, deleted 2018]
- [3.6.3.42 *Transferred entry.  $\beta$ -glucan-transporting ATPase. Now EC 7.5.2.3,  $\beta$ -glucan-transporting ATPase*]  
[EC 3.6.3.42 created 2000, deleted 2018]
- [3.6.3.43 *Transferred entry. peptide-transporting ATPase. Now EC 7.4.2.5, peptide-transporting ATPase*]  
[EC 3.6.3.43 created 2000, deleted 2018]
- [3.6.3.44 *Transferred entry. xenobiotic-transporting ATPase. Now EC 7.6.2.2, ABC-type xenobiotic transporter*]  
[EC 3.6.3.44 created 2000 (EC 3.6.3.45 incorporated 2006), modified 2006, deleted 2018]
- [3.6.3.45 *Deleted entry. steroid-transporting ATPase. Now included with EC 3.6.3.44, xenobiotic-transporting ATPase*]  
[EC 3.6.3.45 created 2000, deleted 2006]
- [3.6.3.46 *Transferred entry. cadmium-transporting ATPase. Now EC 7.2.2.2, ABC-type  $Cd^{2+}$  transporter*]  
[EC 3.6.3.46 created 2000, transferred 2018 to EC 7.2.2.2, deleted 2018]
- [3.6.3.47 *Transferred entry. fatty-acyl-CoA-transporting ATPase. Now EC 7.6.2.4, fatty-acyl-CoA-transporting ATPase*]  
[EC 3.6.3.47 created 2000, deleted 2018]
- [3.6.3.48 *Transferred entry.  $\alpha$ -factor-transporting ATPase. Now EC 7.4.2.7 as  $\alpha$ -factor-pheromone transporting ATPase*]  
[EC 3.6.3.48 created 2000, deleted 2018]
- [3.6.3.49 *Transferred entry. channel-conductance-controlling ATPase. Now EC 5.6.1.6, channel-conductance-controlling ATPase*]  
[EC 3.6.3.49 created 2000, deleted 2018]
- [3.6.3.50 *Transferred entry. protein-secreting ATPase. Now EC 7.4.2.8, protein-secreting ATPase*]  
[EC 3.6.3.50 created 2000, deleted 2018]
- [3.6.3.51 *Transferred entry. mitochondrial protein-transporting ATPase. Now EC 7.4.2.3, mitochondrial protein-transporting ATPase*]

[EC 3.6.3.51 created 2000, deleted 2018]

[3.6.3.52 *Transferred entry. chloroplast protein-transporting ATPase. Now EC 7.4.2.4, chloroplast protein-transporting ATPase*]

[EC 3.6.3.52 created 2000, deleted 2018]

[3.6.3.53 *Transferred entry. Ag<sup>+</sup>-exporting ATPase. Now EC 7.2.2.15, Ag<sup>+</sup>-exporting ATPase*]

[EC 3.6.3.53 created 2000, deleted 2018]

[3.6.3.54 *Transferred entry. Cu<sup>+</sup>-exporting ATPase. Now EC 7.2.2.8, Cu<sup>+</sup>-exporting ATPase*]

[EC 3.6.3.54 created 2013, deleted 2018]

[3.6.3.55 *Transferred entry. tungstate-importing ATPase. Now EC 7.3.2.6, tungstate-importing ATPase*]

[EC 3.6.3.55 created 2013, deleted 2018]

### EC 3.6.4 Acting on acid anhydrides to facilitate cellular and subcellular movement

[3.6.4.1 *Transferred entry. myosin ATPase. Now EC 5.6.1.8, myosin ATPase*]

[EC 3.6.4.1 created 1984 as EC 3.6.1.32, transferred 2000 to EC 3.6.4.1, deleted 2018]

[3.6.4.2 *Transferred entry. dynein ATPase. Now EC 5.6.1.2, dynein ATPase*]

[EC 3.6.4.2 created 1984 as EC 3.6.1.33, transferred 2000 to EC 3.6.4.2, deleted 2018]

[3.6.4.3 *Transferred entry. microtubule-severing ATPase. Now EC 5.6.1.1, microtubule-severing ATPase*]

[EC 3.6.4.3 created 2000 as 3.6.4.3, deleted 2018]

[3.6.4.4 *Transferred entry. plus-end-directed kinesin ATPase. Now EC 5.6.1.3, plus-end-directed kinesin ATPase*]

[EC 3.6.4.4 created 2000, deleted 2018]

[3.6.4.5 *Transferred entry. minus-end-directed kinesin ATPase. Now EC 5.6.1.4, minus-end-directed kinesin ATPase*]

[EC 3.6.4.5 created 2000, deleted 2018]

#### EC 3.6.4.6

**Accepted name:** vesicle-fusing ATPase  
**Reaction:** ATP + H<sub>2</sub>O = ADP + phosphate  
**Systematic name:** ATP phosphohydrolase (vesicle-fusing)  
**Comments:** A large family of ATP-hydrolysing enzymes involved in the heterotypic fusion of membrane vesicles with target membranes and the homotypic fusion of various membrane compartments. They belong to the AAA-type (ATPase associated with a variety of cell activities) ATPase superfamily. They include peroxin, which apparently is involved in Zellweger's syndrome.  
**References:** [472, 1239, 112]

[EC 3.6.4.6 created 2000]

#### EC 3.6.4.7

**Accepted name:** peroxisome-assembly ATPase  
**Reaction:** ATP + H<sub>2</sub>O = ADP + phosphate  
**Other name(s):** peroxisome assembly factor-2  
**Systematic name:** ATP phosphohydrolase (peroxisome-assembling)  
**Comments:** An extremely diversified group of enzymes that use the energy of ATP hydrolysis to import and assemble peroxisome components into the organelle. Their molecular masses range from 25 to 600 kDa.



**References:** [1638, 2982, 3240]

[EC 3.6.4.7 created 2000]

[3.6.4.8 *Transferred entry. proteasome ATPase. Now EC 5.6.1.5, proteasome ATPase*]

[EC 3.6.4.8 created 2000, deleted 2018]

[3.6.4.9 *Transferred entry. chaperonin ATPase. Now EC 5.6.1.7, chaperonin ATPase*]

[EC 3.6.4.9 created 2000, deleted 2018]

#### EC 3.6.4.10

**Accepted name:** non-chaperonin molecular chaperone ATPase  
**Reaction:** ATP + H<sub>2</sub>O = ADP + phosphate  
**Other name(s):** molecular chaperone Hsc70 ATPase  
**Systematic name:** ATP phosphohydrolase (polypeptide-polymerizing)  
**Comments:** This is a highly diverse group of enzymes that perform many functions that are similar to those of chaperonins. They comprise a number of heat-shock-cognate proteins. They are also active in clathrin uncoating and in the oligomerization of actin.  
**References:** [2487, 244, 3140, 2747, 1673]

[EC 3.6.4.10 created 2000]

[3.6.4.11 *Deleted entry. nucleoplasmin ATPase. The activity has been shown not to take place.*]

[EC 3.6.4.11 created 2000, deleted 2018]

#### EC 3.6.4.12

**Accepted name:** DNA helicase  
**Reaction:** ATP + H<sub>2</sub>O = ADP + phosphate  
**Other name(s):** 3' to 5' DNA helicase; 3'-5' DNA helicase; 3'-5' PfdH; 5' to 3' DNA helicase; AvDH1; BACH1 helicase; BcMCM; BLM protein; BRCA1-associated C-terminal helicase; CeWRN-1; Dbp9p; Dm-RECQ5; DNA helicase 120; DNA helicase A; DNA helicase E; DNA helicase II; DNA helicase III; DNA helicase RECQL5β; DNA helicase VI; *dnaB*; DnaB helicase E1; helicase HDH IV; Hel E; helicase DnaB; helicase domain of bacteriophage T<sub>7</sub> gene 4 protein helicase; PcrA helicase; UvrD; hHcsA; Hmi1p; hPif1; MCM helicase; MCM protein; MER3 helicase; MER3 protein; MPH1; PcrA; PcrA helicase; PDH120; PfdH A; Pfh1p; PIF1  
**Systematic name:** ATP phosphohydrolase (DNA helix unwinding)  
**Comments:** DNA helicases utilize the energy from ATP hydrolysis to unwind double-stranded DNA. Some of them unwind duplex DNA with a 3' to 5' polarity [1,3,5,8], others show 5' to 3' polarity [10,11,12,13] or unwind DNA in both directions [2056, 2476]. Some helicases unwind DNA as well as RNA [814, 1279]. May be identical with EC 3.6.4.13 (RNA helicase).  
**References:** [2219, 2885, 2034, 1617, 2273, 202, 2283, 521, 814, 1279, 1280, 3336, 890, 2056, 2476]

[EC 3.6.4.12 created 2009]

#### EC 3.6.4.13

**Accepted name:** RNA helicase  
**Reaction:** ATP + H<sub>2</sub>O = ADP + phosphate  
**Other name(s):** CSFV NS3 helicase; DBP2; DbpA; DDX17; DDX25; DDX3; DDX3X; DDX3Y; DDX4; DDX5; DEAD-box protein DED1; DEAD-box RNA helicase; DEAH-box protein 2; DEAH-box RNA helicase; DED1; Dex(H/D) RNA helicase; EhDEAD1; EhDEAD1 RNA helicase; eIF4A helicase; KOKV helicase; Mtr4p; nonstructural protein 3 helicase; NPH-II; RHA; RNA helicase A; RNA helicase DDX3; RNA helicase Hera; RNA-dependent ATPase; TGBp1 NTPase/helicase domain; VRH1; GRTH/DDX25

**Systematic name:** ATP phosphohydrolase (RNA helix unwinding)  
**Comments:** RNA helicases utilize the energy from ATP hydrolysis to unwind RNA. Some of them unwind RNA with a 3' to 5' polarity [1618], other show 5' to 3' polarity [?]. Some helicases unwind DNA as well as RNA [814, ?]. May be identical with EC 3.6.4.12 (DNA helicase).  
**References:** [487, 2445, 1618, 1670, 3218, 992, 814, ?]

[EC 3.6.4.13 created 2009]

## EC 3.6.5 Acting on GTP to facilitate cellular and subcellular movement

### EC 3.6.5.1

**Accepted name:** heterotrimeric G-protein GTPase  
**Reaction:**  $\text{GTP} + \text{H}_2\text{O} = \text{GDP} + \text{phosphate}$   
**Systematic name:** GTP phosphohydrolase (signalling)  
**Comments:** This group comprises GTP-hydrolysing systems, where GTP and GDP alternate in binding. This group includes stimulatory and inhibitory G-proteins such as  $G_s$ ,  $G_i$ ,  $G_o$  and  $G_{olf}$ , targeting adenylate cyclase and/or  $\text{K}^+$  and  $\text{Ca}^{2+}$  channels;  $G_q$  stimulating phospholipase C; transducin activating cGMP phosphodiesterase; gustducin activating cAMP phosphodiesterase.  $G_{olf}$  is instrumental in odour perception, transducin in vision and gustducin in taste recognition. At least 16 different  $\alpha$  subunits (39-52 kDa), 5  $\beta$  subunits (36 kDa) and 12  $\gamma$  subunits (6-9 kDa) are known.  
**References:** [2064, 2743, 261, 1924]

[EC 3.6.5.1 created 2000 as EC 3.6.1.46, transferred 2003 to EC 3.6.5.1]

### EC 3.6.5.2

**Accepted name:** small monomeric GTPase  
**Reaction:**  $\text{GTP} + \text{H}_2\text{O} = \text{GDP} + \text{phosphate}$   
**Systematic name:** GTP phosphohydrolase (cell-regulating)  
**Comments:** A family of about 50 enzymes with a molecular mass of 21 kDa that are distantly related to the  $\alpha$ -subunit of heterotrimeric G-protein GTPase (EC 3.6.5.1). They are involved in cell-growth regulation (Ras subfamily), membrane vesicle traffic and uncoating (Rab and ARF subfamilies), nuclear protein import (Ran subfamily) and organization of the cytoskeleton (Rho and Rac subfamilies).  
**References:** [270, 1026, 899, 3071]

[EC 3.6.5.2 created 2000 as EC 3.6.1.47, transferred 2003 to EC 3.6.5.2]

### EC 3.6.5.3

**Accepted name:** protein-synthesizing GTPase  
**Reaction:**  $\text{GTP} + \text{H}_2\text{O} = \text{GDP} + \text{phosphate}$   
**Other name(s):** elongation factor (EF); initiation factor (IF); peptide-release or termination factor  
**Systematic name:** GTP phosphohydrolase (mRNA-translation-assisting)  
**Comments:** This enzyme comprises a family of proteins involved in prokaryotic as well as eukaryotic protein synthesis. In the initiation factor complex, it is IF-2b (98 kDa) that binds GTP and subsequently hydrolyses it in prokaryotes. In eukaryotes, it is eIF-2 (150 kDa) that binds GTP. In the elongation phase, the GTP-hydrolysing proteins are the EF-Tu polypeptide of the prokaryotic transfer factor (43 kDa), the eukaryotic elongation factor EF-1 $\alpha$  (53 kDa), the prokaryotic EF-G (77 kDa), the eukaryotic EF-2 (70-110 kDa) and the signal recognition particle that play a role in endoplasmic reticulum protein synthesis (325 kDa). EF-Tu and EF-1 $\alpha$  catalyse binding of aminoacyl-tRNA to the ribosomal A-site, while EF-G and EF-2 catalyse the translocation of peptidyl-tRNA from the A-site to the P-site. GTPase activity is also involved in polypeptide release from the ribosome with the aid of the pRFs and eRFs.  
**References:** [1576, 1469, 2447, 805, 1526]

[EC 3.6.5.3 created 2000 as EC 3.6.1.48, transferred 2003 to EC 3.6.5.3]

#### EC 3.6.5.4

**Accepted name:** signal-recognition-particle GTPase  
**Reaction:**  $\text{GTP} + \text{H}_2\text{O} = \text{GDP} + \text{phosphate}$   
**Systematic name:** GTP phosphohydrolase (protein-synthesis-assisting)  
**Comments:** Activity is associated with the signal-recognition particle (a protein- and RNA-containing structure involved in endoplasmic-reticulum-associated protein synthesis).  
**References:** [479, 480, 1913, 813]

[EC 3.6.5.4 created 2000 as EC 3.6.1.49, transferred 2003 to EC 3.6.5.4]

#### EC 3.6.5.5

**Accepted name:** dynamin GTPase  
**Reaction:**  $\text{GTP} + \text{H}_2\text{O} = \text{GDP} + \text{phosphate}$   
**Systematic name:** GTP phosphohydrolase (vesicle-releasing)  
**Comments:** An enzyme with a molecular mass of about 100 kDa that is involved in endocytosis and is instrumental in pinching off membrane vesicles.  
**References:** [3130, 1863, 2154]

[EC 3.6.5.5 created 2000 as EC 3.6.1.50, transferred 2003 to EC 3.6.5.5]

#### EC 3.6.5.6

**Accepted name:** tubulin GTPase  
**Reaction:**  $\text{GTP} + \text{H}_2\text{O} = \text{GDP} + \text{phosphate}$   
**Systematic name:** GTP phosphohydrolase (microtubule-releasing)  
**Comments:** An intrinsic activity of  $\alpha$ -tubulin involved in tubulin folding, division plane formation in prokaryotic cells and others.  
**References:** [3311, 2921, 2473]

[EC 3.6.5.6 created 2000 as EC 3.6.1.51, transferred 2003 to EC 3.6.5.6]

## EC 3.7 Acting on carbon-carbon bonds

This subclass contains a single sub-subclass for those enzymes that act on carbon-carbon bonds in ketonic substances (EC 3.7.1). There are relatively few carbon-carbon hydrolases and they mostly catalyse the hydrolysis of 3-oxo-carboxylic acids.

### EC 3.7.1 In ketonic substances

#### EC 3.7.1.1

**Accepted name:** oxaloacetase  
**Reaction:**  $\text{oxaloacetate} + \text{H}_2\text{O} = \text{oxalate} + \text{acetate}$   
**Other name(s):** oxalacetic hydrolase  
**Systematic name:** oxaloacetate acetylhydrolase  
**References:** [1085]

[EC 3.7.1.1 created 1961]

#### EC 3.7.1.2

**Accepted name:** fumarylacetoacetase

**Reaction:** 4-fumarylacetoacetate + H<sub>2</sub>O = acetoacetate + fumarate  
**Other name(s):** β-diketonase; fumarylacetoacetate hydrolase  
**Systematic name:** 4-fumarylacetoacetate fumarylhydrolase  
**Comments:** Also acts on other 3,5- and 2,4-dioxo acids.  
**References:** [482, 678, 1880]

[EC 3.7.1.2 created 1961]

#### EC 3.7.1.3

**Accepted name:** kynureninase  
**Reaction:** L-kynurenine + H<sub>2</sub>O = anthranilate + L-alanine  
**Systematic name:** L-kynurenine hydrolase  
**Comments:** A pyridoxal-phosphate protein. Also acts on 3'-hydroxy-L-kynurenine and some other (3-arylcarbonyl)-alanines.  
**References:** [1300, 1299, 1483, 3201]

[EC 3.7.1.3 created 1965]

#### EC 3.7.1.4

**Accepted name:** phloretin hydrolase  
**Reaction:** phloretin + H<sub>2</sub>O = phloretate + phloroglucinol  
**Other name(s):** ErPhy; lactase-phlorerin hydrolase; C-acylphenol hydrolase; 2',4,4',6'-tetrahydroxydehydrochalcone 1,3,5-trihydroxybenzenehydrolase (incorrect)  
**Systematic name:** phloretin acylhydrolase (phloroglucinol forming)  
**Comments:** Also hydrolyses other C-acylated phenols related to phloretin. Isolated from the fungus *Aspergillus niger* and the bacteria *Pantoea agglomerans* and *Eubacterium ramulus*.  
**References:** [403, 1923, 2580]

[EC 3.7.1.4 created 1972, modified 2018]

#### EC 3.7.1.5

**Accepted name:** acylpyruvate hydrolase  
**Reaction:** a 3-acylpyruvate + H<sub>2</sub>O = a carboxylate + pyruvate  
**Systematic name:** 3-acylpyruvate acylhydrolase  
**Comments:** Acts on formylpyruvate, 2,4-dioxopentanoate, 2,4-dioxohexanoate and 2,4-dioxoheptanoate.  
**References:** [3137]

[EC 3.7.1.5 created 1976]

#### EC 3.7.1.6

**Accepted name:** acetylpyruvate hydrolase  
**Reaction:** acetylpyruvate + H<sub>2</sub>O = acetate + pyruvate  
**Systematic name:** 2,4-dioxopentanoate acetylhydrolase  
**Comments:** Highly specific; does not act on pyruvate, oxaloacetate, maleylpyruvate, fumarylpyruvate or acetylacetone.  
**References:** [542]

[EC 3.7.1.6 created 1984]

#### EC 3.7.1.7

**Accepted name:** β-diketone hydrolase  
**Reaction:** nonane-4,6-dione + H<sub>2</sub>O = pentan-2-one + butanoate

**Other name(s):** oxidized PVA hydrolase  
**Systematic name:** nonane-4,6-dione acylhydrolase  
**Comments:** Also acts on the product of the action of EC 1.1.3.18 secondary-alcohol oxidase, on polyvinyl alcohols; involved in the bacterial degradation of polyvinyl alcohol.  
**References:** [2501, 2502]

[EC 3.7.1.7 created 1989]

#### EC 3.7.1.8

**Accepted name:** 2,6-dioxo-6-phenylhexa-3-enoate hydrolase  
**Reaction:** 2,6-dioxo-6-phenylhexa-3-enoate + H<sub>2</sub>O = benzoate + 2-oxopent-4-enoate  
**Other name(s):** HOHPDA hydrolase  
**Systematic name:** 2,6-dioxo-6-phenylhexa-3-enoate benzoylhydrolase  
**Comments:** Cleaves the products from biphenol, 3-isopropylcatechol and 3-methylcatechol produced by EC 1.13.11.39 biphenyl-2,3-diol 1,2-dioxygenase, by ring-fission at a -CO-C bond. Involved in the breakdown of biphenyl-related compounds by *Pseudomonas* sp.  
**References:** [2196]

[EC 3.7.1.8 created 1989]

#### EC 3.7.1.9

**Accepted name:** 2-hydroxymuconate-6-semialdehyde hydrolase  
**Reaction:** 2-hydroxymuconate-6-semialdehyde + H<sub>2</sub>O = formate + 2-oxopent-4-enoate  
**Other name(s):** 2-hydroxy-6-oxohepta-2,4-dienoate hydrolase; 2-hydroxymuconic semialdehyde hydrolase; HMSH; HOD hydrolase; *xyIF* (gene name); 2-hydroxymuconate-semialdehyde formylhydrolase; 2-hydroxymuconate-semialdehyde hydrolase  
**Systematic name:** 2-hydroxymuconate-6-semialdehyde formylhydrolase  
**Comments:** The enzyme is involved in the degradation of catechols.  
**References:** [2508, 1050, 603]

[EC 3.7.1.9 created 1990, modified 2013]

#### EC 3.7.1.10

**Accepted name:** cyclohexane-1,3-dione hydrolase  
**Reaction:** cyclohexane-1,3-dione + H<sub>2</sub>O = 5-oxohexanoate  
**Other name(s):** 1,3-cyclohexanedione hydrolase; cyclohexane-1,3-dione acylhydrolase (decyclizing)  
**Systematic name:** cyclohexane-1,3-dione acylhydrolase (ring-opening)  
**Comments:** Highly specific; does not act on other dione derivatives of cyclohexane, cyclopentane or cycloheptane.  
**References:** [534]

[EC 3.7.1.10 created 1992]

#### EC 3.7.1.11

**Accepted name:** cyclohexane-1,2-dione hydrolase  
**Reaction:** cyclohexane-1,2-dione + H<sub>2</sub>O = 6-oxohexanoate  
**Other name(s):** cyclohexane-1,2-dione acylhydrolase (decyclizing)  
**Systematic name:** cyclohexane-1,2-dione acylhydrolase (ring-opening)  
**Comments:** Highly specific; does not act on cyclohexanone or cyclohexane-1,3-dione as substrate.  
**References:** [1051, 800]

[EC 3.7.1.11 created 2009]

### EC 3.7.1.12

- Accepted name:** cobalt-precorrin 5A hydrolase  
**Reaction:** cobalt-precorrin-5A + H<sub>2</sub>O = cobalt-precorrin-5B + acetaldehyde + 2 H<sup>+</sup>  
**Other name(s):** CbiG  
**Systematic name:** cobalt-precorrin 5A acylhydrolase  
**Comments:** This enzyme hydrolyses the ring A acetate δ-lactone of cobalt-precorrin-5A resulting in the loss of the C-20 carbon and its attached methyl group in the form of acetaldehyde. This is a key reaction in the contraction of the porphyrin-type tetrapyrrole ring and its conversion to a corrin ring in the anaerobic (early cobalt insertion) adenosylcobalamin biosynthesis pathway.  
**References:** [1357, 1958]

[EC 3.7.1.12 created 2010]

### EC 3.7.1.13

- Accepted name:** 2-hydroxy-6-oxo-6-(2-aminophenyl)hexa-2,4-dienoate hydrolase  
**Reaction:** (2E,4E)-6-(2-aminophenyl)-2-hydroxy-6-oxohexa-2,4-dienoate + H<sub>2</sub>O = anthranilate + (2E)-2-hydroxypenta-2,4-dienoate  
**Other name(s):** CarC  
**Systematic name:** (2E,4E)-6-(2-aminophenyl)-2-hydroxy-6-oxohexa-2,4-dienoate acylhydrolase  
**Comments:** This enzyme catalyses the third step in the aerobic degradation pathway of carbazole. The effect of the presence of an amino group or hydroxyl group at the 2-position of the substrate is small. The enzyme has no cofactor requirement [2424].  
**References:** [2105, 2424]

[EC 3.7.1.13 created 2010]

### EC 3.7.1.14

- Accepted name:** 2-hydroxy-6-oxonona-2,4-dienedioate hydrolase  
**Reaction:** (1) (2Z,4E)-2-hydroxy-6-oxonona-2,4-diene-1,9-dioate + H<sub>2</sub>O = (2Z)-2-hydroxypenta-2,4-dienoate + succinate  
(2) (2Z,4E,7E)-2-hydroxy-6-oxonona-2,4,7-triene-1,9-dioate + H<sub>2</sub>O = (2Z)-2-hydroxypenta-2,4-dienoate + fumarate  
**Other name(s):** *mhpC* (gene name)  
**Systematic name:** (2Z,4E)-2-hydroxy-6-oxona-2,4-dienedioate succinylhydrolase  
**Comments:** This enzyme catalyses a step in a pathway of phenylpropanoid compounds degradation. The first step of the enzyme mechanism involves a reversible keto-enol tautomerization [1593].  
**References:** [328, 329, 1592, 1593, 757, 602]

[EC 3.7.1.14 created 2011, modified 2012]

[3.7.1.15 Transferred entry. (+)-caryolan-1-ol synthase. Now EC 4.2.1.138, (+)-caryolan-1-ol synthase]

[EC 3.7.1.15 created 2011, deleted 2013]

[3.7.1.16 Transferred entry. oxepin-CoA hydrolase. Now EC 3.3.2.12, oxepin-CoA hydrolase]

[EC 3.7.1.16 created 2011, deleted 2013]

### EC 3.7.1.17

- Accepted name:** 4,5:9,10-diseco-3-hydroxy-5,9,17-trioxoandrosta-1(10),2-diene-4-oate hydrolase  
**Reaction:** (1E,2Z)-3-hydroxy-5,9,17-trioxo-4,5:9,10-disecoandrosta-1(10),2-dien-4-oate + H<sub>2</sub>O = 3-[(3aS,4S,7aS)-7a-methyl-1,5-dioxo-octahydro-1H-inden-4-yl]propanoate + (2Z,4Z)-2-hydroxyhexa-2,4-dienoate  
**Other name(s):** *tesD* (gene name); *hsaD* (gene name)  
**Systematic name:** 4,5:9,10-diseco-3-hydroxy-5,9,17-trioxoandrosta-1(10),2-diene-4-oate hydrolase ( (2Z,4Z)-2-hydroxyhexa-2,4-dienoate-forming)

**Comments:** The enzyme is involved in the bacterial degradation of the steroid ring structure, and is involved in degradation of multiple steroids, such as testosterone [1179], cholesterol [586], and sitosterol.  
**References:** [1179, 586, 1585, 1586]

[EC 3.7.1.17 created 2012]

#### EC 3.7.1.18

**Accepted name:** 6-oxocamphor hydrolase  
**Reaction:** bornane-2,6-dione + H<sub>2</sub>O = [(1*S*)-4-hydroxy-2,2,3-trimethylcyclopent-3-enyl]acetate  
**Other name(s):** OCH; *camK* (gene name)  
**Systematic name:** bornane-2,6-dione hydrolase  
**Comments:** Isolated from *Rhodococcus* sp. The bornane ring system is cleaved by a retro-Claisen reaction to give the enol of  $\alpha$ -campholonate. When separate from the enzyme the enol is tautomerised to the keto form as a 6:1 mixture of [(1*S*,3*R*)-2,2,3-trimethyl-4-oxocyclopentyl]acetate and [(1*S*,3*S*)-2,2,3-trimethyl-4-oxocyclopentyl]acetate.  
**References:** [990, 3175, 1648]

[EC 3.7.1.18 created 2012]

#### EC 3.7.1.19

**Accepted name:** 2,6-dihydroxypseudooxynicotine hydrolase  
**Reaction:** 1-(2,6-dihydroxypyridin-3-yl)-4-(methylamino)butan-1-one + H<sub>2</sub>O = 2,6-dihydroxypyridine + 4-methylaminobutanoate  
**Systematic name:** 1-(2,6-dihydroxypyridin-3-yl)-4-(methylamino)butan-1-one hydrolase  
**Comments:** The enzyme, characterized from the soil bacterium *Arthrobacter nicotinovorans*, participates in nicotine degradation.  
**References:** [902, 2485]

[EC 3.7.1.19 created 2012]

#### EC 3.7.1.20

**Accepted name:** 3-fumarylpyruvate hydrolase  
**Reaction:** 3-fumarylpyruvate + H<sub>2</sub>O = fumarate + pyruvate  
**Other name(s):** *nagK* (gene name); *naaD* (gene name)  
**Systematic name:** 3-fumarylpyruvate hydrolase  
**Comments:** The enzyme is involved in bacterial degradation of 5-substituted salicylates, including gentisate (5-hydroxysalicylate), 5-nitrosalicylate and 5-halosalicylates.  
**References:** [3337, 2340]

[EC 3.7.1.20 created 2012]

#### EC 3.7.1.21

**Accepted name:** 6-oxocyclohex-1-ene-1-carbonyl-CoA hydratase  
**Reaction:** 6-oxocyclohex-1-ene-1-carbonyl-CoA + 2 H<sub>2</sub>O = 3-hydroxypimeloyl-CoA (overall reaction)  
(1a) 6-oxocyclohex-1-ene-1-carbonyl-CoA + H<sub>2</sub>O = 2-hydroxy-6-oxocyclohexane-1-carbonyl-CoA  
(1b) 2-hydroxy-6-oxocyclohexane-1-carbonyl-CoA + H<sub>2</sub>O = 3-hydroxypimeloyl-CoA  
**Other name(s):** 6-oxocyclohex-1-ene-1-carbonyl-CoA hydrolase; 6-oxocyclohex-1-ene-1-carbonyl-CoA hydrolase (decyclizing)  
**Systematic name:** 6-oxocyclohex-1-ene-1-carbonyl-CoA hydrolase (ring-opening)  
**Comments:** The enzyme, which participates in the anaerobic benzoyl-CoA degradation pathway in certain organisms, catalyses the addition of one molecule of water to the double bond of 6-oxocyclohex-1-ene-1-carbonyl-CoA followed by the hydrolytic C-C cleavage of the alicyclic ring.  
**References:** [1589, 1557]

[EC 3.7.1.21 created 2014]

#### EC 3.7.1.22

**Accepted name:** 3D-(3,5/4)-trihydroxycyclohexane-1,2-dione acylhydrolase (ring-opening)  
**Reaction:** 3D-3,5/4-trihydroxycyclohexa-1,2-dione + H<sub>2</sub>O = 5-deoxy-D-glucuronate  
**Other name(s):** IoID; THcHDO hydrolase; 3D-3,5/4-trihydroxycyclohexa-1,2-dione hydrolase (decyclizing); 3D-(3,5/4)-trihydroxycyclohexane-1,2-dione acylhydrolase (decyclizing)  
**Systematic name:** 3D-3,5/4-trihydroxycyclohexa-1,2-dione hydrolase (ring-opening)  
**Comments:** The enzyme, found in the bacterium *Bacillus subtilis*, is part of the *myo*-inositol degradation pathway leading to acetyl-CoA.  
**References:** [3299]

[EC 3.7.1.22 created 2014, modified 2014]

#### EC 3.7.1.23

**Accepted name:** maleylpyruvate hydrolase  
**Reaction:** 3-maleylpyruvate + H<sub>2</sub>O = maleate + pyruvate  
**Other name(s):** *hbzF* (gene name)  
**Systematic name:** (2Z)-4,6-dioxohept-2-enedioate acylhydrolase  
**Comments:** The enzyme, characterized from the bacterium *Pseudomonas alcaligenes* NCIMB 9867, catalyses the hydrolysis of 3-maleylpyruvate, the ring-cleavage product of gentisate. The enzyme can also act on a number of maleylpyruvate derivatives, such as (2E)-2-methyl-4,6-dioxohept-2-enedioate and (2E)-3-methyl-4,6-dioxohept-2-enedioate. Activated by Mn<sup>2+</sup>. May be identical to EC 3.7.1.5, acylpyruvate hydrolase.  
**References:** [1175, 178, 1722]

[EC 3.7.1.23 created 2016]

## EC 3.8 Acting on halide bonds

This subclass contains enzymes that hydrolyse carbon-halide compounds in a single sub-subclass (EC 3.8.1).

### EC 3.8.1 In carbon-halide compounds

#### EC 3.8.1.1

**Accepted name:** alkylhalidase  
**Reaction:** bromochloromethane + H<sub>2</sub>O = formaldehyde + bromide + chloride  
**Other name(s):** halogenase; haloalkane halidohydrolase; haloalkane dehalogenase  
**Systematic name:** alkyl-halide halidohydrolase  
**References:** [1125]

[EC 3.8.1.1 created 1961]

#### EC 3.8.1.2

**Accepted name:** (S)-2-haloacid dehalogenase  
**Reaction:** (S)-2-haloacid + H<sub>2</sub>O = (R)-2-hydroxyacid + halide  
**Other name(s):** 2-haloacid dehalogenase[ambiguous]; 2-haloacid halidohydrolase [ambiguous][ambiguous]; 2-haloalkanoic acid dehalogenase; 2-haloalkanoic acid halidohydrolase; 2-halocarboxylic acid dehalogenase II; DL-2-haloacid dehalogenase[ambiguous]; L-2-haloacid dehalogenase; L-DEX  
**Systematic name:** (S)-2-haloacid halidohydrolase



**Comments:** Acts on acids of short chain lengths, C<sub>2</sub> to C<sub>4</sub>, with inversion of configuration at C-2. [See also EC 3.8.1.9 (*R*)-2-haloacid dehalogenase, EC 3.8.1.10 2-haloacid dehalogenase (configuration-inverting) and EC 3.8.1.11 2-haloacid dehalogenase (configuration-retaining)]

**References:** [949, 1990, 1471, 611, 1983, 1495, 1987, 1567, 2707]

[EC 3.8.1.2 created 1972, modified 2003]

### EC 3.8.1.3

**Accepted name:** haloacetate dehalogenase  
**Reaction:** haloacetate + H<sub>2</sub>O = glycolate + halide  
**Other name(s):** monohaloacetate dehalogenase  
**Systematic name:** haloacetate halidohydrolase  
**References:** [946, 948]

[EC 3.8.1.3 created 1972]

[3.8.1.4 *Transferred entry. thyroxine deiodinase. Now EC 1.97.1.10, thyroxine 5'-deiodinase*]

[EC 3.8.1.4 created 1984, deleted 2003]

### EC 3.8.1.5

**Accepted name:** haloalkane dehalogenase  
**Reaction:** 1-haloalkane + H<sub>2</sub>O = a primary alcohol + halide  
**Other name(s):** 1-chlorohexane halidohydrolase; 1-haloalkane dehalogenase  
**Systematic name:** 1-haloalkane halidohydrolase  
**Comments:** Acts on a wide range of 1-haloalkanes, haloalcohols, haloalkenes and some haloaromatic compounds.  
**References:** [1431, 2583, 3287]

[EC 3.8.1.5 created 1989]

### EC 3.8.1.6

**Accepted name:** 4-chlorobenzoate dehalogenase  
**Reaction:** 4-chlorobenzoate + H<sub>2</sub>O = 4-hydroxybenzoate + chloride  
**Other name(s):** halobenzoate dehalogenase  
**Systematic name:** 4-chlorobenzoate chlorohydrolase  
**Comments:** Catalyses the first step in the degradation of chlorobenzoate in *Pseudomonas*. In many microorganisms, this activity comprises three separate enzymes, EC 6.2.1.33 (4-chlorobenzoate—CoA ligase), EC 3.8.1.7 (4-chlorobenzoyl-CoA dehalogenase) and EC 3.1.2.23 (4-hydroxybenzoyl-CoA thioesterase).  
**References:** [1998, 1125]

[EC 3.8.1.6 created 1989, modified 1999]

### EC 3.8.1.7

**Accepted name:** 4-chlorobenzoyl-CoA dehalogenase  
**Reaction:** 4-chlorobenzoyl-CoA + H<sub>2</sub>O = 4-hydroxybenzoyl CoA + chloride  
**Systematic name:** 4-chlorobenzoyl CoA chlorohydrolase  
**Comments:** Specific for dehalogenation at the 4-position. Can dehalogenate substrates bearing fluorine, chlorine, bromine and iodine in the 4-position. This enzyme is part of the bacterial 2,4-dichlorobenzoate degradation pathway.  
**References:** [394, 508]

[EC 3.8.1.7 created 1999]

#### EC 3.8.1.8

- Accepted name:** atrazine chlorohydrolase  
**Reaction:** atrazine + H<sub>2</sub>O = hydroxyatrazine + chloride  
**Other name(s):** AtzA  
**Systematic name:** atrazine chlorohydrolase  
**Comments:** Involved in the degradation of the herbicide atrazine, 2-chloro-4-(ethylamino)-6-(isopropylamino)-1,3,5-triazine, in bacteria.  
**References:** [562, 561]

[EC 3.8.1.8 created 2000, modified 2011]

#### EC 3.8.1.9

- Accepted name:** (*R*)-2-haloacid dehalogenase  
**Reaction:** (*R*)-2-haloacid + H<sub>2</sub>O = (*S*)-2-hydroxyacid + halide  
**Other name(s):** 2-haloalkanoic acid dehalogenase[ambiguous]; 2-haloalkanoic acid halidohydrolase[ambiguous]; D-2-haloacid dehalogenase; D-DEX  
**Systematic name:** (*R*)-2-haloacid halidohydrolase  
**Comments:** Acts on acids of short chain lengths, C<sub>2</sub> to C<sub>4</sub>, with inversion of configuration at C-2. [See also EC 3.8.1.2 (*S*)-2-haloacid dehalogenase, EC 3.8.1.10 2-haloacid dehalogenase (configuration-inverting) and EC 3.8.1.11 2-haloacid dehalogenase (configuration-retaining)]  
**References:** [2698, 1641, 2707]

[EC 3.8.1.9 created 2003]

#### EC 3.8.1.10

- Accepted name:** 2-haloacid dehalogenase (configuration-inverting)  
**Reaction:** (1) (*S*)-2-haloacid + H<sub>2</sub>O = (*R*)-2-hydroxyacid + halide  
(2) (*R*)-2-haloacid + H<sub>2</sub>O = (*S*)-2-hydroxyacid + halide  
**Other name(s):** 2-haloalkanoic acid dehalogenase; 2-haloalkanoic acid halidohydrolase; DL-2-haloacid dehalogenase; DL-2-haloacid dehalogenase (inversion of configuration); DL-2-haloacid halidohydrolase (inversion of configuration); DL-DEXi; (*R,S*)-2-haloacid dehalogenase (configuration-inverting)  
**Systematic name:** (*S*)-2-haloacid dehalogenase (configuration-inverting)  
**Comments:** Dehalogenates both (*S*)- and (*R*)-2-haloalkanoic acids to the corresponding (*R*)- and (*S*)-hydroxyalkanoic acids, respectively, with inversion of configuration at C-2. The enzyme from *Pseudomonas* sp. 113 acts on 2-haloalkanoic acids whose carbon chain lengths are five or less. [See also EC 3.8.1.2 (*S*)-2-haloacid dehalogenase, EC 3.8.1.9 (*R*)-2-haloacid dehalogenase and EC 3.8.1.11 2-haloacid dehalogenase (configuration-retaining)]  
**References:** [1987, 1989, 1988, 1567, 1721, 354, 1641, 3149, 2707]

[EC 3.8.1.10 created 2003]

#### EC 3.8.1.11

- Accepted name:** 2-haloacid dehalogenase (configuration-retaining)  
**Reaction:** (1) (*S*)-2-haloacid + H<sub>2</sub>O = (*S*)-2-hydroxyacid + halide  
(2) (*R*)-2-haloacid + H<sub>2</sub>O = (*R*)-2-hydroxyacid + halide  
**Other name(s):** 2-haloalkanoic acid dehalogenase; 2-haloalkanoic acid halidohydrolase; DL-2-haloacid dehalogenase; DL-DEXr  
**Systematic name:** (*S*)-2-haloacid dehalogenase (configuration-retaining)  
**Comments:** Dehalogenates both (*S*)- and (*R*)-2-haloalkanoic acids to the corresponding (*S*)- and (*R*)-hydroxyalkanoic acids, respectively, with retention of configuration at C-2. [See also EC 3.8.1.2 (*S*)-2-haloacid dehalogenase, EC 3.8.1.9 (*R*)-2-haloacid dehalogenase and EC 3.8.1.10 2-haloacid dehalogenase (configuration-inverting)]  
**References:** [3149, 2707]

[EC 3.8.1.11 created 2003]

## EC 3.8.2 In phosphorus-halide compounds (deleted sub-subclass)

[3.8.2.1 *Transferred entry. di-isopropyl-fluorophosphatase. Now EC 3.1.8.2, diisopropyl-fluorophosphatase*]

[EC 3.8.2.1 created 1961, modified 1976, deleted 1992]

## EC 3.9 Acting on phosphorus-nitrogen bonds

This subclass contains a single sub-subclass for enzymes that act on phosphorus-nitrogen bonds (EC 3.9.1).

### EC 3.9.1 Acting on phosphorus-nitrogen bonds (only sub-subclass identified to date)

#### EC 3.9.1.1

**Accepted name:** phosphoamidase  
**Reaction:**  $N$ -phosphocreatine + H<sub>2</sub>O = creatine + phosphate  
**Other name(s):** creatine phosphatase  
**Systematic name:** phosphamide hydrolase  
**Comments:** Also acts on  $N$ -phospho-arginine and other phosphoamides. Possibly identical with EC 3.1.3.9 (glucose-6-phosphatase) or EC 3.1.3.16 (protein-serine/threonine phosphatase).  
**References:** [2242, 2668, 2801]

[EC 3.9.1.1 created 1961]

#### EC 3.9.1.2

**Accepted name:** protein arginine phosphatase  
**Reaction:** a [protein]- $N^{\omega}$ -phospho-L-arginine + H<sub>2</sub>O = a [protein]-L-arginine + phosphate  
**Other name(s):** YwIE  
**Systematic name:** [protein]- $N^{\omega}$ -phospho-L-arginine phosphohydrolase  
**Comments:** The enzyme, characterized from Gram-positive bacteria, hydrolyses the phosphoramidate (P-N) bond of  $N^{\omega}$ -phospho-L-arginine residues in proteins and peptides that were phosphorylated by EC 2.7.14.1, protein-arginine-kinase.  
**References:** [831, 2962, 695]

[EC 3.9.1.2 created 2014]

#### EC 3.9.1.3

**Accepted name:** phosphohistidine phosphatase  
**Reaction:** a [protein]- $N$ -phospho-L-histidine + H<sub>2</sub>O = a [protein]-L-histidine + phosphate  
**Other name(s):** PHPT1 (gene name); protein histidine phosphatase; PHP  
**Systematic name:** [protein]- $N$ -phospho-L-histidine phosphohydrolase  
**Comments:** This eukaryotic enzyme dephosphorylates phosphorylated histidine residues within proteins and peptides. The enzyme acts on phosphate groups attached to both the *pros*- and *tele*-nitrogen atoms, but the *pros*- position is somewhat preferred (by a factor of two at the most) [95]. The substrate specificity depends on the amino acid sequence or structural context of the phosphohistidine in a phosphoprotein. The enzyme is also active on free phosphoramidate [689, 95] and peptide-bound phospholysine [688].  
**References:** [689, 1478, 176, 95, 688]

[EC 3.9.1.3 created 2016]

## EC 3.10 Acting on sulfur-nitrogen bonds

This subclass contains a single sub-subclass for enzymes that act on sulfur-nitrogen bonds (EC 3.10.1).

### EC 3.10.1 Acting on sulfur-nitrogen bonds (only sub-subclass identified to date)

#### EC 3.10.1.1

**Accepted name:** *N*-sulfoglucosamine sulfohydrolase  
**Reaction:** *N*-sulfo-D-glucosamine + H<sub>2</sub>O = D-glucosamine + sulfate  
**Other name(s):** sulfoglucosamine sulfamidase; heparin sulfamidase; 2-desoxy-D-glucoside-2-sulphamate sulphonydrolase (sulphamate sulphonydrolase)  
**Systematic name:** *N*-sulfo-D-glucosamine sulfohydrolase  
**References:** [609, 1775]

[EC 3.10.1.1 created 1972, modified 1981, modified 1982]

#### EC 3.10.1.2

**Accepted name:** cyclamate sulfohydrolase  
**Reaction:** cyclohexylsulfamate + H<sub>2</sub>O = cyclohexylamine + sulfate  
**Other name(s):** cyclamate sulfamatase; cyclamate sulfamidase; cyclohexylsulfamate sulfamidase  
**Systematic name:** cyclohexylsulfamate sulfohydrolase  
**Comments:** Also readily hydrolyses aliphatic sulfamates with 3 to 8 carbons.  
**References:** [2084]

[EC 3.10.1.2 created 1976, modified 1981]

## EC 3.11 Acting on carbon-phosphorus bonds

This subclass contains a single sub-subclass for enzymes that hydrolyse C-phosphono-groups (EC 3.11.1).

### EC 3.11.1 Acting on carbon-phosphorus bonds (only sub-subclass identified to date)

#### EC 3.11.1.1

**Accepted name:** phosphonoacetaldehyde hydrolase  
**Reaction:** phosphonoacetaldehyde + H<sub>2</sub>O = acetaldehyde + phosphate  
**Other name(s):** phosphonatase; 2-phosphonoacetylaldehyde phosphonohydrolase  
**Systematic name:** 2-oxoethylphosphonate phosphonohydrolase  
**Comments:** This enzyme destabilizes the C-P bond, by forming an imine between one of its lysine residues and the carbonyl group of the substrate, thus allowing this, normally stable, bond to be broken. The mechanism is similar to that used by EC 4.1.2.13, fructose-bisphosphate aldolase, to break a C-C bond. Belongs to the haloacetate dehalogenase family.  
**References:** [2060, 2061, 2059, 2193, 124]

[EC 3.11.1.1 created 1972, modified 1976, modified 2001]

#### EC 3.11.1.2

**Accepted name:** phosphonoacetate hydrolase  
**Reaction:** phosphonoacetate + H<sub>2</sub>O = acetate + phosphate  
**Systematic name:** phosphonoacetate phosphonohydrolase

**Comments:** A zinc-dependent enzyme. Belongs to the alkaline phosphatase superfamily of zinc-dependent hydrolases.

**References:** [1871]

[EC 3.11.1.2 created 1999]

### EC 3.11.1.3

**Accepted name:** phosphonopyruvate hydrolase

**Reaction:** 3-phosphonopyruvate + H<sub>2</sub>O = pyruvate + phosphate

**Other name(s):** PPH

**Comments:** Highly specific for phosphonopyruvate as substrate [1546]. The reaction is not inhibited by phosphate but is inhibited by the phosphonates phosphonoformic acid, hydroxymethylphosphonic acid and 3-phosphonopropanoic acid [1546]. The enzyme is activated by the divalent cations Co<sup>2+</sup>, Mg<sup>2+</sup> and Mn<sup>2+</sup>. This enzyme is a member of the phosphoenolpyruvate mutase/isocitrate lyase superfamily [411].

**References:** [2904, 1546, 411]

[EC 3.11.1.3 created 2007]

## EC 3.12 Acting on sulfur-sulfur bonds

This subclass contains a single sub-subclass for enzymes that act on sulfur-sulfur bonds (EC 3.12.1).

### EC 3.12.1 Acting on sulfur-sulfur bonds (only sub-subclass identified to date)

#### EC 3.12.1.1

**Accepted name:** trithionate hydrolase

**Reaction:** trithionate + H<sub>2</sub>O = thiosulfate + sulfate + 2 H<sup>+</sup>

**Systematic name:** trithionate thiosulfohydrolase

**References:** [1746, 2966]

[EC 3.12.1.1 created 1990]

## EC 3.13 Acting on carbon-sulfur bonds

This subclass contains a single sub-subclass for enzymes that act on carbon-sulfur bonds (EC 3.13.1).

### EC 3.13.1 Acting on carbon-sulfur bonds (only sub-subclass identified to date)

#### EC 3.13.1.1

**Accepted name:** UDP-sulfoquinovose synthase

**Reaction:** UDP- $\alpha$ -D-sulfoquinovopyranose + H<sub>2</sub>O = UDP- $\alpha$ -D-glucose + sulfite

**Other name(s):** sulfite:UDP-glucose sulfotransferase; UDPsulfoquinovose synthase; UDP-6-sulfo-6-deoxyglucose sulfohydrolase

**Systematic name:** UDP-6-sulfo-6-deoxy- $\alpha$ -D-glucose sulfohydrolase

**Comments:** Requires NAD<sup>+</sup>, which appears to oxidize UDP- $\alpha$ -D-glucose to UDP-4-dehydroglucose, which dehydrates to UDP-4-dehydro-6-deoxygluc-5-enose, to which sulfite is added. The reaction is completed when the substrate is rehydrogenated at C-4. The enzyme from *Arabidopsis thaliana* is specific for UDP-Glc and sulfite.

**References:** [719, 720, 1993, 2515]

[EC 3.13.1.1 created 2001, modified 2010]

[3.13.1.2 Deleted entry: 5-deoxyribos-5-ylhomocysteinase. The activity is most probably attributable to EC 4.4.1.21, S-ribosylhomocysteine lyase]

[EC 3.13.1.2 created 1972 as EC 3.3.1.3, transferred 2001 to EC 3.2.1.148, transferred 2004 to EC 3.13.1.2, deleted 2005]

#### EC 3.13.1.3

**Accepted name:** 2'-hydroxybiphenyl-2-sulfinate desulfinate  
**Reaction:** 2'-hydroxybiphenyl-2-sulfinate + H<sub>2</sub>O = 2-hydroxybiphenyl + sulfite  
**Other name(s):** gene *dszB*-encoded hydrolase; 2-(2-hydroxyphenyl) benzenesulfinate:H<sub>2</sub>O hydrolase; DszB; HBPSi desulfinate; 2-(2-hydroxyphenyl) benzenesulfinate sulfohydrolase; HPBS desulfinate; 2-(2-hydroxyphenyl)benzenesulfinate hydrolase; 2-(2'-hydroxyphenyl)benzenesulfinate desulfinate; 2-(2-hydroxyphenyl)benzenesulfinate desulfinate  
**Systematic name:** 2'-hydroxybiphenyl-2-sulfinate sulfohydrolase  
**Comments:** The enzyme from *Rhodococcus* sp. strain IGTS8 is encoded by the plasmid-encoded dibenzothiophene-desulfurization (*dsz*) operon. The enzyme has a narrow substrate specificity with biphenyl-2-sulfinate being the only other substrate known to date [2053].  
**References:** [2192, 2053, 3136]

[EC 3.13.1.3 created 2000 as EC 3.1.2.24, transferred 2005 to EC 3.13.1.3]

#### EC 3.13.1.4

**Accepted name:** 3-sulfinopropanoyl-CoA desulfinate  
**Reaction:** 3-sulfinopropanoyl-CoA + H<sub>2</sub>O = propanoyl-CoA + sulfite  
**Other name(s):** 3SP-CoA desulfinate; AcdDPN7; 3-sulfinopropionyl-CoA desulfinate  
**Systematic name:** 3-sulfinopropanoyl-CoA sulfinohydrolase  
**Comments:** The enzyme from the β-proteobacterium *Advenella mimigardefordensis* contains one non-covalently bound FAD per subunit.  
**References:** [2595, 2594]

[EC 3.13.1.4 created 2014]

#### EC 3.13.1.5

**Accepted name:** carbon disulfide hydrolase  
**Reaction:** carbon disulfide + 2 H<sub>2</sub>O = CO<sub>2</sub> + 2 hydrogen sulfide (overall reaction)  
(1a) carbon disulfide + H<sub>2</sub>O = carbonyl sulfide + hydrogen sulfide  
(1b) carbonyl sulfide + H<sub>2</sub>O = CO<sub>2</sub> + hydrogen sulfide  
**Other name(s):** CS<sub>2</sub> hydrolase (misleading); carbon disulfide lyase; CS<sub>2</sub>-converting enzyme; carbon disulphide-lyase (decarboxylating)  
**Systematic name:** carbon-disulfide hydrogen-sulfide-lyase (decarboxylating)  
**Comments:** The enzyme contains Zn<sup>2+</sup>. The hyperthermophilic archaeon *Acidianus* sp. A1-3 obtains energy by the conversion of carbon disulfide to hydrogen sulfide, with carbonyl sulfide as an intermediate.  
**References:** [2695]

[EC 3.13.1.5 created 2013 as EC 4.4.1.27, transferred 2017 to EC 3.13.1.5]

#### EC 3.13.1.6

**Accepted name:** [CysO sulfur-carrier protein]-S-L-cysteine hydrolase  
**Reaction:** [CysO sulfur-carrier protein]-Gly-NH-CH<sub>2</sub>-C(O)-S-L-cysteine + H<sub>2</sub>O = [CysO sulfur-carrier protein]-Gly-NH-CH<sub>2</sub>-COOH + L-cysteine  
**Other name(s):** *mec* (gene name)

**Systematic name:** [CysO sulfur-carrier protein]-S-L-cysteine sulfohydrolase

**Comments:** Requires Zn<sup>2+</sup>. The enzyme, characterized from the bacterium *Mycobacterium tuberculosis*, participates in an L-cysteine biosynthesis pathway. It acts on the product of EC 2.5.1.113, [CysO sulfur-carrier protein]-thiocarboxylate-dependent cysteine synthase.

**References:** [330]

[EC 3.13.1.6 created 2017]

#### EC 3.13.1.7

**Accepted name:** carbonyl sulfide hydrolase

**Reaction:** carbonyl sulfide + H<sub>2</sub>O = hydrogen sulfide + CO<sub>2</sub>

**Other name(s):** COSase; COS hydrolase; *cos* (gene name)

**Systematic name:** carbonyl sulfide hydrogen-sulfide-lyase (decarboxylating)

**Comments:** The enzyme, characterized from the bacterium *Thiobacillus thioparus*, catalyses a step in the degradation pathway of thiocyanate. This activity is also catalysed by the archaeal EC 3.13.1.5, carbon disulfide lyase.

**References:** [2151]

[EC 3.13.1.7 created 2018]

#### EC 3.13.1.8

**Accepted name:** S-adenosyl-L-methionine hydrolase (adenosine-forming)

**Reaction:** S-adenosyl-L-methionine + H<sub>2</sub>O = adenosine + L-methionine

**Other name(s):** SAM hydroxide adenosyltransferase

**Systematic name:** S-adenosyl-L-methionine hydrolase (adenosine-forming)

**Comments:** The enzyme, found in bacteria and archaea, catalyses a nucleophilic attack of water at the C5' carbon of S-adenosyl-L-methionine to generate adenosine and L-methionine. May be involved in regulating SAM levels in the cell. *cf.* EC 3.3.1.2, S-adenosyl-L-methionine hydrolase (L-homoserine-forming).

**References:** [722, 583]

[EC 3.13.1.8 created 2018]

## References

- [1] In E. Reiner, W.N. Aldridge, and C.G. Hoskin, editors, *Enzymes Hydrolysing Organophosphorus Compounds*. Ellis Horwood, Chichester, UK, 1989.
- [2] J. Abdallah, M. Mihoub, V. Gautier, and G. Richarme. The DJ-1 superfamily members YhbO and YajL from *Escherichia coli* repair proteins from glycation by methylglyoxal and glyoxal. *Biochem. Biophys. Res. Commun.*, 470:282–286, 2016.
- [3] K.M. Abdullah, R.Y.C. Lo, and A. Mellors. Cloning, nucleotide sequence, and expression of the *Pasteurella haemolytica* A1 glycoprotease gene. *J. Bacteriol.*, 173:5597–5603, 1991.
- [4] K.M. Abdullah, E.A. Udoh, P.E. Shewen, and A. Mellors. A neutral glycoprotease of *Pasteurella haemolytica* A1 specifically cleaves *O*-sialoglycoproteins. *Infect. Immun.*, 60:56–62, 1992.
- [5] M. Abe, K. Ohno, and R. Sato. Possible identity of lysolecithin acyl-hydrolase with lysolecithin-lysolecithin acyl-transferase in rat-lung soluble fraction. *Biochim. Biophys. Acta*, 369:361–370, 1974.
- [6] N. Abe and S. Tsuiki. Studies on glycogen synthase D phosphatase of rat liver - multiple nature. *Biochim. Biophys. Acta*, 350:383–391, 1974.
- [7] R.M. Abra and P.J. Quinn. A novel pathway for phosphatidylcholine catabolism in rat brain homogenates. *Biochim. Biophys. Acta*, 380:436–441, 1975.
- [8] T. Achstetter, C. Ehmann, and D.H. Wolf. Aminopeptidase Co, a new yeast peptidase. *Biochem. Biophys. Res. Commun.*, 109:341–347, 1982.
- [9] T. Achstetter, C. Ehmann, and D.H. Wolf. Proteinase yscD. Purification and characterization of a new yeast peptidase. *J. Biol. Chem.*, 260:4584–4590, 1985.
- [10] T. Achstetter and D.H. Wolf. Hormone processing and membrane-bound proteinases in yeast. *EMBO J.*, 4:173–177, 1985.
- [11] K.E. Ackermann, B.G. Gish, M.P. Honchar, and W.R. Sherman. Evidence that inositol 1-phosphate in brain of lithium-treated rats results mainly from phosphatidylinositol metabolism. *Biochem. J.*, 242:517–524, 1987.
- [12] A.D., Markland Retzios, , and Jr. A direct-acting fibrinolytic enzyme from the venom of *Agkistrodon contortrix contortrix*: effects on various components of the human blood coagulation and fibrinolysis systems. *Thromb. Res.*, 52:541–552, 1988.
- [13] H. Adachi, M. Tsujimoto, M. Fukasawa, Y. Sato, H. Arai, K. Inoue, and T. Nishimura. cDNA cloning and expression of chicken aminopeptidase H, possessing endopeptidase as well as aminopeptidase activity. *Eur. J. Biochem.*, 245:283–288, 1997.
- [14] W. Adachi, S. Kawamoto, I. Ohno, K. Nishida, S. Kinoshita, K. Matsubara, and K. Okubo. Isolation and characterization of human cathepsin V: a major proteinase in corneal epithelium. *Invest. Ophthalmol. Vis. Sci.*, 39:1789–1796, 1998.
- [15] J.M. Adams. On the release of the formyl group from nascent protein. *J. Mol. Biol.*, 33:571–589, 1968.
- [16] M. Adams, N.K. Richtmyer, and C.S. Hudson. Some enzymes present in highly purified invertase preparations; a contribution to the study of fructofuranosidases, galactosidases, glucosidases and mannosidases. *J. Am. Chem. Soc.*, 65:1369–1380, 1943.
- [17] G.S. Adrian and R.W. Keenan. A dolichyl phosphate-cleaving acid phosphatase from *Tetrahymena pyriformis*. *Biochim. Biophys. Acta*, 575:431–438, 1979.
- [18] P. Aducci, P. Ascenzi, and A. Ballio. Esterolytic properties of leucine-proteinase, the leucine-specific serine proteinase from spinach (*Spinacia oleracea* L.). *Plant Physiol.*, 82:591–593, 1986.
- [19] P. Aducci, P. Ascenzi, M. Pierini, and A. Ballio. Purification and characterization of Leu-proteinase, the leucine specific serine proteinase from spinach (*Spinacia oleracea* L.) leaves. *Plant Physiol.*, 81:812–816, 1986.
- [20] J.A. Affholter, V.A. Fried, and R.A. Roth. Human insulin-degrading enzyme shares structural and functional homologies with *E. coli* protease III. *Science*, 242:1415–1418, 1988.



- [21] J.A. Affholter, C.-L. Hsieh, U. Francke, and R.A. Roth. Insulin-degrading enzyme: stable expression of the human complementary DNA, characterization of its protein product, and chromosomal mapping of the human and mouse genes. *Mol. Endocrinol.*, 4:1125–1135, 1990.
- [22] I. Ahel, U. Rass, S.F. El-Khamisy, S. Katyal, P.M. Clements, P.J. McKinnon, K.W. Caldecott, and S.C. West. The neurodegenerative disease protein aprataxin resolves abortive DNA ligation intermediates. *Nature*, 443:713–716, 2006.
- [23] Y.O. Ahn, M. Mizutani, H. Saino, and K. Sakata. Furcatin hydrolase from *Viburnum furcatum* Blume is a novel disaccharide-specific acuminosidase in glycosyl hydrolase family 1. *J. Biol. Chem.*, 279:23405–23414, 2004.
- [24] K. Akaji, H. Konno, M. Onozuka, A. Makino, H. Saito, and K. Nosaka. Evaluation of peptide-aldehyde inhibitors using R188I mutant of SARS 3CL protease as a proteolysis-resistant mutant. *Bioorg. Med. Chem.*, 16:9400–9408, 2008.
- [25] T.A. Akhtar, R.P. McQuinn, V. Naponelli, J.F. Gregory, Giovannoni 3rd, Hanson J.J., and A.D. Tomato  $\gamma$ -glutamylhydrolases: expression, characterization, and evidence for heterodimer formation. *Plant Physiol.*, 148:775–785, 2008.
- [26] K. Akiyama, T. Nakamura, S. Iwanaga, and M. Hara. The chymotrypsin-like activity of human prostate-specific antigen,  $\gamma$ -seminoprotein. *FEBS Lett.*, 225:168–172, 1987.
- [27] S. Alarico, M. Costa, M.S. Sousa, A. Maranha, E.C. Lourenco, T.Q. Faria, M.R. Ventura, and N. Empadinhas. *Mycobacterium hassiacum* recovers from nitrogen starvation with up-regulation of a novel glucosylglycerate hydrolase and depletion of the accumulated glucosylglycerate. *Sci Rep*, 4:6766–6766, 2014.
- [28] S. Alarico, N. Empadinhas, and M.S. da Costa. A new bacterial hydrolase specific for the compatible solutes  $\alpha$ -D-mannopyranosyl-(1 $\rightarrow$ 2)-D-glycerate and  $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 2)-D-glycerate. *Enzyme Microb. Technol.*, 52:77–83, 2013.
- [29] A.M. Albrecht, E. Boldizar, and D.J. Hutchinson. Carboxypeptidase displaying differential velocity in hydrolysis of methotrexate, 5-methyltetrahydrofolic acid, and leucovorin. *J. Bacteriol.*, 134:506–513, 1978.
- [30] W.N. Aldridge. Serum esterases. I. Two types of esterase (A and B) hydrolysing *p*-nitrophenyl acetate, propionate and butyrate and a method for their determination. *Biochem. J.*, 53:110–117, 1953.
- [31] M. Alexander, L.A. Heppel, and J. Hurwitz. The purification and properties of micrococcal nuclease. *J. Biol. Chem.*, 236:3014–3019, 1961.
- [32] S.E.H. Alexson and J. Nedergaard. A novel type of short- and medium-chain acyl-CoA hydrolases in brown adipose tissue mitochondria. *J. Biol. Chem.*, 263:13564–13571, 1988.
- [33] S.E.H. Alexson, L.T. Svensson, and J. Nedergaard. NADH-sensitive propionyl-CoA hydrolase in brown-adipose-tissue mitochondria of the rat. *Biochim. Biophys. Acta*, 1005:13–19, 1989.
- [34] A. Alhapel, D.J. Darley, N. Wagener, E. Eckel, N. Elsner, and A.J. Pierik. Molecular and functional analysis of nicotinate catabolism in *Eubacterium barkeri*. *Proc. Natl. Acad. Sci. USA*, 103:12341–12346, 2006.
- [35] B. Ali and S. Kaur. Mammalian tissue acetylsalicylic acid esterase(s): identification, distribution and discrimination from other esterases. *J. Pharmacol. Exp. Ther.*, 226:589–594, 1983.
- [36] R.H. Allen and W.B. Jakoby. Tartaric acid metabolism. IX. Synthesis with tartrate epoxidase. *J. Biol. Chem.*, 244:2078–2084, 1969.
- [37] J. Allouch, M. Jam, W. Helbert, T. Barbeyron, B. Kloareg, B. Henrissat, and M. Czjzek. The three-dimensional structures of two  $\beta$ -agarases. *J. Biol. Chem.*, 278:47171–47180, 2003.
- [38] E.S. Alnemri, D.J. Livingston, D.W. Nicholson, G. Salvesen, N.A. Thornberry, W.W. Wong, and J.Y. Yuan. Human ICE/CED-3 protease nomenclature. *Cell*, 87:171–171, 1996.
- [39] G.D. Alonso, A.C. Schoijet, H.N. Torres, and M.M. Flawiá. TcPDE4, a novel membrane-associated cAMP-specific phosphodiesterase from *Trypanosoma cruzi*. *Mol. Biochem. Parasitol.*, 145:40–49, 2006.
- [40] J.M. Martín Alonso, R. Casais, J.A. Boga, and F. Parra. Processing of rabbit hemorrhagic disease virus polyprotein. *J. Virol.*, 70:1261–1265, 1996.

- [41] J. Altenbuchner, M. Siemann-Herzberg, and C. Syltatk. Hydantoinases and related enzymes as biocatalysts for the synthesis of unnatural chiral amino acids. *Curr. Opin. Biotechnol.*, 12:559–563, 2001.
- [42] F. Altmann, B. Kwiatkowski, S. Stirn, L. März, and F.M. Unger. A bacteriophage-associated glycanase cleaving  $\beta$ -pyranosidic linkages of 3-deoxy-D-manno-2-octulosonic acid (KDO). *Biochem. Biophys. Res. Commun.*, 136:329–335, 1986.
- [43] S. Amagase. Digestive enzymes in insectivorous plants. III. Acid proteases in the genus *Nepenthes* and *Drosera peltata*. *J. Biochem. (Tokyo)*, 72:73–81, 1972.
- [44] S. Amagase, S. Nakayama, and A. Tsugita. Acid protease in *Nepenthes*. II. Study on the specificity of nepenthesin. *J. Biochem. (Tokyo)*, 66:431–439, 1969.
- [45] G. Ambrus, P. Gal, M. Kojima, K. Szilagy, J. Balczer, J. Antal, L. Graf, A. Laich, B.E. Moffatt, W. Schwaeble, R.B. Sim, and P. Zavodszky. Natural substrates and inhibitors of mannan-binding lectin-associated serine protease-1 and -2: a study on recombinant catalytic fragments. *J. Immunol.*, 170:1374–1382, 2003.
- [46] B.N. Ames. The biosynthesis of histidine; L-histidinol phosphate phosphatase. *J. Biol. Chem.*, 226:583–593, 1957.
- [47] G. Ammerer, C.P. Hunter, J.H. Rothman, G.C. Saari, L.A. Valls, and T.H. Stevens. PEP4 gene of *Saccharomyces cerevisiae* encodes proteinase A, a vacuolar enzyme required for processing of vacuolar precursors. *Mol. Cell. Biol.*, 6:2490–2499, 1987.
- [48] D.S. An, C.H. Cui, H.G. Lee, L. Wang, S.C. Kim, S.T. Lee, F. Jin, H. Yu, Y.W. Chin, H.K. Lee, W.T. Im, and S.G. Kim. Identification and characterization of a novel *Terrabacter ginsenosidimutans* sp. nov.  $\beta$ -glucosidase that transforms ginsenoside Rb1 into the rare gypenosides XVII and LXXV. *Appl. Environ. Microbiol.*, 76:5827–5836, 2010.
- [49] A. Anastasi, C.G. Knight, and A.J. Barrett. Characterization of the bacterial metalloendopeptidase pitrilysin by use of a continuous fluorescence assay. *Biochem. J.*, 290:601–607, 1993.
- [50] J.M. Anchors and N.L. Karnovsky. Purification of cerebral glucose-6-phosphatase. An enzyme involved in sleep. *J. Biol. Chem.*, 250:6408–6416, 1975.
- [51] B.M. Anderson and C.A. Lang. Nicotinamide-adenine dinucleotide pyrophosphatase in the growing and aging mosquito. *Biochem. J.*, 101:392–396, 1966.
- [52] E.T. Anderson, M.G. Wetherell, L.A. Winter, S.B. Olmsted, P.P. Cleary, and Y.V. Matsuka. Processing, stability, and kinetic parameters of C5a peptidase from *Streptococcus pyogenes*. *Eur. J. Biochem.*, 269:4839–4851, 2002.
- [53] C.E. Andersson, P. Lagerback, and K. Carlson. Structure of bacteriophage T<sub>4</sub> endonuclease II mutant E118A, a tetrameric GIY-YIG enzyme. *J. Mol. Biol.*, 397:1003–1016, 2010.
- [54] J.N. Andexer, S.G. Kendrew, M. Nur e Alam, O. Lazos, T.A. Foster, A.S. Zimmermann, T.D. Warneck, D. Suthar, N.J. Coates, F.E. Koehn, J.S. Skotnicki, G.T. Carter, M.A. Gregory, C.J. Martin, S.J. Moss, P.F. Leadlay, and B. Wilkinson. Biosynthesis of the immunosuppressants FK506, FK520, and rapamycin involves a previously undescribed family of enzymes acting on chorismate. *Proc. Natl. Acad. Sci. USA*, 108:4776–4781, 2011.
- [55] T. Ando. A nuclease specific for heat-denatured DNA isolated from a product of *Aspergillus oryzae*. *Biochim. Biophys. Acta*, 114:158–168, 1966.
- [56] A.J. Andreoli, T.W. Okita, R. Bloom, and T.A. Grover. The pyridine nucleotide cycle: presence of a nicotinamide mononucleotide-specific glycohydrolase in *Escherichia coli*. *Biochem. Biophys. Res. Commun.*, 49:264–269, 1972.
- [57] C.B. Anfinsen, P. Cuatrecasas, and H. Taniuchi. Staphylococcal nuclease, chemical properties and catalysis. In P.D. Boyer, editor, *The Enzymes*, volume 4, pages 177–204. Academic Press, New York, 3rd edition, 1971.
- [58] Y. Anraku. A new cyclic phosphodiesterase having a 3'-nucleotidase activity from *Escherichia coli* B. I. Purification and some properties of the enzyme. *J. Biol. Chem.*, 239:3412–3419, 1964.
- [59] Y. Anraku. A new cyclic phosphodiesterase having a 3'-nucleotidase activity from *Escherichia coli* B. II. Further studies on substrate specificity and mode of action of the enzyme. *J. Biol. Chem.*, 239:3420–3424, 1964.

- [60] H. Anundi, H. Ronne, P.A. Peterson, and L. Rask. Partial amino-acid sequence of the epidermal growth-factor-binding protein. *Eur. J. Biochem.*, 129:365–371, 1982.
- [61] T. Aoki, T. Araki, and M. Kitamikado. Purification and characterization of an endo- $\beta$ -1,3-xylanase from *Vibrio* species. *Nippon Suisan Gakkaishi*, 54:277–281, 1988.
- [62] J.F. Aparicio, C. Lopez-Otin, S. Cal, , and J. A. *Streptomyces glaucescens* endodeoxyribonuclease which shows a strong preference for CC dinucleotide. *Eur. J. Biochem.*, 205:695–699, 1992.
- [63] H.V. Aposhian and G.Y. Tremblay. Deoxythymidylate 5'-nucleotidase. Purification and properties of an enzyme found after infection of *Bacillus subtilis* with phage SP<sub>5</sub>C. *J. Biol. Chem.*, 241:5095–5101, 1966.
- [64] J.J. Aragón and J.M. Lowenstein. A survey of enzymes which generate or use acetoacetyl thioesters in rat liver. *J. Biol. Chem.*, 258:4725–4733, 1983.
- [65] C. Arakawa, M. Kuratsu, K. Furihata, T. Hiratsuka, N. Itoh, H. Seto, and T. Dairi. Diversity of the early step of the futasoline pathway. *Antimicrob. Agents Chemother.*, 55:913–916, 2011.
- [66] T. Araki, N. Inoue, and T. Morishita. Purification and characterization of  $\beta$ -1,3-xylanase from a marine bacterium, *Alcaligenes* sp. XY-234. *J. Gen. Appl. Microbiol.*, 44:269–274, 1998.
- [67] T. Araki and M. Kitamikado. Purification and characterization of a novel exo- $\beta$ -mannanase from *Aeromonas* sp. F-25. *J. Biochem. (Tokyo)*, 91:1181–1186, 1982.
- [68] T. Araki, S. Tani, K. Maeda, S. Hashikawa, H. Nakagawa, and T. Morishita. Purification and characterization of  $\beta$ -1,3-xylanase from a marine bacterium, *Vibrio* sp. XY-214. *Biosci. Biotechnol. Biochem.*, 63:2017–2019, 1999.
- [69] Y. Araki and E. Ito. A pathway of chitosan formation in *Mucor rouxii*: enzymatic deacetylation of chitin. *Biochem. Biophys. Res. Commun.*, 56:669–675, 1974.
- [70] C.A. Arce and H.S. Barra. Association of tubuliny-tyrosine carboxypeptidase with microtubules. *FEBS Lett.*, 157:75–78, 1983.
- [71] I.M. Archer, P.S. Harper, and F.S. Wusteman. Multiple forms of iduronate 2-sulphate sulphatase in human tissues and body fluids. *Biochim. Biophys. Acta*, 708:134–140, 1982.
- [72] J. Arima, Y. Uesugi, M. Iwabuchi, and T. Hatanaka. Study on peptide hydrolysis by aminopeptidases from *Streptomyces griseus*, *Streptomyces septatus* and *Aeromonas proteolytica*. *Appl. Microbiol. Biotechnol.*, 70:541–547, 2006.
- [73] K. Arima, J. Yu, and S. Iwasaki. Milk-clotting enzyme from *Mucor pusillus* var. lindt. *Methods Enzymol.*, 19:446–459, 1970.
- [74] F. Arminjon, M. Guinand, M.-J. Vacheron, and G. Michel. Specificity profiles of the membrane-bound  $\gamma$ -D-glutamyl-(L)*meso*-diaminopimelate endopeptidase and LD-carboxypeptidase from *Bacillus sphaericus* 9602. *Eur. J. Biochem.*, 73:557–565, 1977.
- [75] J.N. Aronson and J.C. Lugay. *N*-Formylmethionine deformylase from *Euglena gracilis*. *Biochem. Biophys. Res. Commun.*, 34:311–314, 1969.
- [76] C. Arsenis and O. Touster. Purification and properties of an acid nucleotidase from rat liver lysosomes. *J. Biol. Chem.*, 243:5702–5708, 1968.
- [77] D. Arthan, P. Kittakoop, A. Esen, and J. Svasti. Furostanol glycoside 26-*O*- $\beta$ -glucosidase from the leaves of *Solanum torvum*. *Phytochemistry*, 67:27–33, 2006.
- [78] G. Arthur, L. Page, T. Mock, and P.C. Choy. The catabolism of plasmenylcholine in the guinea pig heart. *Biochem. J.*, 236:475–480, 1986.
- [79] S. Arvidson. Studies on extracellular proteolytic enzymes from *Staphylococcus aureus*. II. Isolation and characterization of an EDTA-sensitive protease. *Biochim. Biophys. Acta*, 302:149–157, 1973.
- [80] T. Asakura, H. Watanabe, K. Abe, and S. Arai. Rice aspartic proteinase, oryzasin, expressed during seed ripening and germination, has a gene organization distinct from those of animal and microbial aspartic proteinases. *Eur. J. Biochem.*, 232:77–83, 1995.

- [81] S. Asamizu, J. Yang, K.H. Almabruk, and T. Mahmud. Pseudoglycosyltransferase catalyzes nonglycosidic C-N coupling in validamycin A biosynthesis. *J. Am. Chem. Soc.*, 133:12124–12135, 2011.
- [82] Y. Asano, Y. Kato, A. Yamada, and K. Kondo. Structural similarity of D-aminopeptidase to carboxypeptidase DD and  $\beta$ -lactamases. *Biochemistry*, 31:2316–2328, 1992.
- [83] Y. Asano, A. Nakazawa, Y. Kato, and K. Kondo. Properties of a novel D-stereospecific aminopeptidase from *Ochrobactrum anthropi*. *J. Biol. Chem.*, 264:14233–14239, 1989.
- [84] B. Asfaw, D. Schindler, J. Ledvinova, B. Cerny, F. Smid, and E. Conzelmann. Degradation of blood group A glycolipid A-6-2 by normal and mutant human skin fibroblasts. *J. Lipid Res.*, 39:1768–1780, 1998.
- [85] H. Ashida, R. Maki, H. Ozawa, Y. Tani, M. Kiyohara, M. Fujita, A. Imamura, H. Ishida, M. Kiso, and K. Yamamoto. Characterization of two different endo- $\alpha$ -N-acetylgalactosaminidases from probiotic and pathogenic enterobacteria, *Bifidobacterium longum* and *Clostridium perfringens*. *Glycobiology*, 18:727–734, 2008.
- [86] H. Ashida, Y. Saito, C. Kojima, K. Kobayashi, N. Ogasawara, and A. Yokota. A functional link between RuBisCO-like protein of *Bacillus* and photosynthetic RuBisCO. *Science*, 302:286–290, 2003.
- [87] H. Ashida, H. Tamaki, T. Fujimoto, K. Yamamoto, and H. Kumagai. Molecular cloning of cDNA encoding  $\alpha$ -N-acetylgalactosaminidase from *Acremonium* sp. and its expression in yeast. *Arch. Biochem. Biophys.*, 384:305–310, 2000.
- [88] H. Ashida, K. Yamamoto, T. Murata, T. Usui, and H. Kumagai. Characterization of endo- $\alpha$ -N-acetylgalactosaminidase from *Bacillus* sp. and syntheses of neo-oligosaccharides using its transglycosylation activity. *Arch. Biochem. Biophys.*, 373:394–400, 2000.
- [89] N.G. Asp, A. Dahlqvist, and O. Koldovský. Human small-intestinal  $\beta$ -galactosidases. Separation and characterization of one lactase and one hetero  $\beta$ -galactosidase. *Biochem. J.*, 114:351–359, 1969.
- [90] M.T. Assakura, A.P. Reichl, and F.R. Mandelbaum. Comparison of immunological, biochemical and biophysical properties of three hemorrhagic factors isolated from the venom of *Bothrops jararaca* (jararaca). *Toxicon*, 24:943–946, 1986.
- [91] L. Astrachan. The bond hydrolyzed by cardiolipin-specific phospholipase D. *Biochim. Biophys. Acta*, 296:79–88, 1973.
- [92] V.I. Athanasopoulos, K. Niranjana, and R.A. Rastall. The production, purification and characterisation of two novel  $\alpha$ -D-mannosidases from *Aspergillus phoenicis*. *Carbohydr. Res.*, 340:609–617, 2005.
- [93] J. Attias and J.L. Bonnet. A specific alkaline *p*-nitrophenylphosphatase activity from baker's yeast. *Biochim. Biophys. Acta*, 268:422–430, 1972.
- [94] J. Attias and H. Durand. Further characterization of a specific *p*-nitrophenylphosphatase from baker's yeast. *Biochim. Biophys. Acta*, 321:561–568, 1973.
- [95] P.V. Attwood, K. Ludwig, K. Bergander, P.G. Besant, A. Adina-Zada, J. Krieglstein, and S. Klumpp. Chemical phosphorylation of histidine-containing peptides based on the sequence of histone H4 and their dephosphorylation by protein histidine phosphatase. *Biochim. Biophys. Acta*, 1804:199–205, 2010.
- [96] K.G. Au, S. Clark, J.H. Miller, and P. Modrich. *Escherichia coli* *mutY* gene encodes an adenine glycosylase active on G-A mispairs. *Proc. Natl. Acad. Sci. USA*, 86:8877–8881, 1989.
- [97] R.C. Augusteyn, J. de Jersey, E.C. Webb, and B. Zerner. On the homology of the active-site peptides of liver carboxylesterases. *Biochim. Biophys. Acta*, 171:128–137, 1969.
- [98] K.-B. Augustinsson. Cholinesterases. A study in comparative enzymology. *Acta Physiol. Scand.*, 15, Suppl. 2, 1948.
- [99] K.-B. Augustinsson and G. Heimburger. Enzymatic hydrolysis of organophosphorus compounds. I. Occurrence of enzymes hydrolysing dimethyl-amido-ethoxy-phosphoryl cyanide (Tabun). *Acta Chem. Scand.*, 8:753–761, 1954.
- [100] K.-B. Augustinsson and G. Heimburger. Enzymatic hydrolysis of organophosphorus compounds. II. Analysis of reaction products in experiments with Tabun and some properties of blood plasma tabunase. *Acta Chem. Scand.*, 8:762–767, 1954.
- [101] K.-B. Augustinsson and G. Heimburger. Enzymatic hydrolysis of organophosphorus compounds. IV. Specificity studies. *Acta Chem. Scand.*, 8:1533–1541, 1954.

- [102] K.-B. Augustinsson and B. Olsson. Esterases in the milk and blood plasma of swine. 1. Substrate specificity and electrophoresis studies. *Biochem. J.*, 71:477–484, 1959.
- [103] M. Aurrand-Lions, F. Galland, H. Bazin, V.M. Zakharyev, B.A. Imhof, and P. Naquet. Vanin-1, a novel GPI-linked perivascular molecule involved in thymus homing. *Immunity*, 5:391–405, 1996.
- [104] G. Avigad and S. Bauer. Fructan hydrolases. *Methods Enzymol.*, 8:621–628, 1966.
- [105] G. Avigad and R. Zelikson. Cleavage of fructans to levanbiose by a specific hydrolase. *Bull. Res. Counc. Isr.*, 11:253–257, 1963.
- [106] A.C. Awadé, P. Cleuziat, T. Gonzalès, and J. Robert-Baudouy. Pyrrolidone carboxyl peptidase (Pcp): an enzyme that removes pyroglutamic acid (pGlu) from pGlu-peptides and pGlu-proteins. *Proteins: Struct. Funct. Genet.*, 20:34–51, 1994.
- [107] P. Azadi, M.A. O’Neill, C. Bergmann, A.G. Darvill, and P. Albersheim. The backbone of the pectic polysaccharide rhamnogalacturonan I is cleaved by an endohydrolase and an endolyase. *Glycobiology*, 5:783–789, 1995.
- [108] T. Azuma, G. Pals, T.K. Mohandas, J.M. Couvreur, and R.T. Taggart. Human gastric cathepsin E. Predicted sequence, localization to chromosome 1, and sequence homology with other aspartic proteinases. *J. Biol. Chem.*, 264:16748–16753, 1989.
- [109] T. Baba and O. Schneewind. Target cell specificity of a bacteriocin molecule: A C-terminal signal directs lysostaphin to the cell wall of *Staphylococcus aureus*. *EMBO J.*, 15:4789–4797, 1996.
- [110] K.J. Babinski, S.J. Kanjilal, and C.R. Raetz. Accumulation of the lipid A precursor UDP-2,3-diacylglucosamine in an *Escherichia coli* mutant lacking the *lpxH* gene. *J. Biol. Chem.*, 277:25947–25956, 2002.
- [111] K.J. Babinski, A.A. Ribeiro, and C.R. Raetz. The *Escherichia coli* gene encoding the UDP-2,3-diacylglucosamine pyrophosphatase of lipid A biosynthesis. *J. Biol. Chem.*, 277:25937–25946, 2002.
- [112] W. Babst, B. Wendland, E.J. Estepa, and S.D. Emr. The Vps4p AAA ATPase regulates membrane association of a Vps protein complex required for normal endosome function. *EMBO J.*, 17:2982–2983, 1998.
- [113] J. Bach, F. Eisenberg, M. Cantz, and E.C. Neufeld. The defect in the Hunter syndrome: deficiency of sulfiduronate sulfatase. *Proc. Natl. Acad. Sci. USA*, 70:2134–2138, 1973.
- [114] S.J. Bach and J.D. Killip. Studies on the purification and the kinetic properties of arginase from beef, sheep and horse liver. *Biochim. Biophys. Acta*, 47:336–343, 1961.
- [115] M. Bachmann, F. Trautmann, R. Messer, R.K. Zahn, K.H. Meyer zum Büschenfelde, and W.E.G. Müller. Association of a polyuridylylate-specific endoribonuclease with small nuclear ribonucleo-proteins which had been isolated by affinity chromatography using antibodies from a patient with systemic lupus erythematosus. *Eur. J. Biochem.*, 136:447–451, 1983.
- [116] W.W. Bachovchin, A.G. Plaut, G.R. Flentke, M. Lynch, and C.A. Kettner. Inhibition of IgA1 proteinases from *Neisseria gonorrhoeae* and *Hemophilus influenzae* by peptide prolyl boronic acids. *J. Biol. Chem.*, 265:3738–3743, 1990.
- [117] S. Bader, A. Kortholt, H. Snippe, and P.J. Van Haastert. DdPDE4, a novel cAMP-specific phosphodiesterase at the surface of *Dictyostelium* cells. *J. Biol. Chem.*, 281:20018–20026, 2006.
- [118] O.P. Bahl. Glycosidases of *Aspergillus niger*. II. Purification and general properties of 1,2- $\alpha$ -L-fucosidase. *J. Biol. Chem.*, 245:299–304, 1970.
- [119] G.S. Bailey. Rat pancreas kallikrein. *Methods Enzymol.*, 163:115–128, 1989.
- [120] K. Bailey and E.C. Webb. Purification and properties of yeast pyrophosphatase. *Biochem. J.*, 38:394–398, 1944.
- [121] R.W. Bailey and R.T.J. Clarke. A bacterial dextranase. *Biochem. J.*, 72:49–54, 1959.
- [122] R. Bailey-Wood, K.S. Dodgson, and F.A. Rose. A rat liver sulphohydrolase enzyme acting on adenylyl sulphate. *Biochem. J.*, 112:257–258, 1969.

- [123] M. Bajaj and H. Moriyama. Purification, crystallization and preliminary crystallographic analysis of deoxyuridine triphosphate nucleotidohydrolase from *Arabidopsis thaliana*. *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.*, 63:409–411, 2007.
- [124] A.S. Baker, M.J. Ciocci, W.W. Metcalf, J. Kim, P.C. Babbitt, B.L. Wanner, B.M. Martin, and D. Dunaway-Mariano. Insights into the mechanism of catalysis by the P-C bond-cleaving enzyme phosphonoacetaldehyde hydrolase derived from gene sequence analysis and mutagenesis. *Biochemistry*, 37:9305–9315, 1998.
- [125] E.N. Baker, M.J. Boland, P.C. Calder, and M.J. Hardman. The specificity of actinidin and its relationship to the structure of the enzyme. *Biochim. Biophys. Acta*, 616:30–34, 1980.
- [126] E.N. Baker and J. Drenth. The thiol proteases: structure and mechanism. In F.A. Jornak and A. McPherson, editors, *Biological Macromolecules and Assemblies: Active Sites of Enzymes*, volume 3, pages 314–368. John Wiley and Sons, New York, 1987.
- [127] K. Baksi and A.N. Radhakrishnan. Purification and properties of prolidase (imidodipeptidase) from monkey small intestine. *Indian J. Biochem. Biophys.*, 11:7–11, 1974.
- [128] A.S. Balasubramanian and B.K. Bachhawat. Enzymic degradation of active sulphate. *Biochim. Biophys. Acta*, 59:389–397, 1962.
- [129] A.K. Balls, M.K. Walden, and R.R. Thompson. A crystalline  $\beta$ -amylase from sweet potatoes. *J. Biol. Chem.*, 173:9–19, 1948.
- [130] S. Balotra, A.C. Warden, J. Newman, L.J. Briggs, C. Scott, and T.S. Peat. X-ray structure and mutagenesis studies of the *N*-isopropylammelide isopropylaminohydrolase, AtzC. *PLoS One*, 10:e0137700–e0137700, 2015.
- [131] R.M. Bålöw, U. Ragnarsson, and Ö. Zetterqvist. Tripeptidyl aminopeptidase in the extralysosomal fraction of rat liver. *J. Biol. Chem.*, 258:11622–11628, 1983.
- [132] R.M. Bålöw, B. Tomkinson, U. Ragnarsson, and Ö. Zetterqvist. Purification, substrate specificity, and classification of tripeptidyl peptidase II. *J. Biol. Chem.*, 261:2409–2417, 1986.
- [133] K.J. Bame. Heparanases: endoglycosidases that degrade heparan sulfate proteoglycans. *Glycobiology*, 11:91R–98R, 2001.
- [134] A. Banbula, P. Mak, M. Bugno, J. Silberring, A. Dubin, D. Nelson, J. Travis, and J. Potempa. Prolyl tripeptidyl peptidase from *Porphyromonas gingivalis*. A novel enzyme with possible pathological implications for the development of periodontitis. *J. Biol. Chem.*, 274:9246–9252, 1999.
- [135] M.J. Banda and Z. Werb. Mouse macrophage elastase. Purification and characterization as a metalloproteinase. *Biochem. J.*, 193:589–605, 1981.
- [136] E.N. Baramova, J.D. Shannon, J.B. Bjarnason, and J.W. Fox. Identification of the cleavage sites by a hemorrhagic metalloproteinase in type IV collagen. *Matrix*, 10:91–97, 1990.
- [137] T. Barbeyron, G. Michel, P. Potin, B. Henrissat, and B. Kloareg.  $\iota$ -Carrageenases constitute a novel family of glycoside hydrolases, unrelated to that of  $\kappa$ -carrageenases. *J. Biol. Chem.*, 275:35499–35505, 2000.
- [138] F. Barbirato, J.C. Verdoes, J.A.M. de Bont, and M.J. van der Werf. The *Rhodococcus erythropolis* DCL14 limonene-1,2-epoxide hydrolase gene encodes an enzyme belonging to a novel class of epoxide hydrolases. *FEBS Lett.*, 438:293–296, 1998.
- [139] S.L. Bardy and K.F. Jarrell. FlaK of the archaeon *Methanococcus maripaludis* possesses preflagellin peptidase activity. *FEMS Microbiol. Lett.*, 208:53–59, 2002.
- [140] H. Barelli, J.-P. Vincent, and F. Checler. Peripheral inactivation of neurotensin. Isolation and characterization of a metalloproteinase from rat ileum. *Eur. J. Biochem.*, 175:481–489, 1988.
- [141] D.L. Barker and W.P. Jencks. Pig liver esterase. Physical properties. *Biochemistry*, 8:3879–3889, 1969.
- [142] S.A. Barker, P.J. Somers, and M. Stacey. Arrangement of the L-rhamnose units in *Diplococcus pneumoniae* type II polysaccharide. *Carbohydr. Res.*, 1:106–115, 1965.

- [143] V. Barkholt. Amino acid sequence of endothiapepsin. Complete primary structure of the aspartic protease from *Endothia parasitica*. *Eur. J. Biochem.*, 167:327–338, 1987.
- [144] L. Barleben, X. Ma, J. Koepke, G. Peng, H. Michel, and J. Stöckigt. Expression, purification, crystallization and preliminary X-ray analysis of strictosidine glucosidase, an enzyme initiating biosynthetic pathways to a unique diversity of indole alkaloid skeletons. *Biochim. Biophys. Acta*, 1747:89–92, 2005.
- [145] C. Barnes and H.C. Smith. Apolipoprotein B mRNA editing *in vitro* is a zinc-dependent process. *Biochem. Biophys. Res. Commun.*, 197:1410–1414, 1993.
- [146] E.M. Barnes, Wakil Jr., and S.J. Studies on the mechanism of fatty acid synthesis. XIX. Preparation and general properties of palmityl thioesterase. *J. Biol. Chem.*, 243:2955–2962, 1968.
- [147] K. Barnes, J. Ingram, and A.J. Kenny. Proteins of the kidney microvillar membrane. Structural and immunochemical properties of rat endopeptidase-2 and its immunohistochemical localization in tissues of rat and mouse. *Biochem. J.*, 264:335–346, 1989.
- [148] L.D. Barnes and C.A. Culver. Isolation and characterization of diadenosine 5',5'''-P<sup>1</sup>,P<sup>4</sup>-tetraphosphate pyrophosphohydrolase from *Physarum polycephalum*. *Biochemistry*, 21:6123–6128, 1982.
- [149] Y. Barnholz, A. Roitman, and S. Gatt. Enzymatic hydrolysis of sphingolipids. II. Hydrolysis of sphingomyelin by an enzyme from rat brain. *J. Biol. Chem.*, 241:3731–3737, 1966.
- [150] B.K. Barr, Y.L. Hsieh, B. Ganem, and D.B. Wilson. Identification of two functionally different classes of exocellulases. *Biochemistry*, 35:586–592, 1996.
- [151] D.R. Barras, A.E. Moore, and B.A. Stone. Enzyme-substrate relations among  $\beta$ -glucan hydrolases. *Adv. Chem. Ser.*, 95:105–138, 1969.
- [152] D.R. Barras and B.A. Stone.  $\beta$ -1,3-Glucan hydrolases from *Euglena gracilis*. I. The nature of the hydrolases. *Biochim. Biophys. Acta*, 191:329–341, 1969.
- [153] D.R. Barras and B.A. Stone.  $\beta$ -1,3-Glucan hydrolases from *Euglena gracilis*. II. Purification and properties of the  $\beta$ -1,3-glucan exo-hydrolase. *Biochim. Biophys. Acta*, 191:342–353, 1969.
- [154] A.J. Barrett. Cathepsin D and other carboxyl proteinases. In A.J. Barrett, editor, *Proteinases in Mammalian Cells and Tissues*, pages 209–248. North-Holland Publishing Co., Amsterdam, 1977.
- [155] A.J. Barrett. Cathepsin G. *Methods Enzymol.*, 80:561–565, 1981.
- [156] A.J. Barrett. Leukocyte elastase. *Methods Enzymol.*, 80:581–588, 1981.
- [157] A.J. Barrett and M.A. Brown. Chicken liver Pz-peptidase, a thiol-dependent metallo-endopeptidase. *Biochem. J.*, 271:701–706, 1990.
- [158] A.J. Barrett, D.J. Buttle, and R.W. Mason. Lysosomal cysteine proteinases. *ISI Atlas of Science. Biochemistry*, 1:256–260, 1988.
- [159] A.J. Barrett and H. Kirschke. Cathepsin B, cathepsin H and cathepsin L. *Methods Enzymol.*, 80:535–561, 1981.
- [160] D. Barrett and B.F. Edwards. Hatching enzyme of the sea urchin *Strongylocentrotus purpuratus*. *Methods Enzymol.*, 45:354–373, 1976.
- [161] T.E. Barrett, O.D. Scharer, R. Savva, T. Brown, J. Jiricny, G.L. Verdine, and L.H. Pearl. Crystal structure of a thwarted mismatch glycosylase DNA repair complex. *EMBO J.*, 18:6599–6609, 1999.
- [162] I.H. Barrette-Ng, K.K. Ng, B.L. Mark, D. Van Aken, M.M. Cherney, C. Garen, Y. Kolodenco, A.E. Gorbalenya, E.J. Snijder, and M.N. James. Structure of arterivirus nsp4. The smallest chymotrypsin-like proteinase with an  $\alpha/\beta$  C-terminal extension and alternate conformations of the oxyanion hole. *J. Biol. Chem.*, 277:39960–39966, 2002.
- [163] L. Barrientos, J.J. Scott, and P.P. Murthy. Specificity of hydrolysis of phytic acid by alkaline phytase from lily pollen. *Plant Physiol.*, 106:1489–1495, 1994.

- [164] M.L. Barta, K. Thomas, H. Yuan, S. Lovell, K.P. Battaile, V.L. Schramm, and P.S. Hefty. Structural and biochemical characterization of *Chlamydia trachomatis* hypothetical protein CT263 supports that menaquinone synthesis occurs through the futasolose pathway. *J. Biol. Chem.*, 289:32214–32229, 2014.
- [165] B.A. Bartholomew and A.L. Perry. The properties of synovial fluid  $\beta$ -mannosidase activity. *Biochim. Biophys. Acta*, 315:123–127, 1973.
- [166] A.T.M. Bartlett and P.J. White. Species of *Bacillus* that make a vegetative peptidoglycan containing lysine lack diaminopimelate epimerase but have diaminopimelate dehydrogenase. *J. Gen. Microbiol.*, 131:2145–2152, 1985.
- [167] A. Basak, M. Chrétien, and N.G. Seidah. A rapid fluorometric assay for the proteolytic activity of SKI-1/S1P based on the surface glycoprotein of the hemorrhagic fever Lassa virus. *FEBS Lett.*, 514:333–339, 2002.
- [168] M.H. Baslow and J.F. Lenney.  $\alpha$ -N-Acetyl-L-histidine amidohydrolase activity from the brain of the skipjack tuna *Katsuwonus pelamis*. *Can. J. Biochem.*, 45:337–340, 1967.
- [169] R. Basner, H. Kresse, and K. von Figura. N-Acetylglucosamine-6-sulfate sulfatase from human urine. *J. Biol. Chem.*, 254:1151–1158, 1979.
- [170] B.L. Bassler, C. Yu, Y.C. Lee, and S. Roseman. Chitin utilization by marine bacteria. Degradation and catabolism of chitin oligosaccharides by *Vibrio furnissii*. *J. Biol. Chem.*, 266:24276–24286, 1991.
- [171] M. Baudy, S. Foundling, M. Pavlik, T. Blundell, and V. Kostka. Protein chemical characterization of *Mucor pusillus* aspartic proteinase. Amino acid sequence homology with the other aspartic proteinases, disulfide bond arrangement and site of carbohydrate attachment. *FEBS Lett.*, 235:271–274, 1988.
- [172] C.-A. Bauer. Active centers of  $\alpha$ -chymotrypsin and of *Streptomyces griseus* proteases 1 and 3. *Eur. J. Biochem.*, 105:565–570, 1980.
- [173] K. Bauer. Cytosol non-specific dipeptidase. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 1520–1522. Academic Press, London, 1998.
- [174] K. Bauer and P. Nowak. Characterization of a thyrolyberin-degrading serum enzyme catalyzing the hydrolysis of thyrolyberin at the pyroglutamyl-histidine bond. *Eur. J. Biochem.*, 99:239–246, 1979.
- [175] D.J. Baughman. Thrombin assay. *Methods Enzymol.*, 19:145–157, 1970.
- [176] N. Baumer, A. Maurer, J. Krieglstein, and S. Klumpp. Expression of protein histidine phosphatase in *Escherichia coli*, purification, and determination of enzyme activity. *Methods Mol Biol*, 365:247–260, 2007.
- [177] M.E. Bayliss, S.H. Wilkes, and J.M. Prescott. *Aeromonas* neutral protease: specificity toward extended substrates. *Arch. Biochem. Biophys.*, 204:214–219, 1980.
- [178] R.C. Bayly, P.J. Chapman, S. Dagley, and D. Di Berardino. Purification and some properties of maleylpyruvate hydrolase and fumarylpyruvate hydrolase from *Pseudomonas alcaligenes*. *J. Bacteriol.*, 143:70–77, 1980.
- [179] J.F. Bazan and R.J. Fletterick. Viral cysteine proteases are homologous to the trypsin-like family of serine proteases: structural and functional implications. *Proc. Natl Acad. Sci. USA*, 85:7872–7876, 1988.
- [180] A. Becker and J. Hurwitz. The enzymatic cleavage of phosphate termini from polynucleotides. *J. Biol. Chem.*, 242:936–950, 1967.
- [181] A. Becker, I. Schlichting, W. Kabsch, D. Groche, S. Schultz, and A.F. Wagner. Iron center, substrate recognition, and mechanism of peptide deformylase. *Nat. Struct. Biol.*, 5:1053–1058, 1998.
- [182] A. Becker, I. Schlichting, W. Kabsch, S. Schultz, and A.F.V. Wagner. Structure of peptide deformylase and identification of the substrate binding site. *J. Biol. Chem.*, 273:11413–11416, 1998.
- [183] A.B. Becker and R.A. Roth. An unusual active site identified in a family of zinc metalloendopeptidases. *Proc. Natl. Acad. Sci. USA*, 89:3835–3839, 1992.
- [184] K.D. Beebe, J.N. Shin, J. Peng, C. Chaudhury, J. Khera, and D.H. Pei. Substrate recognition through a PDZ domain in tail-specific protease. *Biochemistry*, 39:3149–3155, 2000.



- [185] G.L. Beilhartz, M. Wendeler, N. Baichoo, J. Rausch, S. Le Grice, and M. Gotte. HIV-1 reverse transcriptase can simultaneously engage its DNA/RNA substrate at both DNA polymerase and RNase H active sites: implications for RNase H inhibition. *J. Mol. Biol.*, 388:462–474, 2009.
- [186] M. Belard, R. Cacan, and A. Verbert. Characterization of an oligosaccharide-pyrophosphodolichol pyrophosphatase activity in yeast. *Biochem. J.*, 255:235–242, 1988.
- [187] N. Belhacène, B. Mari, B. Rossi, and P. Auberger. Characterization and purification of T lymphocyte aminopeptidase B: a putative marker of T cell activation. *Eur. J. Immunol.*, 23:1948–1955, 1993.
- [188] G. Bellucci, C. Chiappe, and G. Ingrosso. Kinetics and stereochemistry of the microsomal epoxide hydrolase-catalyzed hydrolysis of *cis*-stilbene oxides. *Chirality*, 6:577–582, 1994.
- [189] A. Ben-Bassat, K. Bauer, S.-Y. Chang, K. Myambo, A. Boosman, and S. Chang. Processing of the initiation methionine from proteins: properties of *Escherichia coli* methionine aminopeptidase and its gene structure. *J. Bacteriol.*, 169:751–757, 1987.
- [190] D. Ben-Meir, A. Spungin, R. Ashkenazi, and S. Blumberg. Specificity of *Streptomyces griseus* aminopeptidase and modulation of activity by divalent metal ion binding and substitution. *Eur. J. Biochem.*, 212:107–112, 1993.
- [191] J. Benach, W.C. Edstrom, I. Lee, K. Das, B. Cooper, R. Xiao, J. Liu, B. Rost, T.B. Acton, G.T. Montelione, and J.F. Hunt. The 2.35 Å structure of the TenA homolog from *Pyrococcus furiosus* supports an enzymatic function in thiamine metabolism. *Acta Crystallogr. D Biol. Crystallogr.*, 61:589–598, 2005.
- [192] H. Bender and K. Wallenfels. Pullulanase (an amylopectin and glycogen debranching enzyme) from *Aerobacter aerogenes*. *Methods Enzymol.*, 8:555–559, 1966.
- [193] R.K. Berge and M. Farstad. Long-chain fatty acyl-CoA hydrolase from rat liver mitochondria. *Methods Enzymol.*, 71:234–242, 1981.
- [194] P. Berger, S. Bonneick, S. Willi, M. Wymann, and U. Suter. Loss of phosphatase activity in myotubularin-related protein 2 is associated with Charcot-Marie-Tooth disease type 4B1. *Hum. Mol. Genet.*, 11:1569–1579, 2002.
- [195] L.E.R. Berghem and L.G. Pettersson. The mechanism of enzymatic cellulose degradation. Purification of a cellulolytic enzyme from *Trichoderma viride* active on highly ordered cellulose. *Eur. J. Biochem.*, 37:21–30, 1973.
- [196] F. Bergmann and S. Rimon. Fractionation of C-esterase from the hog's kidney extract. *Biochem. J.*, 77:209–214, 1960.
- [197] F. Bergmann, S. Rimon, and R. Segal. Effect of pH on the activity of eel esterase towards different substrates. *Biochem. J.*, 68:493–499, 1958.
- [198] R. Bernard, M. El Ghachi, D. Mengin-Lecreulx, M. Chippaux, and F. Denizot. BcrC from *Bacillus subtilis* acts as an undecaprenyl pyrophosphate phosphatase in bacitracin resistance. *J. Biol. Chem.*, 280:28852–28857, 2005.
- [199] A. Bernardi and G. Bernardi. Spleen acid nuclease. In P.D. Boyer, editor, *The Enzymes*, volume 4, pages 329–336. Academic Press, New York, 3rd edition, 1971.
- [200] A. Bernardi, J. Maat, A. de Waard, and G. Bernardi. Preparation and specificity of endonuclease IV induced by bacteriophage T<sub>4</sub>. *Eur. J. Biochem.*, 66:175–179, 1976.
- [201] G. Bernardi. Spleen acid deoxyribonuclease. In G.L. Cantoni and D.R. Davies, editors, *Procedures in Nucleic Acid Research*, pages 102–121. Procedures in Nucleic Acid Research, New York, 1966.
- [202] D.A. Bernstein, M.C. Zittel, and J.L. Keck. High-resolution structure of the *E. coli* RecQ helicase catalytic core. *EMBO J.*, 22:4910–4921, 2003.
- [203] P.S. Bernstein, W.C. Law, and R.R. Rando. Isomerization of *all-trans*-retinoids to 11-*cis*-retinoids in vitro. *Proc. Natl. Acad. Sci. USA*, 84:1849–1853, 1987.
- [204] M.J. Berridge, R.M.C. Dawson, C.P. Downes, J.P. Heslop, and R.F. Irvine. Changes in the levels of inositol phosphates after agonist-dependent hydrolysis of membrane phosphoinositides. *Biochem. J.*, 212:473–482, 1983.
- [205] L.E. Bertani, A. Häggmark, and P. Reichard. Enzymatic synthesis of deoxyribonucleotides. II. Formation and interconversion of deoxyuridine phosphates. *J. Biol. Chem.*, 238:3407–3413, 1963.

- [206] C.L. Berthold, H. Wang, S. Nordlund, and M. Hogbom. Mechanism of ADP-ribosylation removal revealed by the structure and ligand complexes of the dimanganese mono-ADP-ribosylhydrolase DraG. *Proc. Natl. Acad. Sci. USA*, 106:14247–14252, 2009.
- [207] J. Bertram and K. Krisch. Hydrolysis of vitamin A acetate by unspecific carboxylesterases from liver and kidney. *Eur. J. Biochem.*, 11:122–126, 1969.
- [208] R. Bertrand, J. Derancourt, and R. Kassab. Selective cleavage at lysine of the 50 kDa-20 kDa connector loop segment of skeletal myosin S-1 by endoproteinase Arg-C. *FEBS Lett.*, 246:171–176, 1989.
- [209] C. Betzel, A.V. Teplyakov, E.H. Harutyunyan, W. Saenger, and K.S. Wilson. Thermitase and proteinase K: a comparison of the refined three-dimensional structures of the native enzymes. *Protein Engineering*, 3:161–172, 1990.
- [210] R.A. Bever and B.H. Iglewski. Molecular characterization and nucleotide sequence of the *Pseudomonas aeruginosa* elastase structural gene. *J. Bacteriol.*, 170:4309–4314, 1988.
- [211] R.J. Beynon, J.D. Shannon, and J.S. Bond. Purification and characterization of a metallo-endoproteinase from mouse kidney. *Biochem. J.*, 199:591–598, 1981.
- [212] A. Bhumiratana, R.L. Anderson, and R.N. Costilow. Trehalose metabolism by *Bacillus popilliae*. *J. Bacteriol.*, 119:484–493, 1974.
- [213] H. Bierne, C. Garandeau, M.G. Pucciarelli, C. Sabet, S. Newton, F. Garcia del Portillo, P. Cossart, and A. Charbit. Sortase B, a new class of sortase in *Listeria monocytogenes*. *J. Bacteriol.*, 186:1972–1982, 2004.
- [214] J.G. Bieth, S. Dirrig, M.-L. Jung, C. Boudier, E. Papamichael, C. Sakarellos, and J.-L. Dimicoli. Investigation of the active center of rat pancreatic elastase. *Biochim. Biophys. Acta*, 994:64–74, 1989.
- [215] E.K. Bikoff and M.L. Gefter. In vitro synthesis of transfer RNA. I. Purification of required components. *J. Biol. Chem.*, 250:6240–6247, 1975.
- [216] E.K. Bikoff, B.F. La Rue, and M.L. Gefter. In vitro synthesis of transfer RNA. II. Identification of required enzymatic activities. *J. Biol. Chem.*, 250:6248–6255, 1975.
- [217] H. Birkedal-Hansen. Catabolism and turnover of collagens: collagenases. *Methods Enzymol.*, 144:140–171, 1987.
- [218] S.M. Birnbaum. Aminoacylase. Amino acid aminoacylases I and II from hog kidney. *Methods Enzymol.*, 2:115–119, 1955.
- [219] S.M. Birnbaum, L. Levintow, R.B. Kingsley, and J.P. Greenstein. Specificity of amino acid acylases. *J. Biol. Chem.*, 194:455–470, 1952.
- [220] J.B. Bjarnason and J.W. Fox. Proteolytic specificity and cobalt exchange of hemorrhagic toxin e, a zinc protease isolated from the venom of the western diamondback rattlesnake (*Crotalus atrox*). *Biochemistry*, 22:3770–3778, 1983.
- [221] J.B. Bjarnason and J.W. Fox. Characterization of two hemorrhagic zinc proteinases, toxin c and toxin d, from western diamondback rattlesnake (*Crotalus atrox*) venom. *Biochim. Biophys. Acta*, 911:356–363, 1987.
- [222] J.B. Bjarnason and J.W. Fox. Hemorrhagic toxins from snake venoms. *J. Toxicol. Toxin Rev.*, 7:121–209, 1989.
- [223] J.B. Bjarnason, D. Hamilton, and J.W. Fox. Studies on the mechanism of hemorrhage production by five proteolytic hemorrhagic toxins from *Crotalus atrox* venom. *Biol. Chem. Hoppe-Seyler*, 369:121–129, 1988.
- [224] J.B. Bjarnason and A.T. Tu. Hemorrhagic toxins from western diamondback rattlesnake (*Crotalus atrox*) venom: isolation and characterization of five toxins and the role of zinc in hemorrhagic toxin e. *Biochemistry*, 17:3395–3404, 1978.
- [225] K. Bjerregaard-Andersen, T. Sommer, J.K. Jensen, B. Jochimsen, M. Etzerodt, and J.P. Morth. A proton wire and water channel revealed in the crystal structure of isatin hydrolase. *J. Biol. Chem.*, 289:21351–21359, 2014.
- [226] M. Blaber, P.J. Isackson, J.C. Marsters, Burnier Jr., Bradshaw J.P., and R.A. Substrate specificities of growth factor associated kallikreins of the mouse submandibular gland. *Biochemistry*, 28:7813–7819, 1988.
- [227] M.T. Black. Evidence that the catalytic activity of prokaryote leader peptidase depends upon the operation of a serine-lysine catalytic dyad. *J. Bacteriol.*, 175:4957–4961, 1993.

- [228] R.A. Black. Tumor necrosis factor- $\alpha$  converting enzyme. *Int. J. Biochem. Cell Biol.*, 34:1–5, 2002.
- [229] S. Black and N.G. Wright. Enzymatic formation of glyceryl and phosphoglycerol methylthiol esters. *J. Biol. Chem.*, 221:171–180, 1956.
- [230] T.D. Black, B.S. Briggs, R. Evans, W.L. Muth, S. Vangala, , and M.J. o-Phthalyl amidase in the synthesis of Loracarbef: process development using this novel biocatalyst. *Biotechnol. Lett.*, 18:875–880, 1996.
- [231] W.J. Black, F.D. Quinn, and L.S. Tompkins. *Legionella pneumophila* zinc metalloprotease is structurally and functionally homologous to *Pseudomonas aeruginosa* elastase. *J. Bacteriol.*, 172:2608–2613, 1990.
- [232] C.C.F. Blade, L.N. Johnson, G.A. Mair, A.C.T. North, D.C. Phillips, and V.R. Sarma. Crystallographic studies of the activity of hen egg-white lysozyme. *Proc. R. Soc. Lond. B: Biol. Sci.*, 167:378–388, 1967.
- [233] D.E. Blair, A.W. Schuttelkopf, J.I. MacRae, and D.M. van Aalten. Structure and metal-dependent mechanism of peptidoglycan deacetylase, a streptococcal virulence factor. *Proc. Natl. Acad. Sci. USA*, 102:15429–15434, 2005.
- [234] C.C.F. Blake, G.A. Mair, A.C.T. North, D.C. Phillips, and V.R. Sarma. On the conformation of the hen egg-white lysozyme molecule. *Proc. R. Soc. Lond. B: Biol. Sci.*, 167:365–377, 1967.
- [235] J.A. Blakely and S.L. MacKenzie. Purification and properties of a  $\beta$ -hexosidase from *Sporobolomyces singularis*. *Can. J. Biochem.*, 47:1021–1025, 1969.
- [236] R.W. Blakesley, J.B. Dodgson, I.F. Nes, and R.D. Wells. Duplex regions in single-stranded phiX174 DNA are cleaved by a restriction endonuclease from *Haemophilus aegyptius*. *J. Biol. Chem.*, 252:7300–7306, 1977.
- [237] H. Blanchard, M. Donepudi, M. Tschopp, L. Kodandapani, J.C. Wu, and M.G. Grütter. Caspase-8 specificity probed at subsite S(4): crystal structure of the caspase-8-Z-DEVD-cho complex. *J. Mol. Biol.*, 302:9–16, 2000.
- [238] W.S. Blaner, S.R. Das, P. Gouras, and M.T. Flood. Hydrolysis of 11-*cis*- and *all-trans*-retinyl palmitate by homogenates of human retinal epithelial cells. *J. Biol. Chem.*, 262:53–58, 1987.
- [239] W.S. Blaner, J.H. Prystowsky, J.E. Smith, and D.S. Goodman. Rat liver retinyl palmitate hydrolase activity. Relationship to cholesteryl oleate and triolein hydrolase activities. *Biochim. Biophys. Acta*, 794:419–427, 1984.
- [240] M.L. Blank, T.-C. Lee, V. Fitzgerald, and F. Snyder. A specific acetylhydrolase for 1-alkyl-2-acetyl-*sn*-glycero-3-phosphocholine (a hypotensive and platelet-activating lipid). *J. Biol. Chem.*, 256:175–178, 1981.
- [241] M.L. Blank, Z.L. Smith, E.A. Cress, , and F. Characterization of the enzymatic hydrolysis of acetate from alkylacetyl-glycerols in the de novo pathway of PAF biosynthesis. *Biochim. Biophys. Acta*, 1042:153–158, 1990.
- [242] J. Blankenship. Deacetylation of *N*<sup>8</sup>-acetylspermidine by subcellular fractions of rat tissue. *Arch. Biochem. Biophys.*, 189:20–27, 1978.
- [243] M. Blauch, F.C. Koch, and M.E. Hane. A study of xanthine oxidase of rat blood. *J. Biol. Chem.*, 130:471–486, 1939.
- [244] S. Blond-Elquindi, A.M. Fourie, J.F. Sambrook, and M.J. Gething. Peptide-dependent stimulation of the ATPase activity of the molecular chaperone BiP is the result of conversion of oligomers to active monomers. *J. Biol. Chem.*, 268:12730–12735, 1993.
- [245] D.M. Blow. Structure and mechanism of chymotrypsin. *Acc. Chem. Res.*, 9:145–152, 1976.
- [246] S. Blumberg and Z. Tauber. Inhibition of metalloendopeptidases by 2-mercaptoacetyl-dipeptides. *Eur. J. Biochem.*, 136:151–154, 1983.
- [247] A. Blume, D. Ghaderi, V. Liebich, S. Hinderlich, P. Donner, W. Reutter, and L. Lucka. UDP-*N*-acetylglucosamine 2-epimerase/*N*-acetylmannosamine kinase, functionally expressed in and purified from *Escherichia coli*, yeast, and insect cells. *Protein Expr. Purif.*, 35:387–396, 2004.
- [248] G. Boanca, A. Sand, T. Okada, H. Suzuki, H. Kumagai, K. Fukuyama, and J.J. Barycki. Autoprocessing of *Helicobacter pylori*  $\gamma$ -glutamyltranspeptidase leads to the formation of a threonine-threonine catalytic dyad. *J. Biol. Chem.*, 282:534–541, 2007.

- [249] K.M. Boatright, C. Deis, J.B. Denault, D.P. Sutherlin, and G.S. Salvesen. Activation of caspases-8 and -10 by FLIP<sub>1</sub>. *Biochem. J.*, 382:651–657, 2004.
- [250] K.M. Boatright, M. Renatus, F.L. Scott, S. Sperandio, H. Shin, I.M. Pedersen, J.E. Ricci, W.A. Edris, D.P. Sutherlin, D.R. Green, and G.S. Salvesen. A unified model for apical caspase activation. *Mol. Cell.*, 11:529–541, 2003.
- [251] W. Bode, E. Meyer, Powers Jr., and J.C. Human leukocyte and porcine pancreatic elastase: X-ray crystal structures, mechanism, substrate specificity, and mechanism-based inhibitors. *Biochemistry*, 28:1951–1963, 1989.
- [252] J.E. Bodwell and W.L. Meyer. Purification and characterization of carboxypeptidase A from rat skeletal muscle. *Biochemistry*, 20:2767–2777, 1981.
- [253] D.L. Boger, R.A. Fecik, J.E. Patterson, H. Miyauchi, M.P. Patricelli, and B.F. Cravatt. Fatty acid amide hydrolase substrate specificity. *Bioorg. Med. Chem. Lett.*, 10:2613–2616, 2000.
- [254] J.F. Bohnsack, K.W. Mollison, A.M. Buko, J.C. Ashworth, and H.R. Hill. Group B streptococci inactivate complement component C5a by enzymic cleavage at the C-terminus. *Biochem. J.*, 273:635–640, 1991.
- [255] S. Boiteux, T.R. O'Connor, and J. Laval. Formamidopyrimidine-DNA glycosylase of *Escherichia coli*: cloning and sequencing of the fpg structural gene and overproduction of the protein. *EMBO J.*, 6:3177–3183, 1987.
- [256] M.P. Boldin, T.M. Goncharov, Y.V. Goltsev, and D. Wallach. Involvement of MACH, a novel MORT1/FADD-interacting protease, in Fas/APO-1- and TNF receptor-induced cell death. *Cell*, 85:803–815, 1996.
- [257] J.M. Bollinger, D.S. Kwon, G.W. Huisman, R. Kolter, , and C.T. Glutathionylspermidine metabolism in *E. coli*. Purification, cloning, overproduction and characterization of a bifunctional glutathionylspermidine synthetase/amidase. *J. Biol. Chem.*, 270:14031–14041, 1995.
- [258] J.S. Bond and A.J. Barrett. Degradation of fructose-1,6-bisphosphate aldolase by cathepsin B. A further example of peptidyl dipeptidase activity of this proteinase. *Biochem. J.*, 189:17–25, 1980.
- [259] M.D. Bond and H.D. Van Wart. Characterization of the individual collagenases from *Clostridium histolyticum*. *Biochemistry*, 23:3085–3091, 1984.
- [260] M.D. Bond and H.D. Van Wart. Relationship between the individual collagenases of *Clostridium histolyticum*: evidence for evolution by gene duplication. *Biochemistry*, 23:3092–3099, 1984.
- [261] V.A. Bondarenko, M. Deasi, S. Dua, M. Yamazaki, R.H. Amin, K.K. Yousif, T. Kinumi, M. Ohashi, N. Komori, H. Matsumoto, K.W. Jackson, F. Hayashi, J. Usukura, V.M. Lipikin, and A. Yamazaki. Residues within the polycationic region of cGMP phosphodiesterase  $\gamma$  subunit crucial for the interaction with transducin  $\alpha$  subunit. Identification by endogenous ADP-ribosylation and site-directed mutagenesis. *J. Biol. Chem.*, 272:15856–15864, 1997.
- [262] R. Bone, D. Frank, C.A. Kettner, and D.A. Agard. Structural analysis of specificity:  $\alpha$ -lytic protease complexes with analogues of reaction intermediates. *Biochemistry*, 28:7600–7609, 1989.
- [263] A.G. Booth, L.M.L. Hubbard, and A.J. Kenny. Proteins of the kidney microvillar membrane. Immunoelectrophoretic analysis of the membrane hydrolase: identification and resolution of the detergent- and proteinase-solubilized forms. *Biochem. J.*, 179:397–405, 1979.
- [264] P. Borgia and L. Campbell. Properties of two homologous alkaline proteases from *Streptomyces rectus*. *J. Bacteriol.*, 123:1109–1115, 1974.
- [265] M. Borisova, J. Gisin, and C. Mayer. The *N*-acetylmuramic acid 6-phosphate phosphatase MupP completes the *Pseudomonas* peptidoglycan recycling pathway leading to intrinsic fosfomycin resistance. *mBio*, 8:e00092–17–, 2017.
- [266] L.F. Borkenhagen and E.P. Kennedy. The enzymatic exchange of L-serine with *O*-phospho-L-serine catalyzed by a specific phosphatase. *J. Biol. Chem.*, 234:849–853, 1959.
- [267] H.B. Bosmann. Membrane marker enzymes. Characterization of an arylesterase of guinea pig cerebral cortex utilizing *p*-nitrophenyl acetate as substrate. *Biochim. Biophys. Acta*, 276:180–191, 1972.

- [268] M.J. Bossard, T.A. Tomaszek, S.K. Thompson, B.Y. Amegadzie, C.R. Hanning, C. Jones, J.T. Kurdyla, D.E. McNulty, F.H. Drake, M. Gowen, and M.A. Levy. Proteolytic activity of human osteoclast cathepsin K - Expression, purification, activation, and substrate identification. *J. Biol. Chem.*, 271:12517–12524, 1996.
- [269] K.L. Boundy-Mills, M.L. de Souza, R.T. Mandelbaum, L.P. Wackett, and M.J. Sadowsky. The *atzB* gene of *Pseudomonas* sp. strain ADP encodes the second enzyme of a novel atrazine degradation pathway. *Appl. Environ. Microbiol.*, 63:916–923, 1997.
- [270] H.R. Bourne, D.A. Sanders, and F. McCormick. The GTPase superfamily: conserved structure and molecular mechanisms. *Nature*, 349:117–127, 1991.
- [271] J. Bouvier, C. Cordier, H. Vogel, R. Reichelt, and R. Etges. Characterization of the promastigote surface protease of *Leishmania* as a membrane-bound zinc endopeptidase. *Mol. Biochem. Parasitol.*, 37:235–246, 1989.
- [272] J. Bouvier, P. Schneider, R. Etges, and C. Bordier. Peptide substrate specificity of the membrane-bound metalloprotease of *Leishmania*. *Biochemistry*, 29:10113–10119, 1990.
- [273] R.O. Brady, A.E. Gal, R.M. Bradley, and E. Matensson. The metabolism of ceramide trihexosides. I. Purification and properties of an enzyme that cleaves the terminal galactose molecule of galactosylgalactosylglucosylceramide. *J. Biol. Chem.*, 242:1021–1026, 1967.
- [274] R.O. Brady, A.E. Gal, J.N. Kanfer, and R.M. Bradley. The metabolism of glucocerebrosides. 3. Purification and properties of a glucosyl- and galactosylceramide-cleaving enzyme from rat intestinal tissue. *J. Biol. Chem.*, 240:3766–3770, 1965.
- [275] R.O. Brady, J.N. Kanfer, and D. Shapiro. The metabolism of glucocerebrosides. I. Preparation and properties of a glucocerebroside-cleaving enzyme from spleen tissue. *J. Biol. Chem.*, 240:39–43, 1966.
- [276] N. Brakch, H. Boussetta, M. Rholam, and P. Cohen. Processing endoprotease recognizes a structural feature at the cleavage site of peptide prohormones. The *pro*-ocytocin/neurophysin model. *J. Biol. Chem.*, 264:15912–15916, 1989.
- [277] H. Brandt, Z.L. Capulong, and E.Y.C. Lee. Purification and properties of rabbit liver phosphorylase phosphatase. *J. Biol. Chem.*, 250:8038–8044, 1975.
- [278] S.S. Brar, E.J. Sacho, I. Tessmer, D.L. Croteau, D.A. Erie, and M. Diaz. Activation-induced deaminase, AID, is catalytically active as a monomer on single-stranded DNA. *DNA Repair (Amst.)*, 7:77–87, 2008.
- [279] A.G. Braun, M. Radman, and L. Grossman. Enzymic repair of DNA: sites of hydrolysis by the *Escherichia coli* endonuclease specific for pyrimidine dimers (corendonuclease II). *Biochemistry*, 15:4116–4120, 1976.
- [280] H.G. Bray, S.P. James, I.M. Raffan, B.E. Ryman, and W.V. Thorpe. The fate of certain organic acids and amides in the rabbit. 7. An amidase of rabbit liver. *Biochem. J.*, 44:618–625, 1949.
- [281] H.G. Bray, S.P. James, W.V. Thorpe, and M.R. Wasdell. The fate of certain organic acids and amides in the rabbit. 11. Further observations on the hydrolysis of amides by tissue extracts. *Biochem. J.*, 47:294–299, 1950.
- [282] R. Breathnach, L.M. Matrisian, M.-C. Gesnel, and P. Leroy. Sequences coding for part of oncogene-induced transin are highly conserved in a related rat gene. *Nucleic Acids Res.*, 15:1139–1151, 1987.
- [283] K. Breddam. Serine carboxypeptidases. A review. *Carlsberg Res. Commun.*, 51:83–128, 1986.
- [284] K. Breddam and M. Meldal. Substrate preferences of glutamic-acid-specific endopeptidases assessed by synthetic peptide substrates based on intramolecular fluorescence quenching. *Eur. J. Biochem.*, 206:103–107, 1992.
- [285] K. Breddam, S.B. Sørensen, and I. Svendsen. Primary structure and enzymatic properties of carboxypeptidase II from wheat bran. *Carlsberg Res. Commun.*, 52:297–311, 1987.
- [286] J.H. Brehm, J.W. Mellors, and N. Sluis-Cremer. Mechanism by which a glutamine to leucine substitution at residue 509 in the ribonuclease H domain of HIV-1 reverse transcriptase confers zidovudine resistance. *Biochemistry*, 47:14020–14027, 2008.
- [287] M.M. Bresler, S.J. Rosser, A. Basran, and N.C. Bruce. Gene cloning and nucleotide sequencing and properties of a cocaine esterase from *Rhodococcus* sp. strain MB1. *Appl. Environ. Microbiol.*, 66:904–908, 2000.

- [288] C.D. Bridges and R.A. Alvarez. The visual cycle operates via an isomerase acting on *all-trans* retinol in the pigment epithelium. *Science*, 236:1678–1680, 1987.
- [289] W.T. Brigance, C. Barlowe, and T.R. Graham. Organization of the yeast Golgi complex into at least four functionally distinct compartments. *Mol. Biol. Cell*, 11:171–182, 2000.
- [290] B.S. Briggs, , and M.J. Enzyme from microbial source: phthalyl amidase, 1995.
- [291] B.S. Briggs, A.J. Kreuzman, C. Whitesitt, W.K. Yeh, , and M. Discovery, purification, and properties of *o*-phthalyl amidase from *Xanthobacter agilis*. *J. Mol. Catal., B Enzym.*, 2:53–69, 1996.
- [292] R. Brightwell and A.L. Tappel. Lysosomal acid pyrophosphatase and acid phosphatase. *Arch. Biochem. Biophys.*, 124:333–343, 1968.
- [293] J.-M. Brillouet, P. Williams, and M. Moutounet. Purification and some properties of a novel endo- $\beta$ -(1 $\rightarrow$ 6)-D-galactanase from *Aspergillus niger*. *Agric. Biol. Chem.*, 55:1565–1571, 1991.
- [294] A.J. Britt, N.C. Bruce, and C.R. Lowe. Identification of a cocaine esterase in a strain of *Pseudomonas maltophilia*. *J. Bacteriol.*, 174:2087–2094, 1992.
- [295] W.J. Brockbank and K.R. Lynn. Purification and preliminary characterization of two asclepains from the latex of *Asclepias syriaca* L. (milkweed). *Biochim. Biophys. Acta*, 578:13–22, 1979.
- [296] K. Brocklehurst, E. Salih, R. McKee, and H. Smith. Fresh non-fruit latex of *Carica papaya* contains papain, multiple forms of chymopapain A and papaya proteinase . *Biochem. J.*, 228:525–527, 1985.
- [297] K. Brocklehurst, F. Willenbrock, and E. Salih. Cysteine proteinases. In A. Neuberger and K. Brocklehurst, editors, *New Comprehensive Biochemistry: Hydrolytic Enzymes*, pages 39–158. Elsevier, Amsterdam, 1987.
- [298] A.F. Brodie and F. Lipmann. Identification of a gluconolactonase. *J. Biol. Chem.*, 212:677–685, 1955.
- [299] J.W. Brodrick, M.C. Geokas, and C. Largman. Human carboxypeptidase B. II. Purification of the enzyme from pancreatic tissue and comparison with the enzymes present in pancreatic secretion. *Biochim. Biophys. Acta*, 452:468–481, 1976.
- [300] D. Brömme, K. Bescherer, H. Kirschke, and S. Fittkau. Enzyme-substrate interactions in the hydrolysis of peptides by cathepsins B and H from rat liver. *Biochem. J.*, 245:381–385, 1987.
- [301] D. Bromme, J.L. Klaus, K. Okamoto, D. Rasnick, and J.T. Palmer. Peptidyl vinyl sulphones: A new class of potent and selective cysteine protease inhibitors - S2P2 specificity of human cathepsin O<sub>2</sub> in comparison with cathepsins S and L. *Biochem. J.*, 315:85–89, 1996.
- [302] D. Brömme, Z. Li, M. Barnes, and E. Mehler. Human cathepsin V functional expression, tissue distribution, electrostatic surface potential, enzymatic characterization, and chromosomal localization. *Biochemistry*, 38:2377–2385, 1999.
- [303] D. Brömme, A.B. Rossi, S.P. Smeekens, D.C. Anderson, and D.G. Payan. Human bleomycin hydrolase: molecular cloning, sequencing, functional expression, and enzymatic characterization. *Biochemistry*, 35:6706–6714, 1996.
- [304] D. Brömme, A. Steinert, S. Friebe, S. Fittkau, B. Wiederanders, and H. Kirschke. The specificity of bovine spleen cathepsin S. A comparison with rat liver cathepsins L and B. *Biochem. J.*, 264:475–485, 1989.
- [305] K.P. Brooks, E.A. Jones, B.-D. Kim, and E.G. Sander. Bovine liver dihydropyrimidine amidohydrolase: purification, properties, and characterization as a zinc metalloenzyme. *Arch. Biochem. Biophys.*, 226:469–483, 1983.
- [306] F. Brossier, T.J. Jewett, L.D. Sibley, and S. Urban. A spatially localized rhomboid protease cleaves cell surface adhesins essential for invasion by *Toxoplasma*. *Proc. Natl. Acad. Sci. USA*, 102:4146–4151, 2005.
- [307] B. Illingworth Brown and D.H. Brown. The subcellular distribution of enzymes in type II glycogenosis and the occurrence of an oligo- $\alpha$ -1,4-glucan glucohydrolase in human tissues. *Biochim. Biophys. Acta*, 110:124–133, 1965.
- [308] D.H. Brown and B.I. Brown. Enzymes of glycogen debranching: Amylo-1,6-glucosidase (I) and oligo-1,4 $\rightarrow$ 1,4-glucanotransferase (II). *Methods Enzymol.*, 8:515–524, 1966.
- [309] J.L. Brown. Purification and properties of dipeptidase M from *Escherichia coli* B. *J. Biol. Chem.*, 248:409–416, 1973.

- [310] M.S. Brown, J. Ye, R.B. Rawson, and J.L. Goldstein. Regulated intramembrane proteolysis: a control mechanism conserved from bacteria to humans. *Cell*, 100:391–398, 2000.
- [311] P. Browne and G. O’Cuinn. The purification and characterization of a proline dipeptidase from guinea pig brain. *J. Biol. Chem.*, 258:6147–6154, 1983.
- [312] G.J. Broze, Higuchi Jr., and D.A. Coagulation-dependent inhibition of fibrinolysis: Role of carboxypeptidase U and the premature lysis of clots from hemophilic plasma. *Blood*, 88:3815–3823, 1996.
- [313] G.J. Broze, Majerus Jr., and P.W. Human factor VII. *Methods Enzymol.*, 80:228–237, 1981.
- [314] M.J. Brubaker, W.C. Groutas, J.R. Hoidal, and N.V. Rao. Human neutrophil proteinase 3: mapping of the substrate binding site using peptidyl thiobenzyl esters. *Biochem. Biophys. Res. Commun.*, 188:1318–1324, 1992.
- [315] J.S. Bruce, M.W. McLean, W.F. Long, and F.B. Williamson. *Flavobacterium heparinum* 3-O-sulphatase for N-substituted glucosamine 3-O-sulphate. *Eur. J. Biochem.*, 148:359–365, 1985.
- [316] D.F. Bruhn, J. Li, S. Silver, F. Roberto, and B.P. Rosen. The arsenical resistance operon of IncN plasmid R46. *FEMS Microbiol. Lett.*, 139:149–153, 1996.
- [317] A. Brunella, M. Graf, M. Kittelmann, K. Lauma, and O. Ghisalba. Production, purification, and characterization of a highly enantioselective (S)-N-acetyl-1-phenylethyl amidohydrolase from *Rhodococcus*. *Appl. Microbiol. Biotechnol.*, 47:515–520, 1997.
- [318] E.G. Brungraber. Nucleotides involved in the enzymatic conjugation of phenols with sulfate. *J. Biol. Chem.*, 233:472–477, 1958.
- [319] C.B. Bruni, V. Sica, F. Auricchio, and I. Covelli. Further kinetic and structural characterization of the lysosomal  $\alpha$ -D-glucoside glucohydrolase from cattle liver. *Biochim. Biophys. Acta*, 212:470–477, 1970.
- [320] M. Brunner and W. Neupert. Purification and characterization of the mitochondrial processing peptidase of *Neurospora crassa*. *Methods Enzymol.*, 248:717–728, 1994.
- [321] M.B. Brurberg, I.F. Nes, and V.G. Eijsink. Comparative studies of chitinases A and B from *Serratia marcescens*. *Microbiology*, 142:1581–1589, 1996.
- [322] C. Bublitz and A.L. Lehninger. The role of aldonolactonase in the conversion of L-gulonate to L-ascorbate. *Biochim. Biophys. Acta*, 47:288–297, 1961.
- [323] M.J. Buchanan, S.H. Imam, W.A. Eskue, and W.J. Snell. Activation of the cell wall degrading protease, lysin, during sexual signalling in *Chlamydomonas*: the enzyme is stored as an inactive, higher relative molecular mass precursor in the periplasm. *J. Cell. Biol.*, 108:199–207, 1989.
- [324] J. Buchert and L. Viikari. The role of xylonolactone in xylonic acid production by *Pseudomonas fragi*. *Appl. Microbiol. Biotechnol.*, 27:333–336, 1988.
- [325] J. Burch. The purification and properties of horse liver esterase. *Biochem. J.*, 58:415–426, 1954.
- [326] A.W. Burg and G.M. Brown. The biosynthesis of folic acid. 8. Purification and properties of the enzyme that catalyzes the production of formate from carbon atom 8 of guanosine triphosphate. *J. Biol. Chem.*, 243:2349–2358, 1968.
- [327] R.W. Burg and E.E. Snell. The bacterial oxidation of vitamin B<sub>6</sub>. VI. Pyridoxal dehydrogenase and 4-pyridoxolactonase. *J. Biol. Chem.*, 244:2585–2589, 1969.
- [328] R. Burlingame and P.J. Chapman. Catabolism of phenylpropionic acid and its 3-hydroxy derivative by *Escherichia coli*. *J. Bacteriol.*, 155:113–121, 1983.
- [329] R.P. Burlingame, L. Wyman, and P.J. Chapman. Isolation and characterization of *Escherichia coli* mutants defective for phenylpropionate degradation. *J. Bacteriol.*, 168:55–64, 1986.
- [330] K.E. Burns, S. Baumgart, P.C. Dorrestein, H. Zhai, F.W. McLafferty, and T.P. Begley. Reconstitution of a new cysteine biosynthetic pathway in *Mycobacterium tuberculosis*. *J. Am. Chem. Soc.*, 127:11602–11603, 2005.

- [331] K.E. Burns, F.A. Cerda-Maira, T. Wang, H. Li, W.R. Bishai, and K.H. Darwin. "Depupylation" of prokaryotic ubiquitin-like protein from mycobacterial proteasome substrates. *Mol. Cell*, 39:821–827, 2010.
- [332] R.B. Burrows and G.M. Brown. Presence of *Escherichia coli* of a deaminase and a reductase involved in biosynthesis of riboflavin. *J. Bacteriol.*, 136:657–667, 1978.
- [333] R.E. Burton, T.A. Baker, and R.T. Sauer. Nucleotide-dependent substrate recognition by the AAA<sup>+</sup> HslUV protease. *Nat. Struct. Mol. Biol.*, 12:245–251, 2005.
- [334] J. Buschmann, B. Moritz, M. Jeske, H. Lilie, A. Schierhorn, and E. Wahle. Identification of *Drosophila* and human 7-methyl GMP-specific nucleotidases. *J. Biol. Chem.*, 288:2441–2451, 2013.
- [335] D.E. Bussiere, S.D. Pratt, L. Katz, J.M. Severin, T. Holzman, and C.H. Park. The structure of VanX reveals a novel amino-dipeptidase involved in mediating transposon-based vancomycin resistance. *Mol. Cell.*, 2:75–84, 1998.
- [336] M.J. Butler. Snapalysin. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 1134–1135. Handbook of Proteolytic Enzymes, London, 1998.
- [337] P.E. Butler, M.J. McKay, and J.S. Bond. Characterization of meprin, a membrane-bound metalloendopeptidase from mouse kidney. *Biochem. J.*, 241:229–235, 1987.
- [338] J. Butterworth and J.J. Duncan. Carboxypeptidase B activity of cultured skin fibroblasts and relationship to cystic fibrosis. *Clin. Chim. Acta*, 97:39–43, 1979.
- [339] D.J. Buttle, P.M. Dando, P.F. Coe, S.L. Sharp, S.T. Shepherd, and A.J. Barrett. The preparation of fully active chymopain free of contaminating proteinases. *Biol. Chem. Hoppe-Seyler*, 371:1083–1088, 1990.
- [340] D.J. Buttle, A.A. Kembhavi, S.L. Sharp, R.E. Shute, D.H. Rich, and A.J. Barrett. Affinity purification of the novel cysteine proteinase papaya proteinase IV and papain from papaya latex. *Biochem. J.*, 261:469–476, 1989.
- [341] D.J. Buttle, A. Ritonja, P.M. Dando, M. Abrahamson, E.N. Shaw, P. Wikstrom, V. Turk, and A.J. Barrett. Interaction of papaya proteinase IV with inhibitors. *FEBS Lett.*, 262:58–60, 1990.
- [342] D.J. Buttle, A. Ritonja, L.H. Pearl, V. Turk, and A.J. Barrett. Selective cleavage of glycyl bonds by papaya proteinase IV. *FEBS Lett.*, 260:195–197, 1990.
- [343] L.L. Button and W.R. McMaster. Molecular cloning of the major surface antigen of *Leishmania*. *J. Exp. Med.*, 167:724–729, 1988.
- [344] W.L. Byrne. Glucose-6-phosphatase and phosphoserine phosphatase. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 5, pages 73–78. Academic Press, New York, 2nd edition, 1961.
- [345] J. Cabello, C. Basilio, and V. Prajoux. Kinetic properties of erythrocyte- and liver arginase. *Biochim. Biophys. Acta*, 48:148–152, 1961.
- [346] J.A. Cabezas. Some comments on the type references of the official nomenclature (IUB) for  $\beta$ -N-acetylglucosaminidase,  $\beta$ -N-acetylhexosaminidase and  $\beta$ -N-acetylgalactosaminidase. *Biochem. J.*, 261:1059–1060, 1989.
- [347] J.A. Cabezas. Some questions and suggestions on the type references of the official nomenclature (IUB) for sialidase(s) and endosialidase. *Biochem. J.*, 278:311–312, 1991.
- [348] E. Cabib and L.F. Leloir. The biosynthesis of trehalose phosphate. *J. Biol. Chem.*, 231:259–275, 1958.
- [349] S. Cadel, T. Foulon, A. Viron, A. Balogh, S. Midol-Monnet, N. Noel, and P. Cohen. Aminopeptidase B from the rat testis is a bifunctional enzyme structurally related to leukotriene-A<sub>4</sub> hydrolase. *Proc. Natl. Acad. Sci. USA*, 94:2963–2968, 1997.
- [350] S. Cadel, A.R. Pierotti, T. Foulon, C. Créminon, N. Barré, D. Segrétain, and P. Cohen. Aminopeptidase-B in the rat testes: Isolation, functional properties and cellular localization in the seminiferous tubules. *Mol. Cell. Endocrinol.*, 110:149–160, 1995.
- [351] V. Cafardi, M. Biagini, M. Martinelli, R. Leuzzi, J.T. Rubino, F. Cantini, N. Norais, M. Scarselli, D. Serruto, and M. Unnikrishnan. Identification of a novel zinc metalloprotease through a global analysis of *Clostridium difficile* extracellular proteins. *PLoS One*, 8:e81306–e81306, 2013.



- [352] J.J. Caffrey, S.T. Safrany, X. Yang, and S.B. Shears. Discovery of molecular and catalytic diversity among human diphosphoinositol-polyphosphate phosphohydrolases. An expanding Nudt family. *J. Biol. Chem.*, 275:12730–12736, 2000.
- [353] A. Cahour, B. Falgout, and C.-J. Lai. Cleavage of the dengue virus polyprotein at the NS3/NS4A and NS4B/NS5 junctions is mediated by viral protease NS2B-NS3, whereas NS4A/NS4B may be processed by a cellular protease. *J. Virol.*, 66:1535–1542, 1992.
- [354] S.S. Cairns, A. Cornish, and R.A. Cooper. Cloning, sequencing and expression in *Escherichia coli* of two *Rhizobium* sp. genes encoding haloalkanoate dehalogenases of opposite stereospecificity. *Eur. J. Biochem.*, 235:744–749, 1996.
- [355] K.K. Caldwell, D.L. Lips, V.S. Bansal, and P.W. Majerus. Isolation and characterization of two 3-phosphatases that hydrolyze both phosphatidylinositol 3-phosphate and inositol 1,3-bisphosphate. *J. Biol. Chem.*, 266:18378–18386, 1991.
- [356] A.J. Callaghan, J.G. Grossmann, Y.U. Redko, L.L. Ilag, M.C. Moncrieffe, M.F. Symmons, C.V. Robinson, K.J. McDowall, and B.F. Luisi. Quaternary structure and catalytic activity of the *Escherichia coli* ribonuclease E amino-terminal catalytic domain. *Biochemistry*, 42:13848–13855, 2003.
- [357] P. Calvo, A. Reglero, and J.A. Cabezas. Purification and properties of  $\beta$ -*N*-acetylhexosaminidase from the mollusc *Helicella ericetorum* Muller. *Biochem. J.*, 175:743–750, 1978.
- [358] L.A. Camp, , and S.L. Assay and isolation of palmitoyl-protein thioesterase from bovine brain using palmitoylated *H*-Ras as substrate. *Methods Enzymol.*, 250:336–347, 1995.
- [359] B. Campbell, H. Lin, R. Davis, and E. Ballew. The purification and properties of a particulate renal dipeptidase. *Biochim. Biophys. Acta*, 118:371–386, 1966.
- [360] B.J. Campbell. Renal dipeptidase. *Methods Enzymol.*, 19:722–729, 1970.
- [361] L.L. Campbell. Reductive degradation of pyrimidines. 5. Enzymatic conversion of *N*-carbamyl- $\beta$ -alanine to  $\beta$ -alanine, carbon dioxide, and ammonia. *J. Biol. Chem.*, 235:2375–2378, 1960.
- [362] J. Canales, A. Fernández, J.M. Ribeiro, A. Cabezas, J.R. Rodrigues, J.C. Cameselle, and M.J. Costas.  $Mn^{2+}$ -dependent ADP-ribose/CDP-alcohol pyrophosphatase: a novel metallophosphoesterase family preferentially expressed in rodent immune cells. *Biochem. J.*, 413:103–113, 2008.
- [363] J. Canales, A. Fernandez, J.R. Rodrigues, R. Ferreira, J.M. Ribeiro, A. Cabezas, M.J. Costas, and J.C. Cameselle. Hydrolysis of the phosphoanhydride linkage of cyclic ADP-ribose by the  $Mn(2+)$ -dependent ADP-ribose/CDP-alcohol pyrophosphatase. *FEBS Lett.*, 583:1593–1598, 2009.
- [364] J. Canales, R.M. Pinto, M.J. Costas, M.T. Hernández, A. Miró, D. Bernet, A. Fernández, and J.C. Cameselle. Rat liver nucleoside diphosphosugar or diphosphoalcohol pyrophosphatases different from nucleotide pyrophosphatase or phosphodiesterase I: substrate specificities of  $Mg^{2+}$ -and/or  $Mn^{2+}$ -dependent hydrolases acting on ADP-ribose. *Biochim. Biophys. Acta*, 1246:167–177, 1995.
- [365] D.J. Candy and B.A. Kilby. The biosynthesis of trehalose in the locust fat body. *Biochem. J.*, 78:531–536, 1961.
- [366] Z.N. Canellakis, P.K. Bondy, J.A. May, Myers-Robfogel Jr., Sartorelli M.K., and A.C. Identification of a glycosidase activity with apparent specificity for 2-deoxy-D-glucose in glycosidic linkage. *Eur. J. Biochem.*, 143:159–163, 1984.
- [367] J.R. Cantor, E.M. Stone, and G. Georgiou. Expression and biochemical characterization of the human enzyme N-terminal asparagine amidohydrolase. *Biochemistry*, 50:3025–3033, 2011.
- [368] J. Caravaca and S. Grisolia. Enzymatic decarbamylation of carbamyl  $\beta$ -alanine and carbamyl  $\beta$ -aminoisobutyric acid. *J. Biol. Chem.*, 231:357–365, 1958.
- [369] K. Carlson and L.D. Kosturko. Endonuclease II of coliphage T<sub>4</sub>: a recombinase disguised as a restriction endonuclease. *Mol. Microbiol.*, 27:671–676, 1998.
- [370] K. Carlson, L.D. Kosturko, and A.C. Nystrom. Sequence-specific cleavage by bacteriophage T<sub>4</sub> endonuclease II *in vitro*. *Mol. Microbiol.*, 31:1395–1405, 1999.

- [371] K. Carlson, M. Krabbe, A.C. Nystrom, and L.D. Kosturko. DNA determinants of restriction. Bacteriophage T<sub>4</sub> endonuclease II-dependent cleavage of plasmid DNA *in vivo*. *J. Biol. Chem.*, 268:8908–8918, 1993.
- [372] G.M. Carman. Phosphatidate phosphatases and diacylglycerol pyrophosphate phosphatases in *Saccharomyces cerevisiae* and *Escherichia coli*. *Biochim. Biophys. Acta*, 1348:45–55, 1997.
- [373] G.M. Carman and G.S. Han. Phosphatidic acid phosphatase, a key enzyme in the regulation of lipid synthesis. *J. Biol. Chem.*, 284:2593–2597, 2009.
- [374] C. Carmona and G.L. Gray. Nucleotide sequence of the serine protease gene of *Staphylococcus aureus*, strain V8. *Nucleic Acids Res.*, 15:6757–6757, 1987.
- [375] H.R. Carne and E. Onon. Action of *Corynebacterium ovis* exotoxin on endothelial cells of blood vessels. *Nature*, 271:246–248, 1978.
- [376] B.Z. Carter, A.L. Wiseman, R. Orkiszewski, K.D. Ballard, C.N. Ou, and M.W. Lieberman. Metabolism of leukotriene C<sub>4</sub> in  $\gamma$ -glutamyl transpeptidase-deficient mice. *J. Biol. Chem.*, 272:12305–12310, 1997.
- [377] C.E. Carter. Partial purification of a non-phosphorylytic uridine nucleosidase from yeast. *J. Am. Chem. Soc.*, 73:1508–1510, 1951.
- [378] K.M. Carvalho, C. Joudiou, H. Boussetta, A.-M. Leseney, and P. Cohen. A peptide-hormone-inactivating endopeptidase in *Xenopus laevis* skin secretion. *Proc. Natl. Acad. Sci. USA*, 89:84–88, 1992.
- [379] A. Castanares, S.I. McCrae, and T.M. Wood. Purification and properties of a feruloyl/*p*-coumaroyl esterase from the fungus *Penicillium pinophilum*. *Enzyme Microbiol. Technol.*, 14:875–884, 1992.
- [380] F.J. Castellino and J.R. Powell. Human plasminogen. *Methods Enzymol.*, 80:365–378, 1981.
- [381] F.J. Castellino and J.M. Sodetz. Rabbit plasminogen and plasmin isozymes. *Methods Enzymol.*, 45:273–286, 1976.
- [382] N.X. Cawley, H.C. Chen, M.C. Beinfeld, and Y.P. Loh. Specificity and kinetic studies on the cleavage of various prohormone mono- and paired-basic residue sites by yeast aspartic protease 3. *J. Biol. Chem.*, 271:4168–4176, 1996.
- [383] J.J. Cazzulo, V. Stoka, and V. Turk. The major cysteine proteinase of *Trypanosoma cruzi*: a valid target for chemotherapy of Chagas disease. *Curr. Pharm. Des.*, 7:1143–1156, 2001.
- [384] E.S. Cedergren-Zeppezauer, G. Larsson, P.O. Nyman, Z. Dauter, and K.S. Wilson. Crystal structure of a dUTPase. *Nature*, 355:740–743, 1992.
- [385] M.S. Center and F.J. Behal. A cyclic phosphodiesterase with 3'-nucleotidase activity from *Proteus mirabilis*. *J. Biol. Chem.*, 243:138–143, 1968.
- [386] T.B. Cereija, S. Alarico, N. Empadinhas, and P.J.B. Pereira. Production, crystallization and structure determination of a mycobacterial glucosylglycerate hydrolase. *Acta Crystallogr. F Struct. Biol. Commun.*, 73:536–540, 2017.
- [387] N. Cetinbas, M.S. Macauley, K.A. Stubbs, R. Drapala, and D.J. Vocadlo. Identification of Asp<sup>174</sup> and Asp<sup>175</sup> as the key catalytic residues of human O-GlcNAcase by functional analysis of site-directed mutants. *Biochemistry*, 45:3835–3844, 2006.
- [388] B.K. Chaal, R.M. Mould, A.C. Barbrook, J.C. Gray, and C.J. Howe. Characterization of a cDNA encoding the thylakoidal processing peptidase from *Arabidopsis thaliana*. Implications for the origin and catalytic mechanism of the enzyme. *J. Biol. Chem.*, 273:689–692, 1998.
- [389] T.J. Chambers, C.S. Hahn, R. Galler, and C.M. Rice. Flavivirus genome organization, expression, and replication. *Annu. Rev. Microbiol.*, 44:649–688, 1990.
- [390] J.J. Champoux and S.J. Schultz. Ribonuclease H: properties, substrate specificity and roles in retroviral reverse transcription. *FEBS J.*, 276:1506–1516, 2009.
- [391] M.K. Chan, W. Gong, P.T.R. Rajagopalan, B. Hao, C.M. Tsai, and D. Pei. Crystal structure of the *Escherichia coli* peptide deformylase. *Biochemistry*, 36:13904–13909, 1997.

- [392] A.L. Chang, J.R. Tuckerman, G. Gonzalez, R. Mayer, H. Weinhouse, G. Volman, D. Amikam, M. Benziman, and M.A. Gilles-Gonzalez. Phosphodiesterase A1, a regulator of cellulose synthesis in *Acetobacter xylinum*, is a heme-based sensor. *Biochemistry*, 40:3420–3426, 2001.
- [393] H.Y. Chang and X. Yang. Proteases for cell suicide: functions and regulation of caspases. *Microbiol. Mol. Biol. Rev.*, 64:821–846, 2000.
- [394] K.H. Chang, P.H. Liang, W. Beck, J.D. Scholten, , and D. Isolation and characterization of the three polypeptide components of 4-chlorobenzoate dehalogenase from *Pseudomonas* sp. strain CBS-3. *Biochemistry*, 31:5605–5610, 1992.
- [395] P.C. Chang, T-C. Kue, A. Tsugita, and Y.H.W. Lee. Extracellular metalloprotease gene of *Streptomyces cacaoi*: structure, nucleotide sequence and characterization of the cloned gene product. *Gene*, 88:87–95, 1990.
- [396] P.K. Chang, K. Yabe, and J. Yu. The *Aspergillus parasiticus estA*-encoded esterase converts versiconal hemiacetal acetate to versiconal and versiconol acetate to versiconol in aflatoxin biosynthesis. *Appl. Environ. Microbiol.*, 70:3593–3599, 2004.
- [397] W.-J. Chang, S. Horiuchi, K. Takahashi, M. Yamasaki, and Y. Yamada. The structure and function of acid proteases. VI. Effects of acid protease-specific inhibitors on the acid proteases from *Aspergillus niger* var. macrosporus. *J. Biochem. (Tokyo)*, 80:975–981, 1976.
- [398] Y-H. Chang and J.A. Smith. Molecular cloning and sequencing of genomic DNA encoding aminopeptidase I from *Saccharomyces cerevisiae*. *J. Biol. Chem.*, 264:6979–6983, 1989.
- [399] Y.Y. Chang and E.P. Kennedy. Phosphatidyl glycerophosphate phosphatase. *J. Lipid Res.*, 8:456–462, 1967.
- [400] J. Chao and L. Chao. Rat urinary kallikrein. *Methods Enzymol.*, 163:128–143, 1988.
- [401] J.W. Chase and C.C. Richardson. Exonuclease VII of *Escherichia coli*. *J. Biol. Chem.*, 249:4553–4561, 1974.
- [402] J.W. Chase and C.C. Richardson. Ribonuclease VII of *Escherichia coli*. *J. Biol. Chem.*, 249:4545–4552, 1974.
- [403] A.K. Chatterjee and L.N. Gibbins. Metabolism of phloridzin by *Erwinia herbicola*: nature of the degradation products, and the purification and properties of phloretin hydrolase. *J. Bacteriol.*, 100:594–600, 1969.
- [404] S. Chatterjee and N. Ghosh. Neutral sphingomyelinase from human urine. Purification and preparation of monospecific antibodies. *J. Biol. Chem.*, 264:12554–12561, 1989.
- [405] G. Chaudhuri, M. Chaudhuri, A. Pan, and K.-P. Chang. Surface acid proteinase (gp63) of *Leishmania mexicana*. A metalloenzyme capable of protecting liposome-encapsulated proteins from phagolysosomal degradation by macrophages. *J. Biol. Chem.*, 264:7483–7489, 1989.
- [406] M. Chauleau, A. Jacewicz, and S. Shuman. DNA3'pp5'G de-capping activity of aprataxin: effect of cap nucleoside analogs and structural basis for guanosine recognition. *Nucleic Acids Res.*, 43:6075–6083, 2015.
- [407] F. Checler, H. Barelli, and J.-P. Vincent. Tissue distribution of a novel neurotensin-degrading metallopeptidase. An immunological approach using monospecific polyclonal antibodies. *Biochem. J.*, 257:549–554, 1989.
- [408] F. Checler, J.P. Vincent, and P. Kitabgi. Purification and characterization of a novel neurotensin-degrading peptidase from rat brain synaptic membranes. *J. Biol. Chem.*, 261:11274–11281, 1986.
- [409] B.K. Chen, M.T. Overgaard, L.K. Bale, Z.T. Resch, M. Christiansen, C. Oxvig, and C.A. Conover. Molecular regulation of the IGF-binding protein-4 protease system in human fibroblasts: identification of a novel inducible inhibitor. *Endocrinology*, 143:1199–1205, 2002.
- [410] C.B. Chen and R. Wallis. Two mechanisms for mannose-binding protein modulation of the activity of its associated serine proteases. *J. Biol. Chem.*, 279:26058–26065, 2004.
- [411] C.C. Chen, Y. Han, W. Niu, A.N. Kulakova, A. Howard, J.P. Quinn, D. Dunaway-Mariano, and O. Herzberg. Structure and kinetics of phosphonopyruvate hydrolase from *Variovorax* sp. Pal2: new insight into the divergence of catalysis within the PEP mutase/isocitrate lyase superfamily. *Biochemistry*, 45:11491–11504, 2006.
- [412] J.-M. Chen, N.D. Rawlings, R.A.E. Stevens, and A.J. Barrett. Identification of the active site of legumain links it to caspases, clostripain and gingipains in a new clan of cysteine endopeptidases. *FEBS Lett.*, 441:361–365, 1998.

- [413] M. Chen, X. Ma, X. Chen, M. Jiang, H. Song, and Z. Guo. Identification of a hotdog fold thioesterase involved in the biosynthesis of menaquinone in *Escherichia coli*. *J. Bacteriol.*, 195:2768–2775, 2013.
- [414] N. Chen, M.A. Walsh, Y. Liu, R. Parker, and H. Song. Crystal structures of human DcpS in ligand-free and m<sup>7</sup>GDP-bound forms suggest a dynamic mechanism for scavenger mRNA decapping. *J. Mol. Biol.*, 347:707–718, 2005.
- [415] P. Chen, H. Tsuge, R.J. Almassy, C.L. Gribskov, S. Katoh, D.L. Vanderpool, S.A. Margosiak, C. Pinko, D.A. Matthews, and C.C. Kan. Structure of the human cytomegalovirus protease catalytic domain reveals a novel serine protease fold and catalytic triad. *Cell*, 86:477–483, 1996.
- [416] P.R.S. Chen and W.C. Dauterman. Alkylamidase of sheep liver. *Biochim. Biophys. Acta*, 250:216–223, 1971.
- [417] W.P. Chen, M. Matsuo, and Y. Tsuneo. Purification and some properties of  $\beta$ -1,3-xylanase from *Aspergillus terreus* A-07. *Agric. Biol. Chem.*, 50:1183–1194, 1986.
- [418] Y.R. Chen, N. Datta, and S.J. Roux. Purification and partial characterization of a calmodulin-stimulated nucleoside triphosphatase from pea nuclei. *J. Biol. Chem.*, 262:10689–10694, 1987.
- [419] Z. Chen, Y. Li, E. Chen, D. Hall, P. Darke, C. Culberson, J.A. Shafer, and L.A. Kuo. Crystal structure at 1.9-Å resolution of human immunodeficiency virus (HIV) II protease complexed with L-735,524, an orally bioavailable inhibitor of the HIV proteases. *J. Biol. Chem.*, 269:26344–26348, 1994.
- [420] Z. Chen, J. Potempa, A. Polanowski, M. Wikstrom, and J. Travis. Purification and characterization of a 50-kDa cysteine proteinase (gingipain) from *Porphyromonas gingivalis*. *J. Biol. Chem.*, 267:18896–18901, 1992.
- [421] Z.L. Chen, S. Yoshida, K. Kato, Y. Momota, J. Suzuki, T. Tanaka, J. Ito, H. Nishino, S. Aimoto, H. Kiyama, and S. Shiosaka. Expression and activity-dependent changes of a novel limbic-serine protease gene in the hippocampus. *J. Neurosci.*, 15:5088–5097, 1995.
- [422] D. Cheng, P.J. Espenshade, C.A. Slaughter, J.C. Jaen, M.S. Brown, and J.L. Goldstein. Secreted site-1 protease cleaves peptides corresponding to luminal loop of sterol regulatory element-binding proteins. *J. Biol. Chem.*, 274:22805–22812, 1999.
- [423] G. Cheng, N. Shapir, M.J. Sadowsky, and L.P. Wackett. Allophanate hydrolase, not urease, functions in bacterial cyanuric acid metabolism. *Appl. Environ. Microbiol.*, 71:4437–4445, 2005.
- [424] C.J. Chern, A.B. Macdonald, and A.J. Morris. Purification and properties of a nucleoside triphosphate pyrophosphohydrolase from red cells of the rabbit. *J. Biol. Chem.*, 244:5489–5495, 1969.
- [425] V. Chesneau, A.R. Pierotti, N. Barré, C. Créminon, C. Tougard, and P. Cohen. Isolation and characterization of a dibasic selective metalloendopeptidase from rat testes that cleaves at the amino terminus of arginine residues. *J. Biol. Chem.*, 269:2056–2061, 1994.
- [426] A. Chester, A. Somasekaram, M. Tzimina, A. Jarmuz, J. Gisbourne, R. O’Keefe, J. Scott, and N. Navaratnam. The apolipoprotein B mRNA editing complex performs a multifunctional cycle and suppresses nonsense-mediated decay. *EMBO J.*, 22:3971–3982, 2003.
- [427] A. Chester, V. Weinreb, C.W. Carter, Navaratnam Jr., and N. Optimization of apolipoprotein B mRNA editing by APOBEC1 apoenzyme and the role of its auxiliary factor, ACF. *RNA*, 10:1399–1411, 2004.
- [428] C.G.C. Chesters and A.T. Bull. The enzymic degradation of laminarin. 2. The multicomponent nature of fungal laminarinases. *Biochem. J.*, 86:31–38, 1963.
- [429] J.-F. Chich, M.P. Chapot-Chartier, B. Ribadeau-Dumas, and J.-C. Gripon. Identification of the active site serine of the X-prolyl aminopeptidase from *Lactococcus lactis*. *FEBS Lett.*, 314:139–142, 1995.
- [430] J.-F. Chich, J.-C. Gripon, and B. Ribadeau-Dumas. Preparation of bacterial X-prolyl dipeptidyl aminopeptidase and its stabilization by organic cosolvents. *Anal. Biochem.*, 224:245–249, 1995.
- [431] S. Chien, R. Weinburg, S. Li, and Y. Li. Endo- $\beta$ -N-acetylglucosaminidase from fig latex. *Biochem. Biophys. Res. Commun.*, 76:317–323, 1977.

- [432] D.T. Chin, S.A. Goff, T. Webster, T. Smith, and A.L. Goldberg. Sequence of the lon gene in *Escherichia coli*. A heat-shock gene which encodes the ATP-dependent protease La. *J. Biol. Chem.*, 263:11718–11728, 1988.
- [433] J.R. Chin, G. Murphy, and Z. Werb. Stromelysin, a connective tissue-degrading metalloendopeptidase secreted by stimulated rabbit synovial fibroblasts in parallel with collagenase. Biosynthesis, isolation, characterization, and substrates. *J. Biol. Chem.*, 260:12367–12376, 1985.
- [434] M.A. Chinchetru, J.A. Cabezas, and P. Calvo. Characterization and kinetics of  $\beta$ -D-gluco/fuco/galactosidase from sheep liver. *Comp. Biochem. Physiol. B*, 75:719–728, 1983.
- [435] M.A. Chinchetru, J.A. Cabezas, and P. Calvo. Purification and characterization of a broad specificity  $\beta$ -glucosidase from sheep liver. *Int. J. Biochem.*, 21:469–476, 1989.
- [436] J. Cho, J.S. King, X. Qian, A.J. Harwood, and S.B. Shears. Dephosphorylation of 2,3-bisphosphoglycerate by MIPP expands the regulatory capacity of the Rapoport-Luebering glycolytic shunt. *Proc. Natl. Acad. Sci. USA*, 105:5998–6003, 2008.
- [437] K. Cho, T. Tanaka, R.R. Cook, W. Kisiel, K. Fujikawa, K. Kurachi, and J.C. Powers. Active-site mapping of bovine and human blood coagulation serine proteases using synthetic peptide 4-nitroanilide and thio ester substrates. *Biochemistry*, 23:644–650, 1984.
- [438] K.H. Choi and R.A. Laursen. Amino-acid sequence and glycan structures of cysteine proteases with proline specificity from ginger rhizome *Zingiber officinale*. *Eur. J. Biochem.*, 267:1516–1526, 2000.
- [439] K.H. Choi, R.A. Laursen, and K.N. Allen. The 2.1 Å structure of a cysteine protease with proline specificity from ginger rhizome, *Zingiber officinale*. *Biochemistry*, 38:11624–11633, 1999.
- [440] W.K. Chou, S. Hinderlich, W. Reutter, and M.E. Tanner. Sialic acid biosynthesis: stereochemistry and mechanism of the reaction catalyzed by the mammalian UDP-*N*-acetylglucosamine 2-epimerase. *J. Am. Chem. Soc.*, 125:2455–2461, 2003.
- [441] J.T. Christeller and N.E. Tolbert. Phosphoglycolate phosphatase. Purification and properties. *J. Biol. Chem.*, 253:1780–1785, 1978.
- [442] M. Christen, B. Christen, M. Folcher, A. Schauerte, and U. Jenal. Identification and characterization of a cyclic di-GMP-specific phosphodiesterase and its allosteric control by GTP. *J. Biol. Chem.*, 280:30829–30837, 2005.
- [443] A. Christensson, C.-B. Laurell, and H. Lilja. Enzymatic activity of prostate-specific antigen and its reactions with extracellular serine proteinase inhibitors. *Eur. J. Biochem.*, 194:755–763, 1990.
- [444] S. Christoforidis, T. Papamarcaki, D. Galaris, R. Kellner, and O. Tsolas. Purification and properties of human placental ATP diphosphohydrolase. *Eur. J. Biochem.*, 234:66–74, 1995.
- [445] P. Chuankhayan, Y. Hua, J. Svasti, S. Sakdarat, P.A. Sullivan, and J.R. Ketudat Cairns. Purification of an isoflavonoid 7-*O*- $\beta$ -apiosyl-glucoside  $\beta$ -glycosidase and its substrates from *Dalbergia nigrescens* Kurz. *Phytochemistry*, 66:1880–1889, 2005.
- [446] T.M. Chulkova and V.N. Orekhovich. Isolation and properties of aminopeptidase A from bovine kidneys. *Biokhimiya*, 43:964–969, 1978.
- [447] C.H. Chung and A.L. Goldberg. Purification and characterization of protease So, a cytoplasmic serine protease in *Escherichia coli*. *J. Bacteriol.*, 154:231–238, 1983.
- [448] J.H. Chung, J.H. Back, Y.I. Park, and Y.S. Han. Biochemical characterization of a novel hypoxanthine/xanthine dNTP pyrophosphatase from *Methanococcus jannaschii*. *Nucleic Acids Res.*, 29:3099–3107, 2001.
- [449] J.H. Chung, H.Y. Park, J.H. Lee, and Y. Jang. Identification of the dITP- and XTP-hydrolyzing protein from *Escherichia coli*. *J. Biochem. Mol. Biol.*, 35:403–408, 2002.
- [450] S.-T. Chung and K. Aida. Purification and properties of ATP deaminase from *Microsporium audouini*. *J. Biochem. (Tokyo)*, 61:1–9, 1967.

- [451] G. Ciafardini, V. Marsilio, B. Lanza, and N. Pozzi. Hydrolysis of oleuropein by *Lactobacillus plantarum* strains associated with olive fermentation. *Appl. Environ. Microbiol.*, 60:4142–4147, 1994.
- [452] M.A. Cicilini, M.J.F. Ribeiro, E.B. de Oliveira, R.A. Mortara, and A.C.M. de Camargo. Endooligopeptidase A activity in rabbit heart: generation of enkephalin from enkephalin containing peptides. *Peptides (Fayetteville)*, 9:945–955, 1988.
- [453] G. Cilliv and P.T. Ozand. Human erythrocyte acetylcholinesterase purification, properties and kinetic behavior. *Biochim. Biophys. Acta*, 284:136–156, 1972.
- [454] N. Citri. Penicillinase and other  $\beta$ -lactamases. In P.D. Boyer, editor, *The Enzymes*, volume 4, pages 23–46. Academic Press, New York, 3rd edition, 1971.
- [455] D.J. Civello, H.L. Duong, and C.R. Geren. Isolation and characterization of a hemorrhagic proteinase from timber rattlesnake venom. *Biochemistry*, 22:749–755, 1983.
- [456] D.J. Civello, J.B. Moran, and C.R. Geren. Substrate specificity of a hemorrhagic proteinase from timber rattlesnake venom. *Biochemistry*, 22:755–762, 1983.
- [457] C. Clamagirand, C. Creminon, C. Fahy, H. Boussetta, and P. Cohen. Partial purification and functional properties of an endoprotease from bovine neurosecretory granules cleaving proocytosin/neurophysin peptides at the basic amino acid doublet. *Biochemistry*, 26:6018–6023, 1987.
- [458] N.E. Clark and S.C. Garman. The 1.9 Å structure of human  $\alpha$ -N-acetylgalactosaminidase: The molecular basis of Schindler and Kanzaki diseases. *J. Mol. Biol.*, 393:435–447, 2009.
- [459] N. Clarke and R.M.C. Dawson. *rac*-Glycerol 1:2-cyclic phosphate 2-phosphodiesterase, a new soluble phosphodiesterase of mammalian tissues. *Biochem. J.*, 173:579–589, 1978.
- [460] P.H. Clarke. The aliphatic amidases of *Pseudomonas aeruginosa*. *Adv. Microb. Physiol.*, 4:179–222, 1970.
- [461] P.P. Cleary, U. Prahbu, J.B. Dale, D.E. Wexler, and J. Handley. Streptococcal C5a peptidase is a highly specific endopeptidase. *Infect. Immun.*, 60:5219–5223, 1992.
- [462] C. Cobzaru, P. Ganas, M. Mihasan, P. Schleberger, and R. Brandsch. Homologous gene clusters of nicotine catabolism, including a new  $\omega$ -amidase for  $\alpha$ -ketoglutaramate, in species of three genera of Gram-positive bacteria. *Res. Microbiol.*, 162:285–291, 2011.
- [463] J.A. Cohen and M.G.P.J. Warringa. Purification and properties of dialkylfluorophosphatase. *Biochim. Biophys. Acta*, 26:29–39, 1957.
- [464] L.S. Cohen, C. Mikhli, C. Friedman, M. Jankowska-Anyszka, J. Stepinski, E. Darzynkiewicz, and R.E. Davis. Nematode m<sup>7</sup>GpppG and m<sup>3</sup>(2,2,7)GpppG decapping: activities in *Ascaris* embryos and characterization of *C. elegans* scavenger DcpS. *RNA*, 10:1609–1624, 2004.
- [465] S.S. Cohen and H.D. Barner. The conversion of 5-methyldeoxycytidine to thymidine in vitro and in vivo. *J. Biol. Chem.*, 226:631–642, 1957.
- [466] W. Colilla, R.A. Jorgenson, and R.C. Nordlie. Mammalian carbamyl phosphate : glucose phosphotransferase and glucose-6-phosphate phosphohydrolase: extended tissue distribution. *Biochim. Biophys. Acta*, 377:117–117, 1975.
- [467] D. Collen, H. R. Lijnen, and M. Verstraete. The fibrinolytic system and its disorders. In R.I. Handin, S.E. Lux, and J.P. Stossel, editors, *Blood: Principles and Practice of Hematology*. J.B.Lippincott Company, Philadelphia, 2nd edition, 1990.
- [468] I.E. Collier, S.M. Wilhelm, A.Z. Eisen, B.L. Marmer, G.A. Grant, J.L. Seltzer, A. Kronberger, C. He, E.A. Bauer, and G.I. Goldberg. H-ras oncogene-transformed human bronchial epithelial cells (TBE-1) secrete a single metalloprotease capable of degrading basement membrane collagen. *J. Biol. Chem.*, 263:6579–6587, 1988.
- [469] D.G. Comb, , and S. Glucosamine metabolism. IV. Glucosamine-6-phosphate deaminase. *J. Biol. Chem.*, 232:807–827, 1958.
- [470] D. Comfort and R.T. Clubb. A comparative genome analysis identifies distinct sorting pathways in gram-positive bacteria. *Infect. Immun.*, 72:2710–2722, 2004.

- [471] J. Conchie.  $\beta$ -Glucosidase from rumen liquor. Preparation, assay and kinetics of action. *Biochem. J.*, 58:552–560, 1954.
- [472] F. Confalonieri and M. Duguet. A 200-amino acid ATPase module in search of a basic function. *BioEssays*, 17:639–650, 1995.
- [473] K.A. Conklin, K.M. Yamashiro, and G.M. Gray. Human intestinal sucrase-isomaltase. Identification of free sucrase and isomaltase and cleavage of the hybrid into active distinct subunits. *J. Biol. Chem.*, 250:5735–5741, 1975.
- [474] C.A. Conlin and C.G. Miller. Oligopeptidase A and peptidyl-dipeptidase of *Escherichia* and *Salmonella*. *Methods Enzymol.*, 248:567–579, 1995.
- [475] C.A. Conlin, N.J. Trun, T.J. Silhavy, and C.G. Miller. *Escherichia coli prlC* encodes an endopeptidase and is homologous to the *Salmonella typhimurium opdA* gene. *J. Bacteriol.*, 174:5881–5887, 1992.
- [476] C.A. Conlin, E.R. Vimr, and C.G. Miller. Oligopeptidase A is required for normal phage P22 development. *J. Bacteriol.*, 174:5869–5880, 1992.
- [477] T.D. Connell, D.J. Metzger, J. Lynch, and J.P. Folster. Endochitinase is transported to the extracellular milieu by the eps-encoded general secretory pathway of *Vibrio cholerae*. *J. Bacteriol.*, 180:5591–5600, 1998.
- [478] G.E. Conner. Isolation of procathepsin D from mature cathepsin D by pepstatin affinity chromatography. Autocatalytic proteolysis of the zymogen form of the enzyme. *Biochem. J.*, 263:601–604, 1989.
- [479] T. Connolly and R. Gilmore. The signal recognition particle receptor mediates the GTP-dependent displacement of SRP from the signal sequence of the nascent polypeptide. *Cell*, 57:599–610, 1989.
- [480] T. Connolly, P.J. Rapiejko, and R. Gilmore. Requirement of GTP hydrolysis for dissociation of the signal recognition particle from its receptor. *Science*, 252:1171–1173, 1991.
- [481] T.M. Connolly, V.S. Bansal, T.E. Bross, R.F. Irvine, and P.W. Majerus. The metabolism of tris- and tetraphosphates of inositol by 5-phosphomonoesterase and 3-kinase enzymes. *J. Biol. Chem.*, 262:2146–2149, 1987.
- [482] W.M. Connors and E. Stotz. The purification and properties of a triacetic acid-hydrolyzing enzyme. *J. Biol. Chem.*, 178:881–890, 1949.
- [483] A. Contardi and A. Ercoli. The enzymic cleavage of lecithin and lysolecithin. *Biochem. Z.*, 261:275–302, 1933.
- [484] A.M. Cook, P. Beilstein, H. Grossenbacher, and R. Hutter. Ring cleavage and degradative pathway of cyanuric acid in bacteria. *Biochem. J.*, 231:25–30, 1985.
- [485] C. Cooper and D.W. Wilson. Biosynthesis of pyrimidines. *Fed. Proc.*, 13:194–194, 1954.
- [486] J. Cooper, S. Foundling, A. Hemmings, T. Blundell, D.M. Jones, A. Hallett, and M. Szelke. The structure of a synthetic pepsin inhibitor complexed with endothiapepsin. *Eur. J. Biochem.*, 169:215–221, 1987.
- [487] O. Cordin, N.K. Tanner, M. Doere, P. Linder, and J. Banroques. The newly discovered Q motif of DEAD-box RNA helicases regulates RNA-binding and helicase activity. *EMBO J.*, 23:2478–2487, 2004.
- [488] R.S. Cormack, J.L. Genereaux, and G.A. Mackie. RNase E activity is conferred by a single polypeptide: overexpression, purification, and properties of the *ams/rne/hmp1* gene product. *Proc. Natl. Acad. Sci. USA*, 90:9006–9010, 1993.
- [489] G. Correc, J.H. Hehemann, M. Czjzek, and W. Helbert. Structural analysis of the degradation products of porphyrin digested by *Zobellia galactanivorans*  $\beta$ -porphyrinase A. *Carbohydrate Polymers*, 83:277–283, 2011.
- [490] P. Corvol, T.A. Williams, and F. Soubrier. Peptidyl dipeptidase A: angiotensin I-converting enzyme. *Methods Enzymol.*, 248:283–305, 1995.
- [491] D.J. Cosgrove. Ion-exchange chromatography of inositol polyphosphates. *Ann. N.Y. Acad. Sci.*, 165:677–686, 1969.
- [492] D.J. Cosgrove. In *Inositol Phosphates: Their Chemistry, Biochemistry, and Physiology*. Elsevier, Amsterdam, 1980.
- [493] J. Costa, N. Empadinhas, and M.S. da Costa. Glucosylglycerate biosynthesis in the deepest lineage of the bacteria: characterization of the thermophilic proteins GpgS and GpgP from *Persephonella marina*. *J. Bacteriol.*, 189:1648–1654, 2007.

- [494] J. Costa, N. Empadinhas, L. Goncalves, P. Lamosa, H. Santos, and M.S. da Costa. Characterization of the biosynthetic pathway of glucosylglycerate in the archaeon *Methanococoides burtonii*. *J. Bacteriol.*, 188:1022–1030, 2006.
- [495] C. Costello, A. Kreuzman, , and M. Selective deprotection of phthalyl protected proteins. *Tetrahedron Lett.*, 37:7469–7472, 1996.
- [496] N. Côté, A. Fleury, E. Dumont-Blanchette, T. Fukamizo, M. Mitsutomi, and R. Brzezinski. Two exo- $\beta$ -D-glucosaminidases/exochitosanases from actinomycetes define a new subfamily within family 2 of glycoside hydrolases. *Biochem. J.*, 394:675–686, 2006.
- [497] D. Court. RNA processing and degradation by RNase III in control of mRNA stability. In J.G. Belasco and G. Brawerman, editors, *Control of Messenger RNA Stability*, pages 71–116. Academic Press, New York, 1993.
- [498] M.A. Cousin, S. Notermans, P. Hoogerhout, and J.H. Van Boom. Detection of  $\beta$ -galactofuranosidase production by *Penicillium* and *Aspergillus* species using 4-nitrophenyl  $\beta$ -D-galactofuranoside. *J. Appl. Bacteriol.*, 66:311–317, 1989.
- [499] O. Coux, K. Tanaka, and A.L. Goldberg. Structure and functions of the 20S and 26S proteasomes. *Annu. Rev. Biochem.*, 65:801–847, 1996.
- [500] V. Cowling and J. Downward. Caspase-6 is the direct activator of caspase-8 in the cytochrome *c*-induced apoptosis pathway: absolute requirement for removal of caspase-6 prodomain. *Cell Death Differ.*, 9:1046–1056, 2002.
- [501] A. Craxton, J.J. Caffrey, W. Burkhart, S.T. Safrany, and S.B. Shears. Molecular cloning and expression of a rat hepatic multiple inositol polyphosphate phosphatase. *Biochem. J.*, 328:75–81, 1997.
- [502] E.V. Crean. Synthesis and degradation of dolichyl phosphoryl glucose by the cellular slime mold, *Dictyostelium discoideum*. *Biochim. Biophys. Acta*, 792:149–157, 1984.
- [503] C. Créminon, M. Rholam, H. Boussetta, N. Marrakchi, and P. Cohen. Synthetic peptide substrates as models to study a pro-oxytocin/neurophysin converting enzyme. *J. Chromatogr.*, 440:439–448, 1988.
- [504] A.M.V. Crescenzi, K.S. Dodgson, and G.F. White. Purification and some properties of the D-lactate-2-sulphatase of *Pseudomonas syringae* GG. *Biochem. J.*, 223:487–494, 1984.
- [505] H.J. Crombie, S. Chengappa, C. Jarman, C. Sidebottom, and J.S. Reid. Molecular characterisation of a xyloglucan oligosaccharide-acting  $\alpha$ -D-xylosidase from nasturtium (*Tropaeolum majus* L.) cotyledons that resembles plant ‘apoplastic’  $\alpha$ -D-glucosidases. *Planta*, 214:406–413, 2002.
- [506] J.A. Cromlish, N.G. Seidah, M. Marcinkiewicz, J. Hamelin, D.A. Johnson, and M. Chrétien. Human pituitary tryptase: molecular forms, NH<sub>2</sub>-terminal sequence, immunocytochemical localization, and specificity with prohormone and fluorogenic substrates. *J. Biol. Chem.*, 262:1363–1373, 1987.
- [507] A. Cronin, S. Mowbray, H. Dürk, S. Homburg, I. Fleming, B. Fisslthaler, F. Oesch, and M. Arand. The N-terminal domain of mammalian soluble epoxide hydrolase is a phosphatase. *Proc. Natl. Acad. Sci. USA*, 100:1552–1557, 2003.
- [508] G.P. Crooks, , and S.D. Purification and characterization of 4-chlorobenzoyl CoA dehalogenase from *Arthrobacter* sp. strain 4-CB1. *Biochemistry*, 33:11645–11649, 1994.
- [509] L.G. Crossley. C3b inactivator and  $\beta$ 1H. *Methods Enzymol.*, 80:112–124, 1981.
- [510] R. Croteau and F. Karp. Biosynthesis of monoterpenes: hydrolysis of bornyl pyrophosphate, an essential step in camphor biosynthesis, and hydrolysis of geranyl pyrophosphate, the acyclic precursor of camphor, by enzymes from sage (*Salvia officinalis*). *Arch. Biochem. Biophys.*, 198:523–532, 1979.
- [511] R.J. Crouch. Ribonuclease 3 does not degrade deoxyribonucleic acid-ribonucleic acid hybrids. *J. Biol. Chem.*, 249:1314–1316, 1974.
- [512] H. Cudny, R. Zaniewski, and M.P. Deutscher. *Escherichia coli* RNase D. Catalytic properties and substrate specificity. *J. Biol. Chem.*, 256:5633–5637, 1981.
- [513] H. Cudny, R. Zaniewski, and M.P. Deutscher. *Escherichia coli* RNase D. Purification and structural characterization of a putative processing nuclease. *J. Biol. Chem.*, 256:5627–5632, 1981.



- [514] Z. Cui, Y. Maruyama, B. Mikami, W. Hashimoto, and K. Murata. Crystal structure of glycoside hydrolase family 78  $\alpha$ -L-Rhamnosidase from *Bacillus* sp. GL1. *J. Mol. Biol.*, 374:384–398, 2007.
- [515] P.J. Cullen, R.F. Irvine, B.J. Drøbak, and A.P. Dawson. Inositol 1,3,4,5-tetrakisphosphate causes release of  $\text{Ca}^{2+}$  from permeabilized mouse lymphoma L1210 cells by its conversion into inositol 1,4,5-trisphosphate. *Biochem. J.*, 259:931–933, 1989.
- [516] J.A. Cummings, M. Vetting, S.V. Ghodge, C. Xu, B. Hillerich, R.D. Seidel, S.C. Almo, and F.M. Raushel. Prospecting for unannotated enzymes: discovery of a 3',5'-nucleotide bisphosphate phosphatase within the amidohydrolase superfamily. *Biochemistry*, 53:591–600, 2014.
- [517] R. Cunin, N. Glansdorff, A. Pierard, and V. Stalon. Biosynthesis and metabolism of arginine in bacteria. *Microbiol. Rev.*, 50:314–352, 1986.
- [518] R. Cunin, N. Glansdorff, A. Pierard, and V. Stalon. Erratum report: Biosynthesis and metabolism of arginine in bacteria. *Microbiol. Rev.*, 51:178–178, 1987.
- [519] L.W. Cunningham and D.J. Manners. Enzymic degradation of lichenin. *Biochem. J.*, 80:42–42, 1961.
- [520] R.P. Cunningham, S.M. Saporito, S.G. Spitzer, and B. Weiss. Endonuclease IV (nfo) mutant of *Escherichia coli*. *J. Bacteriol.*, 168:1120–1127, 1986.
- [521] E. Curti, S.J. Smerdon, and E.O. Davis. Characterization of the helicase activity and substrate specificity of *Mycobacterium tuberculosis* UvrD. *J. Bacteriol.*, 189:1542–1555, 2007.
- [522] M.A. Curtis, O.J. Aduse, M. Rangarajan, A. Gallagher, J.A. Sterne, C.R. Reid, H.E. Evans, and B. Samuelsson. Attenuation of the virulence of *Porphyromonas gingivalis* by using a specific synthetic Kgp protease inhibitor 2. *Infect. Immun.*, 70:6968–6975, 2002.
- [523] F. Cuskin, A. Basle, S. Ladeveze, A.M. Day, H.J. Gilbert, G.J. Davies, G. Potocki-Veronese, and E.C. Lowe. The GH130 family of mannoside phosphorylases contains glycoside hydrolases that target  $\beta$ -1,2-mannosidic linkages in *Candida* mannan. *J. Biol. Chem.*, 290:25023–25033, 2015.
- [524] F. Cuskin, E.C. Lowe, M.J. Temple, Y. Zhu, E.A. Cameron, N.A. Pudlo, N.T. Porter, K. Urs, A.J. Thompson, A. Cartmell, A. Rogowski, B.S. Hamilton, R. Chen, T.J. Tolbert, K. Piens, D. Bracke, W. Verveckens, Z. Hakki, G. Speciale, J.L. Munoz-Munoz, A. Day, M.J. Pena, R. McLean, M.D. Suits, A.B. Boraston, T. Atherly, C.J. Ziemer, S.J. Williams, G.J. Davies, D.W. Abbott, E.C. Martens, and H.J. Gilbert. Human gut Bacteroidetes can utilize yeast mannan through a selfish mechanism. *Nature*, 517:165–169, 2015.
- [525] M. Czjzek, M. Cicek, V. Zamboni, D.R. Bevan, B. Henrissat, and A. Esen. The mechanism of substrate (aglycone) specificity in  $\beta$ -glucosidases is revealed by crystal structures of mutant maize  $\beta$ -glucosidase-DIMBOA, -DIMBOAGlc, and -dhurrin complexes. *Proc. Natl. Acad. Sci. USA*, 97:13555–13560, 2000.
- [526] T. DaCosta and R.J. Gibbons. Hydrolysis of levan by human plaque streptococci. *Arch. Oral Biol.*, 13:609–617, 1968.
- [527] G.S. Dahler, F. Barras, and N.T. Keen. Cloning of genes encoding extracellular metalloproteases from *Erwinia chrysanthemi* EC16. *J. Bacteriol.*, 172:5803–5815, 1990.
- [528] N. Dahlmann. Human serum thymidine triphosphate nucleotidohydrolase: purification and properties of a new enzyme. *Biochemistry*, 21:6634–6639, 1982.
- [529] A. Dahlqvist. Pig intestinal  $\beta$ -glucosidase activities. I. Relation to  $\beta$ -galactosidase (lactase). *Biochim. Biophys. Acta*, 50:55–61, 1961.
- [530] L.S. Daley and G.A. Strobel.  $\beta$ -Galactofuranosidase activity in *Helminthosporium sacchari* and its relationship to the production of helminthosporoside. *Plant Sci. Lett.*, 30:145–154, 1983.
- [531] J.P. Dalton and P.J. Brindley. Schistosome legumain. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 749–754. Academic Press, London, 1998.
- [532] J.B. Dame, G.R. Reddy, C.A. Yowell, B.M. Dunn, J. Kay, and C. Berry. Sequence, expression and modelled structure of an aspartic proteinase from the human malaria parasite *Plasmodium falciparum*. *Mol. Biochem. Parasitol.*, 64:177–190, 1994.

- [533] J.E. Damen, L. Liu, P. Rosten, R.K. Humphries, A.B. Jefferson, P.W. Majerus, and G. Krystal. The 145-kDa protein induced to associate with Shc by multiple cytokines is an inositol tetrakisphosphate and phosphatidylinositol 3,4,5-triphosphate 5-phosphatase. *Proc. Natl. Acad. Sci. USA*, 93:1689–1693, 1996.
- [534] W. Dangel, A. Tschech, and G. Fuchs. Enzyme-reactions involved in anaerobic cyclohexanol metabolism by a denitrifying *Pseudomonas* species. *Arch. Microbiol.*, 152:273–279, 1989.
- [535] A. D’Aniello and L. Strazullo. Peptidyl-D-amino acid hydrolase from *Loligo vulgaris* Lam. *J. Biol. Chem.*, 259:4237–4243, 1984.
- [536] E.M. Danielsen, O. Norén, H. Sjöström, J. Ingram, and J. Kenny. Proteins of the kidney microvillar membrane. Aspartate aminopeptidase: purification by immunoabsorbent chromatography and properties of the detergent- and proteinase-solubilized forms. *Biochem. J.*, 189:591–603, 1980.
- [537] P.L. Darke. *Herpesvirus assemblin*. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 470–472. Academic Press, London, 1998.
- [538] U. Das, M. Chauleau, H. Ordonez, and S. Shuman. Impact of DNA3’pp5’G capping on repair reactions at DNA 3’ ends. *Proc. Natl. Acad. Sci. USA*, 111:11317–11322, 2014.
- [539] H. DasGupta and D.P. Fan. Purification and characterization of a carboxypeptidase-transpeptidase of *Bacillus megaterium* acting on the tetrapeptide moiety of the peptidoglycan. *J. Biol. Chem.*, 254:5672–5683, 1979.
- [540] P.K. Datta, K.R. Hanson, and D.R. Whitaker. Improved procedures for preparation and characterization of *Myrothecium* cellulase. III. Molecular weight, amino acid composition, terminal residues, and other properties. *Can. J. Biochem. Physiol.*, 41:697–705, 1963.
- [541] D. Dauvillee, I.S. Kinderf, Z. Li, B. Kosar-Hashemi, M.S. Samuel, L. Rampling, S. Ball, and M.K. Morell. Role of the *Escherichia coli* *glgX* gene in glycogen metabolism. *J. Bacteriol.*, 187:1465–1473, 2005.
- [542] J.F. Davey and D.W. Ribbons. Metabolism of resorcinylic compounds by bacteria. Purification and properties of acetylpyruvate hydrolase from *Pseudomonas putida* 01. *J. Biol. Chem.*, 250:3826–3830, 1975.
- [543] F. David, A.-M. Bernard, M. Pierres, and D. Marguet. Identification of serine 624, aspartic acid 702, and histidine 734 as the catalytic triad residues of mouse dipeptidyl-peptidase IV (CD26). A member of a novel family of nonclassical serine hydrolases. *J. Biol. Chem.*, 268:17247–17252, 1993.
- [544] V.A. David, A.H. Deutch, A. Sloma, D. Pawlyk, A. Ally, and D.R. Durham. Cloning, sequencing and expression of the gene encoding the extracellular neutral protease, vibriolysin, of *Vibrio proteolyticus*. *Gene*, 112:107–112, 1992.
- [545] R. Davidson, A. Gertler, and T. Hofmann. *Aspergillus oryzae* acid proteinase. Purification and properties, and formation of  $\pi$ -chymotrypsin. *Biochem. J.*, 147:45–53, 1975.
- [546] E.W. Davie, K. Fujikawa, K. Kurachi, and W. Kisiel. The role of serine proteases in the blood coagulation cascade. *Adv. Enzymol.*, 48:277–318, 1979.
- [547] H.M. Davies, L. Anderson, C. Fan, , and D.J. Developmental induction, purification, and further characterization of 12:0-ACP thioesterase from immature cotyledons of *Umbellularia californica*. *Arch. Biochem. Biophys.*, 290:37–45, 1991.
- [548] J.F. Davies, Hostomska 2nd, Hostomsky Z., Jordan Z., Matthews S.R., and D.A. Crystal structure of the ribonuclease H domain of HIV-1 reverse transcriptase. *Science*, 252:88–95, 1991.
- [549] N.C. Davis and E.L. Smith. Purification and some properties of prolidase of swine kidney. *J. Biol. Chem.*, 224:261–275, 1957.
- [550] R.M.C. Dawson. Liver glycerylphosphorylcholine diesterase. *Biochem. J.*, 62:689–693, 1956.
- [551] R.M.C. Dawson. Studies on the hydrolysis of lecithin by *Penicillium notatum* phospholipase B preparation. *Biochem. J.*, 70:559–570, 1958.
- [552] R.M.C. Dawson and N.G. Clarke. D-myoInositol 1:2-cyclic phosphate 2-phosphohydrolase. *Biochem. J.*, 127:113–118, 1972.

- [553] R.M.C. Dawson and N.G. Clarke. A comparison of D-inositol 1:2-cyclic phosphate 2-phosphohydrolase with other phosphodiesterases of kidney. *Biochem. J.*, 134:59–67, 1973.
- [554] R.M.C. Dawson and N. Hemington. A phosphodiesterase in rat kidney cortex that hydrolyses glycerylphosphorylinositol. *Biochem. J.*, 162:241–245, 1977.
- [555] R.M.C. Dawson, N. Hemington, D.E. Richards, and R.F. Irvine. *sn*-Glycero(3)phosphoinositol glycerophosphohydrolase. A new phosphodiesterase in rat tissues. *Biochem. J.*, 182:39–49, 1979.
- [556] R.M.C. Dawson and W. Thompson. The triphosphoinositide phosphomonoesterase of brain tissue. *Biochem. J.*, 91:244–250, 1964.
- [557] M. Privat de Garilhe and M. Laskowski. Study of the enzymatic degradation of deoxyribonucleic acid by two different deoxyribonucleodepolymerases. *J. Biol. Chem.*, 215:269–276, 1955.
- [558] C.A.D. de Kort and N.A. Granger. Regulation of the juvenile hormone titer. *Annu. Rev. Entomol.*, 26:1–28, 1981.
- [559] G. de la Haba and G.L. Cantoni. The enzymatic synthesis of *S*-adenosyl-L-homocysteine from adenosine and homocysteine. *J. Biol. Chem.*, 234:603–608, 1959.
- [560] A.J. de Ruijter, A.H. van Gennip, H.N. Caron, S. Kemp, and A.B. van Kuilenburg. Histone deacetylases (HDACs): characterization of the classical HDAC family. *Biochem. J.*, 370:737–749, 2003.
- [561] M.L. de Souza, M.J. Sadowsky, and L.P. Wackett. Atrazine chlorohydrolase from *Pseudomonas* sp. strain ADP: gene sequence, enzyme purification, and protein characterization. *J. Bacteriol.*, 178:4894–4900, 1996.
- [562] M.L. de Souza, L.P. Wackett, K.L. Boundy-Mills, R.T. Mandelbaum, and M.J. Sadowsky. Cloning, characterization, and expression of a gene region from *Pseudomonas* sp. strain ADP involved in the dechlorination of atrazine. *Appl. Environ. Microbiol.*, 61:3373–3378, 1995.
- [563] W.J.M. Van de Ven, J.W.M. Creemers, and A.J.M. Roebroek. Furin: the prototype mammalian subtilisin-like proprotein-processing enzyme. Endoproteolytic cleavage at paired basic residues of proproteins of the eukaryotic secretory pathway. *Enzyme*, 45:257–270, 1991.
- [564] W.J.M. Van de Ven, J. Voorberg, R. Fontijn, H. Pannekoek, A.M.W. van den Ouweland, H.L.P. van Duijnhoven, A.J.M. Roebroek, and R.J. Siezen. Furin is a subtilisin-like proprotein processing enzyme in higher eukaryotes. *Mol. Biol. Rep.*, 14:265–275, 1990.
- [565] C.J. Decedue, E.A. Broussard, L. arson II, Braymer A.D., and H.D. Purification and characterization of the extracellular proteinase of *Serratia marcescens*. *Biochim. Biophys. Acta*, 569:293–301, 1979.
- [566] P.A. Deddish, R.A. Skidgel, and E.G. Erdös. Enhanced  $\text{Co}^{2+}$  activation and inhibitor binding of carboxypeptidase M at low pH. *Biochem. J.*, 261:289–291, 1989.
- [567] M.A. Deeg, E.L. Vierman, and M.C. Cheung. GPI-specific phospholipase D associates with an apoA-I- and apoA-IV-containing complex. *J. Lipid Res.*, 42:442–451, 2001.
- [568] R. Deeg, A. Roeder, J. Siedel, H. Gauhl, and J. Ziegenhorn. Process and reagent for the determination of *N*-carbamoylsarcosine with the use of a new enzyme, 1984.
- [569] P. Dehm and A. Nordwig. The cleavage of prolyl peptides by kidney peptidases. Isolation of a microsomal carboxypeptidase from swine kidney. *Eur. J. Biochem.*, 17:372–377, 1970.
- [570] E.E. Dekker, M.J. Schlesinger, and M.J. Coon.  $\beta$ -Hydroxy- $\beta$ -methylglutaryl coenzyme A deacetylase. *J. Biol. Chem.*, 233:434–438, 1958.
- [571] N. Dekker. Omptin. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 212–216. Elsevier, London, 2nd edition, 2004.
- [572] M.M. Sanchez del Pino, G. Lopez-Rodas, R. Sendra, and V. Tordera. Properties of the yeast nuclear histone deacetylase. *Biochem. J.*, 303:723–729, 1994.
- [573] F.P. Delafield, K.E. Cooksey, and M. Doudoroff.  $\beta$ -Hydroxybutyric dehydrogenase and dimer hydrolase of *Pseudomonas lemoignei*. *J. Biol. Chem.*, 240:4023–4028, 1965.

- [574] R.J. Delange and E.L. Smith. Leucine aminopeptidase and other N-terminal exopeptidases. In P.D. Boyer, editor, *The Enzymes*, volume 3, pages 81–118. Academic Press, New York, 3rd edition, 1971.
- [575] E. Delannoy, M. Le Ret, E. Faivre-Nitschke, G.M. Estavillo, M. Bergdoll, N.L. Taylor, B.J. Pogson, I. Small, P. Imbault, and J.M. Gualberto. *Arabidopsis* tRNA adenosine deaminase arginine edits the wobble nucleotide of chloroplast tRNA<sup>Arg</sup>(ACG) and is essential for efficient chloroplast translation. *Plant Cell*, 21:2058–2071, 2009.
- [576] L.T.J. Delbaere and G.D. Brayer. The 1.8 Å structure of the complex between chymostatin and *Streptomyces griseus* protease A. A model for serine protease catalytic tetrahedral intermediates. *J. Mol. Biol.*, 183:89–103, 1985.
- [577] C. Delporte, K.M. Carvalho, A.-M. Leseney, J. Winand, J. Christophe, and P. Cohen. A new metallo-endopeptidase from human neuroblastoma NB-OK-1 cells which inactivates atrial natriuretic peptide by selective cleavage at the Ser<sup>123</sup>-Phe<sup>124</sup> bond. *Biochem. Biophys. Res. Commun.*, 182:158–164, 1992.
- [578] E. DeMoll and T. Auffenberg. Purine metabolism in *Methanococcus vannielii*. *J. Bacteriol.*, 175:5754–5761, 1993.
- [579] W. Van den Ende, S. Clerens, R. Vergauwen, L. Van Riet, A. Van Laere, M. Yoshida, and A. Kawakami. Fructan 1-exohydrolases. β-(2,1)-Trimmers during graminan biosynthesis in stems of wheat? Purification, characterization, mass mapping, and cloning of two fructan 1-exohydrolase isoforms. *Plant Physiol.*, 131:621–631, 2003.
- [580] W. Van den Ende, B. De Coninck, S. Clerens, R. Vergauwen, and A. Van Laere. Unexpected presence of fructan 6-exohydrolases (6-FEHs) in non-fructan plants: characterization, cloning, mass mapping and functional analysis of a novel ‘cell-wall invertase-like’ specific 6-FEH from sugar beet (*Beta vulgaris* L.). *Plant J.*, 36:697–710, 2003.
- [581] W.J.J. Van den Tweel, J.P. Smits, R.L.H.P. Ogg, and J.A.M. de Bont. The involvement of an enantioselective transaminase in the metabolism of D-3- and D-4-hydroxyphenylglycine in *Pseudomonas putida*. *Appl. Microbiol. Biotechnol.*, 29:224–230, 1988.
- [582] J.B. Denault and G.S. Salvesen. Human caspase-7 activity and regulation by its N-terminal peptide. *J. Biol. Chem.*, 278:34042–34050, 2003.
- [583] H. Deng, S.A. McMahon, A.S. Eustaquio, B.S. Moore, J.H. Naismith, and D. O’Hagan. Mechanistic insights into water activation in SAM hydroxide adenosyltransferase (duf-62). *Chembiochem*, 10:2455–2459, 2009.
- [584] R.J. Dennis, E.J. Taylor, M.S. Macauley, K.A. Stubbs, J.P. Turkenburg, S.J. Hart, G.N. Black, D.J. Vocadlo, and G.J. Davies. Structure and mechanism of a bacterial β-glucosaminidase having O-GlcNAcase activity. *Nat. Struct. Mol. Biol.*, 13:365–371, 2006.
- [585] J.A. DePinto and L.L. Campbell. Purification and properties of the cyclodextrinase of *Bacillus macerans*. *Biochemistry*, 7:121–125, 1968.
- [586] R. Van der Geize, K. Yam, T. Heuser, M.H. Wilbrink, H. Hara, M.C. Anderton, E. Sim, L. Dijkhuizen, J.E. Davies, W.W. Mohn, and L.D. Eltis. A gene cluster encoding cholesterol catabolism in a soil actinomycete provides insight into *Mycobacterium tuberculosis* survival in macrophages. *Proc. Natl. Acad. Sci. USA*, 104:1947–1952, 2007.
- [587] M. Desautels and A.L. Goldberg. Demonstration of an ATP-dependent, vanadate-sensitive endoprotease in the matrix of rat liver mitochondria. *J. Biol. Chem.*, 257:11673–11679, 1982.
- [588] M. Desmazeaud. J. and Hermier, J. H. Spécificité de la protéase neutre de *Micrococcus caseolyticus*. *Eur. J. Biochem.*, 19:51–55, 1971.
- [589] M.J. Desmazeaud. Propriétés générales et spécificité d’action d’une endopeptidase neutre intracellulaire de *Streptococcus thermophilus*. *Biochimie*, 56:1173–1181, 1974.
- [590] M.J. Desmazeaud and C. Zevaco. General properties and substrate specificity of an intracellular neutral protease from *Streptococcus diacetilactis*. *Ann. Biol. Anim. Biochem. Biophys.*, 16:851–868, 1976.
- [591] J.J. DeStefano, L.M. Mallaber, P.J. Fay, and R.A. Bambara. Determinants of the RNase H cleavage specificity of human immunodeficiency virus reverse transcriptase. *Nucleic Acids Res.*, 21:4330–4338, 1993.
- [592] H. Deuel, R. Lewuenberger, and G. Huber. Über den enzymatischen Abbau von Carubin, dem Galaktomannan aus *Ceratonia siliqua* L. *Helv. Chim. Acta*, 33:942–946, 1950.

- [593] H. Deuel and E. Stutz. Pectic substances and pectic enzymes. *Adv. Enzymol. Relat. Areas Mol. Biol.*, 20:341–382, 1958.
- [594] A. Deutsch and R. Nilsson. On the dephosphorylation and deamination of adenosine triphosphate by actomyosin gel. *Acta Chem. Scand.*, 8:1898–1906, 1954.
- [595] J. Deutscher, U. Kessler, and W. Hengstenberg. Streptococcal phosphoenolpyruvate: sugar phosphotransferase system: purification and characterization of a phosphoprotein phosphatase which hydrolyzes the phosphoryl bond in seryl-phosphorylated histidine-containing protein. *J. Bacteriol.*, 163:1203–1209, 1985.
- [596] I.K. Dev and P.H. Ray. Signal peptidases and signal peptide hydrolases. *J. Bioenerg. Biomembr.*, 22:271–290, 1990.
- [597] R.P. deVries, Michelsen, B., Poulsen, C.H., Kroon, P.A., van den Heuvel, R.H.H., Faulds, C.B., Williamson, G., van den Homberg, J.P.T.W. and Visser, J. The *faeA* genes from *Aspergillus niger* and *Aspergillus tubingensis* encode ferulic acid esterases involved in degradation of complex cell wall polysaccharides. *Appl. Environ. Microbiol.*, 63:4638–4644, 1997.
- [598] P. M. Dey. Further characterization of  $\beta$ -L-arabinosidase from *Cajanus indicus*. *Biochim. Biophys. Acta*, 746:8–13, 1983.
- [599] P.M. Dey.  $\beta$ -L-Arabinosidase from *Cajanus indicus*: a new enzyme. *Biochim. Biophys. Acta*, 302:393–398, 1973.
- [600] M.V. Dias, F. Huang, D.Y. Chirgadze, M. Tosin, D. Spiteller, E.F. Dry, P.F. Leadlay, J.B. Spencer, and T.L. Blundell. Structural basis for the activity and substrate specificity of fluoroacetyl-CoA thioesterase FIK. *J. Biol. Chem.*, 285:22495–22504, 2010.
- [601] B.E. García Díaz, S. Gauthier, and P.L. Davies.  $\text{Ca}^{2+}$  dependency of calpain 3 (p94) activation. *Biochemistry*, 45:3714–3722, 2006.
- [602] E. Díaz, A. Ferrández, and J.L. García. Characterization of the *hca* cluster encoding the dioxygenolytic pathway for initial catabolism of 3-phenylpropionic acid in *Escherichia coli* K-12. *J. Bacteriol.*, 180:2915–2923, 1998.
- [603] E. Diaz and K.N. Timmis. Identification of functional residues in a 2-hydroxymuconic semialdehyde hydrolase. A new member of the  $\alpha/\beta$  hydrolase-fold family of enzymes which cleaves carbon-carbon bonds. *J. Biol. Chem.*, 270:6403–6411, 1995.
- [604] A.J. Dick, A.T. Matheson, and J.H. Wang. A ribosomal-bound aminopeptidase in *Escherichia coli* B: purification and properties. *Can. J. Biochem.*, 48:1181–1188, 1970.
- [605] T.P. Dick, A.K. Nussbaum, M. Deeg, W. Heinemeyer, M. Groll, M. Schirle, W. Keilholz, S. Stevanovic, D.H. Wolf, R. Huber, H.G. Rammensee, and H. Schild. Contribution of proteasomal  $\beta$ -subunits to the cleavage of peptide substrates analyzed with yeast mutants. *J. Biol. Chem.*, 273:25637–25646, 1998.
- [606] O. Dideberg, P. Charlier, G. Dive, B. Joris, J.M. Frère, and J.M. Ghuyssen. Structure of a  $\text{Zn}^{2+}$ -containing D-alanyl-D-alanine-cleaving carboxypeptidase at 2.5 Å resolution. *Nature*, 299:469–470, 1982.
- [607] B. Diderichsen and L. Christiansen. Cloning of a maltogenic  $\alpha$ -amylase from *Bacillus stearothermophilus*. *FEMS Microbiol. Lett.*, 56:53–59, 1988.
- [608] C.P. Dietrich. Enzymic degradation of heparin. A glucosaminidase and a glycuronidase from *Flavobacterium heparinum*. *Biochemistry*, 8:2089–2094, 1969.
- [609] C.P. Dietrich. Enzymic degradation of heparin. A sulphamidase and a sulphoesterase from *Flavobacterium heparinum*. *Biochem. J.*, 111:91–95, 1969.
- [610] K.J. Dietz, A. Sauter, K. Wichert, D. Messdaghi, and W. Hartung. Extracellular  $\beta$ -glucosidase activity in barley involved in the hydrolysis of ABA glucose conjugate in leaves. *J. Exp. Bot.*, 51:937–944, 2000.
- [611] A. Diez, M.I. Prieto, M.J. Alvarez, J.M. Bautista, J. Garrido, and A. Puyet. Improved catalytic performance of a 2-haloacid dehalogenase from *Azotobacter* sp. by ion-exchange immobilisation. *Biochem. Biophys. Res. Commun.*, 220:828–833, 1996.
- [612] T. Diez and J.A. Cabezas. Properties of two molecular forms of  $\beta$ -glucuronidase from the mollusc *Littorina littorea* L. *Eur. J. Biochem.*, 93:301–311, 1978.

- [613] M. Digby, X.-Y. Zhang, and R.I. Richards. Human prostate specific antigen (PSA) gene: structure and linkage to the kallikrein-like gene, hGK-1. *Nucleic Acids Res.*, 15:2137–2137, 1989.
- [614] D.A. Dillon, X. Chen, G.M. Zeimet, W.I. Wu, D.W. Waggoner, J. Dewald, D.N. Brindley, and G.M. Carman. Mammalian Mg<sup>2+</sup>-independent phosphatidate phosphatase (PAP2) displays diacylglycerol pyrophosphate phosphatase activity. *J. Biol. Chem.*, 272:10361–10366, 1997.
- [615] D.A. Dillon, W.I. Wu, B. Riedel, J.B. Wissing, W. Dowhan, and G.M. Carman. The *Escherichia coli* *pgpB* gene encodes for a diacylglycerol pyrophosphate phosphatase activity. *J. Biol. Chem.*, 271:30548–30553, 1996.
- [616] P. DiNatale and L. Ronsivalle. Identification and partial characterization of two enzyme forms of iduronate sulfatase from human placenta. *Biochim. Biophys. Acta*, 661:106–111, 1981.
- [617] J.Z. Ding, W.J. McGrath, R.M. Sweet, and W.F. Mangel. Crystal structure of the human adenovirus proteinase with its 11 residue cofactor. *EMBO J.*, 15:1778–1783, 1996.
- [618] L. Ding, A.B. Becker, A. Suzuki, and R.A. Roth. Comparison of the enzymatic and biochemical properties of human insulin-degrading enzyme and *Escherichia coli* protease III. *J. Biol. Chem.*, 267:2414–2420, 1992.
- [619] Y. Ding, H. Lapko, I. Ndamukong, Y. Xia, A. Al-Abdallat, S. Lalithambika, M. Sadler, A. Saleh, M. Fromm, J.J. Riethoven, G. Lu, and Z. Avramova. The *Arabidopsis* chromatin modifier ATX1, the myotubularin-like *AtMTM* and the response to drought. *Plant Signal Behav*, 4:1049–1058, 2009.
- [620] N.E. Dixon, C. Gazzola, R.L. Blakeley, and B. Zerner. Metal ions in enzymes using ammonia or amides. *Science*, 191:1144–1150, 1976.
- [621] A. Dmochowska, D. Dignard, D. Henning, D.Y. Thomas, and H. Bussey. Yeast KEX1 gene encodes a putative protease with a carboxypeptidase B-like function involved in killer toxin and  $\alpha$ -factor precursor processing. *Cell*, 50:573–584, 1987.
- [622] P. Dobos. The molecular biology of infectious pancreatic necrosis virus (IPNV). *Annu. Rev. Fish Dis.*, 5:25–54, 1995.
- [623] A.J.P. Docherty and G. Murphy. The tissue metalloproteinase family and the inhibitor TIMP: a study using cDNAs and recombinant proteins. *Ann. Rheum. Dis.*, 49:469–479, 1990.
- [624] K.S. Dodgson. Glycosulphatase: observations on the activity of partially purified preparations towards the sulphate esters of certain monosaccharides and steroids. *Biochem. J.*, 78:324–333, 1961.
- [625] K.S. Dodgson, B. Spencer, and K. Williams. Studies on sulphatases. 13. The hydrolysis of substituted phenyl sulphates by the arylsulphatase of *Alcaligenes metacaligenes*. *Biochem. J.*, 64:216–221, 1956.
- [626] T.L. Doering, W.J. Masteron, P.T. Englund, and G.W. Hart. Biosynthesis of the glycosyl phosphatidylinositol membrane anchor of the trypanosome variant surface glycoprotein. Origin of the non-acetylated glucosamine. *J. Biol. Chem.*, 264:11168–11173, 1989.
- [627] M. Doerr and W.H. Traub. Purification and characterization of two *Serratia marcescens* proteases. *Zentralbl. Bakteriol., Mikrobiol. Hyg. Ser. A*, 257:6–19, 1984.
- [628] H.M. Doery and J.E. Pearson. Haemolysins in venoms of Australian snakes. Observations on the haemolysins of the venoms of some Australian snakes and the separation of phospholipase A from the venom of *Pseudechis porphyriacus*. *Biochem. J.*, 78:820–827, 1961.
- [629] E. Dogru, H. Warzecha, F. Seibel, S. Haebel, F. Lottspeich, and J. Stöckigt. The gene encoding polyneuridine aldehyde esterase of monoterpene indole alkaloid biosynthesis in plants is an ortholog of the hydrolase super family. *Eur. J. Biochem.*, 267:1397–1406, 2000.
- [630] M.D. Doherty and J.F. Morrison. The hydrolysis of adenosine diphosphate ribose by a specific phosphohydrolase of rabbit-muscle extracts. *Biochim. Biophys. Acta*, 65:364–366, 1962.
- [631] A. Domnas and E.C. Catimo. The behavior of amidohydrolases and L-glutamate in synchronized populations of *Blastocladiella emeronii*. *Phytochemistry*, 4:273–284, 1965.

- [632] G. Donelli, A. Fabbri, and C. Fiorentini. *Bacteroides fragilis* enterotoxin induces cytoskeletal changes and surface blebbing in HT-29 cells. *Infect. Immun.*, 64:113–119, 1996.
- [633] D.L. Dong and G.W. Hart. Purification and characterization of an *O*-GlcNAc selective *N*-acetyl- $\beta$ -D-glucosaminidase from rat spleen cytosol. *J. Biol. Chem.*, 269:19321–19330, 1994.
- [634] J.F. Dong, J.L. Moake, L. Nolasco, A. Bernardo, W. Arceneaux, C.N. Shrimpton, A.J. Schade, L.V. McIntire, K. Fujikawa, and J.A. Lopez. ADAMTS-13 rapidly cleaves newly secreted ultralarge von Willebrand factor multimers on the endothelial surface under flowing conditions. *Blood*, 100:4033–4039, 2002.
- [635] T.C. Dong and S.M. Cutting. SpoIVB-mediated cleavage of SpoIVFA could provide the intercellular signal to activate processing of pro- $\sigma^K$  in *Bacillus subtilis*. *Mol. Microbiol.*, 49:1425–1434, 2003.
- [636] Y.H. Dong, A.R. Gusti, Q. Zhang, J.L. Xu, and L.H. Zhang. Identification of quorum-quenching *N*-acyl homoserine lactonases from *Bacillus* species. *Appl. Environ. Microbiol.*, 68:1754–1759, 2002.
- [637] Y.H. Dong, L.H. Wang, J.L. Xu, H.B. Zhang, X.F. Zhang, and L.H. Zhang. Quenching quorum-sensing-dependent bacterial infection by an *N*-acyl homoserine lactonase. *Nature*, 411:813–817, 2001.
- [638] Y.H. Dong, J.L. Xu, X.Z. Li, and L.H. Zhang. AiiA, an enzyme that inactivates the acylhomoserine lactone quorum-sensing signal and attenuates the virulence of *Erwinia carotovora*. *Proc. Natl. Acad. Sci. USA*, 97:3526–3531, 2000.
- [639] M.I. Donnelly and S. Dagley. Production of methanol from aromatic acids by *Pseudomonas putida*. *J. Bacteriol.*, 142:916–924, 1980.
- [640] M.I. Donnelly, J.C. Escalante-Semerena, K.L. Rinehart, Wolfe Jr., and R.S. Methenyl-tetrahydromethanopterin cyclohydrolase in cell extracts of *Methanobacterium*. *Arch. Biochem. Biophys.*, 242:430–439, 1985.
- [641] C. Doumeng and S. Maroux. Aminopeptidase, a cytosol enzyme from rabbit intestinal mucosa. *Biochem. J.*, 177:801–808, 1979.
- [642] C.P. Downes and R.H. Michell. The polyphosphoinositide phosphodiesterase of erythrocyte membranes. *Biochem. J.*, 198:133–140, 1981.
- [643] C.P. Downes, M.C. Mussat, and R.H. Michell. The inositol trisphosphate phosphomonoesterase of the human erythrocyte membrane. *Biochem. J.*, 203:169–177, 1982.
- [644] M.L. Doyle, P.A. Katzman, and E.A. Doisy. Production and properties of bacterial  $\beta$ -glucuronidase. *J. Biol. Chem.*, 217:921–930, 1955.
- [645] D.I. Draganov, J.F. Teiber, A. Speelman, Y. Osawa, R. Sunahara, and B.N. La Du. Human paraoxonases (PON1, PON2, and PON3) are lactonases with overlapping and distinct substrate specificities. *J. Lipid Res.*, 46:1239–1247, 2005.
- [646] G.R. Drapeau. Protease from *Staphylococcus aureus*. *Methods Enzymol.*, 45:469–475, 1976.
- [647] G.R. Drapeau. Role of a metalloprotease in activation of the precursor of staphylococcal protease. *J. Bacteriol.*, 136:607–613, 1978.
- [648] G.R. Drapeau. The primary structure of staphylococcal protease. *Can. J. Biochem.*, 56:534–544, 1978.
- [649] G.R. Drapeau. Substrate specificity of a proteolytic enzyme isolated from a mutant of *Pseudomonas fragi*. *J. Biol. Chem.*, 255:839–840, 1980.
- [650] A.L. Drayer, X. Pesesse, F. De Smedt, R. Woscholski, P. Parker, and C. Erneux. Cloning and expression of a human placenta inositol 1,3,4,5-tetrakisphosphate and phosphatidylinositol 3,4,5-trisphosphate 5-phosphatase. *Biochem. Biophys. Res. Commun.*, 225:243–249, 1996.
- [651] S.L. Drees and S. Fetzner. PqsE of *Pseudomonas aeruginosa* acts as pathway-specific thioesterase in the biosynthesis of alkylquinolone signaling molecules. *Chem. Biol.*, 22:611–618, 2015.
- [652] L.A. Dreyfus and B.H. Iglewski. Purification and characterization of an extracellular protease of *Legionella pneumophila*. *Infect. Immun.*, 70:736–743, 1986.

- [653] C.C. Drinkwater, B.A. Evans, and R.I. Richards. Sequence and expression of mouse  $\gamma$ -renin. *J. Biol. Chem.*, 263:8565–8568, 1988.
- [654] G.I. Drummond, N.T. Iyer, and J. Keith. Hydrolysis of ribonucleoside 2',3'-cyclic phosphates by a diesterase from brain. *J. Biol. Chem.*, 237:3535–3539, 1962.
- [655] G.I. Drummond and J.R. Stern. Enzymes of ketone body metabolism. II. Properties of an acetoacetate-synthesizing enzyme prepared from ox liver. *J. Biol. Chem.*, 235:318–325, 1960.
- [656] K.V. Druzhinina and M.G. Kritzman. [Lecithinase from animal tissues.]. *Biokhimiya*, 17:77–81, 1952.
- [657] G. Dubin, D. Chmiel, P. Mak, M. Rakwalska, M. Rzychon, and A. Dubin. Molecular cloning and biochemical characterisation of proteases from *Staphylococcus epidermidis*. *Biol. Chem.*, 382:1575–1582, 2001.
- [658] T. Dubois, T. Kleinschmidt, A.G. Schnek, Y. Looze, and G. Braunitzer. The thiol proteinases from the latex of *Carica papaya* L. II. The primary structure of proteinase . *Biol. Chem. Hoppe-Seyler*, 369:741–754, 1988.
- [659] M. Duckworth and J.R. Turvey. The action of a bacterial agarase on agarose, porphyran and alkali-treated porphyran. *Biochem. J.*, 113:687–692, 1969.
- [660] W.C. Duckworth. Insulin degradation: mechanisms, products, and significance. *Endocrine Rev.*, 9:319–345, 1988.
- [661] W.C. Duckworth, F.G. Hamel, R. Bennett, M.P. Ryan, and R.A. Roth. Human red blood cell insulin-degrading enzyme and rat skeletal muscle insulin protease share antigenic sites and generate identical products from insulin. *J. Biol. Chem.*, 265:2984–2987, 1990.
- [662] J.A. Duerre. A hydrolytic nucleosidase acting on *S*-adenosylhomocysteine and on 5-methylthioadenosine. *J. Biol. Chem.*, 237:3737–3741, 1962.
- [663] R.B. Duff and D.M. Webley. Metabolism of 6-*O*-acetyl-D-glucopyranose and other monoacetyl-sugars by strains of *Bacillus megaterium* and other soil organisms. *Biochem. J.*, 70:520–528, 1958.
- [664] S.M.G. Duff, D.D. Lefebvre, and W.C. Plaxton. Purification and characterization of a phosphoenolpyruvate phosphatase from *Brassica nigra* suspension cells. *Plant Physiol.*, 90:734–741, 1989.
- [665] I.F. Dumitru. Study of L-arginine amidinohydrolase from vegetable origin. Purification, crystallization and molecular weight. *Acta Vitamin. Enzymol.*, 27:207–210, 1973.
- [666] D. Dunaway-Mariano, , and P.C. On the origins and functions of the enzymes of the 4-chlorobenzoate to 4-hydroxybenzoate converting pathway. *Biodegradation*, 5:259–276, 1994.
- [667] B.M. Dunn. Human immunodeficiency virus 1 retropepsin. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 919–928. Academic Press, London, 1998.
- [668] W.M. Dunne, Buckmire Jr., and F.L.A. Partial purification and characterization of a polymannuronic acid depolymerase produced by a mucoid strain of *Pseudomonas aeruginosa* isolated from a patient with cystic fibrosis. *Appl. Environ. Microbiol.*, 50:562–567, 1985.
- [669] S. Duprè and D. Cavallini. Purification and properties of pantetheinase from horse kidney. *Methods Enzymol.*, 62:262–267, 1979.
- [670] S. Duprè, R. Chiaraluce, M. Nardini, C. Cannella, G. Ricci, and D. Cavallini. Continuous spectrophotometric assay of pantetheinase activity. *Anal. Biochem.*, 142:175–181, 1984.
- [671] P. Dutt, C.N. Spriggs, P.L. Davies, Z. Jia, and J.S. Elce. Origins of the difference in  $\text{Ca}^{2+}$  requirement for activation of  $\mu$ - and *m*-calpain. *Biochem. J.*, 367:263–269, 2002.
- [672] H. Dyckerhoff and R. Armbruster. Zur Kenntnis der Tannase. *Hoppe-Seyler's Z. Physiol. Chem.*, 219:38–56, 1933.
- [673] G.S. Eadie, F. Bernheim, and J.L.C. Bernheim. Metabolism of cytosine, thymidine, uracil and barbituric acid by bacterial enzymes. *J. Biol. Chem.*, 181:449–458, 1949.
- [674] D.L. Eaton, B.E. Malloy, S.P. Tsai, W. Henzel, and D. Drayna. Isolation, molecular cloning, and partial characterization of a novel carboxypeptidase B from human plasma. *J. Biol. Chem.*, 266:21833–21838, 1991.



- [675] R.W. Eaton and J.S. Karns. Cloning and analysis of *s*-triazine catabolic genes from *Pseudomonas* sp. strain NRRLB-12227. *J. Bacteriol.*, 173:1215–1222, 1991.
- [676] R.W. Eaton and J.S. Karns. Cloning and comparison of the DNA encoding ammelide aminohydrolase and cyanuric acid amidohydrolase from three *s*-triazine-degrading bacterial strains. *J. Bacteriol.*, 173:1363–1366, 1991.
- [677] W. Ebeling, N. Hennrich, M. Klockow, H. Metz, H.D. Orth, and H. Lang. Proteinase K from *Tritirachium album* Limber. *Eur. J. Biochem.*, 47:91–97, 1974.
- [678] S.W. Edwards and W.E. Knox. Homogentisate metabolism: the isomerization of maleylacetoacetate by an enzyme which requires glutathione. *J. Biol. Chem.*, 220:79–91, 1956.
- [679] F. Egami and N. Takahashi. Syntheses of adenosinesulfuric acids. *Bull. Chem. Soc. Jpn.*, 28:666–668, 1955.
- [680] T. Egelrud. Purification and preliminary characterization of stratum corneum chymotryptic enzyme: a proteinase that may be involved in desquamation. *J. Invest. Dermatol.*, 101:200–204, 1993.
- [681] T. Egelrud and T. Olivecrona. Purified bovine milk (lipoprotein) lipase: activity against lipid substrates in the absence of exogenous serum factors. *Biochim. Biophys. Acta*, 306:115–127, 1973.
- [682] M.R.W. Ehlers, E.A. Fox, D.J. Strydom, and J.F. Riordan. Molecular cloning of human testicular angiotensin-converting enzyme: the testis enzyme is identical to the C-terminal half of endothelial angiotensin-converting enzyme. *Proc. Natl. Acad. Sci. USA*, 86:7741–7745, 1989.
- [683] C.P. Ehretsmann, A.J. Carpousis, and H.M. Krisch. Specificity of *Escherichia coli* endoribonuclease RNase E: in vivo and in vitro analysis of mutants in a bacteriophage T<sub>4</sub> mRNA processing site. *Genes Dev.*, 6:149–159, 1992.
- [684] D. Eichinger, A. Bacher, M.H. Zenk, and W. Eisenreich. Quantitative assessment of metabolic flux by <sup>13</sup>C NMR analysis. Biosynthesis of anthraquinones in *Rubia tinctorum*. *J. Am. Chem. Soc.*, 121:7469–7475, 1999.
- [685] D.C. Eichler and I.R. Lehman. On the role of ATP in phosphodiester bond hydrolysis catalyzed by the recBC deoxyribonuclease of *Escherichia coli*. *J. Biol. Chem.*, 252:499–503, 1977.
- [686] E. Einset and W.L. Clark. The enzymatically catalyzed release of choline from lecithin. *J. Biol. Chem.*, 231:703–715, 1958.
- [687] Eisenberg and Jr. D-Myoinositol 1-phosphate as product of cyclization of glucose 6-phosphate and substrate for a specific phosphatase in rat testis. *J. Biol. Chem.*, 242:1375–1382, 1967.
- [688] P. Ek, B. Ek, and O. Zetterqvist. Phosphohistidine phosphatase 1 (PHPT1) also dephosphorylates phospholysine of chemically phosphorylated histone H1 and polylysine. *Ups J Med Sci*, 120:20–27, 2015.
- [689] P. Ek, G. Pettersson, B. Ek, F. Gong, J.P. Li, and O. Zetterqvist. Identification and characterization of a mammalian 14-kDa phosphohistidine phosphatase. *Eur. J. Biochem.*, 269:5016–5023, 2002.
- [690] A.M. El-Badry. Hexosediphosphatase from spinach chloroplasts. Purification, crystallization and some properties. *Biochim. Biophys. Acta*, 333:366–377, 1974.
- [691] J.S. Ellingson and W.E.M. Lands. Phospholipid reactivation of plasmalogen metabolism. *Lipids*, 3:111–120, 1968.
- [692] B.W. Elliott and C. Cohen. Isolation and characterization of a lysine-specific protease from *Pseudomonas aeruginosa*. *J. Biol. Chem.*, 261:11259–11265, 1986.
- [693] S.D. Elliott and T.-Y. Liu. Streptococcal proteinase. *Methods Enzymol.*, 19:252–261, 1970.
- [694] A. Elmoujahed, N. Gutman, M. Brillard, and F. Gauthier. Substrate specificity of two kallikrein family gene products isolated from the rat submaxillary gland. *FEBS Lett.*, 265:137–140, 1990.
- [695] A.K. Elsholz, K. Turgay, S. Michalik, B. Hessling, K. Gronau, D. Oertel, U. Mader, J. Bernhardt, D. Becher, M. Hecker, and U. Gerth. Global impact of protein arginine phosphorylation on the physiology of *Bacillus subtilis*. *Proc. Natl. Acad. Sci. USA*, 109:7451–7456, 2012.
- [696] J.J. Elting, W.W. Chen, and J. Lennarz. Characterization of a glucosidase involved in an initial step in the processing of oligosaccharide chains. *J. Biol. Chem.*, 255:2325–2331, 1980.

- [697] T. Emery. Fungal ornithine esterases: relationship to iron transport. *Biochemistry*, 15:2723–2728, 1976.
- [698] S. Emi, D.V. Myers, and G.A. Iacobucci. Purification and properties of the thermostable acid protease of *Penicillium duponti*. *Biochemistry*, 15:842–848, 1976.
- [699] S. Emi and T. Yamamoto. Purification and properties of several galactanases of *Bacillus subtilis* var. *amylosacchariticus*. *Agric. Biol. Chem.*, 36:1945–1954, 1972.
- [700] H. Emonard and J.-A. Grimaud. Matrix metalloproteinase. A review. *Cell. Mol. Biol.*, 36:131–153, 1990.
- [701] N. Empadinhas, J.D. Marugg, N. Borges, H. Santos, and M.S. da Costa. Pathway for the synthesis of mannosylglycerate in the hyperthermophilic archaeon *Pyrococcus horikoshii*. Biochemical and genetic characterization of key-enzymes. *J. Biol. Chem.*, 276:43580–43588, 2001.
- [702] A. Endo, S. Murakawa, H. Shimizu, and Y. Shiraishi. Purification and properties of collagenase from a *Streptomyces* species. *J. Biochem. (Tokyo)*, 102:163–170, 1987.
- [703] M. Endo, M. Mori, S. Akira, and T. Gotoh. C/EBP homologous protein (CHOP) is crucial for the induction of caspase-11 and the pathogenesis of lipopolysaccharide-induced inflammation. *J. Immunol.*, 176:6245–6253, 2006.
- [704] Y. Endo and K. Tsurugi. The RNA *N*-glycosidase activity of ricin A-chain. The characteristics of the enzymatic activity of ricin A-chain with ribosomes and with rRNA. *J. Biol. Chem.*, 263:8735–8739, 1988.
- [705] F.W.H.T. Eng. Dipeptidyl tetrapeptide hydrolase, a new enzyme with dipeptidyl ligase activity. *Can. J. Biochem. Cell. Biol.*, 62:516–528, 1984.
- [706] L. Engström. Studies on calf-intestinal alkaline phosphatase. I. Chromatographic purification, microheterogeneity and some other properties of the purified enzyme. *Biochim. Biophys. Acta*, 52:36–48, 1961.
- [707] B. Epel, K.O. Schafer, A. Quentmeier, C. Friedrich, and W. Lubitz. Multifrequency EPR analysis of the dimanganese cluster of the putative sulfate thiohydrolase SoxB of *Paracoccus pantotrophus*. *J. Biol. Inorg. Chem.*, 10:636–642, 2005.
- [708] D.M. Epstein and P.C. Wensink. The  $\alpha$ -lytic protease gene of *Lysobacter enzymogenes*. The nucleotide sequence predicts a large prepro-peptide with homology to *pro*-peptides of other chymotrypsin-like enzymes. *J. Biol. Chem.*, 263:16586–16590, 1988.
- [709] E.H. Epstein and M.E. Leventhal. Steroid sulfatase of human leukocytes and epidermis and the diagnosis of recessive X-linked ichthyosis. *J. Clin. Invest.*, 67:1257–1262, 1981.
- [710] T.J. Erb, L. Frerichs-Revermann, G. Fuchs, and B.E. Alber. The apparent malate synthase activity of *Rhodobacter sphaeroides* is due to two paralogous enzymes, (3*S*)-methylmalyl-coenzyme A (CoA)/ $\beta$ -methylmalyl-CoA lyase and (3*S*)-methylmalyl-CoA thioesterase. *J. Bacteriol.*, 192:1249–1258, 2010.
- [711] E.G. Erdös and R.A. Skidgel. Neutral endopeptidase 24.11 (enkephalinase) and related regulators of peptide hormones. *FASEB J.*, 3:145–151, 1989.
- [712] A.F.V. Eriksson. Purification and characterisation of a fungal  $\beta$ -mannanase. *Acta Chem. Scand.*, 22:1924–1934, 1968.
- [713] K.E. Eriksson and B. Pettersson. Extracellular enzyme system utilized by the fungus *Sporotrichum pulverulentum* (*Chrysosporium lignorum*) for the breakdown of cellulose. 3. Purification and physico-chemical characterization of an exo-1,4- $\beta$ -glucanase. *Eur. J. Biochem.*, 51:213–218, 1975.
- [714] C. Erneux, M. Lemos, B. Verjans, P. Vanderhaeghen, A. Delvaux, and J.E. Dumont. Soluble and particulate Ins(1,4,5) $P_3$ /Ins(1,3,4,5) $P_4$  5-phosphatase in bovine brain. *Eur. J. Biochem.*, 181:317–322, 1989.
- [715] A.L. Erwin and R.S. Munford. Deacylation of structurally diverse lipopolysaccharides by human acyloxyacyl hydrolase. *J. Biol. Chem.*, 265:16444–16449, 1990.
- [716] C.T. Esmon. The regulation of natural anticoagulant pathways. *Science*, 235:1348–1352, 1987.
- [717] C.T. Esmon. The roles of protein C and thrombomodulin in the regulation of blood coagulation. *J. Biol. Chem.*, 264:4743–4746, 1989.

- [718] P.J. Espenshade, D. Cheng, J.L. Goldstein, and M.S. Brown. Autocatalytic processing of site-1 protease removes propeptide and permits cleavage of sterol regulatory element-binding proteins. *J. Biol. Chem.*, 274:22795–22804, 1999.
- [719] B. Essigmann, S. Gler, R.A. Narang, D. Linke, and C. Benning. Phosphate availability affects the thylakoid lipid composition and the expression of SQD1, a gene required for sulfolipid biosynthesis in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA*, 95:1950–1955, 1998.
- [720] B. Essigmann, B.M. Hesperheide, L.A. Kuhn, and C. Benning. Prediction of the active-site structure and NAD<sup>+</sup> binding in SQD1, a protein essential for sulfolipid biosynthesis in *Arabidopsis*. *Arch. Biochem. Biophys.*, 369:30–41, 1999.
- [721] F.E. Estivariz, N.P. Birch, and Y.P. Loh. Generation of Lys- $\gamma$ 3-melanotropin from *pro*-opiomelanocortin1-77 by a bovine intermediate lobe secretory vesicle membrane-associated aspartic protease and purified *pro*-opiomelanocortin converting enzyme. *J. Biol. Chem.*, 264:17796–17801, 1989.
- [722] A.S. Eustaquio, J. Harle, J.P. Noel, and B.S. Moore. *S*-Adenosyl-L-methionine hydrolase (adenosine-forming), a conserved bacterial and archaeal protein related to SAM-dependent halogenases. *Chembiochem*, 9:2215–2219, 2008.
- [723] B.A. Evans, C.C. Drinkwater, and R.I. Richards. Mouse glandular kallikrein genes. Structure and partial sequence analysis of the kallikrein gene locus. *J. Biol. Chem.*, 262:8027–8034, 1987.
- [724] J.F. Evans, P. Dupuis, and A.W. Ford-Hutchinson. Purification and characterisation of leukotriene A<sub>4</sub> hydrolase from rat neutrophils. *Biochim. Biophys. Acta*, 840:43–50, 1985.
- [725] G. Evensen and E. Seeberg. Adaptation to alkylation resistance involves the induction of a DNA glycosylase. *Nature*, 296:773–775, 1982.
- [726] M.T. Everitt and H. Neurath. Rat peritoneal mast cell carboxypeptidase: localization, purification and enzymatic properties. *FEBS Lett.*, 110:292–296, 1980.
- [727] F.A. Exterkate, A.C. Alting, and P.G. Bruinenberg. Diversity of cell envelope proteinase specificity among strains of *Lactococcus lactis* and its relationship to charge characteristics of the substrate-binding region. *Appl. Environ. Microbiol.*, 59:3640–3647, 1993.
- [728] E.H. Eylar and M. Murakami.  $\beta$ -Aspartyl-*N*-acetylglucosaminidase from epididymis. *Methods Enzymol.*, 8:597–600, 1966.
- [729] J. Ezaki, M. Takeda-Ezaki, K. Oda, and E. Kominami. Characterization of endopeptidase activity of tripeptidyl peptidase-II/CLN2 protein which is deficient in classical late infantile neuronal ceroid lipofuscinosis. *Biochem. Biophys. Res. Commun.*, 268:904–908, 2000.
- [730] J. Ezaki, I. Tanida, N. Kanehagi, and E. Kominami. A lysosomal proteinase, the late infantile neuronal ceroid lipofuscinosis gene (CLN2) product, is essential for degradation of a hydrophobic protein, the subunit c of ATP synthase. *J. Neurochem.*, 72:2573–2582, 1999.
- [731] B.U. Failer, N. Braun, and H. Zimmermann. Cloning, expression, and functional characterization of a Ca(2+)-dependent endoplasmic reticulum nucleoside diphosphatase. *J. Biol. Chem.*, 277:36978–36986, 2002.
- [732] D. Fairbairn. The preparation and properties of a lysophospholipase from *Penicillium notatum*. *J. Biol. Chem.*, 173:705–714, 1948.
- [733] A. Falanga and S.G. Gordon. Isolation and characterization of cancer procoagulant: a cysteine proteinase from malignant tissue. *Biochemistry*, 24:5558–5567, 1985.
- [734] A. Falanga, E. Shaw, M.B. Donati, R. Consonni, T. Barbui, and S. Gordon. Inhibition of cancer procoagulant by peptidyl diazomethyl ketones and peptidyl sulfonium salts. *Thromb. Res.*, 54:389–398, 1989.
- [735] H.J. Fallon and W.L. Byrne. 2-Phosphoglyceric acid phosphatase: identification and properties of the beef-liver enzyme. *Biochim. Biophys. Acta*, 105:43–53, 1965.
- [736] K. Fan, P. Wei, Q. Feng, S. Chen, C. Huang, L. Ma, B. Lai, J. Pei, Y. Liu, J. Chen, and L. Lai. Biosynthesis, purification, and substrate specificity of severe acute respiratory syndrome coronavirus 3C-like proteinase. *J. Biol. Chem.*, 279:1637–1642, 2004.

- [737] B. Fang, P.I. Boross, J. Tozser, and I.T. Weber. Structural and kinetic analysis of caspase-3 reveals role for S5 binding site in substrate recognition. *J. Mol. Biol.*, 360:654–666, 2006.
- [738] P.C. Farley, M.G. Shepherd, and P.A. Sullivan. The purification and properties of yeast proteinase B from *Candida albicans*. *Biochem. J.*, 236:177–184, 1986.
- [739] A.A. Farooqui. The desulphation of hexosamine sulphates by arylsulphatase B. *Experientia*, 32:1242–1244, 1976.
- [740] A.A. Farooqui and A.S. Balasubramanian. Enzymatic dephosphorylation 3'-phosphoadenosine 5'-phosphosulfate to adenosine 5'-phosphosulfate in sheep brain. *Biochim. Biophys. Acta*, 198:56–65, 1970.
- [741] F. Fassy, O. Krebs, H. Rey, B. Komara, C. Gillard, C. Capdevila, C. Yea, C. Faucheu, A.M. Blanchet, C. Miossec, and A. Diu-Hercend. Enzymatic activity of two caspases related to interleukin-1 $\beta$ -converting enzyme. *Eur. J. Biochem.*, 253:76–83, 1998.
- [742] H.R. Fatania, P.A. Patston, and P.J. Randle. Dephosphorylation and reactivation of phosphorylated purified ox-kidney branched-chain dehydrogenase complex by co-purified phosphatase. *FEBS Lett.*, 158:234–238, 1983.
- [743] C. Faucheu, A.M. Blanchet, V. Collard-Dutilleul, J.L. Lalanne, and A. Diu-Hercend. Identification of a cysteine protease closely related to interleukin-1  $\beta$ -converting enzyme. *Eur. J. Biochem.*, 236:207–213, 1996.
- [744] C. Faucheu, A. Diu, A.W. Chan, A.M. Blanchet, C. Miossec, F. Hervé, V. Collard-Dutilleul, Y. Gu, R.A. Aldape, J.A. Lippke, C. Rocher, M.S.-S. Su, D.J. Livingston, T. Hercend, and J.-L. Lalanne. A novel human protease similar to the interleukin-1 $\beta$  converting enzyme induces apoptosis in transfected cells. *EMBO J.*, 14:1914–1922, 1995.
- [745] C.B. Faulds and G. Williamson. The purification and characterisation of 4-hydroxy-3-methoxy-cinnamic (ferulic) acid esterase from *Streptomyces olivochromogenes* (3232). *J. Gen. Microbiol.*, 137:2339–2345, 1991.
- [746] C.B. Faulds and G. Williamson. Purification and characterisation of a ferulic acid esterase (FAE-III) from *Aspergillus niger*. Specificity for the phenolic moiety and binding to microcrystalline cellulose. *Microbiology*, 140:779–787, 1994.
- [747] P. Faulkner. A hexose-1-phosphatase in silkworm blood. *Biochem. J.*, 60:590–596, 1955.
- [748] P.L. Faust, S. Kornfeld, and J.M. Chirgwin. Cloning and sequence analysis of cDNA for human cathepsin D. *Proc. Natl. Acad. Sci. USA*, 82:4910–4914, 1985.
- [749] J. Feder, L. Keay, L.R. Garrett, N. Cirulis, M.H. Moseley, and B.S. Wildi. *Bacillus cereus* neutral protease. *Biochim. Biophys. Acta*, 251:74–78, 1971.
- [750] F.J. Fehrenbach. Reinigung und Kristallisation der NAD-Glykohydrolase aus C-Streptokokken. *Eur. J. Biochem.*, 18:94–102, 1971.
- [751] F. Félix and N. Brouillet. Purification et propriétés de deux peptidases de levure de brasserie. *Biochim. Biophys. Acta*, 122:127–144, 1966.
- [752] J.P. Fellers, G.B. Collins, and A.G. Hunt. The NIa-proteinase of different plant potyviruses provides specific resistance to viral infection. *Crop Sci.*, 38:1309–1319, 1998.
- [753] A.K. Felux, D. Spitteller, J. Klebensberger, and D. Schleheck. Entner-Doudoroff pathway for sulfoquinovose degradation in *Pseudomonas putida* SQ1. *Proc. Natl. Acad. Sci. USA*, 112:E4298–E4305, 2015.
- [754] Y. Feng, T.A. Vickers, and S.N. Cohen. The catalytic domain of RNase E shows inherent 3' to 5' directionality in cleavage site selection. *Proc. Natl. Acad. Sci. USA*, 99:14746–14751, 2002.
- [755] D.M. Fenton and D.E. Eveleigh. Purification and mode of action of a chitosanase from *Penicillium islandicum*. *J. Gen. Microbiol.*, 126:151–165, 1981.
- [756] H. Ferracci and S. Maroux. Rabbit intestinal aminopeptidase N. Purification and molecular properties. *Biochim. Biophys. Acta*, 599:448–463, 1980.
- [757] A. Ferrández, J.L. García, and E. Díaz. Genetic characterization and expression in heterologous hosts of the 3-(3-hydroxyphenyl)propionate catabolic pathway of *Escherichia coli* K-12. *J. Bacteriol.*, 179:2573–2581, 1997.

- [758] A. Ferrandez, B. Minambres, B. Garcia, E.R. Olivera, J.M. Luengo, J.L. Garcia, and E. Diaz. Catabolism of phenylacetic acid in *Escherichia coli*. Characterization of a new aerobic hybrid pathway. *J. Biol. Chem.*, 273:25974–25986, 1998.
- [759] M.L. Ferri-Fioni, M. Fromant, A.P. Bouin, C. Aubard, C. Lazennec, P. Plateau, and S. Blanquet. Identification in archaea of a novel D-Tyr-tRNA<sup>Tyr</sup> deacylase. *J. Biol. Chem.*, 281:27575–27585, 2006.
- [760] M.L. Ferri-Fioni, E. Schmitt, J. Soutourina, P. Plateau, Y. Mechulam, and S. Blanquet. Structure of crystalline D-Tyr-tRNA(Tyr) deacylase. A representative of a new class of tRNA-dependent hydrolases. *J. Biol. Chem.*, 276:47285–47290, 2001.
- [761] A.J. Ferro, A. Barrett, and S.K. Shapiro. Kinetic properties and the effect of substrate analogues on 5'-methylthioadenosine nucleosidase from *Escherichia coli*. *Biochim. Biophys. Acta*, 438:487–494, 1976.
- [762] T.T. Fida, S. Palamuru, G. Pandey, and J.C. Spain. Aerobic biodegradation of 2,4-dinitroanisole by *Nocardioides* sp. strain JS1661. *Appl. Environ. Microbiol.*, 80:7725–7731, 2014.
- [763] F. Fiedler. Effects of secondary interactions on the kinetics of peptide and peptide ester hydrolysis by tissue kallikrein and trypsin. *Eur. J. Biochem.*, 163:303–312, 1987.
- [764] F. Fiedler, E. Fink, H. Tschesche, and H. Fritz. Porcine glandular kallikreins. *Methods Enzymol.*, 80:493–532, 1981.
- [765] C.J. Fielding. Human lipoprotein lipase. I. Purification and substrate specificity. *Biochim. Biophys. Acta*, 206:109–117, 1970.
- [766] G.B. Fields, H.E. Van Wart, and H. Birkedal-Hansen. Sequence specificity of human skin fibroblast collagenase. Evidence for the role of collagen structure in determining the collagenase cleavage site. *J. Biol. Chem.*, 262:6221–6226, 1987.
- [767] S. Fieulaine, J.E. Lunn, F. Borel, and J.L. Ferrer. The structure of a cyanobacterial sucrose-phosphatase reveals the sugar tongs that release free sucrose in the cell. *Plant Cell*, 17:2049–2058, 2005.
- [768] P.W. Finch, R.E. Wilson, K. Brown, I.D. Hickson, and P.T. Emmerson. Complete nucleotide sequence of the *Escherichia coli* ptr gene encoding protease III. *Nucleic Acids Res.*, 14:7695–7703, 1986.
- [769] J. Finne, , and P.H. Cleavage of the polysialosyl units of brain glycoproteins by a bacteriophage endosialidase. Involvement of a long oligosaccharide segment in molecular interactions of polysialic acid. *J. Biol. Chem.*, 260:1265–1270, 1985.
- [770] M.S. Finnin, J.R. Donigian, A. Cohen, V.M. Richon, R.A. Rifkind, P.A. Marks, R. Breslow, and N.P. Pavletich. Structures of a histone deacetylase homologue bound to the TSA and SAHA inhibitors. *Nature*, 401:188–193, 1999.
- [771] E.H. Fischer and E.A. Stein.  $\alpha$ -Amylases. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 4, pages 313–343. Academic Press, New York, 2nd edition, 1960.
- [772] E.H. Fischer and E.A. Stein. Cleavage of *O*- and *S*-glycosidic bonds (survey). In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 4, pages 301–312. Academic Press, New York, 2nd edition, 1960.
- [773] U. Fischer and N. Amrhein. Cyclic nucleotide phosphodiesterase of *Chlamydomonas reinhardtii*. *Biochim. Biophys. Acta*, 341:412–420, 1974.
- [774] U. Fischer, C. Stroh, and K. Schulze-Osthoff. Unique and overlapping substrate specificities of caspase-8 and caspase-10. *Oncogene*, 25:152–159, 2006.
- [775] W.N. Fishbein and S.P. Bessman. Purification and properties of an enzyme in human blood and rat liver microsomes catalyzing the formation and hydrolysis of  $\gamma$ -lactones. I. Tissue localization, stoichiometry, specificity, distinction from esterase. *J. Biol. Chem.*, 241:4835–4841, 1966.
- [776] W.N. Fishbein and S.P. Bessman. Purification and properties of an enzyme in human blood and rat liver microsomes catalyzing the formation and hydrolysis of  $\gamma$ -lactones. II. Metal ion effects, kinetics, and equilibria. *J. Biol. Chem.*, 241:4842–4847, 1966.
- [777] W.H. Fishman. Beta-glucuronidase. *Adv. Enzymol. Relat. Subj. Biochem.*, 16:361–409, 1955.

- [778] N. Fittipaldi, T. Sekizaki, D. Takamatsu, M. de la Cruz Domínguez-Punaro, J. Harel, N.K. Bui, W. Vollmer, and M. Gottschalk. Significant contribution of the *pgdA* gene to the virulence of *Streptococcus suis*. *Mol. Microbiol.*, 70:1120–1135, 2008.
- [779] W.P. Fitzmaurice, L.L. Saari, R.G. Lowery, P.W. Ludden, and G.P. Roberts. Genes coding for the reversible ADP-ribosylation system of dinitrogenase reductase from *Rhodospirillum rubrum*. *Mol. Gen. Genet.*, 218:340–347, 1989.
- [780] J.G. Flaks, M.J. Erwin, and J.M. Buchanan. Biosynthesis of the purines. XVIII. 5-Amino-1-ribosyl-4-imidazolecarboxamide 5'-phosphate transformylase and inosinicase. *J. Biol. Chem.*, 229:603–612, 1957.
- [781] P.R. Flanagan and G.G. Forstner. Purification of rat intestinal maltase/glucoamylase and its anomalous dissociation either by heat or by low pH. *Biochem. J.*, 173:553–563, 1978.
- [782] G. Fleminger, A. Carmel, and A. Yaron. Fluorogenic substrates for bacterial aminopeptidase P and its analogs detected in human serum and calf lung. *Eur. J. Biochem.*, 125:609–615, 1982.
- [783] T.S. Fletcher, M. Alhadeff, C.S. Craik, and C. Largman. Isolation and characterization of a cDNA encoding rat cationic trypsinogen. *Biochemistry*, 26:3081–3086, 1987.
- [784] T.S. Fletcher, W.-F. Shen, and C. Largman. Primary structure of human pancreatic elastase 2 determined by sequence analysis of the cloned mRNA. *Biochemistry*, 26:7256–7261, 1987.
- [785] M.J. Flipphi, H. Panneman, P. van der Veen, J. Visser, and L.H. de Graaff. Molecular cloning, expression and structure of the endo-1,5- $\alpha$ -L-arabinase gene of *Aspergillus niger*. *Appl. Microbiol. Biotechnol.*, 40:318–326, 1993.
- [786] M. Florkin and G. Duchateau-Bosson. Microdosage photométrique de l'allantoïne en solutions pures et dans l'urine. *Enzymologia*, 9:5–9, 1940.
- [787] J.M. Fogg, M.J. Schofield, M.F. White, and D.M. Lilley. Sequence and functional-group specificity for cleavage of DNA junctions by RuvC of *Escherichia coli*. *Biochemistry*, 38:11349–11358, 1999.
- [788] J.E. Folk. Carboxypeptidase B (porcine pancreas). *Methods Enzymol.*, 19:504–508, 1970.
- [789] J.E. Folk. Chymotrypsin C (porcine pancreas). *Methods Enzymol.*, 19:109–112, 1970.
- [790] B. Foltmann. A review of prorennin and rennin. *C. R. Trav. Lab. Carlsberg*, 35:143–231, 1966.
- [791] B. Foltmann. Gastric proteinases - structure, function, evolution and mechanism of action. *Essays Biochem.*, 17:52–84, 1981.
- [792] B. Foltmann and A.L. Jensen. Human progastricsin - analysis of intermediates during activation into gastricsin and determination of the amino-acid sequence of the propart. *Eur. J. Biochem.*, 128:63–70, 1982.
- [793] R. Foltmann. Gastric proteinases -structure, function, evolution and mechanism of action. *Essays Biochem.*, 17:52–84, 1981.
- [794] M.L. Fonda. Purification and characterization of vitamin B<sub>6</sub>-phosphate phosphatase from human erythrocytes. *J. Biol. Chem.*, 267:15978–15983, 1992.
- [795] M.L. Fonda and Y.N. Zhang. Kinetic mechanism and divalent metal activation of human erythrocyte pyridoxal phosphatase. *Arch. Biochem. Biophys.*, 320:345–352, 1995.
- [796] W.S. Fones and M. Lee. Hydrolysis of *N*-acyl derivatives of alanine and phenylalanine by acylase I and carboxypeptidase. *J. Biol. Chem.*, 201:847–856, 1953.
- [797] F. Foor and G.M. Brown. Purification and properties of guanosine triphosphate cyclohydrolase II from *Escherichia coli*. *J. Biol. Chem.*, 250:3545–3551, 1975.
- [798] J.G. Foulkes, R.F. Howard, and A. Ziemiecki. Detection of a novel mammalian protein phosphatase with activity for phosphotyrosine. *FEBS Lett.*, 130:197–200, 1981.
- [799] J.W. Fox, R. Campbell, L. Beggerly, and J.B. Bjarnason. Substrate specificities and inhibition of two hemorrhagic zinc proteases Ht-c and Ht-d from *Crotalus atrox* venom. *Eur. J. Biochem.*, 156:65–72, 1986.

- [800] S. Fraas, A.K. Steinbach, A. Tabbert, J. Harder, U. Ermler, K. Tittmann, A., and Kroneck P.M.H. Cyclohexane-1,2-dione hydrolase: A new tool to degrade alicyclic compounds. *J. Mol. Catalysis B: Enzymatic*, 61:47–49, 2009.
- [801] H. Fraenkel-Conrat and J. Fraenkel-Conrat. Inactivation of crotoxin by group-specific reagents. *Biochim. Biophys. Acta*, 5:98–104, 1950.
- [802] O. Francetic, C. Badaut, S. Rimsky, and A.P. Pugsley. The ChiA (YheB) protein of *Escherichia coli* K-12 is an endochitinase whose gene is negatively controlled by the nucleoid-structuring protein H-NS. *Mol. Microbiol.*, 35:1506–1517, 2000.
- [803] S.E. Francis, I.Y. Gluzman, A. Oksman, A. Knickerbocker, R. Mueller, M.L. Bryant, D.R. Sherman, D.G. Russell, and D.E. Goldberg. Molecular characterization and inhibition of a *Plasmodium falciparum* aspartic hemoglobinase. *EMBO J.*, 13:306–317, 1994.
- [804] G. Fredrikson, P. Stralfors, N.O. Nilsson, and P. Belfrage. Hormone-sensitive lipase of rat adipose tissue. Purification and some properties. *J. Biol. Chem.*, 256:6311–6320, 1981.
- [805] D.V. Freistroffer, M.Y. Pavlov, J. MacDougall, R.H. Buckingham, and M. Ehrenberg. Release factor RF3 in *E. coli* accelerates the dissociation of release factors RF1 and RF2 from the ribosome in a GTP-dependent manner. *EMBO J.*, 16:4126–4133, 1997.
- [806] J.O. Freitas, Termignoni Jr., Guimaraes C., and J.A. Methionine aminopeptidase associated with liver mitochondria and microsomes. *Int. J. Biochem.*, 17:1285–1291, 1985.
- [807] D. French.  $\beta$ -Amylases. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 4, pages 345–368. Academic Press, New York, 2nd edition, 1960.
- [808] D. French and D.W. Knapp. The maltase of *Clostridium acetobutylicum*. *J. Biol. Chem.*, 187:463–471, 1950.
- [809] J.B. French, T.P. Begley, and S.E. Ealick. Structure of trifunctional THI20 from yeast. *Acta Crystallogr. D Biol. Crystallogr.*, 67:784–791, 2011.
- [810] J.M. Frère and B. Joris. Penicillin-sensitive enzymes in peptidoglycan biosynthesis. *CRC Crit. Rev. Microbiol.*, 11:306–331, 1985.
- [811] K. Freter, J.C. Rabinowitz, and B. Witkop. Labile Stoffwechselprodukte. V. Zur Biogenese des Formiminoglycins aus 4(5H)-imidazolone. *Ann. Chem.*, 607:174–187, 1957.
- [812] A.J. Fretland and C.J. Omiecinski. Epoxide hydrolases: biochemistry and molecular biology. *Chem. Biol. Interact.*, 129:41–59, 2000.
- [813] D.M. Freymann, R.J. Keenan, R.M. Stroud, and P. Walter. Structure of the conserved GTPase domain of the signal recognition particle. *Nature*, 385:361–364, 1997.
- [814] D.N. Frick. The hepatitis C virus NS3 protein: a model RNA helicase and potential drug target. *Curr. Issues Mol. Biol.*, 9:1–20, 2007.
- [815] D.N. Frick and M.J. Bessman. Cloning, purification, and properties of a novel NADH pyrophosphatase. Evidence for a nucleotide pyrophosphatase catalytic domain in MutT-like enzymes. *J. Biol. Chem.*, 270:1529–1534, 1995.
- [816] L.D. Fricker. Carboxypeptidase E. *Annu. Rev. Physiol.*, 50:309–321, 1988.
- [817] L.D. Fricker. Methods for studying carboxypeptidase E. *Methods Neurosci.*, 23:237–250, 1995.
- [818] L.D. Fricker. Carboxypeptidase E/H. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 1341–1344. Academic Press, London, 1998.
- [819] E.C. Friedberg and D.A. Goldthwait. Endonuclease II of *E. coli*. I. Isolation and purification. *Proc. Natl. Acad. Sci. USA*, 62:934–940, 1969.
- [820] E.C. Friedberg, S.-M. Hadi, and D.A. Goldthwait. Endonuclease II of *Escherichia coli*. II. Enzyme properties and studies on the degradation of alkylated and native deoxyribonucleic acid. *J. Biol. Chem.*, 244:5879–5889, 1969.

- [821] C.G. Friedrich and G. Mitrenga. Utilization of aliphatic amides and formation of two different amidases by *Alcaligenes eutrophus*. *J. Gen. Microbiol.*, 125:367–374, 1981.
- [822] C.G. Friedrich, D. Rother, F. Bardischewsky, A. Quentmeier, and J. Fischer. Oxidation of reduced inorganic sulfur compounds by bacteria: emergence of a common mechanism. *Appl. Environ. Microbiol.*, 67:2873–2882, 2001.
- [823] Y.S. Frohwein and S. Gatt. Isolation of  $\beta$ -*N*-acetylhexosaminidase,  $\beta$ -*N*-acetylglucosaminidase, and  $\beta$ -*N*-acetylgalactosaminidase from calf brain. *Biochemistry*, 6:2775–2782, 1967.
- [824] I. Fruchey, N. Shapir, M.J. Sadowsky, and L.P. Wackett. On the origins of cyanuric acid hydrolase: purification, substrates, and prevalence of AtzD from *Pseudomonas* sp. strain ADP. *Appl. Environ. Microbiol.*, 69:3653–3657, 2003.
- [825] L. Fruhaufová, E. Suska-Brezinska, T. Barth, and I. Rychlik. Rat liver enzyme inactivating oxytocin and its deaminocarba analogues. *Coll. Czech. Chem. Commun.*, 38:2793–2798, 1973.
- [826] J.S. Fruton. Aspartyl proteinases. In A. Neuberger and K. Brocklehurst, editors, *New Comprehensive Biochemistry: Hydrolytic Enzymes*, volume 16, pages 1–38. Elsevier, Amsterdam, 1987.
- [827] K.B. Fryxell, K. O'Donoghue, R.M. Graeff, H.C. Lee, and W.D. Branton. Functional expression of soluble forms of human CD38 in *Escherichia coli* and *Pichia pastoris*. *Protein Expr. Purif.*, 6:329–336, 1995.
- [828] T.B. Fu and J. Taylor. When retroviral reverse transcriptases reach the end of their RNA templates. *J. Virol.*, 66:4271–4278, 1992.
- [829] Y. Fu, H. Yu, S.H. Tang, X. Hu, Y. Wang, B. Liu, C. Yu, and F. Jin. New dioscin-glycosidase hydrolyzing multi-glycosides of dioscin from *Absidia* strain. *J Microbiol Biotechnol*, 20:1011–1017, 2010.
- [830] R. Fuchs, W. Machleidt, and H.G. Gassen. Molecular cloning and sequencing of a cDNA coding for mature human kidney cathepsin H. *Biol. Chem. Hoppe-Seyler*, 369:469–475, 1988.
- [831] J. Fuhrmann, B. Mierzwa, D.B. Trentini, S. Spiess, A. Lehner, E. Charpentier, and T. Clausen. Structural basis for recognizing phosphoarginine and evolving residue-specific protein phosphatases in gram-positive bacteria. *Cell Rep*, 3:1832–1839, 2013.
- [832] N. Fujii, K. Kimura, T. Yashiki, K. Tsuzuki, K. Moriishi, N. Yokosawa, B. Syuto, and K. Oguma. A zinc-protease specific domain in botulinum and tetanus neurotoxins. *Microbiol. Intern.*, 36:213–220, 1992.
- [833] K. Fujikawa. Bovine Hageman factor and its fragments. *Methods Enzymol.*, 163:54–68, 1988.
- [834] K. Fujikawa, D.W. Chung, L.E. Hendrickson, and E.W. Davie. Amino acid sequence of human factor XI, a blood coagulation factor with four tandem repeats that are highly homologous with plasma prekallikrein. *Biochemistry*, 25:2417–2424, 1986.
- [835] K. Fujikawa and E.W. Davie. Bovine factor IX (Christmas factor). *Methods Enzymol.*, 45:74–83, 1976.
- [836] K. Fujikawa and E.W. Davie. Bovine factor X (Stuart factor). *Methods Enzymol.*, 45:89–95, 1976.
- [837] K. Fujikawa and E.W. Davie. Human factor XII (Hageman factor). *Methods Enzymol.*, 80:198–211, 1981.
- [838] K. Fujikawa, H. Kamiya, H. Yakushiji, Y. Fujii, Y. Nakabeppu, and H. Kasai. The oxidized forms of dATP are substrates for the human MutT homologue, the hMTH1 protein. *J. Biol. Chem.*, 274:18201–18205, 1999.
- [839] K. Fujikawa, H. Kamiya, H. Yakushiji, Y. Nakabeppu, and H. Kasai. Human MTH1 protein hydrolyzes the oxidized ribonucleotide, 2-hydroxy-ATP. *Nucleic Acids Res.*, 29:449–454, 2001.
- [840] K. Fujikawa and H. Kasai. The oxidized pyrimidine ribonucleotide, 5-hydroxy-CTP, is hydrolyzed efficiently by the *Escherichia coli* recombinant Orf135 protein. *DNA Repair (Amst.)*, 1:571–576, 2002.
- [841] K. Fujikawa, H. Suzuki, B. McMullen, and D. Chung. Purification of human von Willebrand factor-cleaving protease and its identification as a new member of the metalloproteinase family. *Blood*, 98:1662–1666, 2001.
- [842] D. Fujimoto, T. Koyama, and N. Tamiya. *N*-Acetyl- $\beta$ -alanine deacetylase in hog kidney. *Biochim. Biophys. Acta*, 167:407–413, 1968.



- [843] S. Fujimura, O. Ueda, Y. Shibata, and K. Hirai. Isolation and properties of a tripeptidyl peptidase from a periodontal pathogen *Prevotella nigrescens*. *FEMS Microbiol. Lett.*, 219:305–309, 2003.
- [844] M. Fujinaga and M.N.G. James. Rat submaxillary gland serine protease, tonin. Structure solution and refinement at 1.8 Å resolution. *J. Mol. Biol.*, 195:373–396, 1987.
- [845] M. Fujinaga, R.J. Read, A. Sielecki, W. Ardelt, M. Laskowski, James Jr., and M.N.G. Refined crystal structure of the molecular complex of *Streptomyces griseus* protease B, a serine protease, with the third domain of the ovomucoid inhibitor from turkey. *Proc. Natl Acad. Sci. USA*, 79:4868–4872, 1982.
- [846] T. Fujino, N. Navaratnam, A. Jarmuz, A. von Haeseler, and J. Scott. C→U editing of apolipoprotein B mRNA in marsupials: identification and characterisation of APOBEC-1 from the American opossum *Monodelphus domestica*. *Nucleic Acids Res.*, 27:2662–2671, 1999.
- [847] M. Fujisaki and K. Sugawara. Properties of peptidylarginine deiminase from the epidermis of newborn rats. *J. Biochem. (Tokyo)*, 89:257–263, 1981.
- [848] Y. Fujisawa, H. Shirafuji, M. Kida, and K. Nara. New findings on cephalosporin C biosynthesis. *Nat. New Biol.*, 246:154–155, 1973.
- [849] A. Fujita, Y. Nose, and K. Kuratani. Second type of bacterial thiaminase. *J. Vitaminol. (Kyoto)*, 1:1–7, 1954.
- [850] K. Fujita, F. Oura, N. Nagamine, T. Katayama, J. Hiratake, K. Sakata, H. Kumagai, and K. Yamamoto. Identification and molecular cloning of a novel glycoside hydrolase family of core 1 type *O*-glycan-specific endo- $\alpha$ -*N*-acetylglucosaminidase from *Bifidobacterium longum*. *J. Biol. Chem.*, 280:37415–37422, 2005.
- [851] K. Fujita, S. Sakamoto, Y. Ono, M. Wakao, Y. Suda, K. Kitahara, and T. Sukanuma. Molecular cloning and characterization of a  $\beta$ -L-Arabinobiosidase in *Bifidobacterium longum* that belongs to a novel glycoside hydrolase family. *J. Biol. Chem.*, 286:5143–5150, 2011.
- [852] K. Fujita, Y. Takashi, E. Obuchi, K. Kitahara, and T. Sukanuma. Characterization of a novel  $\beta$ -L-arabinofuranosidase in *Bifidobacterium longum*: functional elucidation of a DUF1680 family member. *J. Biol. Chem.*, 286:38079–38085, 2011.
- [853] T. Fukamizo, A. Fleury, N. Côté, M. Mitsutomi, and R. Brzezinski. Exo- $\beta$ -D-glucosaminidase from *Amycolatopsis orientalis*: catalytic residues, sugar recognition specificity, kinetics, and synergism. *Glycobiology*, 16:1064–1072, 2006.
- [854] K. Fukasawa, K.M. Fukasawa, H. Iwamoto, J. Hirose, and M. Harada. The HELLGH motif of rat liver dipeptidyl peptidase III is involved in zinc coordination and the catalytic activity of the enzyme. *Biochemistry*, 38:8299–8303, 1999.
- [855] K.M. Fukasawa, K. Fukasawa, M. Kanai, S. Fujii, and M. Harada. Molecular cloning and expression of rat liver aminopeptidase B. *J. Biol. Chem.*, 271:30731–30735, 1996.
- [856] H. Fukatsu, Y. Hashimoto, M. Goda, H. Higashibata, and M. Kobayashi. Amine-synthesizing enzyme *N*-substituted formamide deformylase: screening, purification, characterization, and gene cloning. *Proc. Natl. Acad. Sci. USA*, 101:13726–13731, 2004.
- [857] H. Fukuda, S. Iwade, and A. Kimura. A new enzyme: long acyl aminoacylase from *Pseudomonas diminuta*. *J. Biochem. (Tokyo)*, 91:1731–1738, 1982.
- [858] M.N. Fukuda and G. Matsumara. Endo- $\beta$ -galactosidase of *Escherichia freundii*. Purification and endoglycosidic action on keratan sulfates, oligosaccharides, and blood group active glycoprotein. *J. Biol. Chem.*, 251:6218–6225, 1976.
- [859] S. Fukui, T. Suzuki, K. Kitahara, and T. Miwa.  $\beta$ -1,3'-Xylanase. *J. Gen. Appl. Microbiol.*, 6:270–282, 1960.
- [860] T. Fukumura, G. Talbot, H. Misono, Y. Teremura, K., Soda Kato, properties of a novel enzyme K. Purification, and L.  $\alpha$ -amino- $\epsilon$ -caprolactamase from *Cryptococcus laurentii*. *FEBS Lett.*, 89:298–300, 1978.
- [861] R.S. Fuller. Yapsin 2. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 908–909. Academic Press, London, 1998.
- [862] R.S. Fuller, A. Brake, and J. Thorner. Yeast prohormone processing enzyme (KEX2 gene product) is a  $\text{Ca}^{2+}$ -dependent serine protease. *Proc. Natl. Acad. Sci. USA*, 86:1434–1438, 1989.

- [863] Y. Fundoiano-Hershcovitz, L. Rabinovitch, Y. Langut, V. Reiland, G. Shoham, and Y. Shoham. Identification of the catalytic residues in the double-zinc aminopeptidase from *Streptomyces griseus*. *FEBS Lett.*, 571:192–196, 2004.
- [864] B.C. Furie and B. Furie. Coagulant protein of Russell's viper venom. *Methods Enzymol.*, 45:191–205, 1976.
- [865] S. Fushinobu, M. Hidaka, Y. Honda, T. Wakagi, H. Shoun, and M. Kitaoka. Structural basis for the specificity of the reducing end xylose-releasing exo-oligoxylanase from *Bacillus halodurans* C-125. *J. Biol. Chem.*, 280:17180–17186, 2005.
- [866] M. Futai and D. Mizuno. A new phosphodiesterase forming nucleoside 5'-monophosphate from rat liver. Its partial purification and substrate specificity for nicotinamide adenine dinucleotide and oligonucleotides. *J. Biol. Chem.*, 242:5301–5307, 1967.
- [867] S.B. Gabelli, M.A. Bianchet, W. Xu, C.A. Dunn, Z.D. Niu, L.M. Amzel, and M.J. Bessman. Structure and function of the *E. coli* dihydroneopterin triphosphate pyrophosphatase: a Nudix enzyme involved in folate biosynthesis. *Structure*, 15:1014–1022, 2007.
- [868] C. Gagnon, G. Harbour, and R. Camato. Purification and characterization of protein methylesterase from rat kidney. *J. Biol. Chem.*, 259:10212–10215, 1984.
- [869] H. Gainer, J.T. Russell, and Y.P. Loh. An aminopeptidase activity in bovine pituitary secretory vesicles that cleaves the N-terminal arginine from  $\beta$ -lipotropin(60-65). *FEBS Lett.*, 175:135–139, 1984.
- [870] B. Gallis, P. Bornstein, and D.L. Brautigan. Tyrosylprotein kinase and phosphatase activities in membrane vesicles from normal and Rous sarcoma virus-transformed rat cells. *Proc. Natl. Acad. Sci. USA*, 78:6689–6693, 1981.
- [871] D. Gao, D.L. Narasimhan, J. Macdonald, R. Brim, M.C. Ko, D.W. Landry, J.H. Woods, R.K. Sunahara, and C.G. Zhan. Thermostable variants of cocaine esterase for long-time protection against cocaine toxicity. *Mol. Pharmacol.*, 75:318–323, 2009.
- [872] X.D. Gao, V. Kaigorodov, and Y. Jigami. YND1, a homologue of GDA1, encodes membrane-bound apyrase required for Golgi N- and O-glycosylation in *Saccharomyces cerevisiae*. *J. Biol. Chem.*, 274:21450–21456, 1999.
- [873] Y. Gao, L. Wells, F.I. Comer, G.J. Parker, and G.W. Hart. Dynamic O-glycosylation of nuclear and cytosolic proteins: cloning and characterization of a neutral, cytosolic  $\beta$ -N-acetylglucosaminidase from human brain. *J. Biol. Chem.*, 276:9838–9845, 2001.
- [874] B. García, E.R. Olivera, B. Mi nambres, Ca nedo Fernández-Valverde, Prieto L.M., García M.A., Martínez J.L., Luengo M., and J.M. Novel biodegradable aromatic plastics from a bacterial source. Genetic and biochemical studies on a route of the phenylacetyl-CoA catabolon. *J. Biol. Chem.*, 274:29228–29241, 1999.
- [875] N. Garcia-Alvarez, U. Teichert, and D.H. Wolf. Proteinase yscD mutants of yeast. Isolation and characterization. *Eur. J. Biochem.*, 163:339–346, 1987.
- [876] R. Garcia-Lepe, O.M. Nuero, F. Reyes, and F. Santamaria. Lipases in autolysed cultures of filamentous fungi. *Lett. Appl. Microbiol.*, 25:127–130, 1997.
- [877] A. Garcia-Sastre, E. Villar, J.C. Manuguerra, C. Hannoun, and J.A. Cabezas. Activity of influenza C virus O-acetylsterase with O-acetyl-containing compounds. *Biochem. J.*, 273:435–441, 1991.
- [878] S.J. Gardell, C.S. Craik, E. Clauser, E.J. Goldsmith, C.B. Stewart, M. Graf, and W.J. Rutter. A novel rat carboxypeptidase, CPA2: characterization, molecular cloning, and evolutionary implications on substrate specificity in the carboxypeptidase gene family. *J. Biol. Chem.*, 263:17828–17836, 1988.
- [879] G.K. Garg and T.K. Virupaksha. Acid protease from germinated sorghum. 2. Substrate specificity with synthetic peptides and ribonuclease A. *Eur. J. Biochem.*, 17:4–12, 1970.
- [880] M. Garnier, M.-J. Vacheron, M. Guinard, and G. Michel. Purification and partial characterization of the extracellular  $\gamma$ -D-glutamyl-(L)meso-diaminopimelate endopeptidase I, from *Bacillus sphaericus* NCTC 9602. *Eur. J. Biochem.*, 148:539–543, 1985.
- [881] A.R. Garrett, L.A. Johnson, , and I.R. Isolation, molecular characterization and expression of the ushB gene of *Salmonella typhimurium* which encodes a membrane bound UDP-sugar hydrolase. *Mol. Microbiol.*, 3:177–186, 1989.

- [882] F.T. Gates and S. Linn. Endonuclease V of *Escherichia coli*. *J. Biol. Chem.*, 252:1647–1653, 1977.
- [883] S. Gatt. Purification and properties of phospholipase A-1 from rat and calf brain. *Biochim. Biophys. Acta*, 159:304–316, 1968.
- [884] G.M. Gaucher and K.J. Stevenson. Thermomycin. *Methods Enzymol.*, 45:415–433, 1976.
- [885] N.S. Gee and A.J. Kenny. Proteins of the kidney microvillar membrane. The 130kDa protein in pig kidney, recognised by monoclonal antibody GK5C1, is an ectoenzyme with aminopeptidase activity. *Biochem. J.*, 230:753–764, 1985.
- [886] N.S. Gee and A.J. Kenny. Proteins of the kidney microvillar membrane. Enzymic and molecular properties of aminopeptidase W. *Biochem. J.*, 246:97–102, 1987.
- [887] N.S. Gee, C.I. Ragan, K.J. Watling, S. Aspley, R.G. Jackson, G.G. Reid, D. Gani, and J.K. Shute. The purification and properties of *myo*-inositol monophosphatase from bovine brain. *Biochem. J.*, 249:883–889, 1988.
- [888] K. Geider and H. Hofmann-Berling. Proteins controlling the helical structure of DNA. *Annu. Rev. Biochem.*, 50:233–260, 1981.
- [889] R. Geiger and W. Miska. Human tissue kallikrein. *Methods Enzymol.*, 163:102–115, 1988.
- [890] T. George, Q. Wen, R. Griffiths, A. Ganesh, M. Meuth, and C.M. Sanders. Human Pif1 helicase unwinds synthetic DNA structures resembling stalled DNA replication forks. *Nucleic Acids Res.*, 37:6491–6502, 2009.
- [891] R.D. Gerard and R.S. Meidell. Regulation of tissue plasminogen activator expression. *Annu. Rev. Physiol.*, 51:245–262, 1989.
- [892] A. Gerber, H. Grosjean, T. Melcher, and W. Keller. Tad1p, a yeast tRNA-specific adenosine deaminase, is related to the mammalian pre-mRNA editing enzymes ADAR1 and ADAR2. *EMBO J.*, 17:4780–4789, 1998.
- [893] G.A. Gerencser and K.R. Purushotham. Reconstituted Cl<sup>-</sup> pump protein: a novel Cl<sup>-</sup>-motive ATPase. *J. Bioenerget. Biomembr.*, 28:459–469, 1996.
- [894] J. Gergely, P. Hele, and C.V. Ramakrishnan. Succinyl and acetyl coenzyme A deacylases. *J. Biol. Chem.*, 198:323–334, 1952.
- [895] H.K. Gershenfeld, R.J. Hershberger, T.B. Shows, and I.L. Weissman. Cloning and chromosomal assignment of a human cDNA encoding a T cell- and natural killer cell-specific trypsin-like serine protease. *Proc. Natl. Acad. Sci. USA*, 85:1184–1188, 1988.
- [896] F. Geu-Flores, M.E. Møldrup, C. Böttcher, C.E. Olsen, D. Scheel, and B.A. Halkier. Cytosolic  $\gamma$ -glutamyl peptidases process glutathione conjugates in the biosynthesis of glucosinolates and camalexin in *Arabidopsis*. *Plant Cell*, 23:2456–2469, 2011.
- [897] B. Geueke, T. Heck, M. Limbach, V. Nesatyy, D. Seebach, and H.P. Kohler. Bacterial  $\beta$ -peptidyl aminopeptidases with unique substrate specificities for  $\beta$ -oligopeptides and mixed  $\beta,\alpha$ -oligopeptides. *FEBS J.*, 273:5261–5272, 2006.
- [898] B. Geueke, K. Namoto, D. Seebach, and H.P. Kohler. A novel  $\beta$ -peptidyl aminopeptidase (BapA) from strain 3-2W4 cleaves peptide bonds of synthetic  $\beta$ -tri- and  $\beta$ -dipeptides. *J. Bacteriol.*, 187:5910–5917, 2005.
- [899] M. Geyer and A. Wittinghofer. GEFs, GAPs, GDIs and effectors: taking a closer (3D) look at the regulation of Ras-related GTP-binding proteins. *Curr. Opin. Struct. Biol.*, 7:786–792, 1997.
- [900] M. El Ghachi, A. Bouhss, D. Blanot, and D. Mengin-Lecreulx. The *bacA* gene of *Escherichia coli* encodes an undecaprenyl pyrophosphate phosphatase activity. *J. Biol. Chem.*, 279:30106–30113, 2004.
- [901] M. El Ghachi, A. Derbise, A. Bouhss, and D. Mengin-Lecreulx. Identification of multiple genes encoding membrane proteins with undecaprenyl pyrophosphate phosphatase (UppP) activity in *Escherichia coli*. *J. Biol. Chem.*, 280:18689–18695, 2005.
- [902] R.L. Gherna, S.H. Richardson, and S.C. Rittenberg. The bacterial oxidation of nicotine. VI. The metabolism of 2,6-dihydroxypseudoxyntine. *J. Biol. Chem.*, 240:3669–3674, 1965.

- [903] O. Ghisalpa and M. Küenzi. Biodegradation and utilization of monomethyl sulfate by specialized methylotrophs. *Experientia*, 39:1257–1263, 1983.
- [904] S.V. Ghodge, J.A. Cummings, H.J. Williams, and F.M. Raushel. Discovery of a cyclic phosphodiesterase that catalyzes the sequential hydrolysis of both ester bonds to phosphorus. *J. Am. Chem. Soc.*, 135:16360–16363, 2013.
- [905] J. Ghosh, P.J. Anderson, S. Chandrasekaran, and M.G. Caparon. Characterization of *Streptococcus pyogenes*  $\beta$ -NAD<sup>+</sup> glycohydrolase: re-evaluation of enzymatic properties associated with pathogenesis. *J. Biol. Chem.*, 285:5683–5694, 2010.
- [906] R.K. Ghosh and M.P. Deutscher. Identification of an *Escherichia coli* nuclease acting on structurally altered transfer RNA molecules. *J. Biol. Chem.*, 253:997–1000, 1978.
- [907] S. Ghosh and U. DasGupta. Studies with endonuclease X: a new deoxyendonuclease of *E. coli* that preferentially cleaves supercoiled plasmid DNA. *Curr. Trends Life Sci.*, 12:79–88, 1984.
- [908] J.-M. Ghuysen, L. Dierickx, J. Coyette, M. Leyh-Bouille, M. Guinand, and J.N. Campbell. An improved technique for the preparation of *Streptomyces* peptidases and *N*-acetylmuramyl-L-alanine amidase active on bacterial wall peptidoglycans. *Biochemistry*, 8:213–222, 1969.
- [909] J.-M. Ghuysen, J.-M. Frère, M. Leyh-Bouille, M. Nguyen-Distèche, J. Coyette, J. Dusart, B. Joris, C. Duez, O. Dideberg, P. Charlier, G. Dive, and J. Lamotte-Brasseur. Bacterial wall peptidoglycan, DD-peptidases and  $\beta$ -lactam antibiotics. *Scand. J. Infect. Dis. Suppl.*, 42:17–37, 1984.
- [910] B. Giabbai and M. Degano. Cloning, purification, crystallization and X-ray analysis of the *Escherichia coli* pyrimidine nucleoside hydrolase YeiK. *Acta Crystallogr. D Biol. Crystallogr.*, 60:524–527, 2004.
- [911] B. Giabbai and M. Degano. Crystal structure to 1.7 Å of the *Escherichia coli* pyrimidine nucleoside hydrolase YeiK, a novel candidate for cancer gene therapy. *Structure*, 12:739–749, 2004.
- [912] D. Gibson and G.H. Dixon. Chymotrypsin-like proteases from the sea anemone, *Metridium senile*. *Nature*, 222:753–756, 1969.
- [913] D.M. Gibson, P. Ayengar, and D.R. Sanadi. A phosphatase specific for nucleoside diphosphates. *Biochim. Biophys. Acta*, 16:536–538, 1955.
- [914] F. Giffhorn and G. Gottschalk. Inactivation of citrate lyase from *Rhodopseudomonas gelatinosa* by a specific deacetylase and inhibition of this inactivation by L-(+)-glutamate. *J. Bacteriol.*, 124:1052–1061, 1975.
- [915] F. Giffhorn, H. Rode, A. Kuhn, and G. Gottschalk. Citrate lyase deacetylase of *Rhodopseudomonas gelatinosa*. Isolation of the enzyme and studies on the inhibition by L-glutamate. *Eur. J. Biochem.*, 111:461–471, 1980.
- [916] C. Giglione, M. Pierre, and T. Meinnel. Peptide deformylase as a target for new generation, broad spectrum antimicrobial agents. *Mol. Microbiol.*, 36:1197–1205, 2000.
- [917] G. Gil and F.G. Hegardt. Some properties of purified 3-hydroxy-3-methylglutaryl coenzyme A reductase phosphatases from rat liver. *Arch. Biochem. Biophys.*, 214:192–198, 1982.
- [918] G. Gil, M. Sitges, and F.G. Hegardt. Purification and properties of rat liver hydroxymethylglutaryl coenzyme A reductase phosphatases. *Biochim. Biophys. Acta*, 663:211–221, 1981.
- [919] J.V. Gilbert, A.G. Plaut, Y. Fishman, and A. Wright. Cloning of the gene encoding streptococcal immunoglobulin A protease and its expression in *Escherichia coli*. *Infect. Immun.*, 56:1961–1966, 1988.
- [920] J.V. Gilbert, A.G. Plaut, and A. Wright. Analysis of the immunoglobulin A protease gene of *Streptococcus sanguis*. *Infect. Immun.*, 59:7–17, 1991.
- [921] R. Gilboa, H.M. Greenblatt, M. Perach, A. Spungin-Bialik, U. Lessel, G. Wohlfahrt, D. Schomburg, S. Blumberg, and G. Shoham. Interactions of *Streptomyces griseus* aminopeptidase with a methionine product analogue: a structural study at 1.53 Å resolution. *Acta Crystallogr. D Biol. Crystallogr.*, 56:551–558, 2000.
- [922] A.-M. Gilles, J.-M. Imhoff, and B. Keil.  $\alpha$ -Clostripain. Chemical characterization, activity, and thiol content of the highly active form of clostripain. *J. Biol. Chem.*, 254:1462–1468, 1979.

- [923] A.-M. Gilles, A. Lecroisey, and B. Keil. Primary structure of  $\alpha$ -clostripain light chain. *Eur. J. Biochem.*, 145:469–476, 1984.
- [924] L.E. Giroir and W.A. Deutsch. Drosophila deoxyuridine triphosphatase. Purification and characterization. *J. Biol. Chem.*, 262:130–134, 1987.
- [925] S. Giuriato, B. Payrastra, A.L. Drayer, M. Plantavid, R. Woscholski, P. Parker, C. Erneux, and H. Chap. Tyrosine phosphorylation and relocation of SHIP are integrin-mediated in thrombin-stimulated human blood platelets. *J. Biol. Chem.*, 272:26857–26863, 1997.
- [926] L. Glaser. The synthesis of teichoic acid. IV. On the regulation of cytidine 5'-diphosphateglycerol concentration. *Biochim. Biophys. Acta*, 101:6–15, 1965.
- [927] L. Glaser, A. Melo, , and R. Uridine diphosphate sugar hydrolase. Purification of enzyme and protein inhibitor. *J. Biol. Chem.*, 242:1944–1954, 1987.
- [928] P.A. Glaze, D.C. Watson, N.M. Young, and M.E. Tanner. Biosynthesis of CMP-*N,N'*-diacetyllegionaminic acid from UDP-*N,N'*-diacetylbaicillosamine in *Legionella pneumophila*. *Biochemistry*, 47:3272–3282, 2008.
- [929] G.G. Glenner, P.J. McMillan, and J.E. Folk. A mammalian peptidase specific for the hydrolysis of N-terminal  $\alpha$ -L-glutamyl and aspartyl residues. *Nature*, 194:867–867, 1962.
- [930] D. Glick, S. Glaubach, and D.H. Moore. Azolesterase activities of electrophoretically separated proteins of serum. *J. Biol. Chem.*, 144:525–528, 1942.
- [931] J. Glössl and H. Kresse. Impaired degradation of keratan sulphate by Morquio A fibroblasts. *Biochem. J.*, 203:335–338, 1982.
- [932] P. Gluschankof, S. Gomez, A. Morel, and P. Cohen. Enzymes that process somatostatin precursors. A novel endoprotease that cleaves before the arginine-lysine doublet is involved in somatostatin-28 convertase activity of rat brain cortex. *J. Biol. Chem.*, 262:9615–9620, 1987.
- [933] I.Y. Gluzman, S.E. Francis, A. Oksman, C.E. Smith, K.L. Duffin, and D.E. Goldberg. Order and specificity of the *Plasmodium falciparum* hemoglobin degradation pathway. *J. Clin. Invest.*, 93:1602–1608, 1994.
- [934] A.M. Goble, Y. Feng, F.M. Raushel, and J.E. Cronan. Discovery of a cAMP deaminase that quenches cyclic AMP-dependent regulation. *ACS Chem. Biol.*, 8:2622–2629, 2013.
- [935] A.M. Goble, R. Toro, X. Li, A. Ornelas, H. Fan, S. Eswaramoorthy, Y. Patskovsky, B. Hillerich, R. Seidel, A. Sali, B.K. Shoichet, S.C. Almo, S. Swaminathan, M.E. Tanner, and F.M. Raushel. Deamination of 6-aminodeoxyfutasoline in menaquinone biosynthesis by distantly related enzymes. *Biochemistry*, 52:6525–6536, 2013.
- [936] H.M. Goda, K. Ushigusa, H. Ito, N. Okino, H. Narimatsu, and M. Ito. Molecular cloning, expression, and characterization of a novel endo- $\alpha$ -*N*-acetylgalactosaminidase from *Enterococcus faecalis*. *Biochem. Biophys. Res. Commun.*, 375:441–446, 2008.
- [937] E.R. Goedken and S. Marqusee. Metal binding and activation of the ribonuclease H domain from moloney murine leukemia virus. *Protein Eng.*, 12:975–980, 1999.
- [938] D.H. Goetz, Y. Choe, E. Hansell, Y.T. Chen, M. McDowell, C.B. Jonsson, W.R. Roush, J. McKerrow, and C.S. Craik. Substrate specificity profiling and identification of a new class of inhibitor for the major protease of the SARS coronavirus. *Biochemistry*, 46:8744–8752, 2007.
- [939] L.V. Gofshtein-Gandman, A. Keynan, and Y. Milner. Bacteria of the genus *Bacillus* have a hydrolase stereospecific to the D isomer of benzoyl-arginine-*p*-nitroanilide. *J. Bacteriol.*, 170:5895–5900, 1988.
- [940] E. Gohda and H.C. Pitot. A new thiol proteinase from rat liver. *J. Biol. Chem.*, 256:2567–2572, 1981.
- [941] E. Gohda and H.C. Pitot. Purification and characterization of a new thiol proteinase from rat kidney. *Biochim. Biophys. Acta*, 659:114–122, 1981.
- [942] M. Golczak, P.D. Kiser, D.T. Lodowski, A. Maeda, and K. Palczewski. Importance of membrane structural integrity for RPE65 retinoid isomerization activity. *J. Biol. Chem.*, 285:9667–9682, 2010.

- [943] A.L. Goldberg, K.H.S. Swamy, C.H. Chung, and F.S. Larimore. Proteases in *Escherichia coli*. *Methods Enzymol.*, 80:680–702, 1981.
- [944] D.E. Goldberg, A.F.G. Slater, R. Beavis, B. Chait, A. Cerami, and G.B. Henderson. Hemoglobin degradation in the human malaria pathogen *Plasmodium falciparum*: a catabolic pathway initiated by a specific aspartic protease. *J. Exp. Med.*, 173:961–969, 1991.
- [945] G.I. Goldberg, S.M. Wilhelm, A. Kronberger, E.A. Bauer, G.A. Grant, and A.Z. Eisen. Human fibroblast collagenase. Complete primary structure and homology to an oncogene transformation-induced rat protein. *J. Biol. Chem.*, 261:6600–6605, 1986.
- [946] P. Goldman. The enzymatic cleavage of the carbon-fluorine bond in fluoroacetate. *J. Biol. Chem.*, 240:3434–3438, 1965.
- [947] P. Goldman and C.C. Levy. Carboxypeptidase G: purification and properties. *Proc. Natl Acad. Sci. USA*, 58:1299–1306, 1967.
- [948] P. Goldman and G.W.A. Milne. Carbon-fluorine bond cleavage. II. Studies on the mechanism of the defluorination of fluoroacetate. *J. Biol. Chem.*, 241:5557–5559, 1966.
- [949] P. Goldman, G.W.A. Milne, and D.B. Keister. Carbon-halogen bond cleavage. 3. Studies on bacterial halohydrases. *J. Biol. Chem.*, 243:428–434, 1968.
- [950] R. Goldman and J.L. Strominger. Purification and properties of C<sub>55</sub>-isoprenylpyrophosphate phosphatase from *Micrococcus lysodeikticus*. *J. Biol. Chem.*, 247:5116–5122, 1972.
- [951] P.J. Goldmark and S. Liun. Purification and properties of the recBC DNase of *Escherichia coli* K-12. *J. Biol. Chem.*, 247:1849–1860, 1972.
- [952] G.A. Gomez, C. Morisseau, B.D. Hammock, and D.W. Christianson. Structure of human epoxide hydrolase reveals mechanistic inferences on bifunctional catalysis in epoxide and phosphate ester hydrolysis. *Biochemistry*, 43:4716–4723, 2004.
- [953] S. Gomez, P. Gluschankof, A. Morel, and P. Cohen. The somatostatin-28 convertase of rat brain cortex is associated with secretory granule membranes. *J. Biol. Chem.*, 260:10541–10545, 1985.
- [954] G. Gomori. Hexosediphosphatase. *J. Biol. Chem.*, 148:139–149, 1943.
- [955] F. Gong, P. Jemth, M.L. Escobar Galvis, I. Vlodaysky, A. Horner, U. Lindahl, and J.P. Li. Processing of macromolecular heparin by heparanase. *J. Biol. Chem.*, 278:35152–35158, 2003.
- [956] I. Goodman, J.R. Fouts, E. Bresnick, R. Menegas, and G.H. Hitchings. A mammalian thioglucosidase. *Science*, 130:450–451, 1959.
- [957] C.M. Gorbea, P. Marchand, W. Jiang, N.G. Copeland, D.J. Gilbert, N.A. Jenkins, and J.S. Bond. Cloning, expression, and chromosomal localization of the mouse meprin  $\beta$  subunit. *J. Biol. Chem.*, 268:21035–21043, 1993.
- [958] R.W. Gordon, D.J. Manners, and J.R. Stark. The limit dextrinase of the broad bean (*Vicia faba*). *Carbohydr. Res.*, 42:125–134, 1975.
- [959] S.D. Gorham and M. Cantz. Arylsulphatase B, an exo-sulphatase for chondroitin 4-sulphate tetrasaccharide. *Hoppe-Seyler's Z. Physiol. Chem.*, 359:1811–1814, 1978.
- [960] S. Gottesman, W.P. Clark, and M.R. Maurizi. The ATP-dependent Clp protease of *Escherichia coli*. Sequence of clpA and identification of a Clp-specific substrate. *J. Biol. Chem.*, 265:7886–7893, 1990.
- [961] D.B. Grabarczyk and B.C. Berks. Intermediates in the Sox sulfur oxidation pathway are bound to a sulfane conjugate of the carrier protein SoxYZ. *PLoS One*, 12:e0173395–e0173395, 2017.
- [962] D.E. Graham, M. Graupner, H. Xu, and R.H. White. Identification of coenzyme M biosynthetic 2-phosphosulfolactate phosphatase. *Eur. J. Biochem.*, 268:5176–5188, 2001.
- [963] D.E. Graham, H. Xu, and R.H. White. A member of a new class of GTP cyclohydrolases produces formylaminopyrimidine nucleotide monophosphates. *Biochemistry*, 41:15074–15084, 2002.

- [964] C.R. Grant and T. ap Rees. Sorbitol metabolism by apple seedlings. *Phytochemistry*, 20:1505–1511, 1981.
- [965] G.A. Grant, A.Z. Eisen, and R.A. Bradshaw. Collagenolytic protease from fiddler crab (*Uca pugilator*). *Methods Enzymol.*, 80:722–734, 1981.
- [966] D.J. Graves, E.H. Fischer, and E.G. Krebs. Specificity studies on muscle phosphorylase phosphatase. *J. Biol. Chem.*, 235:805–809, 1960.
- [967] C.W. Gray, R.V. Ward, E. Karran, S. Turconi, A. Rowles, D. Viglienghi, C. Southan, A. Barton, K.G. Fantom, A. West, J. Savopoulos, N.J. Hassan, H. Clinkenbeard, C. Hanning, B. Amegadzie, J.B. Davis, C. Dingwall, G.P. Livi, and C.L. Creasy. Characterization of human HtrA2, a novel serine protease involved in the mammalian cellular stress response. *Eur. J. Biochem.*, 267:5699–5710, 2000.
- [968] G.M. Gray and N.A. Santiago. Intestinal surface amino-oligopeptidases. I. Isolation of two weight isomers and their subunits from rat brush border. *J. Biol. Chem.*, 252:4922–4928, 1977.
- [969] N.C.C. Gray and K.P. Strickland. On the specificity of a phospholipase A<sub>2</sub> purified from the 106,000 X g pellet of bovine brain. *Lipids*, 17:91–96, 1982.
- [970] N.C.C. Gray and K.P. Strickland. The purification and characterization of a phospholipase A<sub>2</sub> activity from the 106,000 x g pellet (microsomal fraction) of bovine brain acting on phosphatidylinositol. *Can. J. Biochem.*, 60:108–117, 1982.
- [971] D.M. Greenberg. Arginase. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 4, pages 257–267. Academic Press, New York, 2nd edition, 1960.
- [972] D.M. Greenberg, A.E. Bagot, and O.A. Roholt. Liver arginase. III. Properties of highly purified arginase. *Arch. Biochem. Biophys.*, 62:446–453, 1956.
- [973] D.M. Greenberg and T. Winnick. Plant proteases. I. Activation-inhibition reactions. *J. Biol. Chem.*, 135:761–773, 1940.
- [974] G.R. Greenberg and R.L. Somerville. Deoxyuridylylase activity and deoxyuridinetriphosphatase in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA*, 48:247–257, 1962.
- [975] H.M. Greenblatt, C.A. Ryan, and M.N.G. James. Structure of the complex of *Streptomyces griseus* proteinase B and polypeptide chymotrypsin inhibitor-1 from Russet Burbank potato tubers at 2.1 Å resolution. *J. Mol. Biol.*, 205:201–228, 1989.
- [976] K.J. Gregg and A.B. Boraston. Cloning, recombinant production, crystallization and preliminary X-ray diffraction analysis of a family 101 glycoside hydrolase from *Streptococcus pneumoniae*. *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.*, 65:133–135, 2009.
- [977] H. Greten, R.I. Levy, H. Fales, and D.S. Fredrickson. Hydrolysis of diglyceride and glyceryl monoester diethers with lipoprotein lipase. *Biochim. Biophys. Acta*, 210:39–45, 1970.
- [978] J.K. Griffiths and C.E. Sansom. In *The Transporter Factsbook*. Academic Press, San Diego, 1998.
- [979] S. Grigoryev, A.E. Stewart, Y.T. Kwon, S.M. Arfin, R.A. Bradshaw, N.A. Jenkins, N.G. Copeland, and A. Varshavsky. A mouse amidase specific for N-terminal asparagine. The gene, the enzyme, and their function in the N-end rule pathway. *J. Biol. Chem.*, 271:28521–28532, 1996.
- [980] G.B. Grindey and C.A. Nichol. Mammalian deoxyuridine 5'-triphosphate pyrophosphatase. *Biochim. Biophys. Acta*, 240:180–183, 1971.
- [981] L.S. Grinna and P.W. Robbins. Glycoprotein biosynthesis. Rat liver microsomal glucosidases which process oligosaccharides. *J. Biol. Chem.*, 254:8814–8818, 1979.
- [982] L.S. Grinna and P.W. Robbins. Substrate specificities of rat liver microsomal glucosidases which process glycoproteins. *J. Biol. Chem.*, 255:2255–2258, 1980.
- [983] J.-C. Gripon and J. Hermier. Le système protéolytique de *Penicillium roqueforti*. III. Purification, propriétés et spécificité d'une protéase inhibée par l'E.D.T.A. *Biochimie*, 56:1324–1332, 1974.
- [984] J.C. Gripon, B. Auberger, and J. Lenoir. Metalloproteases from *Penicillium caseicolum* and *P. roqueforti*: comparison of specificity and chemical characterization. *Int. J. Biochem.*, 12:451–455, 1980.

- [985] S.G. Grishutin, A.V. Gusakov, A.V. Markov, B.B. Ustinov, M.V. Semenova, and A.P. Sinityn. Specific xyloglucanases as a new class of polysaccharide-degrading enzymes. *Biochim. Biophys. Acta*, 1674:268–281, 2004.
- [986] D. Groche, A. Becker, I. Schlichting, W. Kabsch, S. Schultz, and A.F.V. Wagner. Isolation and crystallization of functionally competent *Escherichia coli* peptide deformylase forms containing either iron or nickel in the active site. *Biochem. Biophys. Res. Commun.*, 246:342–346, 1998.
- [987] L.L. Grochowski, H. Xu, K. Leung, and R.H. White. Characterization of an Fe<sup>2+</sup>-dependent archaeal-specific GTP cyclohydrolase, MptA, from *Methanocaldococcus jannaschii*. *Biochemistry*, 46:6658–6667, 2007.
- [988] L.L. Grochowski, H. Xu, and R.H. White. An iron(II) dependent formamide hydrolase catalyzes the second step in the archaeal biosynthetic pathway to riboflavin and 7,8-didemethyl-8-hydroxy-5-deazariboflavin. *Biochemistry*, 48:4181–4188, 2009.
- [989] J. Grodberg, M.D. Lundrigan, D.L. Toledo, W.F. Mangel, and J.J. Dunn. Complete nucleotide sequence and deduced amino acid sequence of the *ompT* gene of *Escherichia coli* K-12. *Nucleic Acids Res.*, 16:1209–1209, 1988.
- [990] G. Grogan, G.A. Roberts, D. Bougioukou, N.J. Turner, and S.L. Flitsch. The desymmetrization of bicyclic  $\beta$ -diketones by an enzymatic retro-Claisen reaction. A new reaction of the crotonase superfamily. *J. Biol. Chem.*, 276:12565–12572, 2001.
- [991] M. Groll, L. Ditzel, J. Löwe, D. Stock, M. Bochtler, H.D. Bartunik, and R. Huber. Structure of 20S proteasome from yeast at 2.4Å resolution. *Nature*, 386:463–471, 1997.
- [992] C.H. Gross and S. Shuman. The nucleoside triphosphatase and helicase activities of vaccinia virus NPH-II are essential for virus replication. *J. Virol.*, 72:4729–4736, 1998.
- [993] D. Grossman and K. Lang. Anorganische Poly- und Metaphosphatasen sowie Polyphosphate im tierischen Zellkern. *Biochem. Z.*, 336:351–370, 1962.
- [994] M. Grunberg-Manago. Messenger RNA stability and its role in control of gene expression in bacteria and phages. *Annu. Rev. Genet.*, 33:193–227, 1999.
- [995] P.S. Grushoff, S. Shany, and A.W. Bernheimer. Purification and properties of streptococcal nicotinamide adenine dinucleotide glycohydrolase. *J. Bacteriol.*, 122:599–605, 1975.
- [996] A.L. Guan, A.D. Retzios, G.N., Markland Henderson, , and Jr. Purification and characterization of a fibrinolytic enzyme from venom of southern copperhead snake (*Agkistrodon contortrix contortrix*). *Arch. Biochem. Biophys.*, 289:197–207, 1991.
- [997] A. Guarné, B. Hampoelz, W. Glaser, X. Carpena, J. Torma, I. Fita, and T. Skern. Structural and biochemical features distinguish the foot-and-mouth disease virus leader proteinase from other papain-like enzymes. *J. Mol. Biol.*, 302:1227–1240, 2000.
- [998] M.D. Guillou, M. Camier, and C. Clamagirand. Evidence for the presence of *pro*-oxytocin/neurophysin converting enzyme in the human ovary. *J. Endocrinol.*, 142:345–352, 1994.
- [999] J.M. Gulland and E.M. Jackson. 5-Nucleotidase. *Biochem. J.*, 32:597–601, 1938.
- [1000] J. Gunawan and H. Debuch. Liberation of free aldehyde from 1-(1-alkenyl)-*sn*-glycero-3-phosphoethanolamine (lysoplasmalogen) by rat liver microsomes. *Hoppe-Seyler's Z. Physiol. Chem.*, 362:445–452, 1981.
- [1001] P. Guo, B. Wang, B. Hang, L. Li, W. Ali, J. He, and S. Li. Pyrethroid-degrading *Sphingobium* sp. JZ-2 and the purification and characterization of a novel pyrethroid hydrolase. *Int. Biodeter. Biodegrad.*, 63:1107–1112, 2009.
- [1002] A. Guranowski, H. Jakubowski, and E. Holler. Catabolism of diadenosine 5',5'''-P<sup>1</sup>,P<sup>4</sup>-tetrphosphate in procaryotes. Purification and properties of diadenosine 5',5'''-P<sup>1</sup>,P<sup>4</sup>-tetrphosphate (symmetrical) pyrophosphohydrolase from *Escherichia coli* K12. *J. Biol. Chem.*, 258:14784–14789, 1983.
- [1003] A.B. Guranowski, P.K. Chiang, and G.L. Cantoni. 5'-Methylthioadenosine nucleosidase. Purification and characterization of the enzyme from *Lupinus luteus* seeds. *Eur. J. Biochem.*, 114:293–299, 1981.



- [1004] M. Gurevitz, N. Watson, and D. Apirion. A cleavage site of ribonuclease F. A putative processing endoribonuclease from *Escherichia coli*. *Eur. J. Biochem.*, 124:553–559, 1982.
- [1005] S. Gus-Mayer, H. Brunner, H.A. Schneider-Poetsch, F. Lottspeich, C. Eckerskorn, R. Grimm, and W. Rudiger. The amino acid sequence previously attributed to a protein kinase or a TCP1-related molecular chaperone and co-purified with phytochrome is a  $\beta$ -glucosidase. *FEBS Lett.*, 347:51–54, 1994.
- [1006] S. Gus-Mayer, H. Brunner, H.A. Schneider-Poetsch, and W. Rudiger. Avenacosidase from oat: purification, sequence analysis and biochemical characterization of a new member of the BGA family of  $\beta$ -glucosidases. *Plant Mol. Biol.*, 26:909–921, 1994.
- [1007] M.I. Gutierrez-Roman, M.F. Dunn, R. Tinoco-Valencia, F. Holguin-Melendez, G. Huerta-Palacios, and K. Guillen-Navarro. Potentiation of the synergistic activities of chitinases ChiA, ChiB and ChiC from *Serratia marcescens* CFFSUR-B2 by chitobiase (Chb) and chitin binding protein (CBP). *World J Microbiol Biotechnol*, 30:33–42, 2014.
- [1008] F. Gutierrez-Rosales, M.P. Romero, M. Casanovas, M.J. Motilva, and M.I. Minguez-Mosquera.  $\beta$ -Glucosidase involvement in the formation and transformation of oleuropein during the growth and development of olive fruits (*Olea europaea* L. cv. Arbequina) grown under different farming practices. *J. Agric. Food Chem.*, 60:4348–4358, 2012.
- [1009] J. Gutkowska, P. Corvol, A.F. Figueiredo, T. Inagami, J. Bouhnik, and J. Genest. Kinetic studies of rat renin and tonin on purified rat angiotensinogen. *Can. J. Biochem. Cell Biol.*, 62:137–142, 1984.
- [1010] P. Gutwein, S. Mechttersheimer, S. Riedle, A. Stoeck, D. Gast, S. Joumaa, H. Zentgraf, M. Fogel, and D.P. Altevogt. ADAM10-mediated cleavage of L1 adhesion molecule at the cell surface and in released membrane vesicles. *FASEB J.*, 17:292–294, 2003.
- [1011] H.-W., Sandermann Krell, , and Jr. Plant biochemistry of xenobiotics. Purification and properties of a wheat esterase hydrolyzing the plasticizer chemical, bis(2-ethylhexyl)phthalate. *Eur. J. Biochem.*, 143:57–62, 1984.
- [1012] K. Haakansson, A.H.J. Wang, and C.G. Miller. The structure of aspartyl dipeptidase reveals a unique fold with a Ser-His-Glu catalytic triad. *Proc. Natl. Acad. Sci. USA*, 97:14097–14102, 2000.
- [1013] I. Haase, S. Sarge, B. Illarionov, D. Laudert, H.P. Hohmann, A. Bacher, and M. Fischer. Enzymes from the haloacid dehalogenase (HAD) superfamily catalyse the elusive dephosphorylation step of riboflavin biosynthesis. *Chembiochem*, 14:2272–2275, 2013.
- [1014] R.C. Haberkern and G.L. Cantoni. Studies on a calf thymus ribonuclease specific for ribonucleic acid-deoxyribonucleic acid hybrids. *Biochemistry*, 12:2389–2395, 1973.
- [1015] M.B. Habibi-Najafi and B.H. Lee. Purification and characterization of X-prolyl dipeptidyl peptidase from *Lactobacillus casei* subsp. *casei* LLG. *Appl. Microbiol. Biotechnol.*, 42:280–286, 1994.
- [1016] S.M. Hadi and D.A. Goldthwait. Endonuclease II of *Escherichia coli*. Degradation of partially depurinated deoxyribonucleic acid. *Biochemistry*, 10:4986–4993, 1971.
- [1017] J. Haeggström, J. Meijer, and O. Rådmark. Leukotriene A<sub>4</sub>. Enzymatic conversion into 5,6-dihydroxy-7,9,11,14-eicosatetraenoic acid by mouse liver cytosolic epoxide hydrolase. *J. Biol. Chem.*, 261:6332–6337, 1986.
- [1018] M. Hagemann and N. Erdmann. Activation and pathway of glucosylglycerol biosynthesis in the cyanobacterium *Synechocystis* sp. PCC 6803. *Microbiology*, 140:1427–1431, 1994.
- [1019] M. Hagemann, S. Richter, E. Zuther, and A. Schoor. Characterization of a glucosylglycerol-phosphate-accumulating salt-sensitive mutant of the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Arch. Microbiol.*, 166:83–91, 1996.
- [1020] M. Hagemann, A. Schoor, R. Jeanjean, E. Zuther, and F. Joset. The gene *stpA* from *Synechocystis* sp. strain PCC 6803 encodes for the glucosylglycerol-phosphate phosphatase involved in cyanobacterial salt adaptation. *J. Bacteriol.*, 179:1727–1733, 1997.
- [1021] F.S. Hagen, F.J. Grant, J.L. Kuijper, C.A. Slaughter, C.R. Moomaw, K. Orth, P.J. O’Hara, and R.S. Munford. Expression and characterization of recombinant human acyloxyacyl hydrolase, a leukocyte enzyme that deacylates bacterial lipopolysaccharides. *Biochemistry*, 30:8415–8423, 1991.

- [1022] D.D. Hagerman, H. Rosenberg, A.H. Ennor, P. Schiff, and S. Inove. The isolation and properties of chicken kidney serine ethanolamine phosphate phosphodiesterase. *J. Biol. Chem.*, 240:1108–1012, 1965.
- [1023] Z. Hakki, A.J. Thompson, S. Bellmaine, G. Speciale, G.J. Davies, and S.J. Williams. Structural and kinetic dissection of the endo- $\alpha$ -1,2-mannanase activity of bacterial GH99 glycoside hydrolases from *Bacteroides* spp. *Chemistry*, 21:1966–1977, 2015.
- [1024] S. Halak, T. Basta, S. Burger, M. Contzen, V. Wray, D.H. Pieper, and A. Stolz. 4-sulfomuconolactone hydrolases from *Hydrogenophaga intermedia* S1 and *Agrobacterium radiobacter* S2. *J. Bacteriol.*, 189:6998–7006, 2007.
- [1025] E.E. Haley.  $\beta$ -Aspartyl peptidase from rat liver. *Methods Enzymol.*, 19:737–741, 1970.
- [1026] A. Hall. Small GTP-binding proteins and the regulation of actin cytoskeleton. *Annu. Rev. Cell Biol.*, 10:31–54, 1994.
- [1027] R.S. Hall, A.A. Fedorov, R. Marti-Arbona, E.V. Fedorov, P. Kolb, J.M. Sauder, S.K. Burley, B.K. Shoichet, S.C. Almo, and F.M. Raushel. The hunt for 8-oxoguanine deaminase. *J. Am. Chem. Soc.*, 132:1762–1763, 2010.
- [1028] L.M. Hallcher and W.R. Sherman. The effects of lithium ion and other agents on the activity of *myo*-inositol-1-phosphatase from bovine brain. *J. Biol. Chem.*, 255:10896–10901, 1980.
- [1029] P.C. Hallenbeck, E.R. Vimr, F. Yu, B. Bassler, and F.A. Troy. Purification and properties of a bacteriophage-induced endo-*N*-acetylneuraminidase specific for poly- $\alpha$ -2,8-sialosyl carbohydrate units. *J. Biol. Chem.*, 262:3553–3561, 1987.
- [1030] G. Halliwell, M. Griffin, and R. Vincent. The role of component C<sub>1</sub> in cellulolytic systems. *Biochem. J.*, 127:43P–43P, 1972.
- [1031] Y.S. Halpern and N. Grossowicz. Hydrolysis of amides by extracts from mycobacteria. *Biochem. J.*, 65:716–720, 1957.
- [1032] C.H. Halsted, E.-H. Ling, R. Luthi-Carter, J.A. Villanueva, J.M. Gardner, , and J.T. Folylpoly- $\gamma$ -glutamate carboxypeptidase from pig jejunum: molecular characterization and relation to glutamate carboxypeptidase II. *J. Biol. Chem.*, 273:20417–20424, 1998.
- [1033] Y. Hamano, N. Matsuura, M. Kitamura, and H. Takagi. A novel enzyme conferring streptothricin resistance alters the toxicity of streptothricin D from broad-spectrum to bacteria-specific. *J. Biol. Chem.*, 281:16842–16848, 2006.
- [1034] H. Hamazaki and M.H. Hamazaki. Catalytic site of human protein-glucosylgalactosylhydroxylysine glucosidase: Three crucial carboxyl residues were determined by cloning and site-directed mutagenesis. *Biochem. Biophys. Res. Commun.*, 469:357–362, 2016.
- [1035] H. Hamazaki and K. Hotta. Purification and characterization of an  $\alpha$ -glucosidase specific for hydroxylysine-linked disaccharide of collagen. *J. Biol. Chem.*, 254:9682–9687, 1979.
- [1036] H. Hamazaki and K. Hotta. Enzymatic hydrolysis of disaccharide unit of collagen. Isolation of 2-*O*- $\alpha$ -D-glucopyranosyl-*O*- $\beta$ -D-galactopyranosyl-hydroxylysine glucohydrolase from rat spleens. *Eur. J. Biochem.*, 111:587–591, 1980.
- [1037] S.R. Hamilton, H. Li, H. Wischnewski, A. Prasad, J.S. Kerley-Hamilton, T. Mitchell, A.J. Walling, R.C. Davidson, S. Wildt, and T.U. Gerngross. Intact  $\alpha$ -1,2-endomannosidase is a typical type II membrane protein. *Glycobiology*, 15:615–624, 2005.
- [1038] E. Hammond, C.P. Li, and V. Ferro. Development of a colorimetric assay for heparanase activity suitable for kinetic analysis and inhibitor screening. *Anal. Biochem.*, 396:112–116, 2010.
- [1039] B. Han, G. Luo, Z.Z. Shi, R. Barrios, D. Atwood, W. Liu, G.M. Habib, R.N. Sifers, D.B. Corry, and M.W. Lieberman.  $\gamma$ -glutamyl leukotrienase, a novel endothelial membrane protein, is specifically responsible for leukotriene D<sub>4</sub> formation *in vivo*. *Am J Pathol*, 161:481–490, 2002.
- [1040] G.S. Han, C.N. Johnston, X. Chen, K. Athenstaedt, G. Daum, and G.M. Carman. Regulation of the *Saccharomyces cerevisiae* DPP1-encoded diacylglycerol pyrophosphate phosphatase by zinc. *J. Biol. Chem.*, 276:10126–10133, 2001.
- [1041] K. Hanada, T. Mizutani, M. Yamagishi, H. Tsuji, Misaki, and J. The isolation of collagenase and its enzymological and physico-chemical properties. *Agric. Biol. Chem.*, 37:1771–1781, 1973.
- [1042] D.J. Hanahan, H. Brockerhoff, and E.J. Barron. The site of attack of phospholipase (lecithinase) A on lecithin: a re-evaluation. Position of fatty acids on lecithins and triglycerides. *J. Biol. Chem.*, 235:1917–1923, 1960.

- [1043] D.J. Hanahan and I.L. Chaikoff. On the nature of the phosphorus-containing lipides of cabbage leaves and their relation to a phospholipide-splitting enzyme contained in these leaves. *J. Biol. Chem.*, 172:191–198, 1948.
- [1044] M.H. Hanigan and W.A. Ricketts. Extracellular glutathione is a source of cysteine for cells that express  $\gamma$ -glutamyl transpeptidase. *Biochemistry*, 32:6302–6306, 1993.
- [1045] C. Hanke, J. Hess, G. Schumacher, and W. Goebel. Processing by OmpT of fusion proteins carrying the HlyA transport signal during secretion by the *Escherichia coli* hemolysin transport system. *Mol. Gen. Genet.*, 233:42–48, 1992.
- [1046] L. Hansson, M. Strömqvist, A. Bäckman, P. Wallbrandt, A. Carlstein, and T. Egelrud. Cloning, expression, and characterization of stratum corneum chymotryptic enzyme. A skin-specific human serine proteinase. *J. Biol. Chem.*, 269:19420–19426, 1994.
- [1047] A. Hara and J. Sy. Guanosine 5'-triphosphate, 3'-diphosphate 5'-phosphohydrolase. Purification and substrate specificity. *J. Biol. Chem.*, 258:1678–1683, 1983.
- [1048] J. Hara, K. Higuchi, R. Okamoto, M. Kawashima, and G. Imokawa. High-expression of sphingomyelin deacylase is an important determinant of ceramide deficiency leading to barrier disruption in atopic dermatitis. *J. Invest. Dermatol.*, 115:406–413, 2000.
- [1049] I. Hara-Nishimura. Asparaginyl endopeptidase. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 746–749. Academic Press, London, 1998.
- [1050] S. Harayama, M. Rekik, A. Wasserfallen, and A. Bairoch. Evolutionary relationships between catabolic pathways for aromatics: conservation of gene order and nucleotide sequences of catechol oxidation genes of pWW0 and NAH7 plasmids. *MGG Mol. Gen. Genet.*, 210:241–247, 1987.
- [1051] J. Harder. Anaerobic degradation of cyclohexane-1,2-diol by a new *Azoarcus* species. *Arch. Microbiol.*, 168:199–204, 1997.
- [1052] B. Hardt, C. Volker, S. Mundt, M. Salska-Navarro, M. Hauptmann, and E. Bause. Human endo- $\alpha$ 1,2-mannosidase is a Golgi-resident type II membrane protein. *Biochimie*, 87:169–179, 2005.
- [1053] D.M. Hardy and J.L. Hedrick. Oviductin. Purification and properties of the oviductal protease that processes the molecular weight 43,000 glycoprotein of the *Xenopus laevis* egg envelope. *Biochemistry*, 31:4466–4472, 1992.
- [1054] D.R. Harkness. Studies on human placental alkaline phosphatase. II. Kinetic properties and studies on the apoenzyme. *Arch. Biochem. Biophys.*, 126:513–523, 1968.
- [1055] V. Harmat, P. Gal, J. Kardos, K. Szilagyi, G. Ambrus, B. Vegh, G. Naray-Szabo, and P. Zavodszky. The structure of MBL-associated serine protease-2 reveals that identical substrate specificities of C1s and MASP-2 are realized through different sets of enzyme-substrate interactions. *J. Mol. Biol.*, 342:1533–1546, 2004.
- [1056] N. Harms, J. Ras, W.N. Reijnders, R.J. van Spanning, and A.H. Stouthamer. S-Formylglutathione hydrolase of *Paracoccus denitrificans* is homologous to human esterase D: a universal pathway for formaldehyde detoxification? *J. Bacteriol.*, 178:6296–6299, 1996.
- [1057] N. Harpaz and H. Schachter. Control of glycoprotein synthesis. Processing of asparagine-linked oligosaccharides by one or more rat liver Golgi  $\alpha$ -D-mannosidases dependent on the prior action of UDP-N-acetylglucosamine:  $\alpha$ -D-mannoside  $\beta$ 2-N-acetylglucosaminyltransferase I. *J. Biol. Chem.*, 255:4894–4902, 1980.
- [1058] D.B. Harper. Microbial metabolism of aromatic nitriles. Enzymology of C-N cleavage by *Nocardia* sp. (rhodochrous group) N.C.I.B. 11216. *Biochem. J.*, 165:309–319, 1977.
- [1059] J.W. Harper, R.R. Cook, C.J. Roberts, B.J. McLaughlin, and J.C. Powers. Active site mapping of the serine proteases human leukocyte elastase, cathepsin G, porcine pancreatic elastase, rat mast cell proteases I and II, bovine chymotrypsin A $\alpha$ , and *Staphylococcus aureus* protease V-8 using tripeptide thiobenzyl ester substrates. *Biochemistry*, 23:2995–3002, 1984.
- [1060] J. Harris, D.E. Mason, J. Li, K.W. Burdick, B.J. Backes, T. Chen, A. Shipway, G. Van Heeke, L. Gough, A. Ghaemmaghami, F. Shakib, F. Debaene, and N. Winssinger. Activity profile of dust mite allergen extract using substrate libraries and functional proteomic microarrays. *Chem. Biol.*, 11:1361–1372, 2004.

- [1061] R.B. Harris and I.B. Wilson. Atrial tissue contains a metallo dipeptidyl carboxylhydrolase not present in ventricular tissue: partial purification and characterization. *Arch. Biochem. Biophys.*, 233:667–675, 1984.
- [1062] R.B. Harris and I.B. Wilson. Conversion of atriopeptin II to atriopeptin I by atrial dipeptidyl carboxy hydrolase. *Peptides (Fayetteville)*, 6:393–396, 1985.
- [1063] T.J.R. Harris, P.A. Lowe, A. Lyons, P.G. Thomas, M.A.W. Eaton, T.A. Millican, T.P. Patel, C.C. Bose, N.H. Carey, and M.T. Doel. Molecular cloning and nucleotide sequence of cDNA coding for calf preprochymosin. *Nucleic Acids Res.*, 10:2177–2187, 1982.
- [1064] M. Harun-Or-Rashid, T. Matsuzawa, Y. Satoh, T. Shiraishi, M. Ando, G. Sadik, and Y. Uda. Purification and characterization of  $\alpha$ -N-acetylgalactosaminidases I and II from the starfish *Asterina amurensis*. *Biosci. Biotechnol. Biochem.*, 74:256–261, 2010.
- [1065] I.T. Harvima, N.M. Schechter, R.J. Harvima, and J.E. Fräki. Human skin tryptase: purification, partial characterization and comparison with human lung tryptase. *Biochim. Biophys. Acta*, 957:71–80, 1988.
- [1066] H. Hasegawa, R.D. Bennett, and C.P. Verdon. Metabolism of limonoids via a deoxylimonoid pathway in *Citrus*. *Phytochemistry*, 19:1445–1447, 1980.
- [1067] H. Hasegawa and C.W. Nagel. Isolation of an oligogalacturonate hydrolase from a *Bacillus* species. *Arch. Biochem. Biophys.*, 124:513–520, 1968.
- [1068] S. Hasegawa, J.H. Nordin, and S. Kirkwood. Enzymes that hydrolyze fungal cell wall polysaccharides. I. Purification and properties of an endo- $\alpha$ -D-(1-3)-glucanase from *Trichoderma*. *J. Biol. Chem.*, 244:5460–5470, 1969.
- [1069] W.A. Haseltine, L.K. Gordon, C.P. Lindan, R.H. Grafstrom, N.L. Shaper, and L. Grossman. Cleavage of pyrimidine dimers in specific DNA sequences by a pyrimidine dimer DNA-glycosylase of *M. luteus*. *Nature*, 285:634–641, 1980.
- [1070] W. Hashimoto, E. Kobayashi, H. Nankai, N. Sato, T. Miya, S. Kawai, and K. Murata. Unsaturated glucuronyl hydrolase of *Bacillus* sp. GL1: novel enzyme prerequisite for metabolism of unsaturated oligosaccharides produced by polysaccharide lyases. *Arch. Biochem. Biophys.*, 368:367–374, 1999.
- [1071] Y. Hashitani and J.R. Cooper. The partial purification of thiamine triphosphatase from rat brain. *J. Biol. Chem.*, 247:2117–2199, 1972.
- [1072] K. Hasty. A., Jeffrey, J. J., Hibbs, M. S. and Welgus, H. G. The collagen substrate specificity of human neutrophil collagenase. *J. Biol. Chem.*, 262:10048–10052, 1987.
- [1073] K. Hasty. A., Pourmotabbed, T. F., Goldberg, G. I., Thompson, J. P., Spinella, D. G., Stevens, R. M. and Mainardi, C. L. Human Neutrophil Collagenase. A distinct gene product with homology to other matrix metalloproteinases. *J. Biol. Chem.*, 265:11421–11424, 1990.
- [1074] T. Hata, R. Hayashi, and E. Dot. Purification of yeast proteinases. Part III. Isolation and physicochemical properties of yeast proteinase A and C. *Agric. Biol. Chem.*, 31:357–367, 1967.
- [1075] C. Hatanaka and J. Ozawa. Enzymic degradation of pectic acid. XIII. New exopolygalacturonase producing digalacturonic acid from pectic acid. *J. Agric. Chem. Soc. Jpn.*, 43:764–772, 1968.
- [1076] C. Hatanaka and J. Ozawa. *Ber. des O'Hara Inst.*, 15:47–47, 1971.
- [1077] R. Hatfield and D.J. Nevins. Purification and properties of an endoglucanase isolated from the cell walls of *Zea mays* seedlings. *Carbohydr. Res.*, 148:265–278, 1986.
- [1078] R. Hatfield and D.J. Nevins. Hydrolytic activity and substrate specificity of an endoglucanase from *Zea mays* seedling cell walls. *Plant Physiol.*, 83:203–207, 1987.
- [1079] K. Hatsuzawa, K. Murakami, and K. Nakayama. Molecular and enzymatic properties of furin, a Kex2-like endoprotease involved in precursor cleavage at Arg-X-Lys/Arg-Arg sites. *J. Biochem. (Tokyo)*, 111:296–301, 1992.
- [1080] H.-P. Hauri, A. Quaroni, and K.J. Isselbacher. Biogenesis of intestinal plasma membrane: posttranslational route and cleavage of sucrase-isomaltase. *Proc. Natl. Acad. Sci. USA*, 76:5183–5186, 1979.

- [1081] J.S. Hawker and M.D. Hatch. A specific sucrose phosphatase from plant tissues. *Biochem. J.*, 99:102–107, 1966.
- [1082] R.E. Hawtin, T. Zarkowska, K. Arnold, C.J. Thomas, G.W. Gooday, L.A. King, J.A. Kuzio, and R.D. Possee. Liquefaction of *Autographa californica* nucleopolyhedrovirus-infected insects is dependent on the integrity of virus-encoded chitinase and cathepsin genes. *Virology*, 238:243–253, 1997.
- [1083] O. Hayaishi and A. Kornberg. Metabolism of cytosine, thymine, uracil, and barbituric acid by bacterial enzymes. *J. Biol. Chem.*, 197:717–723, 1952.
- [1084] O. Hayaishi and A. Kornberg. Metabolism of phospholipides by bacterial enzymes. *J. Biol. Chem.*, 206:647–663, 1954.
- [1085] O. Hayaishi, H. Shimazono, M. Katagiri, and Y. Saito. Enzymatic formation of oxalate and acetate from oxaloacetate. *J. Am. Chem. Soc.*, 78:5126–5127, 1956.
- [1086] O. Hayaishi and R.Y. Stanier. The bacterial oxidation of tryptophan. III. Enzymatic activities of cell-free extracts from bacteria employing the aromatic pathway. *J. Bacteriol.*, 62:691–709, 1951.
- [1087] O. Hayaishi, H. Tabor, and T. Hayaishi. *N*-Formimino-L-aspartic acid as an intermediate in the enzymic conversion of imidazoleacetic acid to formylaspartic acid. *J. Biol. Chem.*, 227:161–180, 1957.
- [1088] T. Hayano, K. Sogawa, Y. Ichihara, Y. Fujii-Kuriyama, and K. Takahasi. Primary structure of human pepsinogen C gene. *J. Biol. Chem.*, 263:1382–1385, 1988.
- [1089] K. Hayashi and M. Terada. Some characteristics of hydrolysis of synthetic substrates and proteins by the alkaline proteases from *Aspergillus sojae*. *Agric. Biol. Chem.*, 36:1755–1765, 1972.
- [1090] G.W. Haywood and P.J. Large. 4-Acetamidobutyrate deacetylase in the yeast *Candida boidinii* grown on putrescine or spermidine as sole nitrogen source and its probable role in polyamine catabolism. *J. Gen. Microbiol.*, 132:7–14, 1986.
- [1091] S.M. He, M. Wathier, K. Podzelinska, M. Wong, F.R. McSorley, A. Asfaw, B. Hove-Jensen, Z. Jia, and D.L. Zechel. Structure and mechanism of PhnP, a phosphodiesterase of the carbon-phosphorus lyase pathway. *Biochemistry*, 50:8603–8615, 2011.
- [1092] Z. He and J.C. Spain. Studies of the catabolic pathway of degradation of nitrobenzene by *Pseudomonas pseudoalcaligenes* JS45: removal of the amino group from 2-aminomuconic semialdehyde. *Appl. Environ. Microbiol.*, 63:4839–4843, 1997.
- [1093] Z. He and J.C. Spain. A novel 2-aminomuconate deaminase in the nitrobenzene degradation pathway of *Pseudomonas pseudoalcaligenes* JS45. *J. Bacteriol.*, 180:2502–2506, 1998.
- [1094] T. Heck, H.P. Kohler, M. Limbach, O. Flgel, D. Seebach, and B. Geueke. Enzyme-catalyzed formation of  $\beta$ -peptides:  $\beta$ -peptidyl aminopeptidases BapA and DmpA acting as  $\beta$ -peptide-synthesizing enzymes. *Chem. Biodivers.*, 4:2016–1030, 2007.
- [1095] T. Heck, M. Limbach, B. Geueke, M. Zacharias, J. Gardiner, H.P. Kohler, and D. Seebach. Enzymatic degradation of  $\beta$ - and mixed  $\alpha,\beta$ -oligopeptides. *Chem. Biodivers.*, 3:1325–1348, 2006.
- [1096] S. Hedeager-Sorensen and A.J. Kenny. Proteins of the kidney microvillar membrane. Purification and properties of carboxypeptidase P from pig kidneys. *Biochem. J.*, 229:251–257, 1985.
- [1097] D. Heese, S. Berger, and K.H. Rohm. Nuclear magnetic relaxation studies of the role of the metal ion in  $Mn^{2+}$ -substituted aminoacylase I. *Eur. J. Biochem.*, 188:175–180, 1990.
- [1098] J.H. Hehemann, G. Correc, T. Barbeyron, W. Helbert, M. Czjzek, and G. Michel. Transfer of carbohydrate-active enzymes from marine bacteria to Japanese gut microbiota. *Nature*, 464:908–912, 2010.
- [1099] E.J. Hehre, T. Sawai, C.F. Brewer, M. Nakano, and T. Kanda. Trehalase: stereocomplementary hydrolytic and glucosyl transfer reactions with  $\alpha$ - and  $\beta$ -D-glucosyl fluoride. *Biochemistry*, 21:3090–3097, 1982.
- [1100] R.L. Heimark and E.W. Davie. Bovine and human plasma prekallikrein. *Methods Enzymol.*, 80:157–172, 1981.
- [1101] M.-C. Heindl, S. Fermandjian, and B. Keil. Circular dichroism comparative studies of two bacterial collagenases and thermolysin. *Biochim. Biophys. Acta*, 624:51–59, 1980.

- [1102] E.-A. Heinemeyer and D. Richter. Characterization of the guanosine 5'-triphosphate 3'-diphosphate and guanosine 5'-diphosphate 3'-diphosphate degradation reaction catalyzed by a specific pyrophosphorylase from *Escherichia coli*. *Biochemistry*, 17:5368–5372, 1978.
- [1103] S. Heinel, D. Hartinger, M. Thamhesl, E. Vekiru, R. Krska, G. Schatzmayr, W.D. Moll, and R. Grabherr. Degradation of fumonisin B1 by the consecutive action of two bacterial enzymes. *J. Biotechnol.*, 145:120–129, 2010.
- [1104] V.E. Held and E. Buddecke. Nachweis, Reinigung und Eigenschaften einer Chondroitin-4-Sulfatase aus der Aorta des Rindes. *Hoppe-Seyler's Z. Physiol. Chem.*, 348:1047–1060, 1967.
- [1105] D.M. Helfman and J.F. Kuo. A homogeneous cyclic CMP phosphodiesterase hydrolyzes both pyrimidine and purine cyclic 2':3'- and 3':5'-nucleotides. *J. Biol. Chem.*, 257:1044–1047, 1982.
- [1106] D.M. Helfman, M. Shoji, and J.F. Kuo. Purification to homogeneity and general properties of a novel phosphodiesterase hydrolyzing cyclic CMP and cyclic AMP. *J. Biol. Chem.*, 256:6327–6334, 1981.
- [1107] M. Heller and B. Shapiro. Enzymic hydrolysis of sphingomyelin by rat liver. *Biochem. J.*, 98:763–769, 1966.
- [1108] P.J. Helmsing. Purification and properties of galactolipase. *Biochim. Biophys. Acta*, 178:519–533, 1969.
- [1109] T. Hemscheidt and M.H. Zenk. Glucosidases involved in indole alkaloid biosynthesis of *Catharanthus* cell cultures. *FEBS Lett.*, 110:187–191, 1980.
- [1110] G. Henderson, P. Krygsmann, C.J. Liu, C.C. Davey, and L.T. Malek. Characterization and structure of genes for proteases A and B from *Streptomyces griseus*. *J. Bacteriol.*, 169:3778–3784, 1987.
- [1111] W. Hengstenberg, W.K. Penberthy, and M.L. Morse. Purification of the staphylococcal 6-phospho- $\beta$ -D-galactosidase. *Eur. J. Biochem.*, 14:27–32, 1970.
- [1112] T.D. Hennessey and M.H. Richmond. The purification and some properties of a  $\beta$ -lactamase (cephalosporinase) synthesized by *Enterobacter cloacae*. *Biochem. J.*, 109:469–473, 1968.
- [1113] H.R. Henney and G. Tavana. Purification and some properties of an intracellular acid (carboxyl) proteinase from differentiating haploid cells of *Physarum flavicomum*. *Exp. Mycol.*, 6:161–170, 1982.
- [1114] B. Henrich, S. Becker, U. Schroeder, and R. Plapp. *dcp* gene of *Escherichia coli*: cloning, sequencing, transcript mapping, and characterization of the gene product. *J. Bacteriol.*, 175:7290–7300, 1993.
- [1115] P.J. Hensbergen, O.I. Klychnikov, D. Bakker, I. Dragan, M.L. Kelly, N.P. Minton, J. Corver, E.J. Kuijper, J.W. Drijfhout, and H.C. van Leeuwen. *Clostridium difficile* secreted Pro-Pro endopeptidase PPEP-1 (ZMP1/CD2830) modulates adhesion through cleavage of the collagen binding protein CD2831. *FEBS Lett.*, 589:3952–3958, 2015.
- [1116] P.J. Hensbergen, O.I. Klychnikov, D. Bakker, V.J. van Winden, N. Ras, A.C. Kemp, R.A. Cordfunke, I. Dragan, A.M. Deelder, E.J. Kuijper, J. Corver, J.W. Drijfhout, and H.C. van Leeuwen. A novel secreted metalloprotease (CD2830) from *Clostridium difficile* cleaves specific proline sequences in LPXTG cell surface proteins. *Mol. Cell. Proteomics*, 13:1231–1244, 2014.
- [1117] J. Henseling and K.H. Rohm. Aminoacylase I from hog kidney: anion effects and the pH dependence of kinetic parameters. *Biochim. Biophys. Acta*, 959:370–377, 1988.
- [1118] D. Hensen, D. Sperling, H.G. Truper, D.C. Brune, and C. Dahl. Thiosulphate oxidation in the phototrophic sulphur bacterium *Allochromatium vinosum*. *Mol. Microbiol.*, 62:794–810, 2006.
- [1119] C.A. Henson and D.P. Livingston. , III. Purification and characterization of an oat fructan exohydrolase that preferentially hydrolyzes  $\beta$ -2,6-fructans. *Plant Physiol.*, 110:639–644, 1996.
- [1120] N.L. Hepowitz, S. Uthandi, H.V. Miranda, M. Toniutti, L. Prunetti, O. Olivarez, I.M. De Vera, G.E. Fanucci, S. Chen, and J.A. Maupin-Furlow. Archaeal JAB1/MPN/MOV34 metalloenzyme (HvJAMM1) cleaves ubiquitin-like small archaeal modifier proteins (SAMPs) from protein-conjugates. *Mol. Microbiol.*, 86:971–987, 2012.
- [1121] L.A. Heppel and R.J. Hilme. Purification and properties of 5-nucleotidase. *J. Biol. Chem.*, 188:665–676, 1951.
- [1122] L.A. Heppel and R.J. Hilme. Phosphorolysis and hydrolysis of purine ribosides from yeast. *J. Biol. Chem.*, 198:683–694, 1952.

- [1123] L.A. Heppel and R.J. Hilmo. Mechanism of enzymatic hydrolysis of adenosinetriphosphate. *J. Biol. Chem.*, 202:217–226, 1953.
- [1124] L.A. Heppel, J. Hurwitz, and B.L. Horecker. Adenine deaminase of *Azotobacter vinelandii*. *J. Am. Chem. Soc.*, 79:630–633, 1957.
- [1125] L.A. Heppel and V.T. Porterfield. Enzymatic dehalogenation of certain brominated and chlorinated compounds. *J. Biol. Chem.*, 176:763–769, 1948.
- [1126] D.R. Herbold and L. Glaser. *Bacillus subtilis* N-acetylmuramic acid L-alanine amidase. *J. Biol. Chem.*, 250:1676–1682, 1975.
- [1127] D.R. Herbold and L. Glaser. Interaction of N-acetylmuramic acid L-alanine amidase with cell wall polymers. *J Biol Chem*, 250:7231–7238, 1975.
- [1128] M. Herlan, C. Bornhovd, K. Hell, W. Neupert, and A.S. Reichert. Alternative topogenesis of Mgm1 and mitochondrial morphology depend on ATP and a functional import motor. *J. Cell Biol.*, 165:167–173, 2004.
- [1129] M. Herlan, F. Vogel, C. Bornhovd, W. Neupert, and A.S. Reichert. Processing of Mgm1 by the rhomboid-type protease Pcp1 is required for maintenance of mitochondrial morphology and of mitochondrial DNA. *J. Biol. Chem.*, 278:27781–27788, 2003.
- [1130] J.C. Hermann, R. Marti-Arbona, A.A. Fedorov, E. Fedorov, S.C. Almo, B.K. Shoichet, and F.M. Raushel. Structure-based activity prediction for an enzyme of unknown function. *Nature*, 448:775–779, 2007.
- [1131] M. Hernick, H.A. Gennadios, D.A. Whittington, K.M. Rusche, D.W. Christianson, and C.A. Fierke. UDP-3-O-((R)-3-hydroxymyristoyl)-N-acetylglucosamine deacetylase functions through a general acid-base catalyst pair mechanism. *J. Biol. Chem.*, 280:16969–16978, 2005.
- [1132] L.B. Hersh. 5-Hydroxy-N-methylpyroglutamate synthetase. Purification and mechanism of action. *J. Biol. Chem.*, 245:3526–3535, 1970.
- [1133] L.B. Hersh, L. Tsai, and E.R. Stadtman. The enzymatic synthesis of 5-hydroxy-N-methylpyroglutamic acid. *J. Biol. Chem.*, 244:4677–4683, 1969.
- [1134] W.D.W. Heston. Characterization and glutamyl preferring carboxypeptidase function of prostate specific membrane antigen: a novel folate hydrolase. *Urology*, 49:104–112, 1997.
- [1135] R. Heyworth and P.G. Walker. Almond-emulsin  $\beta$ -D-glucosidase and  $\beta$ -D-galactosidase. *Biochem. J.*, 83:331–335, 1962.
- [1136] M.S. Hibbs, J.R. Hoidal, and A.H. Kang. Expression of a metalloproteinase that degrades native type V collagen and denatured collagens by cultured human alveolar macrophages. *J. Clin. Invest.*, 80:1644–1650, 1987.
- [1137] K.A. Hicks and S.E. Ealick. Biochemical and structural characterization of *Klebsiella pneumoniae* oxamate amidohydrolase in the uric acid degradation pathway. *Acta Crystallogr D Struct Biol*, 72:808–816, 2016.
- [1138] C.F. Higgins. ABC transporters: from microorganisms to man. *Annu. Rev. Cell Biol.*, 8:67–113, 1992.
- [1139] K. Higuchi, J. Hara, R. Okamoto, M. Kawashima, and G. Imokawa. The skin of atopic dermatitis patients contains a novel enzyme, glucosylceramide sphingomyelin deacylase, which cleaves the N-acyl linkage of sphingomyelin and glucosylceramide. *Biochem. J.*, 350:747–756, 2000.
- [1140] D.L. Hill and P. Chambers. The biosynthesis of proline by *Tetrahymena pyriformis*. *Biochim. Biophys. Acta*, 148:435–447, 1967.
- [1141] J. Hill, L. Tyas, L.H. Phylip, J. Kay, B.M. Dunn, and C. Berry. High level expression and characterisation of plasmepsin II, an aspartic proteinase from *Plasmodium falciparum*. *FEBS Lett.*, 352:155–158, 1994.
- [1142] S.R. Himmelhoch. Leucine aminopeptidase from swine kidney. *Methods Enzymol.*, 19:508–513, 1970.
- [1143] Y. Hirabayashi, M. Kimura, M. Matsumoto, K. Yamamoto, S. Kadowaki, and T. Tochikura. A novel glycosphingolipid hydrolyzing enzyme, glycosphingolipid ceramide deacylase, which cleaves the linkage between the fatty acid and sphingosine base in glycosphingolipids. *J. Biochem. (Tokyo)*, 103:1–4, 1988.

- [1144] S. Hiraizumi, U. Spohr, and R.G. Spiro. Ligand affinity chromatographic purification of rat liver Golgi endomannosidase. *J. Biol. Chem.*, 269:4697–4700, 1994.
- [1145] M. Hiramaru, T. Uchida, and F. Egami. Studies on two nucleases and a ribonuclease from *Physarum polycephalum*. Purification and mode of action. *J. Biochem. (Tokyo)*, 65:701–708, 1969.
- [1146] A. Hiramatsu and T. Ouchi. A neutral proteinase from *Streptomyces naraensis*. *J. Biochem. (Tokyo)*, 71:767–781, 1972.
- [1147] N. Hirano, H. Ohshima, and H. Takahashi. Biochemical analysis of the substrate specificity and sequence preference of endonuclease IV from bacteriophage T<sub>4</sub>, a dC-specific endonuclease implicated in restriction of dC-substituted T4 DNA synthesis. *Nucleic Acids Res.*, 34:4743–4751, 2006.
- [1148] T. Hirano, K. Kadokura, T. Ikegami, Y. Shigeta, Y. Kumaki, W. Hakamata, T. Oku, and T. Nishio. Heterodisaccharide 4-*O*-(*N*-acetyl- $\beta$ -D-glucosaminy)-D-glucosamine is a specific inducer of chitinolytic enzyme production in *Vibrios* harboring chitin oligosaccharide deacetylase genes. *Glycobiology*, 19:1046–1053, 2009.
- [1149] T. Hiratsuka, K. Furihata, J. Ishikawa, H. Yamashita, N. Itoh, H. Seto, and T. Dairi. An alternative menaquinone biosynthetic pathway operating in microorganisms. *Science*, 321:1670–1673, 2008.
- [1150] O. Hirayama, H. Matsuda, H. Takeda, K. Maenaka, and H. Takatsuka. Purification and properties of a lipid acyl-hydrolase from potato tubers. *Biochim. Biophys. Acta*, 384:127–137, 1975.
- [1151] I.N. Hirshfeld, H.J. Rosenfeld, Z. Leifer, and W.K. Maas. Isolation and characterization of a mutant of *Escherichia coli* blocked in the synthesis of putrescine. *J. Bacteriol.*, 101:725–730, 1970.
- [1152] G.H. Hitchings and E.A. Falco. The identification of guanine in extracts of *Girella nigricans*. The specificity of guanase. *Proc. Natl. Acad. Sci. USA*, 30:294–297, 1944.
- [1153] M. Hiyoshi, K. Takamune, K. Mita, H. Kubo, Y. Sugimoto, and C. Katagiri. Oviductin, the oviductal protease that mediates gamete interaction by affecting the vitelline coat in *Bufo japonicus*: its molecular cloning and analyses of expression and posttranslational activation. *Dev. Biol.*, 243:176–184, 2002.
- [1154] P.P.K. Ho, B.H. Frank, and P.J. Burek. Crystalline L-asparaginase from *Escherichia coli* B. *Science*, 165:510–512, 1969.
- [1155] N.T. Hoa, J.A. Brannigan, and S.M. Cutting. The PDZ domain of the SpoIVB serine peptidase facilitates multiple functions. *J. Bacteriol.*, 183:4364–4373, 2001.
- [1156] N.T. Hoa, J.A. Brannigan, and S.M. Cutting. The *Bacillus subtilis* signaling protein SpoIVB defines a new family of serine peptidases. *J. Bacteriol.*, 184:191–199, 2002.
- [1157] B. Hofmann, H.J. Hecht, M. Kiess, and D. Schomburg. Crystal structure of a thiol proteinase from *Staphylococcus aureus* V8 in the E-64 inhibitor complex. *Acta Crystallogr. Sect. A (Suppl.)*, 49:102–102, 1993.
- [1158] T. Hofmann. Penicillopepsin. *Methods Enzymol.*, 45:434–450, 1976.
- [1159] P.A. Hohn, N.C. Popescu, R.D. Hanson, G. Salvesen, and T.J. Ley. Genomic organization and chromosomal localization of the human cathepsin G gene. *J. Biol. Chem.*, 264:13412–13419, 1989.
- [1160] Y. Hojima, J. McKenzie, M. van der Rest, and D.J. Prockop. Type I procollagen *N*-proteinase from chick embryo tendons. Purification of a new 500-kDa form of the enzyme and identification of the catalytically active polypeptides. *J. Biol. Chem.*, 264:11336–11345, 1989.
- [1161] Y. Hojima, M. van der Rest, and D.J. Prockop. Type I procollagen carboxyl-terminal proteinase from chick embryo tendons. Purification and characterization. *J. Biol. Chem.*, 260:15996–16003, 1985.
- [1162] M. Holden. The breakdown of chlorophyll by chlorophyllase. *Biochem. J.*, 78:359–364, 1961.
- [1163] C. Holm, T. Osterlund, H. Laurell, and J.A. Contreras. Molecular mechanisms regulating hormone-sensitive lipase and lipolysis. *Annu. Rev. Nutr.*, 20:365–393, 2000.
- [1164] C.E. Holmlund and R.H. Blank. Preparation and properties of a steroid lactonase. *Arch. Biochem. Biophys.*, 109:29–35, 1965.



- [1165] B. Holmquist and B.L. Vallee. Esterase activity of zinc neutral proteases. *Biochemistry*, 15:101–107, 1976.
- [1166] Y. Honda and M. Kitaoka. A family 8 glycoside hydrolase from *Bacillus halodurans* C-125 (BH<sub>2</sub>105) is a reducing end xylose-releasing exo-oligoxyranase. *J. Biol. Chem.*, 279:55097–55103, 2004.
- [1167] H. Hong, C.H. Cui, J.K. Kim, F.X. Jin, S.C. Kim, and W.T. Im. Enzymatic Biotransformation of Ginsenoside Rb1 and Gypenoside XVII into Ginsenosides Rd and F2 by Recombinant  $\beta$ -glucosidase from *Flavobacterium johnsoniae*. *J Ginseng Res*, 36:418–424, 2012.
- [1168] L. Hong, R.T. Turner, G. Koelsch, D. Shin, A.K. Ghosh, and J. Tang. Crystal structure of memapsin 2 ( $\beta$ -secretase) in complex with an inhibitor OM00-3. *Biochemistry*, 41:10963–10967, 2002.
- [1169] M. Honma and T. Shimomura. Metabolism of 1-aminocyclopropane-1-carboxylic acid. *Agric. Biol. Chem.*, 42:1825–1831, 1978.
- [1170] G.J.M. Hoogwinkel, W.A. Veltkamp, B. Overdijk, and J.W. Lisman. Electrophoretic separation of  $\beta$ -N-acetylhexosaminidases of human and bovine brain and liver and of Tay-Sachs brain tissue. *Hoppe-Seylers Z. Physiol. Chem.*, 353:839–841, 1972.
- [1171] R.H. Hook and W.G. Robinson. Ricinine nitrilase. II. Purification and properties. *J. Biol. Chem.*, 239:4263–4267, 1964.
- [1172] N.M. Hooper, J. Hryszko, and A.J. Turner. Purification and characterization of pig kidney aminopeptidase P. *Biochem. J.*, 267:509–515, 1990.
- [1173] N.M. Hooper, J.N. Keen, and A.J. Turner. Characterization of the glycosyl-phosphatidylinositol-anchored human renal dipeptidase reveals that it is more extensively glycosylated than the pig enzyme. *Biochem. J.*, 265:429–433, 1990.
- [1174] D.A. Hopkinson, M.A. Mestriner, J. Cortner, and H. Harris. Esterase D: a new human polymorphism. *Ann. Hum. Genet.*, 37:119–137, 1973.
- [1175] D.J. Hopper, P.J. Chapman, and S. Dagley. The enzymic degradation of alkyl-substituted gentisates, maleates and malates. *Biochem. J.*, 122:29–40, 1971.
- [1176] B.L. Horecker, J. Hurwitz, and L.A. Heppel. The synthesis of ribose 5-pyrophosphate and ribose 5-triphosphate. *J. Am. Chem. Soc.*, 79:701–702, 1957.
- [1177] D.J. Horgan, J.K. Stoops, E.C. Webb, and B. Zerner. Carboxylesterases (EC 3.1.1). A large-scale purification of pig liver carboxylesterase. *Biochemistry*, 8:2000–2006, 1969.
- [1178] T. Horii, T. Ogawa, and H. Ogawa. Nucleotide sequence of the LexA gene of *E. coli*. *Cell*, 23:689–697, 1981.
- [1179] M. Horinouchi, T. Hayashi, H. Koshino, T. Kurita, and T. Kudo. Identification of 9,17-dioxo-1,2,3,4,10,19-hexanorandrostan-5-oic acid, 4-hydroxy-2-oxohexanoic acid, and 2-hydroxyhexa-2,4-dienoic acid and related enzymes involved in testosterone degradation in *Comamonas testosteroni* TA441. *Appl. Environ. Microbiol.*, 71:5275–5281, 2005.
- [1180] S. Hörtensteiner. Chlorophyll degradation during senescence. *Annu. Rev. Plant Biol.*, 57:55–77, 2006.
- [1181] W. Hösel and W. Barz.  $\beta$ -Glucosidases from *Cicer arietinum* L. Purification and Properties of isoflavone-7-O-glucoside-specific  $\beta$ -glucosidases. *Eur. J. Biochem.*, 57:607–616, 1975.
- [1182] W. Hösel, E. Surholt, and E. Borgmann. Characterization of  $\beta$ -glucosidase isoenzymes possibly involved in lignification from chick pea (*Cicer arietinum* L.) cell suspension cultures. *Eur. J. Biochem.*, 84:487–492, 1978.
- [1183] D.J. Hosfield, Y. Guan, B.J. Haas, R.P. Cunningham, and J.A. Tainer. Structure of the DNA repair enzyme endonuclease IV and its DNA complex: double-nucleotide flipping at abasic sites and three-metal-ion catalysis. *Cell*, 98:397–408, 1999.
- [1184] T. Hoshino, C. Nakano, T. Ootsuka, Y. Shinohara, and T. Hara. Substrate specificity of Rv3378c, an enzyme from *Mycobacterium tuberculosis*, and the inhibitory activity of the bicyclic diterpenoids against macrophage phagocytosis. *Org. Biomol. Chem.*, 9:2156–2165, 2011.
- [1185] L.C. Hoskins, E.T. Boulding, and G. Larson. Purification and characterization of blood group A-degrading isoforms of  $\alpha$ -N-acetylgalactosaminidase from *Ruminococcus torques* strain IX-70. *J. Biol. Chem.*, 272:7932–7939, 1997.

- [1186] C.T. Hou and D. Perlman. Microbial transformations of peptide antibiotics. V. Purification and properties of the actinomycin lactonase from *Actinoplanes missouriensis*. *J. Biol. Chem.*, 245:1289–1295, 1970.
- [1187] R.F. Hough and B.L. Bass. Purification of the *Xenopus laevis* double-stranded RNA adenosine deaminase. *J. Biol. Chem.*, 269:9933–9939, 1994.
- [1188] M.-L. Hourdou, M. Guinand, M.-J. Vacheron, G. Michel, L. Denoroy, C. Duez, S. Englebert, B. Joris, G. Weber, and J.-M. Ghuysen. Characterization of the sporulation-related  $\gamma$ -D-glutamyl-(L)-*meso*-diaminopimelic-acid-hydrolysing peptidase I of *Bacillus sphaericus* NCTC 9602 as a member of the metallo(zinc) carboxypeptidase A family. Modular design of the protein. *Biochem. J.*, 292:563–570, 1993.
- [1189] B. Houston, E. Seawright, D. Jefferies, E. Hoogland, D. Lester, C. Whitehead, and C. Farquharson. Identification and cloning of a novel phosphatase expressed at high levels in differentiating growth plate chondrocytes. *Biochim. Biophys. Acta*, 1448:500–506, 1999.
- [1190] B. Hove-Jensen, F.R. McSorley, and D.L. Zechel. Physiological role of *phnP*-specified phosphoribosyl cyclic phosphodiesterase in catabolism of organophosphonic acids by the carbon-phosphorus lyase pathway. *J. Am. Chem. Soc.*, 133:3617–3624, 2011.
- [1191] A. Howard, M.J. Kostura, N. Thornberry, G.J. Ding, G. Limjuco, J. Weidner, J.P. Salley, K.A. Hogquist, D.D. Chaplin, R.A. Mumford, J.A. Schmidt, and M.J. Tocci. IL-1 converting enzyme requires aspartic acid residues for processing of the IL-1 $\beta$  precursor at two distinct sites and does not cleave 31-kDa IL-1 $\alpha$ . *J. Immunol.*, 147:2964–2969, 1991.
- [1192] B.H. Howard, G. Jones, and M.R. Purdom. The pentosanases of some rumen bacteria. *Biochem. J.*, 74:173–180, 1960.
- [1193] M. Howard, J.C. Grimaldi, J.F. Bazan, F.E. Lund, L. Santos-Argumedo, R.M. Parkhouse, T.F. Walseth, and H.C. Lee. Formation and hydrolysis of cyclic ADP-ribose catalyzed by lymphocyte antigen CD38. *Science*, 262:1056–1059, 1993.
- [1194] S. Howell, R.J. Barnaby, T. Rowe, C.I. Ragan, and N.S. Gee. Evidence for at least four different inositol bisphosphatases in bovine brain. *Eur. J. Biochem.*, 183:169–172, 1989.
- [1195] H.L. Hsieh, C.G. Tong, C. Thomas, and S.J. Roux. Light-modulated abundance of an mRNA encoding a calmodulin-regulated, chromatin-associated NTPase in pea. *Plant Mol. Biol.*, 30:135–147, 1996.
- [1196] J.M. Hsieh, K. Tsurulnikov, M.R. Sawaya, N. Magilnick, N. Abuladze, I. Kurtz, J. Abramson, and A. Pushkin. Structures of aminoacylase 3 in complex with acetylated substrates. *Proc. Natl. Acad. Sci. USA*, 107:17962–17967, 2010.
- [1197] I.-N. Hsu, L.T.J. Delbaere, M.N.G. James, and T. Hofmann. Penicillopepsin from *Penicillium janthinellum* crystal structure at 2.8 Å and sequence homology with porcine pepsin. *Nature*, 266:140–144, 1977.
- [1198] J. Hu, Y. Xue, S. Lee, and Y. Ha. The crystal structure of GXGD membrane protease FlaK. *Nature*, 475:528–531, 2011.
- [1199] L. Hu, L.M. Mulfinger, and A.T. Phillips. Purification and properties of formylglutamate amidohydrolase from *Pseudomonas putida*. *J. Bacteriol.*, 169:4696–4702, 1987.
- [1200] Y.J. Hu, Y. Wei, Y. Zhou, P.T.R. Rajagopalan, and D. Pei. Determination of substrate specificity for peptide deformylase through the screening of a combinatorial peptide library. *Biochemistry*, 38:643–650, 1999.
- [1201] A.H.C. Huang, R.A. Moreau, and K.D.F. Liu. Development and properties of a wax ester hydrolase in the cotyledons of jojoba seedlings. *Plant Physiol.*, 61:339–341, 1978.
- [1202] F. Huang, S.F. Haydock, D. Spiteller, T. Mironenko, T.L. Li, D. O'Hagan, P.F. Leadlay, and J.B. Spencer. The gene cluster for fluorometabolite biosynthesis in *Streptomyces cattleya*: a thioesterase confers resistance to fluoroacetyl-coenzyme A. *Chem. Biol.*, 13:475–484, 2006.
- [1203] H. Huang, R. Chopra, G.L. Verdine, and S.C. Harrison. Structure of a covalently trapped catalytic complex of HIV-1 reverse transcriptase: implications for drug resistance. *Science*, 282:1669–1675, 1998.
- [1204] X. Huang, R. Zeng, X. Ding, X. Mao, Y. Ding, Z. Rao, Y. Xie, W. Jiang, and G. Zhao. Affinity alkylation of the Trp-B4 residue of the  $\beta$ -subunit of the glutaryl 7-aminocephalosporanic acid acylase of *Pseudomonas* sp. 130. *J. Biol. Chem.*, 277:10256–10264, 2002.
- [1205] R. Huber and W. Bode. Structural basis of the activation and action of trypsin. *Acc. Chem. Res.*, 11:114–122, 1978.

- [1206] F.M. Huennekens. Enzymatic deacylation of  $N^{10}$ -formyltetrahydrofolic acid. *Fed. Proc.*, 16:199–199, 1957.
- [1207] E.L. Hult, F. Katouno, T. Uchiyama, T. Watanabe, and J. Sugiyama. Molecular directionality in crystalline  $\beta$ -chitin: hydrolysis by chitinases A and B from *Serratia marcescens* 2170. *Biochem. J.*, 388:851–856, 2005.
- [1208] M.C. Hunt, K. Solaas, B.F. Kase, and S.E. Alexson. Characterization of an acyl-coA thioesterase that functions as a major regulator of peroxisomal lipid metabolism. *J. Biol. Chem.*, 277:1128–1138, 2002.
- [1209] J. Hur, S.Y. Kim, H. Kim, S. Cha, M.S. Lee, and K. Suk. Induction of caspase-11 by inflammatory stimuli in rat astrocytes: lipopolysaccharide induction through p38 mitogen-activated protein kinase pathway. *FEBS Lett.*, 507:157–162, 2001.
- [1210] N. Hurion, H. Fromentin, and B. Keil. Specificity of the collagenolytic enzyme from the fungus *Entomophthora coronata*: comparison with the bacterial collagenase from *Achromobacter iophagus*. *Arch. Biochem. Biophys.*, 192:438–445, 1979.
- [1211] J. Hurwitz, L.A. Heppel, and B.L. Horecker. The enzymatic cleavage of adenylic acid to adenine and ribose 5-phosphate. *J. Biol. Chem.*, 226:525–540, 1957.
- [1212] M.S. Huynh and E.E. Snell. Enzymes of vitamin B<sub>6</sub> degradation. Purification and properties of two *N*-acetylamidohydrolases. *J. Biol. Chem.*, 260:2379–2383, 1985.
- [1213] K.Y. Hwang, J.H. Chung, S.H. Kim, Y.S. Han, and Y. Cho. Structure-based identification of a novel NTPase from *Methanococcus jannaschii*. *Nat. Struct. Biol.*, 6:691–696, 1999.
- [1214] S.A. Hyland, S.S. Eveland, and M.S. Anderson. Cloning, expression, and purification of UDP-3-*O*-acyl-GlcNAc deacetylase from *Pseudomonas aeruginosa*: a metalloamidase of the lipid A biosynthesis pathway. *J. Bacteriol.*, 179:2029–2037, 1997.
- [1215] J.W. Hylin and K. Sawai. The enzymatic hydrolysis of *Leucaena glauca* galactomannan. Isolation of crystalline galactomannan depolymerase. *J. Biol. Chem.*, 239:990–992, 1964.
- [1216] J. Hyun, H. Kothari, E. Herm, J. Mortenson, C.R. Treadwell, and G.V. Vahouny. Purification and properties of pancreatic juice cholesterol esterase. *J. Biol. Chem.*, 244:1937–1945, 1969.
- [1217] S.A. Ibrahim, P.J. Lea, and L. Fowden. Preparation and properties of 4-methyleneglutaminase from the leaves of peanut (*Arachis hypogaea*). *Phytochemistry*, 23:1545–1549, 1984.
- [1218] S. Ichihara and F. Uchino. The specificity of acid proteinase from *Acrocylindrium*. *Agric. Biol. Chem.*, 39:423–428, 1975.
- [1219] E. Ichishima, H. Kumagai, and K. Tomoda. Substrate specificity of carboxyl proteinase from *Pycnoporus coccineus*, a wood-deteriorating fungus. *Curr. Microbiol.*, 3:333–337, 1980.
- [1220] E. Ichishima, Y. Yamagata, H. Chiba, K. Sawaguchi, and T. Tanaka. Soluble and bound forms of aminopeptidase in hens egg-yolk. *Agric. Biol. Chem.*, 53:1867–1872, 1989.
- [1221] H. Ide, K. Tedzuka, H. Shimzu, Y. Kimura, A.A. Purmal, S.S. Wallace, and Y.W. Kow. Alpha-deoxyadenosine, a major anoxic radiolysis product of adenine in DNA, is a substrate for *Escherichia coli* endonuclease IV. *Biochemistry*, 33:7842–7847, 1994.
- [1222] N. Iglesias, J.A. Abelenda, M. Rodino, J. Sampedro, G. Revilla, and I. Zarra. Apoplastic glycosidases active against xyloglucan oligosaccharides of *Arabidopsis thaliana*. *Plant Cell Physiol.*, 47:55–63, 2006.
- [1223] M. Ihara, H. Koyama, Y. Uchimura, H. Saitoh, and A. Kikuchi. Noncovalent binding of small ubiquitin-related modifier (SUMO) protease to SUMO is necessary for enzymatic activities and cell growth. *J. Biol. Chem.*, 282:16465–16475, 2007.
- [1224] E. Iida, K. Satou, M. Mishima, C. Kojima, H. Harashima, and H. Kamiya. Amino acid residues involved in substrate recognition of the *Escherichia coli* Orf135 protein. *Biochemistry*, 44:5683–5689, 2005.
- [1225] Y. Iijima, D.R. Gang, E. Fridman, E. Lewinsohn, and E. Pichersky. Characterization of geraniol synthase from the peltate glands of sweet basil. *Plant Physiol.*, 134:370–379, 2004.

- [1226] K. Iio and M. Yamasaki. Specificity of acid proteinase A from *Aspergillus niger* var. macrosporus towards B-chain of performic acid oxidized bovine insulin. *Biochim. Biophys. Acta*, 429:912–924, 1976.
- [1227] Y. Ijima, K. Ogawa, N. Watanabe, T. Usui, M. Ohnishi-Kameyama, T. Nagata, and K. Sakata. Characterization of  $\beta$ -primeverosidase, being concerned with alcoholic aroma formation in tea leaves to be processed into black tea, and preliminary observations on its substrate specificity. *J. Agric. Food Chem.*, 46:1712–1718, 1998.
- [1228] M. Ike, K. Isami, Y. Tanabe, M. Nogawa, W. Ogasawara, H. Okada, and Y. Morikawa. Cloning and heterologous expression of the exo- $\beta$ -D-glucosaminidase-encoding gene (*gls93*) from a filamentous fungus, *Trichoderma reesei* PC-3-7. *Appl. Microbiol. Biotechnol.*, 72:687–695, 2006.
- [1229] F. Ikegami, K. Matsunae, M. Hisamitsu, T. Kurihara, T. Yamamoto, and I. Murakoshi. Purification and properties of a plant  $\beta$ -D-glucuronidase from *Scutellaria* root. *Biol. Pharm. Bull.*, 18:1531–1534, 1995.
- [1230] Y. Ikehara, S. Ogata, and Y. Misumi. Dipeptidyl-peptidase IV from rat liver. *Methods Enzymol.*, 244:215–227, 1994.
- [1231] H. Ikehata. Purification of thiaminase II. *J. Gen. Appl. Microbiol.*, 6:30–39, 1960.
- [1232] H. Ikemura, H. Takagi, and M. Inouye. Requirement of *pro*-sequence for the production of active subtilisin E in *Escherichia coli*. *J. Biol. Chem.*, 262:7859–7864, 1987.
- [1233] A. Imada. Degradation of pyrimidine nucleotides by enzyme systems of *Streptomyces*. II. Pyrimidine 5'-nucleotide phosphoribo(deoxyribo) hydrolase of *Streptomyces virginiae*. *J. Gen. Appl. Microbiol.*, 13:267–278, 1967.
- [1234] A. Imada, M. Kuno, and S. Igarasi. Degradation of pyrimidine nucleotides by enzyme systems of *Streptomyces*. I. Ribose-5-phosphate formation from pyrimidine nucleotides. *J. Gen. Appl. Microbiol.*, 13:255–265, 1967.
- [1235] T. Imai. Purification and properties of nicotinamide mononucleotide amidohydrolase from *Azotobacter vinelandii*. *J. Biochem. (Tokyo)*, 73:139–153, 1973.
- [1236] T. Imai. Isolation and properties of a glycohydrolase specific for nicotinamide mononucleotide from *Azotobacter vinelandii*. *J. Biochem.*, 85:887–899, 1979.
- [1237] T. Imai. Properties of allosteric nicotinamide mononucleotide glycohydrolase from *Azotobacter vinelandii*: activation and inhibition. *J. Biochem.*, 101:163–173, 1987.
- [1238] T. Imai. Purification and characterization of a pyridine nucleotide glycohydrolase from rabbit spleen. *J. Biochem.*, 106:928–937, 1989.
- [1239] A. Imamura, S. Tamura, N. Shimoyawa, Y. Suzuki, Z. Zhang, T. Tsukamoto, T. Orii, N. Kondo, T. Osumi, and Y. Fujiki. Temperature-sensitive mutation in PEX1 moderates the phenotypes of peroxisome deficiency disorders. *Hum. Mol. Genet.*, 7:2089–2094, 1998.
- [1240] R. Imamura, K. Yamanaka, T. Ogura, S. Hiraga, N. Fujita, A. Ishihama, and H. Niki. Identification of the *cpdA* gene encoding cyclic 3',5'-adenosine monophosphate phosphodiesterase in *Escherichia coli*. *J. Biol. Chem.*, 271:25423–25429, 1996.
- [1241] T. Imamura, H. Nitta, Y. Wada, H. Kobayashi, and K. Okamoto. Impaired plasma clottability induction through fibrinogen degradation by ASP, a serine protease released from *Aeromonas sobria*. *FEMS Microbiol. Lett.*, 284:35–42, 2008.
- [1242] T. Imanaka and T. Kuriki. Pattern of action of *Bacillus stearothermophilus* neopullulanase on pullulan. *J. Bacteriol.*, 171:369–374, 1989.
- [1243] M.J. Imber, L.R. Glasgow, and S.V. Pizzo. Purification of an almond emulsin fucosidase on Cibacron blue-sepharose and demonstration of its activity toward fucose-containing glycoproteins. *J. Biol. Chem.*, 257:8205–8210, 1982.
- [1244] J.M. Inacio, I.L. Correia, and I. de Sa-Nogueira. Two distinct arabinofuranosidases contribute to *arabino*-oligosaccharide degradation in *Bacillus subtilis*. *Microbiology*, 154:2719–2729, 2008.
- [1245] C. Inagaki, M. Hara, and X.T. Zeng. A Cl<sup>-</sup> pump in rat brain neurons. *J. Exp. Zool.*, 275:262–268, 1996.
- [1246] T. Inagami. Structure and function of renin. *J. Hypertension*, 2):S3–S8, 1989.

- [1247] T. Inagami and K. Murakami. Pure renin. Isolation from hog kidney and characterization. *J. Biol. Chem.*, 252:2978–2983, 1977.
- [1248] T. Inaoka, G. Bilbe, O. Ishibashi, K. Tezuka, M. Kumegawa, and T. Kokubo. Molecular cloning of human cDNA for cathepsin K: Novel cysteine proteinase predominantly expressed in bone. *Biochem. Biophys. Res. Commun.*, 206:89–96, 1995.
- [1249] T.S. Ingebritsen and P. Cohen. The protein phosphatases involved in cellular regulation. I. Classification and substrate specificities. *Eur. J. Biochem.*, 132:255–261, 1983.
- [1250] D. Ingrosso, A.V. Fowler, J. Bleibaum, and S. Clarke. Specificity of endoproteinase Asp-N (*Pseudomonas fragi*): cleavage at glutamyl residues in two proteins. *Biochem. Biophys. Res. Commun.*, 162:1528–1534, 1989.
- [1251] R.C. Inhorn and P.W. Majerus. Inositol polyphosphate 1-phosphatase from calf brain. Purification and inhibition by  $\text{Li}^+$ ,  $\text{Ca}^{2+}$ , and  $\text{Mn}^{2+}$ . *J. Biol. Chem.*, 262:15946–15952, 1987.
- [1252] M. Inohue, K. Hayashgi, and D.J. Nevins. Polypeptide characteristics and immunological properties of exo- and endoglucanases purified from maize coleoptile cell walls. *J. Plant Physiol.*, 154:334–340, 1999.
- [1253] K. Inoue, A.J. Baldwin, R.L. Shipman, K. Matsui, S.M. Theg, and M. Ohme-Takagi. Complete maturation of the plastid protein translocation channel requires a type I signal peptidase. *J. Cell Biol.*, 171:425–430, 2005.
- [1254] K. Inoue and Y. Ebizuka. Purification and characterization of furostanol glycoside 26-O- $\beta$ -glucosidase from *Costus speciosus* rhizomes. *FEBS Lett.*, 378:157–160, 1996.
- [1255] G.C.J. Irving and D.J. Cosgrove. Inositol phosphate phosphatases of microbiological origin: the inositol pentaphosphate products of *Aspergillus ficuum* phytases. *J. Bacteriol.*, 112:434–438, 1972.
- [1256] G. Isaya, F. Kalousek, and L.E. Rosenberg. Amino-terminal octapeptides function as recognition signals for the mitochondrial intermediate peptidase. *J. Biol. Chem.*, 267:7904–7910, 1992.
- [1257] G. Isaya, F. Kalousek, and L.E. Rosenberg. Sequence analysis of rat mitochondrial intermediate peptidase: similarity to zinc metallopeptidases and to a putative yeast homologue. *Proc. Natl. Acad. Sci. USA*, 89:8317–8321, 1992.
- [1258] M. Ishibashi, J. Arikawa, R. Okamoto, M. Kawashima, Y. Takagi, K. Ohguchi, and G. Imokawa. Abnormal expression of the novel epidermal enzyme, glucosylceramide deacylase, and the accumulation of its enzymatic reaction product, glucosylsphingosine, in the skin of patients with atopic dermatitis. *Lab. Invest.*, 83:397–408, 2003.
- [1259] T. Ishibashi, H. Hayakawa, R. Ito, M. Miyazawa, Y. Yamagata, and M. Sekiguchi. Mammalian enzymes for preventing transcriptional errors caused by oxidative damage. *Nucleic Acids Res.*, 33:3779–3784, 2005.
- [1260] T. Ishibashi, H. Hayakawa, and M. Sekiguchi. A novel mechanism for preventing mutations caused by oxidation of guanine nucleotides. *EMBO Rep.*, 4:479–483, 2003.
- [1261] M. Ishihara and K. Shimizu.  $\alpha$ -(1 $\rightarrow$ 2)-Glucuronidase in the enzymatic saccharification of hardwood xylan .1. Screening of  $\alpha$ -glucuronidase producing fungi. *Mokuzai Gakkaishi*, 34:58–64, 1988.
- [1262] Y. Ishii, Y. Saito, T. Fujimura, H. Sasaki, Y. Noguchi, H. Yamada, M. Niwa, and K. Shimomura. High-level production, chemical modification and site-directed mutagenesis of a cephalosporin C acylase from *Pseudomonas* strain N176. *Eur. J. Biochem.*, 230:773–778, 1995.
- [1263] T. Ishimizu, A. Sasaki, S. Okutani, M. Maeda, M. Yamagishi, and S. Hase. Endo- $\beta$ -mannosidase, a plant enzyme acting on *N*-glycan. Purification, molecular cloning, and characterization. *J. Biol. Chem.*, 279:38555–38562, 2004.
- [1264] Y. Ishihida, H. Shirafiji, M. Kida, and M. Yoneda. Studies on the guanosine degrading system in bacterial cell. III. Preparation and properties of guanosine deaminase. *Agric. Biol. Chem.*, 33:384–390, 1969.
- [1265] W. Ismail, M. El-Said Mohamed, B.L. Wanner, K.A. Datsenko, W. Eisenreich, F. Rohdich, A. Bacher, and G. Fuchs. Functional genomics by NMR spectroscopy. Phenylacetate catabolism in *Escherichia coli*. *Eur. J. Biochem.*, 270:3047–3054, 2003.
- [1266] M. Ito and T. Yamagata. A novel glycosphingolipid-degrading enzyme cleaves the linkage between the oligosaccharide and ceramide of neutral and acidic glycosphingolipids. *J. Biol. Chem.*, 261:14278–14282, 1986.

- [1267] R. Ito, H. Hayakawa, M. Sekiguchi, and T. Ishibashi. Multiple enzyme activities of *Escherichia coli* MutT protein for sanitization of DNA and RNA precursor pools. *Biochemistry*, 44:6670–6674, 2005.
- [1268] R. Ito, M. Sekiguchi, D. Setoyama, Y. Nakatsu, Y. Yamagata, and H. Hayakawa. Cleavage of oxidized guanine nucleotide and ADP sugar by human NUDT5 protein. *J. Biochem.*, 149:731–738, 2011.
- [1269] N. Itoh, N. Tanaka, I. Funakoshi, T. Kawasaki, S. Mihashi, and I. Yamashina. Organisation of the gene for batroxobin, a thrombin-like snake venom enzyme. Homology with the trypsin/kallikrein gene family. *J. Biol. Chem.*, 263:7628–7631, 1988.
- [1270] R. Itoh. Purification and some properties of an IMP-specific 5'-nucleotidase from yeast. *Biochem. J.*, 298:593–598, 1994.
- [1271] R. Itoh, C. Saint-Marc, S. Chaignepain, R. Katahira, J.M. Schmitter, and B. Daignan-Fornier. The yeast ISN1 (YOR155c) gene encodes a new type of IMP-specific 5'-nucleotidase. *BMC Biochem.*, 4:4–4, 2003.
- [1272] T. Itoh, S. Akao, W. Hashimoto, B. Mikami, and K. Murata. Crystal structure of unsaturated glucuronyl hydrolase, responsible for the degradation of glycosaminoglycan, from *Bacillus* sp. GL1 at 1.8 Å resolution. *J. Biol. Chem.*, 279:31804–31812, 2004.
- [1273] T. Itoh, W. Hashimoto, B. Mikami, and K. Murata. Substrate recognition by unsaturated glucuronyl hydrolase from *Bacillus* sp. GL1. *Biochem. Biophys. Res. Commun.*, 344:253–262, 2006.
- [1274] T. Itoh, A. Ochiai, B. Mikami, W. Hashimoto, and K. Murata. A novel glycoside hydrolase family 105: the structure of family 105 unsaturated rhamnogalacturonyl hydrolase complexed with a disaccharide in comparison with family 88 enzyme complexed with the disaccharide. *J. Mol. Biol.*, 360:573–585, 2006.
- [1275] T. Itoh, A. Ochiai, B. Mikami, W. Hashimoto, and K. Murata. Structure of unsaturated rhamnogalacturonyl hydrolase complexed with substrate. *Biochem. Biophys. Res. Commun.*, 347:1021–1029, 2006.
- [1276] Y. Itoh. Cloning and characterization of the *aru* genes encoding enzymes of the catabolic arginine succinyltransferase pathway in *Pseudomonas aeruginosa*. *J. Bacteriol.*, 179:7280–7290, 1997.
- [1277] Y. Itoh, A. Takamura, N. Ito, Y. Maru, H. Sato, N. Suenaga, T. Aoki, and M. Seiki. Homophilic complex formation of MT<sub>1</sub>-MMP facilitates proMMP-2 activation on the cell surface and promotes tumor cell invasion. *EMBO J.*, 20:4782–4793, 2001.
- [1278] L.A. Ivanoff, T. Towatari, J. Ray, B.D. Korant, and S.R. Petteway. Expression and site-specific mutagenesis of the poliovirus 3C protease in *Escherichia coli*. *Proc. Natl Acad. Sci. USA*, 83:5392–5396, 1986.
- [1279] K.A. Ivanov and J. Ziebuhr. Human coronavirus 229E nonstructural protein 13: characterization of duplex-unwinding, nucleoside triphosphatase, and RNA 5'-triphosphatase activities. *J. Virol.*, 78:7833–7838, 2004.
- [1280] A.S. Ivessa, J.Q. Zhou, V.P. Schulz, E.K. Monson, and V.A. Zakian. *Saccharomyces* Rrm3p, a 5' to 3' DNA helicase that promotes replication fork progression through telomeric and subtelomeric DNA. *Genes Dev.*, 16:1383–1396, 2002.
- [1281] T. Iwai, S. Kuramitsu, and R. Masui. The Nudix hydrolase Ndx1 from *Thermus thermophilus* HB8 is a diadenosine hexaphosphate hydrolase with a novel activity. *J. Biol. Chem.*, 279:21732–21739, 2004.
- [1282] A. Iwamura, T. Fukami, R. Higuchi, M. Nakajima, and T. Yokoi. Human  $\alpha/\beta$  hydrolase domain containing 10 (ABHD10) is responsible enzyme for deglucuronidation of mycophenolic acid acyl-glucuronide in liver. *J. Biol. Chem.*, 287:9240–9249, 2012.
- [1283] A. Iwayama, T. Kimura, O. Adachi, and M. Ameyama. Crystallization and characterization of a novel aminopeptidase from *Trichosporon cutaneum*. *Agric. Biol. Chem.*, 47:2483–2493, 1983.
- [1284] T. Iyama, N. Abolhassani, D. Tsuchimoto, M. Nonaka, and Y. Nakabeppu. NUDT16 is a (deoxy)inosine diphosphatase, and its deficiency induces accumulation of single-strand breaks in nuclear DNA and growth arrest. *Nucleic Acids Res.*, 38:4834–4843, 2010.
- [1285] K. Izaki and J.L. Strominger. Biosynthesis of the peptidoglycan of bacterial cell walls. XIV. Purification and properties of two D-alanine carboxypeptidases from *Escherichia coli*. *J. Biol. Chem.*, 243:3193–3201, 1968.

- [1286] M. Izume, S. Nagae, H. Kawagishi, M. Mitsutomi, and A. Ohtakara. Action pattern of *Bacillus* sp. No. 7-M chitosanase on partially *N*-acetylated chitosan. *Biosci. Biotechnol. Biochem.*, 56:448–453, 1992.
- [1287] H.L. Jackman, P.W. Morris, P.A. Deddish, R.A. Skidgel, and E.G. Erdös. Inactivation of endothelin I by deamidase (lysosomal protective protein). *J. Biol. Chem.*, 267:2872–2875, 1992.
- [1288] J.E. Jackman, C.R. Raetz, and C.A. Fierke. UDP-3-*O*-(*R*-3-hydroxymyristoyl)-*N*-acetylglucosamine deacetylase of *Escherichia coli* is a zinc metalloenzyme. *Biochemistry*, 38:1902–1911, 1999.
- [1289] C.M. Jackson and Y. Nemerson. Blood coagulation. *Annu. Rev. Biochem.*, 49:765–811, 1980.
- [1290] M.C. Jackson, C.M. Kucera, and J.F. Lenney. Purification and properties of human serum carnosinase. *Clin. Chim. Acta*, 196:193–206, 1991.
- [1291] J.B. Jacobson and N.O. Kaplan. A reduced pyridine nucleotide pyrophosphatase. *J. Biol. Chem.*, 226:427–437, 1957.
- [1292] N.J. Jacques, J.G. Morrey-Jones, and G.J. Walker. Inducible and constitutive formation of fructanase in batch and continuous cultures of *Streptococcus mutans*. *J. Gen. Microbiol.*, 131:1625–1633, 1985.
- [1293] A. Jacquet, T. Kleinschmidt, A.G. Schnek, Y. Looze, and G. Braunitzer. The thiol proteinases from the latex of *Carica papaya* L. III. The primary structure of chymopapain. *Biol. Chem. Hoppe-Seyler*, 370:425–434, 1989.
- [1294] L. Jaenicke, W. Kunhe, R. Spessert, U. Wahle, and S. Waffenschmidt. Cell-wall lytic enzymes (autolysins) of *Chlamydomonas reinhardtii* are (hydroxy)proline-specific proteases. *Eur. J. Biochem.*, 170:485–491, 1987.
- [1295] W.G. Jaffé. A new vegetable proteolytic enzyme of the papain class. *Rev. Brasil Biol.*, 3:149–157, 1943.
- [1296] W.G. Jaffé. Hurain, a new plant protease from *Hura crepitans*. *J. Biol. Chem.*, 149:1–7, 1943.
- [1297] T. Jahns, R. Schepp, , and H. Purification and characterisation of an enzyme from a strain of *Ochrobactrum anthroπ* that degrades condensation products of urea and formaldehyde (ureaform). *Can. J. Microbiol.*, 43:1111–1117, 1997.
- [1298] W.B. Jakoby. Kynurenine formamidase from *Neurospora*. *J. Biol. Chem.*, 207:657–663, 1954.
- [1299] W.B. Jakoby and D.M. Bonner. Kynureninase from *Neurospora*: interactions of enzyme with substrates, coenzyme and amines. *J. Biol. Chem.*, 205:709–715, 1953.
- [1300] W.B. Jakoby and D.M. Bonner. Kynureninase from *Neurospora*: purification and properties. *J. Biol. Chem.*, 205:699–707, 1953.
- [1301] W.B. Jakoby and T.A. Fjellstedt. Epoxidases. In P.D. Boyer, editor, *The Enzymes*, volume 7, pages 199–212. Academic Press, New York, 3rd edition, 1972.
- [1302] H. Jakubowski and A. Guranowski. Enzymes hydrolyzing ApppA and/or AppppA in higher plants. Purification and some properties of diadenosine triphosphatase, diadenosine tetraphosphatase, and phosphodiesterase from yellow lupin (*Lupinus luteus*) seeds. *J. Biol. Chem.*, 258:9982–9989, 1983.
- [1303] M. Jam, D. Flament, J. Allouch, P. Potin, L. Thion, B. Kloareg, M. Czjzek, W. Helbert, G. Michel, and T. Barbeyron. The endo- $\beta$ -agarases AgaA and AgaB from the marine bacterium *Zobellia galactanivorans*: two paralogue enzymes with different molecular organizations and catalytic behaviours. *Biochem. J.*, 385:703–713, 2005.
- [1304] N. Jambunathan and R. Mahalingam. Analysis of *Arabidopsis* growth factor gene 1 (GFG1) encoding a nudix hydrolase during oxidative signaling. *Planta*, 224:1–11, 2006.
- [1305] M.N.G. James and A.R. Sielecki. Molecular structure of an aspartic proteinase zymogen, porcine pepsinogen, at 1.8 Å resolution. *Nature*, 319:33–38, 1986.
- [1306] M.N.G. James, A.R. Sielecki, G.D. Brayer, L.T.J. Delbaere, and C.-A. Bauer. Structures of product and inhibitor complexes of *Streptomyces griseus* protease A at 1.8 Å resolution. *J. Mol. Biol.*, 144:43–88, 1980.
- [1307] Y.M. Jang, D.W. Kim, T.C. Kang, M.H. Won, N.I. Baek, B.J. Moon, S.Y. Choi, and O.S. Kwon. Human pyridoxal phosphatase. Molecular cloning, functional expression, and tissue distribution. *J. Biol. Chem.*, 278:50040–50046, 2003.

- [1308] E.F. Jansen, M.-D.F. Nutting, and A.K. Balls. The reversible inhibition of acetyl esterase by diisopropyl fluorophosphate and tetraethyl pyrophosphate. *J. Biol. Chem.*, 175:975–987, 1948.
- [1309] A. Jansson, J. Niemi, P. Mantsala, and G. Schneider. Crystal structure of aclacinomycin methylesterase with bound product analogues: implications for anthracycline recognition and mechanism. *J. Biol. Chem.*, 278:39006–39013, 2003.
- [1310] K.-D. Jany, G. Lederer, and B. Mayer. Amino acid sequence of proteinase K from the mold *Tritirachium album* Limber. *FEBS Lett.*, 199:139–144, 1986.
- [1311] J.C., Brown Pascall, , and RHBDL. 2. *Biochem. Biophys. Res. Commun.*, 317:244–252, 2004.
- [1312] F. Jean, A. Basak, J. Dimasio, N.G. Seidah, and C. Lazure. An internally quenched fluorogenic substrate of prohormone convertase 1 and furin leads to a potent prohormone convertase inhibitor. *Biochem. J.*, 307:689–695, 1995.
- [1313] R. Jeanningros, N. Creuzet-Sigal, C. Frixon, and J. Cattaneo. Purification and properties of a debranching enzyme from *Escherichia coli*. *Biochim. Biophys. Acta*, 438:186–199, 1976.
- [1314] P.L. Jeffrey, D.H. Brown, and B.I. Brown. Studies of lysosomal  $\alpha$ -glucosidase. I. Purification and properties of the rat liver enzyme. *Biochemistry*, 9:1403–1415, 1970.
- [1315] P.A. Jekel, W.J. Weijer, and J.J. Beintema. Use of endoproteinase Lys-C from *Lysobacter enzymogenes* in protein sequence analysis. *Anal. Biochem.*, 134:347–354, 1983.
- [1316] D. Jendrossek. Microbial degradation of polyesters. *Adv. Biochem. Eng./Biotechnol.*, 71:293–325, 2001.
- [1317] A.H. Jenkins, G. Schyns, S. Potot, G. Sun, and T.P. Begley. A new thiamin salvage pathway. *Nat. Chem. Biol.*, 3:492–497, 2007.
- [1318] A.L. Jenkins, Y. Zhang, S.E. Ealick, and T.P. Begley. Mutagenesis studies on TenA: a thiamin salvage enzyme from *Bacillus subtilis*. *Bioorg. Chem.*, 36:29–32, 2008.
- [1319] R.E. Jensen and M.P. Yaffe. Import of proteins into yeast mitochondria: the nuclear MAS2 gene encodes a component of the processing protease that is homologous to the MAS1-encoded subunit. *EMBO J.*, 7:3863–3871, 1988.
- [1320] J. Jesty and Y. Nemerson. The activation of bovine coagulation factor X. *Methods Enzymol.*, 45:95–107, 1976.
- [1321] M.-F. Jett, L. Hue, and H.-G. Hers. Pyruvate kinase phosphatase. *FEBS Lett.*, 132:183–186, 1981.
- [1322] J.I. Jimenez, A. Canales, J. Jimenez-Barbero, K. Ginalski, L. Rychlewski, J.L. Garcia, and E. Diaz. Deciphering the genetic determinants for aerobic nicotinic acid degradation: the *nic* cluster from *Pseudomonas putida* KT2440. *Proc. Natl. Acad. Sci. USA*, 105:11329–11334, 2008.
- [1323] X.F. Jin, H.S. Yu, D.M. Wang, T.Q. Liu, C.Y. Liu, D.S. An, W.T. Im, S.G. Kim, and F.X. Jin. Kinetics of a cloned special ginsenosidase hydrolyzing 3-*O*-glucoside of multi-protopanaxadiol-type ginsenosides, named ginsenosidase type III. *J Microbiol Biotechnol*, 22:343–351, 2012.
- [1324] A. Johnner, S. Kunz, M. Linder, Y. Shakur, and T. Seebeck. Cyclic nucleotide specific phosphodiesterases of *Leishmania major*. *BMC Microbiol.*, 6:25–25, 2006.
- [1325] G.D. Johnson and L.B. Hersh. Expression of meprin subunit precursors. Membrane anchoring through the  $\beta$  subunit and mechanism of zymogen activation. *J. Biol. Chem.*, 269:7682–7688, 1994.
- [1326] J.E. Johnson and A. Schneemann. Nodavirus endopeptidase. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 964–967. Handbook of Proteolytic Enzymes, London, 1998.
- [1327] L.A. Johnson, K.E. Moon, and M. Eisenberg. Purification to homogeneity of the human skin chymotryptic proteinase "chymase". *Anal. Biochem.*, 155:358–364, 1986.
- [1328] L.F. Johnson and M.E. Tate. The structure of *myo*-inositol pentaphosphates. *Ann. N.Y. Acad. Sci.*, 165:526–532, 1969.
- [1329] M. Johnson, M.A.G. Kaye, R. Hems, and H.A. Krebs. Enzymic hydrolysis of adenosine phosphates by cobra venom. *Biochem. J.*, 54:625–629, 1953.
- [1330] M.J. Johnson. Isolation and properties of a pure yeast polypeptidase. *J. Biol. Chem.*, 137:575–586, 1941.



- [1331] P. Johnson and L.B. Smillie. The disulfide bridge sequences of a serine protease of wide specificity from *Streptomyces griseus*. *Can. J. Biochem.*, 49:548–562, 1971.
- [1332] M. Johnston, S. Andrews, R. Brinkman, J. Cooper, H. Ding, J. Dover, Z. Du, A. Favello, L. Fulton, S. Gattung, C. Geisel, J. Kirsten, T. Kucaba, L. Hillier, M. Jier, L. Johnston, Y. Langston, P. Latreille, E.J. Louis, Macri, and M. , St.Peter, H., Trevaskis, E., Vaughan, K., Vignati, D., Wilcox, L., Wohldman, P., Waterston, R., Wilson, R., Vaudin, M. Complete nucleotide sequence of *Saccharomyces cerevisiae* chromosome VIII. *Science*, 265:2077–2082, 1994.
- [1333] S.C. Johnston, C.N. Larsen, W.J. Cook, K.D. Wilkinson, and C.P. Hill. Crystal structure of a deubiquitinating enzyme (human UCH-L3) at 1.8Å resolution. *EMBO J.*, 16:3787–3796, 1997.
- [1334] P. Jollès. Lysozyme. In P.D. Boyer, H. Lardy, and K. Myrback, editors, *The Enzymes*, volume 4, pages 431–445. Academic Press, New York, 2nd edition, 1960.
- [1335] C.H. Jones, P. Dexter, A.K. Evans, C. Liu, S.J. Hultgren, and D.E. Hruby. *Escherichia coli* DegP protease cleaves between paired hydrophobic residues in a natural substrate: the PapA pilin. *J. Bacteriol.*, 184:5762–5771, 2002.
- [1336] G.H. Jones and C.E. Ballou. Studies on the structure of yeast mannan. I. Purification and some properties of an  $\alpha$ -mannosidase from an *Arthrobacter* species. *J. Biol. Chem.*, 244:1043–1051, 1969.
- [1337] G.H. Jones and C.E. Ballou. Studies on the structure of yeast mannan. II. Mode of action of the *Arthrobacter*  $\alpha$ -mannosidase on yeast mannan. *J. Biol. Chem.*, 244:1052–1059, 1969.
- [1338] N.R. Jones. The free amino acids of fish. 1-Methylhistidine and  $\beta$ -alanine liberation by skeletal muscle anserinase of codling (*Gadus callarias*). *Biochem. J.*, 60:81–87, 1955.
- [1339] J.A. Jorge, S.G. Kinney, and J.L. Reissig. Purification and characterization of *Neurospora crassa* N-acetyl galactosaminoglycan deacetylase. *Braz. J. Med. Biol. Res.*, 15:29–34, 1982.
- [1340] B. Joris, J. Van Beeumen, F. Casagrande, C. Gerday, J.-M. Frère, and J.-M. Ghuysen. The complete amino acid sequence of the Zn<sup>2+</sup>-containing D-alanyl-D-alanine-cleaving carboxypeptidase of *Streptomyces albus* G. *Eur. J. Biochem.*, 130:53–69, 1983.
- [1341] L.J. Joseph, L.C. Chang, D. Stamenkovich, and V.P. Sukhatme. Complete nucleotide and deduced amino acid sequences of human and murine preprocathepsin L. An abundant transcript induced by transformation of fibroblasts. *J. Clin. Invest.*, 81:1621–1629, 1988.
- [1342] J.-P. Jost and R.M. Bock. Enzymatic hydrolysis of N-substituted aminoacyl transfer ribonucleic acid in yeast. *J. Biol. Chem.*, 244:5866–5873, 1969.
- [1343] C. Joudiou, K.M. Carvalho, G. Camarao, H. Boussetta, and P. Cohen. Characterization of the thermolysin-like cleavage of biologically active peptides by *Xenopus laevis* peptide hormone inactivating enzyme. *Biochemistry*, 32:5959–5966, 1993.
- [1344] G.W. Jourdain, A. Swanson, D. Watson, and S. Roseman. N-Acetylneuraminic (sialic) acid 9-phosphatase. *Methods Enzymol.*, 8:205–208, 1966.
- [1345] B.K. Joyce and S. Grisolia. Purification and properties of a nonspecific acid phosphatase from wheat germ. *J. Biol. Chem.*, 235:2278–2281, 1960.
- [1346] D. Julius, A. Brake, L. Blair, R. Kunisawa, and J. Thorner. Isolation of the putative structural gene for the lysine-arginine-cleaving endopeptidase required for processing of yeast prepro- $\alpha$ -factor. *Cell*, 37:1075–1089, 1984.
- [1347] M.A. Junaid, G.X. Wu, and R.K. Pullarkat. Purification and characterization of bovine brain lysosomal pepstatin-insensitive proteinase, the gene product deficient in the human late-infantile neuronal ceroid lipofuscinosis. *J. Neurochem.*, 74:287–294, 2000.
- [1348] P. Juneja, F. Hubrich, K. Diederichs, W. Welte, and J.N. Andexer. Mechanistic implications for the chorismatase FkbO based on the crystal structure. *J. Mol. Biol.*, 426:105–115, 2014.
- [1349] R.A. Jupp, A.D. Richards, J. Kay, B.M. Dunn, J.B. Wyckoff, I.M. Samloff, and K. Yamamoto. Identification of the aspartic proteinases from human erythrocyte membranes and gastric mucosa (slow-moving proteinase) as catalytically equivalent to cathepsin E. *Biochem. J.*, 254:895–898, 1988.

- [1350] L. Jurasek, D. Fackre, and L.B. Smillie. Remarkable homology about the disulfide bridges of a trypsin-like enzyme from *Streptomyces griseus*. *Biochem. Biophys. Res. Commun.*, 37:99–105, 1969.
- [1351] Y. Kabuyama, N. Nakatsu, Y. Homma, , and Y. Purification and characterization of phosphatidyl inositol-3,4,5-trisphosphate phosphatase in bovine thymus. *Eur. J. Biochem.*, 238:350–356, 1996.
- [1352] K. Kadokura, A. Rokutani, M. Yamamoto, T. Ikegami, H. Sugita, S. Itoi, W. Hakamata, T. Oku, and T. Nishio. Purification and characterization of *Vibrio parahaemolyticus* extracellular chitinase and chitin oligosaccharide deacetylase involved in the production of heterodisaccharide from chitin. *Appl. Microbiol. Biotechnol.*, 75:357–365, 2007.
- [1353] K. Kainuma, K. Wako, A. Kobayashi, A. Nogami, and S. Suzuki. Purification and some properties of a novel maltohexaose-producing exo-amylase from *Aerobacter aerogenes*. *Biochim. Biophys. Acta*, 410:333–346, 1975.
- [1354] A. Kaji and T. Saheki. Endo-arabinanase from *Bacillus subtilis* F-11. *Biochim. Biophys. Acta*, 410:354–360, 1975.
- [1355] A. Kaji and K. Tagawa. Purification, crystallization and amino acid composition of  $\alpha$ -L-arabinofuranosidase from *Aspergillus niger*. *Biochim. Biophys. Acta*, 207:456–464, 1970.
- [1356] A. Kaji and O. Yoshihara. Properties of purified  $\alpha$ -L-arabinofuranosidase from *Corticium rolfsii*. *Biochim. Biophys. Acta*, 250:367–371, 1971.
- [1357] Y. Kajiwara, P.J. Santander, C.A. Roessner, L.M. Perez, and A.I. Scott. Genetically engineered synthesis and structural characterization of cobalt-precorrin 5A and -5B, two new intermediates on the anaerobic pathway to vitamin B<sub>12</sub>: definition of the roles of the CbiF and CbiG enzymes. *J. Am. Chem. Soc.*, 128:9971–9978, 2006.
- [1358] S. Kakudo, S. Negoro, I. Urabe, and H. Okada. Nylon oligomer degradation gene, *nylC*, on plasmid pOAD2 from a *Flavobacterium* strain encodes endo-type 6-aminohexanoate oligomer hydrolase: purification and characterization of the *nylC* gene product. *Appl. Environ. Microbiol.*, 59:3978–3980, 1993.
- [1359] S. Kakugawa, P.F. Langton, M. Zebisch, S.A. Howell, T.H. Chang, Y. Liu, T. Feizi, G. Bineva, N. O'Reilly, A.P. Snijders, E.Y. Jones, and J.P. Vincent. Notum deacylates Wnt proteins to suppress signalling activity. *Nature*, 2015.
- [1360] T. Kakuma, J. Nishida, T. Tsuzuki, and M. Sekiguchi. Mouse MTH1 protein with 8-oxo-7,8-dihydro-2'-deoxyguanosine 5'-triphosphatase activity that prevents transversion mutation. cDNA cloning and tissue distribution. *J. Biol. Chem.*, 270:25942–25948, 1995.
- [1361] H.M. Kalckar. Differential spectrophotometry of purine compounds by means of specific enzymes. III. Studies of the enzymes of purine metabolism. *J. Biol. Chem.*, 167:461–475, 1947.
- [1362] H.M. Kalckar. Biosynthetic aspects of nucleosides and nucleic acids. *Pubbl. Staz. Zool. (Napoli)*, 23:87–103, 1951.
- [1363] G.F. Kalf and S.V. Rieder. The preparation and properties of trehalase. *J. Biol. Chem.*, 230:691–698, 1958.
- [1364] M. Kalinowska and Z.A. Wojciechowski. Purification and some properties of steryl  $\beta$ -D-glucoside hydrolase from *Sinapis alba* seedlings. *Phytochemistry*, 17:1533–1537, 1978.
- [1365] F. Kalousek, W. Neupert, T. Omura, G. Schatz, and U.K. Schmitz. Uniform nomenclature for the mitochondrial peptidases cleaving precursors of mitochondrial proteins. *Trends Biochem. Sci.*, 18:249–249, 1993.
- [1366] N.A. Kalsheker, S. Deam, L. Chambers, S. Sreedharan, K. Brocklehurst, and D.A. Lomas. The house dust mite allergen Der p1 catalytically inactivates  $\alpha$ 1-antitrypsin by specific reactive centre loop cleavage: a mechanism that promotes airway inflammation and asthma. *Biochem. Biophys. Res. Commun.*, 221:59–61, 1996.
- [1367] C.-M. Kam, J.E. Kerrigan, K.M. Dolman, R. Goldschmeding, A.E.G.K. von dem Borne, and J.C. Powers. Substrate and inhibitor studies on proteinase 3. *FEBS Lett.*, 297:119–123, 1992.
- [1368] M. Kamada, K. Oda, and S. Murao. The purification of the extracellular acid protease of *Rhodotorula glutinis* K-24 and its general properties. *Agric. Biol. Chem.*, 36:1095–1101, 1972.
- [1369] S. Kamada, Y. Funahashi, and Y. Tsujimoto. Caspase-4 and caspase-5, members of the ICE/CED-3 family of cysteine proteases, are CrmA-inhibitable proteases. *Cell Death Differ.*, 4:473–478, 1997.
- [1370] S.S. Kamat, H.J. Williams, and F.M. Raushel. Intermediates in the transformation of phosphonates to phosphate by bacteria. *Nature*, 480:570–573, 2011.

- [1371] J. Kamens, M. Paskind, M. Hugunin, R.V. Talanian, H. Allen, D. Banach, N. Bump, M. Hackett, C.G. Johnston, P. Li, J.A. Mankovich, M. Terranova, and T. Ghayur. Identification and characterization of ICH-2, a novel member of the interleukin-1 $\beta$ -converting enzyme family of cysteine proteases. *J. Biol. Chem.*, 270:15250–15256, 1995.
- [1372] E. Kaminskas, Y. Kimhi, and B. Magasanik. Urocanase and *N*-formimino-L-glutamate formiminohydrolase of *Bacillus subtilis*, two enzymes of the histidine degradation pathway. *J. Biol. Chem.*, 245:3536–3544, 1970.
- [1373] H. Kamiya, M. Hori, T. Arimori, M. Sekiguchi, Y. Yamagata, and H. Harashima. NUDT5 hydrolyzes oxidized deoxyribonucleoside diphosphates with broad substrate specificity. *DNA Repair (Amst)*, 8:1250–1254, 2009.
- [1374] H. Kamiya, E. Iida, and H. Harashima. Important amino acids in the phosphohydrolase module of *Escherichia coli* Orf135. *Biochem. Biophys. Res. Commun.*, 323:1063–1068, 2004.
- [1375] I.G. Kamphuis, J. Drenth, and E.N. Baker. Thiol proteases. Comparative studies based on the high-resolution structures of papain and actinidin, and on amino acid sequence information for cathepsins B and H, and stem bromelain. *J. Mol. Biol.*, 182:317–329, 1985.
- [1376] A. Kanai, A. Sato, Y. Fukuda, K. Okada, T. Matsuda, T. Sakamoto, Y. Muto, S. Yokoyama, G. Kawai, and M. Tomita. Characterization of a heat-stable enzyme possessing GTP-dependent RNA ligase activity from a hyperthermophilic archaeon, *Pyrococcus furiosus*. *RNA*, 15:420–431, 2009.
- [1377] T. Kanamori, N. Kanou, S. Kusakabe, H. Atomi, and T. Imanaka. Allophanate hydrolase of *Oleomonas sagaranensis* involved in an ATP-dependent degradation pathway specific to urea. *FEMS Microbiol. Lett.*, 245:61–65, 2005.
- [1378] A. Kanatani, T. Masuda, T. Shimoda, F. Misoka, X.S. Lin, T. Yoshimoto, and D. Tsuru. Protease II from *Escherichia coli*: sequencing and expression of the enzyme gene and characterization of the expressed enzyme. *J. Biochem. (Tokyo)*, 110:315–320, 1991.
- [1379] M. Kaneda, H. Ohmine, H. Yonezawa, and N. Tominaga. Amino acid sequence around the reactive site of cucumisin from melon fruit. *J. Biochem. (Tokyo)*, 95:825–829, 1984.
- [1380] M. Kaneda and N. Tominaga. Isolation and characterization of a proteinase from the sarcocarp of melon fruit. *J. Biochem. (Tokyo)*, 78:1287–1296, 1975.
- [1381] N. Kaneda, Y. Minematsu, J.C. Powers, and N. Tominaga. Specificity of cucumisin in hydrolysis of peptide thiobenzyl esters. *Agric. Biol. Chem.*, 50:1075–1076, 1986.
- [1382] M. Kanemori, K. Nishihara, H. Yanagi, and T. Yura. Synergistic roles of HslVU and other ATP-dependent proteases in controlling *in vivo* turnover of  $\sigma^{32}$  and abnormal proteins in *Escherichia coli*. *J. Bacteriol.*, 179:7219–7225, 1997.
- [1383] J.N. Kanfer, O.M. Young, D. Shapiro, and R.O. Brady. The metabolism of sphingomyelin. I. Purification and properties of a sphingomyelin-cleaving enzyme from rat liver tissue. *J. Biol. Chem.*, 241:1081–1084, 1966.
- [1384] B.H. Kang, E. Ko, O.K. Kwon, and K.Y. Choi. The structure of procaspase 6 is similar to that of active mature caspase 6. *Biochem. J.*, 364:629–634, 2002.
- [1385] E.J. Kang, S.O. Lee, J.D. Lee, T.H. Lee, and T.H. Lee. Purification and characterization of a levanbiose-producing levanase from *Pseudomonas* sp. No. 43. *Biotechnol. Appl. Biochem.*, 29:263–268, 1999.
- [1386] S.J. Kang, S. Wang, H. Hara, E.P. Peterson, S. Namura, S. Amin-Hanjani, Z. Huang, A. Srinivasan, K.J. Tomaselli, N.A. Thornberry, M.A. Moskowitz, and J. Yuan. Dual role of caspase-11 in mediating activation of caspase-1 and caspase-3 under pathological conditions. *J. Cell. Biol.*, 149:613–622, 2000.
- [1387] Y.J. Kao, W.L. Starnes, and F.J. Behal. Human kidney alanine aminopeptidase: physical and kinetic properties of a sialic acid containing glycoprotein. *Biochemistry*, 17:2990–2994, 1978.
- [1388] N.O. Kaplan, S.P. Colowick, and M.M. Ciotti. Enzymatic deamination of cytosine nucleosides. *J. Biol. Chem.*, 194:579–591, 1952.
- [1389] J.S. Karns. Gene sequence and properties of an *s*-triazine ring-cleavage enzyme from *Pseudomonas* sp. strain NRRLB-12227. *Appl. Environ. Microbiol.*, 65:3512–3517, 1999.

- [1390] P. Karran, T. Hjelmgren, and T. Lindahl. Induction of a DNA glycosylase for *N*-methylated purines is part of the adaptive response to alkylating agents. *Nature*, 296:770–773, 1982.
- [1391] P. Karran and T. Lindahl. Enzymatic excision of free hypoxanthine from polydeoxynucleotides and DNA containing deoxyinosine monophosphate residues. *J. Biol. Chem.*, 253:5877–5879, 1978.
- [1392] K. Kasai, T. Uchida, F. Egami, K. Yoshida, and M. Nomoto. Purification and crystallization of ribonuclease N<sub>1</sub> from *Neurospora crassa*. *J. Biochem. (Tokyo)*, 66:389–396, 1969.
- [1393] K.D. Kasschau and J.C. Carrington. Requirement for HC-Pro processing during genome amplification of tobacco etch potyvirus. *Virology*, 209:268–273, 1995.
- [1394] Y. Katayama, Y. Matsushita, M. Kaneko, M. Kondo, T. Mizuno, and H. Nyunoya. Cloning of genes coding for the three subunits of thiocyanate hydrolase of *Thiobacillus thioparus* THI 115 and their evolutionary relationships to nitrile hydratase. *J. Bacteriol.*, 180:2583–2589, 1998.
- [1395] Y. Katayama, Y. Narahara, Y. Inoue, F. Amano, T. Kanagawa, and H. Kuraishi. A thiocyanate hydrolase of *Thiobacillus thioparus*. *J. Biol. Chem.*, 267:9170–9175, 1992.
- [1396] W.M. Kati, K.A. Johnson, L.F. Jerva, and K.S. Anderson. Mechanism and fidelity of HIV reverse transcriptase. *J. Biol. Chem.*, 267:25988–25997, 1992.
- [1397] H. Kato, K. Enjyoji, T. Miyata, I. Hayashi, S. Oh-Ishi, and S. Iwanaga. Demonstration of arginyl-bradykinin moiety in rat HMW kininogen: direct evidence for liberation of bradykinin by rat glandular kallikreins. *Biochem. Biophys. Res. Commun.*, 127:289–295, 1985.
- [1398] H. Kato, E. Nakanishi, K. Enjyoji, I. Hayashi, S. Oh-ishi, and S. Iwanaga. Characterization of serine proteinases isolated from rat submaxillary gland: with special reference to the degradation of rat kininogens by these enzymes. *J. Biochem. (Tokyo)*, 102:1389–1404, 1987.
- [1399] K. Kato, K. Kawahara, T. Takahashi, and A. Kakinuma. Purification of an  $\alpha$ -amino acid ester hydrolase from *Xanthomonas citri*. *Agric. Biol. Chem.*, 44:1069–1074, 1980.
- [1400] K. Kato, K. Kawahara, T. Takahashi, and A. Kakinuma. Substrate specificity of an  $\alpha$ -amino acid ester hydrolase from *Xanthomonas citri*. *Agric. Biol. Chem.*, 44:1075–1081, 1980.
- [1401] S. Kato, H. Ueda, S. Nonomura, and C. Tatsumi. [Degradation of dehydroacetic acid by microorganisms. III. Properties of triacetic acid lactone hydrolase.]. *Nippon Nogei Kagaku Kaishi*, 42:596–600, 1968.
- [1402] Y. Kato, J. Matsushita, T. Kubodera, and K. Matsuda. A novel enzyme producing isoprimeverose from oligoxyloglucans of *Aspergillus oryzae*. *J. Biochem. (Tokyo)*, 97:801–810, 1985.
- [1403] H. Kato-Noguchi and Y. Tanaka. Effect of ABA- $\beta$ -D-glucopyranosyl ester and activity of ABA- $\beta$ -D-glucosidase in *Arabidopsis thaliana*. *J. Plant Physiol.*, 165:788–790, 2008.
- [1404] S. Kauppinen, S. Christgau, L.V. Kofod, T. Halkier, K. Dorreich, and H. Dalboge. Molecular cloning and characterization of a rhamnogalacturonan acetyltransferase from *Aspergillus aculeatus*. Synergism between rhamnogalacturonan degrading enzymes. *J. Biol. Chem.*, 270:27172–27178, 1995.
- [1405] M. Kawada, Y. Kagawa, H. Takiguchi, and N. Shimazono. Purification of 6-phosphogluconolactonase from rat liver and yeast; its separation from gluconolactonase. *Biochim. Biophys. Acta*, 57:404–407, 1962.
- [1406] K. Kawamoto, I. Horibe, and K. Uchida. Purification and characterization of a new hydrolase for conjugated bile acids, chenodeoxycholytaurine hydrolase, from *Bacteroides vulgatus*. *J. Biochem. (Tokyo)*, 106:1049–1053, 1989.
- [1407] I. Kawashima, T. Tani, K. Shimoda, and Y. Takiguchi. Characterization of pancreatic elastase II cDNAs: two elastase II mRNAs are expressed in human pancreas. *DNA*, 6:163–172, 1987.
- [1408] Y. Kazama, T. Hamamoto, D.C. Foster, and W. Kisiel. Hepsin, a putative membrane-associated serine protease, activates human factor VII and initiates a pathway of blood coagulation on the cell surface leading to thrombin formation. *J. Biol. Chem.*, 270:66–72, 1995.

- [1409] E.L. Kean and K.J. Bighouse. Cytidine 5'-monophosphosialic acid hydrolase. Subcellular location and properties. *J. Biol. Chem.*, 249:7813–7823, 1974.
- [1410] L.P. Keegan, A.P. Gerber, J. Brindle, R. Leemans, A. Gallo, W. Keller, and M.A. O'Connell. The properties of a tRNA-specific adenosine deaminase from *Drosophila melanogaster* support an evolutionary link between pre-mRNA editing and tRNA modification. *Mol. Cell Biol.*, 20:825–833, 2000.
- [1411] M.R. Kehry, T.G. Doak, and F.W. Dahlquist. Stimulus-induced changes in methyltransferase activity during chemotaxis in *Escherichia coli*. *J. Biol. Chem.*, 259:11828–11835, 1984.
- [1412] K.C. Keiler and R.T. Sauer. Tsp protease. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 460–461. Handbook of Proteolytic Enzymes, London, 1998.
- [1413] S. Keime, I.M. Adham, and W. Engel. Nucleotide sequence and exon-intron organisation of the human proacrosin gene. *Eur. J. Biochem.*, 190:195–200, 1990.
- [1414] G.M. Kellerman. Isolation and characteristics of the enzyme acyl 5'-nucleotidase. *Biochim. Biophys. Acta*, 33:101–105, 1959.
- [1415] R.B. Kelley, M.R. Atkinson, J.A. Huberman, and A. Kornberg. Excision of thymine dimers and other mismatched sequences by DNA polymerases of *Escherichia coli*. *Nature*, 224:495–501, 1969.
- [1416] J.A. Kelly, E.L. Neidle, and A. Neidle. An aminopeptidase from mouse brain cytosol that cleaves N-terminal acidic amino acid residues. *J. Neurochem.*, 40:1727–1734, 1983.
- [1417] J.J. Kelly and D.H. Alpers. Properties of human intestinal glucoamylase. *Biochim. Biophys. Acta*, 315:113–122, 1973.
- [1418] S.J. Kelly, D.E. Dardinger, and L.G. Butler. Hydrolysis of phosphonate esters catalyzed by 5'-nucleotide phosphodiesterase. *Biochemistry*, 14:4983–4988, 1975.
- [1419] J.M. Kemner, S. Liang, and E.W. Nester. The *Agrobacterium tumefaciens* virulence gene *chvE* is part of a putative ABC-type sugar transport operon. *J. Bacteriol.*, 179:2452–2458, 1997.
- [1420] M.A. Kerr. The second component of human complement. *Methods Enzymol.*, 80:54–64, 1980.
- [1421] M.A. Kerr. Human factor B. *Methods Enzymol.*, 80:102–112, 1981.
- [1422] P.J. Kersten, S. Dagley, J.W. Whittaker, D.M. Arciero, and J.D. Lipscomb. 2-Pyrone-4,6-dicarboxylic acid, a catabolite of gallic acids in *Pseudomonas* species. *J. Bacteriol.*, 152:1154–1162, 1982.
- [1423] J. Kervinen, P. Sarkkinen, N. Kalkkinen, L. Mikola, and M. Saarna. Hydrolytic specificity of the barley grain aspartic proteinase. *Phytochemistry*, 32:799–803, 1993.
- [1424] J. Kervinen, K. Törmäkangas, P. Runeberg-Roos, K. Guruprasad, T. Blundell, and T.H. Teeri. Structure and possible function of aspartic proteinases in barley and other plants. *Adv. Exp. Med. Biol.*, 362:241–254, 1995.
- [1425] M. Kessel, Maurizi, M.R., Kim, B., Kocsis, E., Trus, B., Singh, S.K. and Steven, A.C. Homology in structural organization between *E. coli* ClpAP protease and the eukaryotic 26 S proteasome. *J. Mol. Biol.*, 250:587–594, 1995.
- [1426] E. Kessler and R. Adar. Type I procollagen C-proteinase from mouse fibroblasts. Purification and demonstration of a 55-kDa enhancer glycoprotein. *Eur. J. Biochem.*, 186:115–121, 1989.
- [1427] E. Kessler and A. Yaron. A novel aminopeptidase from *Clostridium histolyticum*. *Biochem. Biophys. Res. Commun.*, 50:405–412, 1973.
- [1428] E. Kessler and A. Yaron. An extracellular aminopeptidase from *Clostridium histolyticum*. *Eur. J. Biochem.*, 63:271–287, 1976.
- [1429] E. Kessler and A. Yaron. Extracellular aminopeptidase from *Clostridium histolyticum*. *Methods Enzymol.*, 45:544–552, 1976.
- [1430] C. Kettner, E. Shaw, R. White, and A. Janoff. The specificity of macrophage elastase on the insulin B-chain. *Biochem. J.*, 195:369–372, 1981.

- [1431] S. Keuning, D.B. Janssen, and B. Witholt. Purification and characterization of hydrolytic haloalkane dehalogenase from *Xanthobacter autotrophicus* GJ10. *J. Bacteriol.*, 163:635–639, 1985.
- [1432] N.O. Keyhani and S. Roseman. The chitin catabolic cascade in the marine bacterium *Vibrio furnissii*. Molecular cloning, isolation, and characterization of a periplasmic chitodextrinase. *J. Biol. Chem.*, 271:33414–33424, 1996.
- [1433] N.A. Khan and N.R. Eaton. Purification and characterization of maltase and  $\alpha$ -methyl glucosidase from yeast. *Biochim. Biophys. Acta*, 146:173–180, 1967.
- [1434] O. Khersonsky and D.S. Tawfik. Structure-reactivity studies of serum paraoxonase PON1 suggest that its native activity is lactonase. *Biochemistry*, 44:6371–6382, 2005.
- [1435] G.H. Khorana. Phosphodiesterases. In P.D. Boyer, H. Lardy, and K. Myrback, editors, *The Enzymes*, volume 5, pages 79–94. Academic Press, New York, 2nd edition, 1961.
- [1436] H. Kido, N. Fukusen, and N. Katunuma. Chymotrypsin- and trypsin-type serine proteases in rat mast cells: properties and functions. *Arch. Biochem. Biophys.*, 239:436–443, 1985.
- [1437] W.W. Kielley. Myosin adenosine triphosphatase. In P.D. Boyer, H. Lardy, and K. Myrback, editors, *The Enzymes*, volume 5, pages 159–168. Academic Press, New York, 2nd edition, 1961.
- [1438] W.W. Kielley and L.B. Bradley. Glutathione thiolesterase. *J. Biol. Chem.*, 206:327–338, 1954.
- [1439] Y. Kikkawa, N. Yamanaka, J. Tada, N. Kanamori, K. Tsumura, and K. Hosoi. Prorenin processing and restricted endoproteolysis by mouse tissue kallikrein family enzymes (mK1, mK9, mK13, and mK22). *Biochim. Biophys. Acta*, 1382:55–64, 1998.
- [1440] M. Kikuchi, H. Hayashida, E. Nakano, and K. Sakaguchi. Peptidoglutaminase. Enzymes for selective deamidation of  $\gamma$ -amide of peptide-bound glutamine. *Biochemistry*, 10:1222–1229, 1971.
- [1441] R.D. Kilker, B. Saunier, J.S. Tkacz, and A. Herscovics. Partial purification from *Saccharomyces cerevisiae* of a soluble glucosidase which removes the terminal glucose from the oligosaccharide Glc<sub>3</sub>Man<sub>9</sub>GlcNAc<sub>2</sub>. *J. Biol. Chem.*, 256:5299–5603, 1981.
- [1442] B. Kim and J.W. Little. LexA and CI repressors as enzymes: specific cleavage in an intermolecular reaction. *Cell*, 73:1165–1173, 1993.
- [1443] D.-H. Kim and K.Y. Choi. Potyvirus NIa protease. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 721–723. Academic Press, London, 1998.
- [1444] D.-H. Kim, Y.-S. Yang, and W.B. Jakoby. Aspirin hydrolyzing esterases from rat liver cytosol. *Biochem. Pharmacol.*, 40:481–487, 1990.
- [1445] D.-H. Kim, Y.-S. Yang, and W.B. Jakoby. Nonserine esterases from rat liver cytosol. *Protein Expr. Purif.*, 1:19–27, 1990.
- [1446] D.H. Kim, D.C. Hwang, B.H. Kang, J. Lew, J.S. Han, B.O.D. Song, and K.Y. Choi. Effects of internal cleavages and mutations in the C-terminal region of NIa protease of turnip mosaic potyvirus on the catalytic activity. *Virology*, 226:183–190, 1996.
- [1447] E.J. Kim, D.O. Kang, D.C. Love, and J.A. Hanover. Enzymatic characterization of O-GlcNAcase isoforms using a fluorogenic GlcNAc substrate. *Carbohydr. Res.*, 341:971–982, 2006.
- [1448] G.A. Kim, M.S. Lee, Y. Sun, B.D. Lee, J.I. Lee, J.H. Lee, and S.T. Kwon. Characterization of cold-active uracil-DNA glycosylase from *Bacillus* sp. HJ171 and its use for contamination control in PCR. *Appl. Microbiol. Biotechnol.*, 80:785–794, 2008.
- [1449] J.K. Kim, I.S. Yang, S. Rhee, Z. Dauter, Y.S. Lee, S.S. Park, and K.H. Kim. Crystal structures of glutaryl 7-aminocephalosporanic acid acylase: insight into autoproteolytic activation. *Biochemistry*, 42:4084–4093, 2003.
- [1450] J.L. Kim, K.A. Morgenstern, C. Lin, T. Fox, M.D. Dwyer, J.A. Landro, S.P. Chambers, W. Markland, C.A. Lepre, E.T. O'Malley, S.L. Harbeson, C.M. Rice, M.A. Murcko, P.R. Caron, and J.A. Thomson. Crystal structure of the hepatitis C virus NS3 protease domain complexed with a synthetic NS4A cofactor peptide. *Cell*, 87:343–355, 1996.

- [1451] J.M. Kim, S. Shimizu, and H. Yamada. Amidohydrolysis of *N*-methylhydantoin coupled with ATP hydrolysis. *Biochem. Biophys. Res. Commun.*, 142:1006–1012, 1987.
- [1452] K.S. Kim, J.G. Pelton, W.B. Inwood, U. Andersen, S. Kustu, and D.E. Wemmer. The Rut pathway for pyrimidine degradation: novel chemistry and toxicity problems. *J. Bacteriol.*, 192:4089–4102, 2010.
- [1453] M.H. Kim, W.C. Choi, H.O. Kang, J.S. Lee, B.S. Kang, K.J. Kim, Z.S. Derewenda, T.K. Oh, C.H. Lee, and J.K. Lee. The molecular structure and catalytic mechanism of a quorum-quenching *N*-acyl-L-homoserine lactone hydrolase. *Proc. Natl. Acad. Sci. USA*, 102:17606–17611, 2005.
- [1454] R.Q. Kim, W.A. Offen, G.J. Davies, and K.A. Stubbs. Structural enzymology of *Helicobacter pylori* methylthioadenosine nucleosidase in the futasoline pathway. *Acta Crystallogr. D Biol. Crystallogr.*, 70:177–185, 2014.
- [1455] W.S. Kim, K. Hatsuzawa, Y. Ishizuka, K. Hashiba, K. Murakami, and K. Nakayama. A processing enzyme for prorenin in mouse submandibular gland. Purification and characterization. *J. Biol. Chem.*, 265:5930–5933, 1990.
- [1456] Y. Kim, K.-H. Yoon, Y. Khang, S. Turley, and W.G.J. Hol. The 2.0 Å crystal structure of cephalosporin acylase. *Structure*, 8:1059–1068, 2000.
- [1457] Y.K. Kim, M. Kitaoka, K. Hayashi, C.H. Kim, and G.L. Cote. Purification and characterization of an intracellular cycloalternan-degrading enzyme from *Bacillus* sp. NRRL B-21195. *Carbohydr. Res.*, 339:1179–1184, 2004.
- [1458] Y.S. Kim and E.J. Brophy. Rat intestinal brush border membrane peptidases. I. Solubilization, purification of two different forms of enzyme. *J. Biol. Chem.*, 251:3199–3205, 1976.
- [1459] S.H. Kindler and C. Gilvarg. *N*-Succinyl-L-2,6-diaminopimelic acid deacylase. *J. Biol. Chem.*, 235:3532–3535, 1960.
- [1460] S. Kinoshita, S. Negoro, M. Muramatsu, V.S. Bisaria, S. Sawada, and H. Okada. 6-Aminohexanoic acid cyclic dimer hydrolase. A new cyclic amide hydrolase produced by *Achromobacter guttatus* KI74. *Eur. J. Biochem.*, 80:489–495, 1977.
- [1461] S. Kinoshita, T. Terada, T. Taniguchi, Y. Takeney, S. Masuda, N. Matsunaga, and H. Okada. Purification and characterization of 6-aminohexanoic-acid-oligomer hydrolase of *Flavobacterium* sp. KI72. *Eur. J. Biochem.*, 116:547–551, 1981.
- [1462] T. Kinoshita, T. Tada, Y. Saito, Y. Ishii, A. Sato, and M. Murata. Crystallization and preliminary X-ray analysis of cephalosporin C acylase from *Pseudomonas* sp. strain N176. *Acta Crystallogr. D Biol. Crystallogr.*, 56:458–459, 2000.
- [1463] H. Kirschke, B. Wiederanders, D. Brömme, and A. Rinne. Cathepsin S from bovine spleen. Purification, distribution, intracellular localization and action on proteins. *Biochem. J.*, 264:467–473, 1989.
- [1464] H. Kirschke, P. Wikstrom, and E. Shaw. Active center differences between cathepsins L and B: the S1 binding region. *FEBS Lett.*, 228:128–130, 1988.
- [1465] L. Kirszbaum, C. Sotiropoulos, C. Jackson, S. Cleal, N. Slakeski, and E.C. Reynolds. Complete nucleotide sequence of a gene prtR of *Porphyromonas gingivalis* W50 encoding a 132 kDa protein that contains an arginine-specific thiol endopeptidase domain and a haemagglutinin domain. *Biochem. Biophys. Res. Commun.*, 207:424–431, 1995.
- [1466] T. Kishi, S.M. Cloutier, C. Kündig, D. Deperthes, and E.P. Diamandis. Activation and enzymatic characterization of recombinant human kallikrein 8. *Biol. Chem.*, 387:723–731, 2006.
- [1467] M. Kishida, T. Nikai, N. Mori, S. Kohmura, and H. Sugihara. Characterization of mucrotoxin A from the venom of *Trimeresurus mucrosquamatus* (the Chinese habu snake). *Toxicon*, 23:637–645, 1985.
- [1468] W. Kisiel and W. Canfield. M. Snake venom proteases that activate blood-coagulation factor V. *Methods Enzymol.*, 80:275–285, 1981.
- [1469] L.L. Kisselev and L.Yu. Frolova. Termination of translation in eukaryotes. *Biochem. Cell Biol.*, 73:1079–1086, 1995.
- [1470] K. Kitakima, S. Inoue, Y. Inoue, and F.A. Troy. Use of a bacteriophage-derived endo-*N*-acetylneuraminidase and an equine antipolysialyl antibody to characterize the polysialyl residues in salmonid fish egg polysialoglycoproteins. Substrate and immunospecificity studies. *J. Biol. Chem.*, 263:18269–18276, 1988.

- [1471] M. Klages, S. Krauss, and F. Lingens. 2-Haloacid dehalogenase from a 4-chlorobenzoate-degrading *Pseudomonas* spec. CBS 3. *Hoppe-Seyler's Z. Physiol. Chem.*, 364:529–535, 1983.
- [1472] S.M. Klaus, A. Wegkamp, W. Sybesma, J. Hugenholtz, J.F. Gregory, Hanson 3rd, and A.D. A nudix enzyme removes pyrophosphate from dihydroneopterin triphosphate in the folate synthesis pathway of bacteria and plants. *J. Biol. Chem.*, 280:5274–5280, 2005.
- [1473] A.O. Klein and W. Vishniac. Activity and partial purification of chlorophyllase in aqueous systems. *J. Biol. Chem.*, 236:2544–2547, 1961.
- [1474] R. Kleine. Properties of thermitase, a thermostable serine protease from *Thermoactinomyces vulgaris*. *Acta Biol. Med. Ger.*, 41:89–102, 1982.
- [1475] O.A. Klimova, S.I. Borukhov, N.I. Solovyeva, T.O. Balaevskaya, and A.Y. Strongin. The isolation and properties of collagenolytic proteases from crab hepatopancreas. *Biochem. Biophys. Res. Commun.*, 166:1411–1420, 1990.
- [1476] J.J. Kling, R.L. Wright, J.S. Moncrief, and T.D. Wilkins. Cloning and characterization of the gene for the metalloprotease enterotoxin of *Bacteroides fragilis*. *FEMS Microbiol. Lett.*, 146:279–284, 1997.
- [1477] C. Klinger, M. Roszbach, R. Howe, and M. Kaufmann. Thermophile-specific proteins: the gene product of aq\_1292 from *Aquifex aeolicus* is an NTPase. *BMC Biochem.*, 4:12–12, 2003.
- [1478] S. Klumpp, J. Hermesmeier, D. Selke, R. Baumeister, R. Kellner, and J. Krieglstein. Protein histidine phosphatase: a novel enzyme with potency for neuronal signaling. *J Cereb Blood Flow Metab*, 22:1420–1424, 2002.
- [1479] J. Knappe, W. Brümer, and K. Biederbick. Reinigung und Eigenschaften der Biotinidase aus Schweinenieren und *Lactobacillus casei*. *Biochem. Z.*, 338:599–613, 1963.
- [1480] V. Knäuper, S. Krämer, H. Reinke, and H. Tschesche. Characterization and activation of procollagenase from human polymorphonuclear leucocytes. N-terminal sequence and determination of the proenzyme and various proteolytically activated forms. *Eur. J. Biochem.*, 189:295–300, 1990.
- [1481] T. Knaus, M. Schober, B. Kepplinger, M. Faccinelli, J. Pitzer, K. Faber, P. Macheroux, and U. Wagner. Structure and mechanism of an inverting alkylsulfatase from *Pseudomonas* sp. DSM6611 specific for secondary alkyl sulfates. *FEBS J.*, 279:4374–4384, 2012.
- [1482] B. Kneidinger, C. Marolda, M. Graninger, A. Zamyatina, F. McArthur, P. Kosma, M.A. Valvano, and P. Messner. Biosynthesis pathway of ADP-L-glycero- $\beta$ -D-manno-heptose in *Escherichia coli*. *J. Bacteriol.*, 184:363–369, 2002.
- [1483] W.E. Knox. The relation of liver kynureninase to tryptophan metabolism in pyridoxine deficiency. *Biochem. J.*, 53:379–385, 1953.
- [1484] K. Kobashi, S. Takebe, and T. Sakai. Urethane-hydrolyzing enzyme from *Citrobacter* sp. *Chem. Pharm. Bull.*, 38:1326–1328, 1990.
- [1485] H. Kobayashi, I. Kusakabe, and K. Murakami. Substrate specificity of a carboxyl proteinase from *Irpex lacteus*. *Agric. Biol. Chem.*, 47:1921–1923, 1983.
- [1486] H. Kobayashi, I. Kusakabe, and K. Murakami. Purification and characterization of a pepstatin-insensitive carboxyl proteinase from *Polyporus tulipiferae* (*Irpex lacteus*). *Agric. Biol. Chem.*, 49:2393–2397, 1985.
- [1487] H. Kobayashi, S. Sekibata, H. Shibuya, S. Yoshida, I. Kusakabe, and K. Murakami. Cloning and sequence analysis of cDNA for *Irpex lacteus* aspartic proteinase. *Agric. Biol. Chem.*, 53:1927–1933, 1989.
- [1488] H. Kobayashi, E. Takahashi, K. Oguma, Y. Fujii, H. Yamanaka, T. Negishi, S. Arimoto-Kobayashi, T. Tsuji, and K. Okamoto. Cleavage specificity of the serine protease of *Aeromonas sobria*, a member of the kexin family of subtilases. *FEMS Microbiol. Lett.*, 256:165–170, 2006.
- [1489] H. Kobayashi, H. Utsunomiya, H. Yamanaka, Y. Sei, N. Katunuma, K. Okamoto, and H. Tsuge. Structural basis for the kexin-like serine protease from *Aeromonas sobria* as sepsis-causing factor. *J. Biol. Chem.*, 284:27655–27663, 2009.
- [1490] K. Kobayashi and J.A. Smith. Acyl-peptide hydrolase from rat liver. Characterization of enzyme reaction. *J. Biol. Chem.*, 262:11435–11437, 1987.



- [1491] M. Kobayashi, N. Yanaka, T. Nagasawa, , and H. Primary structure of an aliphatic nitrile-degrading enzyme, aliphatic nitrilase from *Rhodococcus rhodochromus* K22 and expression of its gene and identification of its active site residue. *Biochemistry*, 31:9000–9007, 1992.
- [1492] A.L. Koch. Some enzymes of nucleoside metabolism of *Escherichia coli*. *J. Biol. Chem.*, 223:535–549, 1956.
- [1493] G.B. Koelle. Cholinesterases of the tissues and sera of rabbits. *Biochem. J.*, 53:217–226, 1953.
- [1494] L.V. Kofod, S. Kauppinen, S. Christgau, L.N. Andersen, H.P. Heldt-Hansen, K. Dorreich, and H. Dalboge. Cloning and characterization of two structurally and functionally divergent rhamnogalacturonases from *Aspergillus aculeatus*. *J. Biol. Chem.*, 269:29182–29189, 1994.
- [1495] R. Köhler, A. Brokamp, R. Schwarze, R.H. Reiting, and F.R.J. Schmidt. Characteristics and DNA-sequence of a cryptic haloalkanoic acid dehalogenase from *Agrobacterium tumefaciens* RS5. *Curr. Microbiol.*, 36:96–101, 1998.
- [1496] L.D. Kohn, C. Iserky, J. Zupnik, A. Lenaers, G. Lee, and C.M. Lapière. Calf tendon procollagen peptidase: its purification and endopeptidase mode of action. *Proc. Natl. Acad. Sci. USA*, 71:40–44, 1974.
- [1497] M. Kohno and I. Yamashina. Purification and properties of 4-L-aspartylglycosylamine amidohydrolase from hog kidney. *Biochim. Biophys. Acta*, 258:600–617, 1972.
- [1498] N. Koide and T. Muramatsu. Endo- $\beta$ -*N*-acetylglucosaminidase acting on carbohydrate moieties of glycoproteins. Purification and properties of the enzyme from *Diplococcus pneumoniae*. *J. Biol. Chem.*, 249:4897–4904, 1974.
- [1499] J. Kolinska and J. Kraml. Separation and characterization of sucrose-isomaltase and of glucoamylase of rat intestine. *Biochim. Biophys. Acta*, 284:235–247, 1972.
- [1500] I.K. Kolos and A.F. Makarchikov. [Identification of thiamine monophosphate hydrolyzing enzymes in chicken liver]. *Ukr Biochem J*, 86:39–49, 2014.
- [1501] H. Komatsuzawa, M. Sugai, K. Ohta, T. Fujiwara, S. Nakashima, J. Suzuki, C.Y. Lee, and H. Suginaka. Cloning and characterization of the *fmt* gene which affects the methicillin resistance level and autolysis in the presence of triton X-100 in methicillin-resistant *Staphylococcus aureus*. *Antimicrob. Agents Chemother.*, 41:2355–2361, 1997.
- [1502] H. Komeda, H. Harada, S. Washika, T. Sakamoto, M. Ueda, and Y. Asano. A novel *R*-stereoselective amidase from *Pseudomonas* sp. MCI3434 acting on piperazine-2-*tert*-butylcarboxamide. *Eur. J. Biochem.*, 271:1580–1590, 2004.
- [1503] H. Komeda, H. Harada, S. Washika, T. Sakamoto, M. Ueda, and Y. Asano. *S*-stereoselective piperazine-2-*tert*-butylcarboxamide hydrolase from *Pseudomonas* azotoformans IAM 1603 is a novel L-amino acid amidase. *Eur. J. Biochem.*, 271:1465–1475, 2004.
- [1504] Y. Komeda, M. Tanaka, and T. Nishimune. A *th-1* mutant of *Arabidopsis thaliana* is defective for a thiamin-phosphate-synthesizing enzyme: thiamin phosphate pyrophosphorylase. *Plant Physiol.*, 88:248–250, 1988.
- [1505] E. Kominami, H. Hoffschulte, L. Leuschel, K. Maier, and H. Holzer. The substrate specificity of proteinase B from baker's yeast. *Biochim. Biophys. Acta*, 661:136–141, 1981.
- [1506] T. Komiyama, T.L. Bigler, N. Yoshida, K., Laskowski Noda, , and Jr. Replacement of P1 Leu<sup>18</sup> by Glu<sup>18</sup> in the reactive site of turkey ovomucoid third domain converts it into a strong inhibitor of Glu-specific Streptomyces griseus Proteinase (GluSGP). *J. Biol. Chem.*, 266:10727–10730, 1991.
- [1507] P. Konczalik and J. Moss. Identification of critical, conserved vicinal aspartate residues in mammalian and bacterial ADP-ribosylarginine hydrolases. *J. Biol. Chem.*, 274:16736–16740, 1999.
- [1508] H. König and B. Rosenwirth. Purification and partial characterization of poliovirus protease 2A by means of a functional assay. *J. Virol.*, 62:1243–1250, 1988.
- [1509] E.V. Koonin, K.S. Makarova, I.B. Rogozin, L. Davidovic, M.C. Letellier, and L. Pellegrini. The rhomboids: a nearly ubiquitous family of intramembrane serine proteases that probably evolved by multiple ancient horizontal gene transfers. *Genome Biol.*, 4:R19–R19, 2003.

- [1510] M. Korczynska, D.F. Xiang, Z. Zhang, C. Xu, T. Narindoshvili, S.S. Kamat, H.J. Williams, S.S. Chang, P. Kolb, B. Hillerich, J.M. Sauder, S.K. Burley, S.C. Almo, S. Swaminathan, B.K. Shoichet, and F.M. Raushel. Functional annotation and structural characterization of a novel lactonase hydrolyzing D-xylono-1,4-lactone-5-phosphate and L-arabino-1,4-lactone-5-phosphate. *Biochemistry*, 53:4727–4738, 2014.
- [1511] E.D. Korn and T.W. Quigley. Lipoprotein lipase of chicken adipose tissue. *J. Biol. Chem.*, 226:833–839, 1957.
- [1512] A. Kornberg and W.E. Pricer. Nucleotide pyrophosphatase. *J. Biol. Chem.*, 182:763–778, 1950.
- [1513] S.R. Kornberg. Tripolyphosphate and trimetaphosphate in yeast extracts. *J. Biol. Chem.*, 218:23–31, 1956.
- [1514] S.R. Kornberg, I.R. Lehman, M.J. Bessman, E.S. Simms, and A. Kornberg. Enzymatic cleavage of deoxyguanosine triphosphate to deoxyguanosine and tripolyphosphate. *J. Biol. Chem.*, 233:159–162, 1958.
- [1515] S.J. Kornfeld and A.G. Plaut. Secretory immunity and the bacterial IgA proteases. *Rev. Infect. Dis.*, 3:521–534, 1981.
- [1516] S.S. Koshy, M.H. Montrose, and C.L. Sears. Human intestinal epithelial cells swell and demonstrate actin rearrangement in response to the metalloprotease toxin of *Bacteroides fragilis*. *Infect. Immun.*, 64:5022–5028, 1996.
- [1517] T. Kosuge and E.E. Conn. The metabolism of aromatic compounds in higher plants. V. Purification and properties of dihydrocoumarin hydrolase of *Melilotus alba*. *J. Biol. Chem.*, 237:1653–1656, 1962.
- [1518] T. Kotake, N. Hirata, Y. Degi, M. Ishiguro, K. Kitazawa, R. Takata, H. Ichinose, S. Kaneko, K. Igarashi, M. Samejima, and Y. Tsumuraya. Endo- $\beta$ -1,3-galactanase from winter mushroom *Flammulina velutipes*. *J. Biol. Chem.*, 286:27848–27854, 2011.
- [1519] T. Kotake, S. Kaneko, A. Kubomoto, M.A. Haque, H. Kobayashi, and Y. Tsumuraya. Molecular cloning and expression in *Escherichia coli* of a *Trichoderma viride* endo- $\beta$ -(1 $\rightarrow$ 6)-galactanase gene. *Biochem. J.*, 377:749–755, 2004.
- [1520] K. Koudounas, G. Banilas, C. Michaelidis, C. Demoliou, S. Rigas, and P. Hatzopoulos. A defence-related *Olea europaea*  $\beta$ -glucosidase hydrolyses and activates oleuropein into a potent protein cross-linking agent. *J. Exp. Bot.*, 66:2093–2106, 2015.
- [1521] K. Koudounas, M. Thomopoulou, C. Michaelidis, E. Zevgiti, G. Papakostas, P. Tserou, G. Daras, and P. Hatzopoulos. The C-domain of oleuropein  $\beta$ -glucosidase assists in protein folding and sequesters the enzyme in nucleus. *Plant Physiol.*, 174:1371–1383, 2017.
- [1522] M.Z. Kounnas, R.L. Wolz, C.M. Gorbea, and J.S. Bond. Meprin-A and -B. Cell surface endopeptidases of the mouse kidney. *J. Biol. Chem.*, 266:17350–17357, 1991.
- [1523] D. Koutsoulis, D. Landry, and E.P. Guthrie. Novel endo- $\alpha$ -N-acetylgalactosaminidases with broader substrate specificity. *Glycobiology*, 18:799–805, 2008.
- [1524] R.J. Kovachy, S.D. Copley, and R.H. Allen. Recognition, isolation, and characterization of rat liver D-methylmalonyl coenzyme A hydrolase. *J. Biol. Chem.*, 258:11415–11421, 1983.
- [1525] G.G. Kovaleva, M.P. Shimanskaya, and V.M. Stepanov. The site of diazoacetyl inhibitor attachment to acid proteinase of *Aspergillus awamori* - an analog of penicillopepsin and pepsin. *Biochem. Biophys. Res. Commun.*, 49:1075–1082, 1972.
- [1526] I.M. Krab and A. Parmeggiani. EF-Tu, a GTPase odyssey. *Biochim. Biophys. Acta*, 1443:1–22, 1998.
- [1527] G.R. Krakower and K.-H. Kim. Purification and properties of acetyl-CoA carboxylase phosphatase. *J. Biol. Chem.*, 256:2408–2413, 1980.
- [1528] R.A. Kramer, L. Vandeputte-Rutten, G.J. de Roon, P. Gros, N. Dekker, and M.R. Egmond. Identification of essential acidic residues of outer membrane protease OmpT supports a novel active site. *FEBS Lett.*, 505:426–430, 2001.
- [1529] E. Krauhs, H. Dörsam, M. Little, R. Zwilling, and H. Ponstingl. A protease from *Astacus fluviatilis* as an aid in protein sequencing. *Anal. Biochem.*, 119:153–157, 1982.
- [1530] E. Kraus, H.H. Klitz, and U.F. Fembert. The specificity of proteinase K against oxidized insulin B chain. *Hoppe-Seyler's Z. Physiol. Chem.*, 357:233–237, 1976.

- [1531] H.-G. Kräusslich and E. Wimmer. Viral proteinases. *Annu. Rev. Biochem.*, 57:701–754, 1988.
- [1532] J. Kream and E. Chargaff. On the cytosine deaminase of yeast. *J. Am. Chem. Soc.*, 74:5157–5160, 1952.
- [1533] J.F. Krebs, A. Srinivasan, A.M. Wong, K.J. Tomaselli, L.C. Fritz, and J.C. Wu. Heavy membrane-associated caspase 3: identification, isolation, and characterization. *Biochemistry*, 39:16056–16063, 2000.
- [1534] H. Kresse, W. Fuchs, J. Glössl, D. Holtfrerich, and W. Gilberg. Liberation of *N*-acetylglucosamine-6-sulfate by human  $\beta$ -*N*-acetylhexosaminidase A. *J. Biol. Chem.*, 256:12926–12932, 1981.
- [1535] D.E. Krieger, R. Levine, R.B. Merrifield, G. Vidali, and V.G. Allfrey. Chemical studies of histone acetylation. Substrate specificity of a histone deacetylase from calf thymus nuclei. *J. Biol. Chem.*, 249:332–334, 1974.
- [1536] P.S. Krishnan. Apyrase, pyrophosphatase and metaphosphatase of *Penicillium chrysogenum*. *Arch. Biochem. Biophys.*, 37:224–234, 1952.
- [1537] T. Krojer, M. Garrido-Franco, R. Huber, M. Ehrmann, and T. Clausen. Crystal structure of DegP (HtrA) reveals a new protease-chaperone machine. *Nature*, 416:455–459, 2002.
- [1538] P.A. Kroon, C.B. Faulds, and G. Williamson. Purification and characterisation of a novel ferulic acid esterase induced by growth of *Aspergillus niger* on sugarbeet pulp. *Biotechnol. Appl. Biochem.*, 23:255–262, 1996.
- [1539] H. Kropp, J.G. Sundelof, R. Hajdu, and F.M. Kahan. Metabolism of thienamycin and related carbapenem antibiotics by renal dipeptidase, dehydropeptidase-I. *Antimicrob. Agents Chemother.*, 22:62–70, 1982.
- [1540] M.S. Krug and S.L. Berger. Ribonuclease H activities associated with viral reverse transcriptases are endonucleases. *Proc. Natl. Acad. Sci. USA*, 86:3539–3543, 1989.
- [1541] G. Kuan, E. Dassa, N. Saurin, M., Saier Hofnung, , and Jr. Phylogenetic analyses of the ATP-binding constituents of bacterial extracytoplasmic receptor-dependent ABC-type nutrient uptake permeases. *Res. Microbiol.*, 146:271–278, 1995.
- [1542] S.A. Kuby and H.A. Lardy. Purification and kinetics of  $\beta$ -D-galactosidase from *Escherichia coli*, strain K-12. *J. Am. Chem. Soc.*, 75:890–896, 1953.
- [1543] F. Kudo, T. Fujii, S. Kinoshita, and T. Eguchi. Unique *O*-ribosylation in the biosynthesis of butirosin. *Bioorg. Med. Chem.*, 15:4360–4368, 2007.
- [1544] M. Kukkonen, K. Lähteenmäki, M. Suomalainen, N. Kalkkinen, L. Emödy, H. Laang, and T.K. Korhonen. Protein regions important for plasminogen activation and inactivation of  $\alpha$ 2-antiplasmin in the surface protease Pla of *Yersinia pestis*. *Mol. Microbiol.*, 40:1097–1111, 2001.
- [1545] I.S. Kulaev, G.I. Konoshenko, and A.M. Umnov. Localization of polyphosphatase hydrolyzing polyphosphates to orthophosphate in subcellular structures of *Neurospora crassa*. *Biochemistry (Moscow)*, 37:190–194, 1972.
- [1546] A.N. Kulakova, G.B. Wisdom, L.A. Kulakov, and J.P. Quinn. The purification and characterization of phosphonopyruvate hydrolase, a novel carbon-phosphorus bond cleavage enzyme from *Variovorax* sp. Pal2. *J. Biol. Chem.*, 278:23426–23431, 2003.
- [1547] T. Kumagai, Y. Koyama, K. Oda, M. Noda, Y. Matoba, and M. Sugiyama. Molecular cloning and heterologous expression of a biosynthetic gene cluster for the antitubercular agent D-cycloserine produced by *Streptomyces lavendulae*. *Antimicrob. Agents Chemother.*, 54:1132–1139, 2010.
- [1548] T. Kumagai, K. Takagi, Y. Koyama, Y. Matoba, K. Oda, M. Noda, and M. Sugiyama. Heme protein and hydroxyarginase necessary for biosynthesis of D-cycloserine. *Antimicrob. Agents Chemother.*, 56:3682–3689, 2012.
- [1549] N. Kumar and M. Flavin. Preferential action of a brain detyrosinating carboxypeptidase on polymerized tubulin. *J. Biol. Chem.*, 256:7678–7686, 1981.
- [1550] S. Kumar, M. Kinoshita, M. Noda, N.G. Copeland, and N.A. Jenkins. Induction of apoptosis by the mouse Nedd2 gene, which encodes a protein similar to the product of the *Caenorhabditis elegans* cell death gene *ced-3* and the mammalian IL-1 $\beta$ -converting enzyme. *Genes Dev.*, 8:1613–1626, 1994.

- [1551] S.A. Kumar, N.A. Rao, and C.S. Vaidyanathan. Nucleotidases in plants. I. Partial purification and properties of the enzyme hydrolyzing flavine adenine dinucleotide from mung bean seedlings (*Phaseolus radiatus*). *Arch. Biochem. Biophys.*, 111:646–652, 1965.
- [1552] A. Kumon, Y. Matsuoka, Y. Kakimoto, T. Nakajima, and I. Sano. A peptidase that hydrolyzes  $\alpha$ -N-( $\gamma$ -aminobutyryl)lysine. *Biochim. Biophys. Acta*, 200:466–474, 1970.
- [1553] H.-F. Kung and C. Wagner.  $\gamma$ -Glutamylmethylamide. A new intermediate in the metabolism of methylamine. *J. Biol. Chem.*, 244:4136–4140, 1969.
- [1554] A. Kuninaka. 5'-Inosinic acid-N-ribosidase in *Aspergillus*. *Koso Kagaka Shinojiumu*, 12:65–69, 1957.
- [1555] M. Kunitz. Isolation of crystalline deoxyribonuclease from beef pancreas. *Science*, 108:19–20, 1948.
- [1556] M. Kunitz. Crystalline inorganic pyrophosphatase isolated from baker's yeast. *J. Gen. Physiol.*, 35:423–450, 1952.
- [1557] K. Kuntze, Y. Shinoda, H. Moutakki, M.J. McInerney, C. Vogt, H.H. Richnow, and M. Boll. 6-Oxocyclohex-1-ene-1-carbonyl-coenzyme A hydrolases from obligately anaerobic bacteria: characterization and identification of its gene as a functional marker for aromatic compounds degrading anaerobes. *Environ Microbiol*, 10:1547–1556, 2008.
- [1558] A. Kunzmann, A. Brennicke, and A. Marchfelder. 5' end maturation and RNA editing have to precede tRNA 3' processing in plant mitochondria. *Proc. Natl. Acad. Sci. USA*, 95:108–113, 1998.
- [1559] C.H. Kuo and W.W. Wells.  $\beta$ -Galactosidase from rat mammary gland. Its purification, properties, and role in the biosynthesis of 6 $\beta$ -O-D-galactopyranosyl myo-inositol. *J. Biol. Chem.*, 253:3550–3556, 1978.
- [1560] L.C. Kuo and J.A. Shafer. (eds) Retroviral Proteases. *Methods Enzymol.*, 241:1–431, 1994.
- [1561] M.-H. Kuo and H.J. Blumenthal. Purification and properties of an acid phosphomonoesterase from *Neurospora crassa*. *Biochim. Biophys. Acta*, 52:13–29, 1961.
- [1562] W.-L. Kuo, B.D. Gehm, and M.R. Rosner. Cloning and expression of the cDNA for a *Drosophila* insulin-degrading enzyme. *Mol. Endocrinol.*, 4:1580–1591, 1990.
- [1563] K. Kurachi and E.W. Davie. Human factor XI (plasma thromboplastin antecedent). *Methods Enzymol.*, 80:211–220, 1981.
- [1564] M. Kuratani, R. Ishii, Y. Bessho, R. Fukunaga, T. Sengoku, M. Shirouzu, S. Sekine, and S. Yokoyama. Crystal structure of tRNA adenosine deaminase (TadA) from *Aquifex aeolicus*. *J. Biol. Chem.*, 280:16002–16008, 2005.
- [1565] T. Kurecki, M. Laskowski, Kress Sr., and L.F. Purification and some properties of two proteinases from *Crotalus adamanteus* venom that inactivate human  $\alpha_1$ -proteinase inhibitor. *J. Biol. Chem.*, 253:8340–8345, 1978.
- [1566] S. Kurihara, S. Oda, K. Kato, H.G. Kim, T. Koyanagi, H. Kumagai, and H. Suzuki. A novel putrescine utilization pathway involves  $\gamma$ -glutamylated intermediates of *Escherichia coli* K-12. *J. Biol. Chem.*, 280:4602–4608, 2005.
- [1567] T. Kurihara, N. Esaki, and K. Soda. Bacterial 2-haloacid dehalogenases: structures and reaction mechanisms. *J. Mol. Catal., B Enzym.*, 10:57–65, 2000.
- [1568] T. Kurihara, Y. Nishizawa, Y. Takahashi, and S. Odani. Chemical, immunological and catalytic properties of 2':3'-cyclic nucleotide 3'-phosphodiesterase purified from brain white matter. *Biochem. J.*, 195:153–157, 1981.
- [1569] G. Kurisu, Y. Kai, and S. Harada. Structure of the zinc-binding site in the crystal structure of a zinc endoprotease from *Streptomyces caespitosus* at 1 Å resolution. *J. Inorg. Biochem.*, 82:225–228, 2000.
- [1570] G. Kurisu, A. Sugimoto, S. Harada, M. Takagi, T. Imanaka, and Y. Kai. Characterization of a small metalloprotease from *Streptomyces caespitosus* with high specificity to aromatic residues. *J. Ferment. Bioeng.*, 83:590–592, 1997.
- [1571] H. Kuroda, A. Miyadera, A. Imura, and A. Suzuki. Partial purification, and some properties and reactivities of cetraxate benzyl ester hydrochloride-hydrolyzing enzyme. *Chem. Pharm. Bull.*, 37:2929–2932, 1989.
- [1572] G. Kuroki, P.A. Lizotte, and J.E. Poulton. L- $\beta$ -Glycosidases from *Prunus serotina* EHRH and *Davallia trichomanoides*. *Z. Naturforsch. C: Biosci.*, 39:232–239, 1984.

- [1573] K. Kuroki, F. Eng, T. Ishikawa, C. Turck, F. Harada, and D. Ganem. gp180, a host cell glycoprotein that binds duck hepatitis B virus particles, is encoded by a member of the carboxypeptidase gene family. *J. Biol. Chem.*, 270:15022–15028, 1995.
- [1574] Y. Kurono, M. Chidimatsu, K. Horikoshi, and Y. Ikeda. Isolation of a protease from a *Rhizopus* product. *Agric. Biol. Chem.*, 35:1668–1675, 1971.
- [1575] Y. Kurosawa, K. Ikeda, and F. Egami.  $\alpha$ -L-rhamnosidases of the liver of *Turbo cornutus* and *Aspergillus niger*. *J. Biochem.*, 73:31–37, 1973.
- [1576] T.V. Kurzchalia, U.A. Bommer, G.T. Babkina, and G.G. Karpova. GTP interacts with the  $\gamma$ -subunit of eukaryotic initiation factor EIF-2. *FEBS Lett.*, 175:313–316, 1984.
- [1577] K. Kusumoto and D.P. Hsieh. Purification and characterization of the esterases involved in aflatoxin biosynthesis in *Aspergillus parasiticus*. *Can. J. Microbiol.*, 42:804–810, 1996.
- [1578] S. Kuwabara. Purification and properties of two extracellular  $\beta$ -lactamases from *Bacillus cereus* 569-H. *Biochem. J.*, 118:457–465, 1970.
- [1579] E. Kuznetsova, M. Proudfoot, C.F. Gonzalez, G. Brown, M.V. Omelchenko, I. Borozan, L. Carmel, Y.I. Wolf, H. Mori, A.V. Savchenko, C.H. Arrowsmith, E.V. Koonin, A.M. Edwards, and A.F. Yakunin. Genome-wide analysis of substrate specificities of the *Escherichia coli* haloacid dehalogenase-like phosphatase family. *J. Biol. Chem.*, 281:36149–36161, 2006.
- [1580] B. Kwiatkowski, B. Boscheck, H. Thiele, and S. Stirm. Endo-N-acetylneuraminidase associated with bacteriophage particles. *J. Virol.*, 43:697–704, 1982.
- [1581] T.H. Kwon, S. Rhee, Y.S. Lee, S.S. Park, and K.H. Kim. Crystallization and preliminary X-Ray diffraction analysis of glutaryl-7-aminocephalosporanic acid acylase from *Pseudomonas* sp. GK16. *J. Struct. Biol.*, 131:79–81, 2000.
- [1582] J. Labadie and M.-C. Montel. Purification et étude de quelques propriétés d'une collagénase produite par *Empedobacter collagenolyticum*. *Biochimie*, 64:49–54, 1982.
- [1583] J.M. Labavitch, L.E. Freeman, and P. Albersheim. Structure of plant cell walls. Purification and characterization of a  $\beta$ -1,4-galactanase which degrades a structural component of the primary cell walls of dicots. *J. Biol. Chem.*, 251:5904–5910, 1976.
- [1584] C. Labbaye, P. Musette, and Y.E. Cayre. Wegener autoantigen and myeloblastin are encoded by a single mRNA. *Proc. Natl. Acad. Sci. USA*, 88:9253–9256, 1991.
- [1585] N. Lack, E.D. Lowe, J. Liu, L.D. Eltis, M.E. Noble, E. Sim, and I.M. Westwood. Structure of HsaD, a steroid-degrading hydrolase, from *Mycobacterium tuberculosis*. *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.*, 64:2–7, 2008.
- [1586] N.A. Lack, K.C. Yam, E.D. Lowe, G.P. Horsman, R.L. Owen, E. Sim, and L.D. Eltis. Characterization of a carbon-carbon hydrolase from *Mycobacterium tuberculosis* involved in cholesterol metabolism. *J. Biol. Chem.*, 285:434–443, 2010.
- [1587] G.M. Lacourciere and R.N. Armstrong. The catalytic mechanism of microsomal epoxide hydrolase involves an ester intermediate. *J. Am. Chem. Soc.*, 115:10466–10456, 1993.
- [1588] R.D. Ladner, D.E. McNulty, S.A. Carr, G.D. Roberts, and S.J. Caradonna. Characterization of distinct nuclear and mitochondrial forms of human deoxyuridine triphosphate nucleotidohydrolase. *J. Biol. Chem.*, 271:7745–7751, 1996.
- [1589] D. Laempe, M. Jahn, and G. Fuchs. 6-Hydroxycyclohex-1-ene-1-carbonyl-CoA dehydrogenase and 6-oxocyclohex-1-ene-1-carbonyl-CoA hydrolase, enzymes of the benzoyl-CoA pathway of anaerobic aromatic metabolism in the denitrifying bacterium *Thauera aromatica*. *Eur. J. Biochem.*, 263:420–429, 1999.
- [1590] W.A. Laing, S. Bulley, M. Wright, J. Cooney, D. Jensen, D. Barraclough, and E. MacRae. A highly specific L-galactose-1-phosphate phosphatase on the path to ascorbate biosynthesis. *Proc. Natl. Acad. Sci. USA*, 101:16976–16981, 2004.
- [1591] J. Laliberte and R.L. Momparler. Human cytidine deaminase: purification of enzyme, cloning, and expression of its complementary DNA. *Cancer Res.*, 54:5401–5407, 1994.

- [1592] W. W. Y Lam and T. D. H. Bugg. Chemistry of extradiol aromatic ring cleavage: isolation of a stable dienol ring fission intermediate and stereochemistry of its enzymatic hydrolytic cleavage. *J. Chem. Soc., Chem. Commun.*, 10:1163–1164, 1994.
- [1593] W.W. Lam and T.D. Bugg. Purification, characterization, and stereochemical analysis of a C-C hydrolase: 2-hydroxy-6-keto-nona-2,4-diene-1,9-dioic acid 5,6-hydrolase. *Biochemistry*, 36:12242–12251, 1997.
- [1594] D.W. Lambert, N.M. Hooper, and A.J. Turner. Angiotensin-converting enzyme 2 and new insights into the renin-angiotensin system. *Biochem. Pharmacol.*, 75:781–786, 2008.
- [1595] J.A. Lambrecht, J.M. Flynn, and D.M. Downs. Conserved YjgF protein family deaminates reactive enamine/imine intermediates of pyridoxal 5'-phosphate (PLP)-dependent enzyme reactions. *J. Biol. Chem.*, 287:3454–3461, 2012.
- [1596] O.E. Landman. Properties and induction of  $\beta$ -galactosidase in *Bacillus megaterium*. *Biochim. Biophys. Acta*, 23:558–569, 1957.
- [1597] M. Landt and L.G. Butler. 5'-Nucleotide phosphodiesterase: isolation of covalently bound 5'-adenosine monophosphate, an intermediate in the catalytic mechanism. *Biochemistry*, 17:4130–4135, 1978.
- [1598] C.F. LaPointe and R.K. Taylor. The type 4 prepilin peptidases comprise a novel family of aspartic acid proteases. *J. Biol. Chem.*, 275:1502–1510, 2000.
- [1599] C. Lapresle, V. Puizdar, C. Porchon-Bertolotto, E. Joukoff, and V. Turk. Structural differences between rabbit cathepsin E and cathepsin D. *Biol. Chem. Hoppe-Seyler*, 367:523–526, 1986.
- [1600] M. Larijani and A. Martin. Single-stranded DNA structure and positional context of the target cytidine determine the enzymatic efficiency of AID. *Mol. Cell Biol.*, 27:8038–8048, 2007.
- [1601] M. Larijani, A.P. Petrov, O. Kolenchenko, M. Berru, S.N. Krylov, and A. Martin. AID associates with single-stranded DNA with high affinity and a long complex half-life in a sequence-independent manner. *Mol. Cell Biol.*, 27:20–30, 2007.
- [1602] F.S. Larimore, L. Waxman, and A.L. Goldberg. Studies of the ATP-dependent proteolytic enzyme, protease La, from *Escherichia coli*. *J. Biol. Chem.*, 257:4187–4195, 1982.
- [1603] J. Larner. Other glucosidases. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 4, pages 369–378. Academic Press, New York, 2nd edition, 1960.
- [1604] J. Larsbrink, A. Izumi, F. Ibatullin, A. Nakhai, H.J. Gilbert, G.J. Davies, and H. Brumer. Structural and enzymatic characterisation of a glycoside hydrolase family 31  $\alpha$ -xylosidase from *Cellvibrio japonicus* involved in xyloglucan saccharification. *Biochem. J.*, 436:567–580, 2011.
- [1605] N.A. Larsen, J.M. Turner, J. Stevens, S.J. Rosser, A. Basran, R.A. Lerner, N.C. Bruce, and I.A. Wilson. Crystal structure of a bacterial cocaine esterase. *Nat. Struct. Biol.*, 9:17–21, 2002.
- [1606] T.J. Larson, M. Ehrmann, and W. Boos. Periplasmic glycerophosphodiester phosphodiesterase of *Escherichia coli*, a new enzyme of the glp regulon. *J. Biol. Chem.*, 258:5428–5432, 1983.
- [1607] M. Laskowski. , Sr. Pancreatic deoxyribonuclease I. In G.L. Cantoni and D.R. Davies, editors, *Procedures in Nucleic Acid Research*, pages 85–101. Procedures in Nucleic Acid Research, New York, 1966.
- [1608] M. Laskowski. , Sr. Venom exonuclease. In P.D. Boyer, editor, *The Enzymes*, volume 4, pages 313–328. Academic Press, New York, 3rd edition, 1971.
- [1609] R.A.L. Lassy and C.G. Miller. Peptidase E, a peptidase specific for N-terminal aspartic dipeptides, is a serine hydrolase. *J. Bacteriol.*, 182:2536–2543, 2000.
- [1610] S. Latt. A., Holmquist, B. and Vallee, B. L. Thermolysin: a zinc metalloenzyme. *Biochem. Biophys. Res. Commun.*, 37:333–339, 1969.
- [1611] J.B. Lawrence, C. Oxvig, M.T. Overgaard, L. Sottrup-Jensen, , and G.J. , Hays L.G., Yates, J.R. 3rd, and Conover, C.A. The insulin-like growth factor (IGF)-dependent IGF binding protein-4 protease secreted by human fibroblasts is pregnancy-associated plasma protein-A. *Proc. Natl. Acad. Sci. USA*, 96:3149–3153, 1999.

- [1612] T.F. Leal and I. de Sa-Nogueira. Purification, characterization and functional analysis of an endo-arabinanase (AbnA) from *Bacillus subtilis*. *FEMS Microbiol. Lett.*, 241:41–48, 2004.
- [1613] A. Lecroisey, C. Boulard, and B. Keil. Chemical and enzymatic characterization of the collagenase from the insect *Hypoderma lineatum*. *Eur. J. Biochem.*, 101:385–393, 1979.
- [1614] A. Lecroisey, A.-M. Gilles, A. De Wolf, and B. Keil. Complete amino acid sequence of the collagenase from the insect *Hypoderma lineatum*. *J. Biol. Chem.*, 262:7546–7551, 1987.
- [1615] A. Lecroisey and B. Keil. Specificity of the collagenase from the insect *Hypoderma lineatum*. *Eur. J. Biochem.*, 152:123–130, 1985.
- [1616] I.G. Leder. A novel 3-O sulfatase from human urine acting on methyl-2-deoxy-2-sulfamino- $\alpha$ -D-glucopyranoside 3-sulfate. *Biochem. Biophys. Res. Commun.*, 94:1183–1189, 1980.
- [1617] C. Lee and Y.S. Seo. Isolation and characterization of a processive DNA helicase from the fission yeast *Schizosaccharomyces pombe* that translocates in a 5'-to-3' direction. *Biochem. J.*, 334:377–386, 1998.
- [1618] C.G. Lee and J. Hurwitz. A new RNA helicase isolated from HeLa cells that catalytically translocates in the 3' to 5' direction. *J. Biol. Chem.*, 267:4398–4407, 1992.
- [1619] D. Lee and A.P. Ryle. Pepsin D. A minor component of commercial pepsin preparations. *Biochem. J.*, 104:742–748, 1967.
- [1620] D. Lee and A.P. Ryle. Pepsinogen D. A fourth proteolytic zymogen from pig gastric mucosa. *Biochem. J.*, 104:735–741, 1967.
- [1621] E.Y.C. Lee, J.H. Carter, L.D. Nielsen, and E.H. Fischer. Purification and properties of yeast amylo-1,6-glucosidase-oligo-1,4 leads to 1,4-glucantransferase. *Biochemistry*, 9:2347–2355, 1970.
- [1622] E.Y.C. Lee and W.J. Whelan. Glycogen and starch debranching enzymes. In P.D. Boyer, editor, *The Enzymes*, volume 5, pages 191–234. Academic Press, New York, 3rd edition, 1972.
- [1623] F.T. Lee, J.B. Adams, A.J. Garton, and S.J. Yeaman. Hormone-sensitive lipase is involved in the hydrolysis of lipoidal derivatives of estrogens and other steroid hormones. *Biochim. Biophys. Acta*, 963:258–264, 1988.
- [1624] K.H. Lee, H.L. Piao, H.Y. Kim, S.M. Choi, F. Jiang, W. Hartung, I. Hwang, J.M. Kwak, I.J. Lee, and I. Hwang. Activation of glucosidase via stress-induced polymerization rapidly increases active pools of abscisic acid. *Cell*, 126:1109–1120, 2006.
- [1625] K.S. Lee, S. Yasumasu, K. Nomura, and I. Iuchi. HCE, a constituent of the hatching enzymes of *Oryzias latipes* embryos, releases unique proline-rich polypeptides from its natural substrate, the hardened chorion. *FEBS Lett.*, 339:281–284, 1993.
- [1626] M.S. Lee, G.A. Kim, M.S. Seo, J.H. Lee, and S.T. Kwon. Characterization of heat-labile uracil-DNA glycosylase from *Psychrobacter* sp. HJ147 and its application to the polymerase chain reaction. *Biotechnol. Appl. Biochem.*, 52:167–175, 2009.
- [1627] S.-H. Lee, E. Minagawa, H. Taguchi, H. Matsuzawa, T. Ohta, S. Kaminogawa, and K. Yamauchi. Purification and characterization of a thermostable carboxypeptidase (carboxypeptidase Taq) from *Thermus aquaticus* YT-1. *Biosci. Biotechnol. Biochem.*, 56:1839–1844, 1992.
- [1628] S.-H. Lee, H. Taguchi, E. Yoshimura, E. Minagawa, S. Kaminogawa, T. Ohta, and H. Matsuzawa. Carboxypeptidase Taq, a thermostable zinc enzyme, from *Thermus aquaticus* YT-1: molecular cloning, sequencing, and expression of the encoding gene in *Escherichia coli*. *Biosci. Biotechnol. Biochem.*, 58:1490–1495, 1994.
- [1629] S.C. Lee, J. Chan, M.V. Clement, and S. Pervaiz. Functional proteomics of resveratrol-induced colon cancer cell apoptosis: caspase-6-mediated cleavage of lamin A is a major signaling loop. *Proteomics*, 6:2386–2394, 2006.
- [1630] S.J. Lee, S.Y. Park, J.J. Lee, D.Y. Yum, B.T. Koo, and J.K. Lee. Genes encoding the *N*-acyl homoserine lactone-degrading enzyme are widespread in many subspecies of *Bacillus thuringiensis*. *Appl. Environ. Microbiol.*, 68:3919–3924, 2002.

- [1631] S.L. Lee, R.B. Dickson, and C.Y. Lin. Activation of hepatocyte growth factor and urokinase/plasminogen activator by matriptase, an epithelial membrane serine protease. *J. Biol. Chem.*, 275:36720–36725, 2000.
- [1632] T.-C. Lee, B. Malone, and F. Snyder. A new de novo pathway for the formation of 1-alkyl-2-acetyl-*sn*-glycerols, precursors of platelet activating factor. Biochemical characterization of 1-alkyl-2-lyso-*sn*-glycero-3-P:acetyl-CoA acetyltransferase in rat spleen. *J. Biol. Chem.*, 261:5373–5377, 1986.
- [1633] W.H. Lee, Y.K. Kim, K.H. Nam, A. Priyadarshi, E.H. Lee, E.E. Kim, Y.H. Jeon, C. Cheong, and K.Y. Hwang. Crystal structure of the tRNA-specific adenosine deaminase from *Streptococcus pyogenes*. *Proteins*, 68:1016–1019, 2007.
- [1634] Y.-P. Lee. 5'-Adenylic acid deaminase. I. Isolation of the crystalline enzyme from rabbit skeletal muscle. *J. Biol. Chem.*, 227:987–992, 1957.
- [1635] Y.-P. Lee. 5'-Adenylic acid deaminase. II. Homogeneity and physicochemical properties. *J. Biol. Chem.*, 227:993–998, 1957.
- [1636] Y.-P. Lee. 5'-Adenylic acid deaminase. III. Properties and kinetic studies. *J. Biol. Chem.*, 227:999–1007, 1957.
- [1637] Y.-P. Lee, J.R. Sowokinos, and M.J. Erwin. Sugar phosphate phosphohydrolase. I. Substrate specificity, intracellular localization, and purification from *Neisseria meningitidis*. *J. Biol. Chem.*, 242:2264–2271, 1967.
- [1638] Y.J. Lee and R.B. Wickner. AFG1, a new member of the SEC18-NSF, PAS1, CDC48-VCP, TBP family of ATPases. *Yeast*, 8:787–790, 1992.
- [1639] H.J. Leese and G. Semenza. On the identity between the small intestinal enzymes phlorizin hydrolase and glycosylceramidase. *J. Biol. Chem.*, 248:8170–8173, 1973.
- [1640] I.R. Lehman and A.L. Nussbaum. The deoxyribonucleases of *Escherichia coli*. V. On the specificity of exonuclease I (phosphodiesterase). *J. Biol. Chem.*, 239:2628–2636, 1964.
- [1641] J.A. Leigh, A.J. Skinner, and R.A. Cooper. Partial purification, stereospecificity and stoichiometry of three dehalogenases from a *Rhizobium* species. *FEMS Microbiol. Lett.*, 49:353–356, 1988.
- [1642] S. Lemak, N. Beloglazova, B. Nocek, T. Skarina, R. Flick, G. Brown, A. Popovic, A. Joachimiak, A. Savchenko, and A.F. Yakunin. Toroidal structure and DNA cleavage by the CRISPR-associated [4Fe-4S] cluster containing Cas4 nuclease SSO0001 from *Sulfolobus solfataricus*. *J. Am. Chem. Soc.*, 135:17476–17487, 2013.
- [1643] M.K. Lemberg, J. Menendez, A. Misik, M. Garcia, C.M. Koth, and M. Freeman. Mechanism of intramembrane proteolysis investigated with purified rhomboid proteases. *EMBO J.*, 24:464–472, 2005.
- [1644] Y. Lemoine, A. Wach, and J.M. Jeltsch. To be free or not: the fate of pimelate in *Bacillus sphaericus* and in *Escherichia coli*. *Mol. Microbiol.*, 19:645–647, 1996.
- [1645] J.F. Lenney, M.H. Baslow, and G.H. Sugiyama. Similarity of tuna *N*-acetylhistidine deacetylase and cod fish anserinase. *Comp. Biochem. Physiol. B Comp. Biochem.*, 61:253–258, 1978.
- [1646] J.F. Lenney, R.P. George, A.M. Weiss, C.M. Kucera, P.W.H. Chan, and G.S. Rinzler. Human serum carnosinase: characterization, distinction from cellular carnosinase, and activation by cadmium. *Clin. Chim. Acta*, 123:221–231, 1982.
- [1647] O. Lenz, J. ter Meulen, H.D. Klenk, N.G. Seidah, and W. Garten. The Lassa virus glycoprotein precursor GP-C is proteolytically processed by subtilase SKI-1/S1P. *Proc. Natl. Acad. Sci. USA*, 98:12701–12705, 2001.
- [1648] P.M. Leonard and G. Grogan. Structure of 6-oxo camphor hydrolase H122A mutant bound to its natural product, (2*S*,4*S*)- $\alpha$ -campholinic acid: mutant structure suggests an atypical mode of transition state binding for a crotonase homolog. *J. Biol. Chem.*, 279:31312–31317, 2004.
- [1649] T. Lepage and C. Gache. Purification and characterization of the sea urchin embryo hatching enzyme. *J. Biol. Chem.*, 264:4787–4793, 1989.
- [1650] T. Lepage and C. Gache. Early expression of a collagenase-like hatching enzyme gene in the sea urchin embryo. *EMBO J.*, 9:3003–3012, 1990.
- [1651] N.R. Leslie, A.G. McLennan, and S.T. Safrany. Cloning and characterisation of hAps1 and hAps2, human diadenosine polyphosphate-metabolising Nudix hydrolases. *BMC Biochem.*, 3:20–20, 2002.



- [1652] M. Letarte, S. Vera, R. Tran, J.B.L. Addis, R.J. Onizuka, , and E. J., Jongneel, C.V. and McInnes, R.R. Common acute lymphocytic leukemia antigen is identical to neutral endopeptidase. *J. Exp. Med.*, 168:1247–1253, 1988.
- [1653] W. Leuzinger, A.L. Baker, and E. Cauvin. Acetylcholinesterase. II. Crystallization, absorption spectra, isoionic point. *Proc. Natl. Acad. Sci. USA*, 59:620–623, 1968.
- [1654] B. Levenberg and O. Hayaishi. A bacterial pterin deaminase. *J. Biol. Chem.*, 234:955–961, 1959.
- [1655] W. Levin, D.P. Michaud, P.E. Thomas, and D.M. Jerina. Distinct rat hepatic microsomal epoxide hydrolases catalyze the hydration of cholesterol 5,6  $\alpha$ -oxide and certain xenobiotic alkene and arene oxides. *Arch. Biochem. Biophys.*, 220:485–494, 1983.
- [1656] Y. Levin, R.A. Skidgel, and E.G. Erdös. Isolation and characterization of the subunits of human plasma carboxypeptidase N (kininase I). *Proc. Natl. Acad. Sci. USA*, 79:4618–4622, 1982.
- [1657] G.A. Levvy and C.A. Marsh.  $\beta$ -Glucuronidase. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 4, pages 397–407. Academic Press, New York, 2nd edition, 1960.
- [1658] G.A. Levvy and A. McAllan. Mammalian fucosidases. 2.  $\alpha$ -L-Fucosidase. *Biochem. J.*, 80:435–439, 1961.
- [1659] C.C. Levy and P. Goldman. Bacterial peptidases. *J. Biol. Chem.*, 244:4467–4472, 1969.
- [1660] M. Lewis and S. Weissman. Properties of a soluble nucleoside triphosphatase activity in mammalian liver. *Arch. Biochem. Biophys.*, 109:490–498, 1965.
- [1661] W.G. Lewis, J.M. Basford, and P.L. Walton. Specificity and inhibition studies of *Armillaria mellea* protease. *Biochim. Biophys. Acta*, 522:551–560, 1978.
- [1662] S.P. Leytus, K. Kurachi, K.S. Sakariassen, and E.W. Davie. Nucleotide sequence of the cDNA coding for human complement C1r. *Biochemistry*, 25:4855–4863, 1986.
- [1663] H. Li, L. Bergeron, V. Cryns, M.S. Pasternack, H. Zhu, L. Shi, A. Greenberg, and J. Yuan. Activation of caspase-2 in apoptosis. *J. Biol. Chem.*, 272:21010–21017, 1997.
- [1664] H. Li, H. Xu, D.E. Graham, and R.H. White. The *Methanococcus jannaschii* dCTP deaminase is a bifunctional deaminase and diphosphatase. *J. Biol. Chem.*, 278:11100–11106, 2003.
- [1665] J.Y. Li, K. Hollfelder, K.S. Huang, and M.G. Low. Structural features of GPI-specific phospholipase D revealed by fragmentation and  $\text{Ca}^{2+}$  binding studies. *J. Biol. Chem.*, 269:28963–28971, 1994.
- [1666] S. Li. L., Norioka, S. and Sakiyama, F. Molecular cloning and nucleotide sequence of the  $\beta$ -lytic protease gene from *Achromobacter lyticus*. *J. Bacteriol.*, 172:6506–6511, 1990.
- [1667] S.-C. Li and Y.-T. Li. Studies on the glycosidases of jack bean meal. 3. Crystallization and properties of  $\beta$ -N-acetylhexosaminidase. *J. Biol. Chem.*, 245:5153–5160, 1970.
- [1668] S.-J. Li and M. Hochstrasser. A new protease required for cell-cycle progression in yeast. *Nature*, 398:246–251, 1999.
- [1669] S.-J. Li and M. Hochstrasser. The Ulp1 SUMO isopeptidase: distinct domains required for viability, nuclear envelope localization, and substrate specificity. *J. Cell Biol.*, 160:1069–1081, 2003.
- [1670] S.C. Li, M.C. Chung, and C.S. Chen. Cloning and characterization of a DEAD box RNA helicase from the viable seedlings of aged mung bean. *Plant Mol. Biol.*, 47:761–770, 2001.
- [1671] W. Li, S.M. Srinivasula, J. Chai, P. Li, J.W. Wu, Z. Zhang, E.S. Alnemri, and Y. Shi. Structural insights into the pro-apoptotic function of mitochondrial serine protease HtrA2/Omi. *Nat. Struct. Biol.*, 9:436–441, 2002.
- [1672] X. Li, D. Apel, E.C. Gaynor, and M.E. Tanner. 5'-methylthioadenosine nucleosidase is implicated in playing a key role in a modified futasosine pathway for menaquinone biosynthesis in *Campylobacter jejuni*. *J. Biol. Chem.*, 286:19392–19398, 2011.
- [1673] X. Li, R.T. Su, H.T. Hsu, and H. Sze. The molecular chaperone calnexin associated with the vacuolar  $\text{H}^+$ -ATPase from oat seedlings. *Plant Cell*, 10:119–130, 1998.

- [1674] X.D. Li, L.F. Huergo, A. Gasperina, F.O. Pedrosa, M. Merrick, and F.K. Winkler. Crystal structure of dinitrogenase reductase-activating glycohydrolase (DraG) reveals conservation in the ADP-ribosylhydrolase fold and specific features in the ADP-ribose-binding pocket. *J. Mol. Biol.*, 390:737–746, 2009.
- [1675] Y.-T. Li. Presence of  $\alpha$ -D-mannosidic linkage in glycoproteins. Liberation of D-mannose from various glycoproteins by  $\alpha$ -mannosidase isolated from jack bean meal. *J. Biol. Chem.*, 241:1010–1012, 1966.
- [1676] Y.T. Li, L.X. Wang, N.V. Pavlova, S.C. Li, and Y.C. Lee.  $\alpha$ -KDOase activity in oyster and synthesis of  $\alpha$ - and  $\beta$ -4-methylumbelliferyl ketosides of 3-deoxy-D-manno-octulosonic acid (KDO). *J. Biol. Chem.*, 272:26419–26424, 1997.
- [1677] W.Q. Liang, Z.Y. Wang, H. Li, P.C. Wu, J.M. Hu, N. Luo, L.X. Cao, and Y.H. Liu. Purification and characterization of a novel pyrethroid hydrolase from *Aspergillus niger* ZD11. *J. Agric. Food Chem.*, 53:7415–7420, 2005.
- [1678] D.-I. Liao, K. Breddam, R.M. Sweet, T. Bullock, and S.J. Remington. Refined atomic model of wheat serine carboxypeptidase II at 2.2-Å resolution. *Biochemistry*, 31:9796–9812, 1992.
- [1679] D.I. Liao, J. Qian, D.A. Chisholm, D.B. Jordan, and B.A. Diner. Crystal structures of the photosystem II D1 C-terminal processing protease. *Nat. Struct. Biol.*, 7:749–753, 2000.
- [1680] P.R. Libby. Properties of an acetylspermidine deacetylase from rat liver. *Arch. Biochem. Biophys.*, 188:360–363, 1978.
- [1681] C. Liébecq, A. Lallemand, and M.-J. Degueldre-Guillaume. [Partial purification and properties of potato apyrase.]. *Bull. Soc. Chim. Biol.*, 45:573–594, 1963.
- [1682] I. Lieberman and A. Kornberg. Enzymatic synthesis and breakdown of a pyrimidine, orotic acid. II. Dihydroorotic acid, ureidosuccinic acid, and 5-carboxymethylhydantoin. *J. Biol. Chem.*, 207:911–924, 1954.
- [1683] I. Lieberman and A. Kornberg. Enzymatic synthesis and breakdown of a pyrimidine, orotic acid. III. Ureidosuccinase. *J. Biol. Chem.*, 212:909–920, 1955.
- [1684] I.E. Liener and B. Friedenson. Ficin. *Methods Enzymol.*, 19:261–273, 1970.
- [1685] A. Light and H. Janska. Enterokinase (enteropeptidase): comparative aspects. *Trends Biochem. Sci.*, 14:110–112, 1989.
- [1686] H.R. Lijnen, B. Van Hoef, L. Nelles, and D. Collen. Plasminogen activation with single-chain urokinase-type plasminogen activator (scu-PA). Studies with active site mutagenized plasminogen (Ser<sup>740</sup>→Glu). *J. Biol. Chem.*, 265:5232–5236, 1990.
- [1687] C.T. Lim and A.L. Horwitz. Purification and properties of human *N*-acetylgalactosamine-6-sulfate sulfatase. *Biochim. Biophys. Acta*, 657:344–355, 1981.
- [1688] P.E. Lim and M.E. Tate. The phytases. II. Properties of phytase fractions F1 and F2 from wheat bran and the *myo*-inositol phosphates produced by fraction F2. *Biochim. Biophys. Acta*, 302:316–328, 1973.
- [1689] C.D. Lima. Ulp1 endopeptidase. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 1340–1344. Elsevier, London, 2nd edition, 2004.
- [1690] C. Lin, S.M. Amberg, T.J. Chambers, and C.M. Rice. Cleavage at a novel site in the NS4A region by the yellow fever virus NS2B-3 proteinase is a prerequisite for processing at the downstream 4A/4B signalase site. *J. Virol.*, 67:2327–2335, 1993.
- [1691] C.Y. Lin, J. Anders, M. Johnson, Q.A. Sang, and R.B. Dickson. Molecular cloning of cDNA for matriptase, a matrix-degrading serine protease with trypsin-like activity. *J. Biol. Chem.*, 274:18231–18236, 1999.
- [1692] L. Lin, I. Sohar, H. Lackland, and P. Lobel. The human CLN2 protein/tripeptidyl-peptidase I is a serine protease that autoactivates at acidic pH. *J. Biol. Chem.*, 276:2249–2255, 2001.
- [1693] S. Lin, R.E. Hanson, and J.E. Cronan. Biotin synthesis begins by hijacking the fatty acid synthetic pathway. *Nat. Chem. Biol.*, 6:682–688, 2010.
- [1694] W. Lin, J.C. Ame, N. Aboul-Ela, E.L. Jacobson, and M.K. Jacobson. Isolation and characterization of the cDNA encoding bovine poly(ADP-ribose) glycohydrolase. *J. Biol. Chem.*, 272:11895–11901, 1997.

- [1695] X. Lin and J. Tang. Thermopsin. *Methods Enzymol.*, 248:156–168, 1995.
- [1696] X.Y. Lin, M.S. Choi, and A.G. Porter. Expression analysis of the human caspase-1 subfamily reveals specific regulation of the *CASP5* gene by lipopolysaccharide and interferon- $\gamma$ . *J. Biol. Chem.*, 275:39920–39926, 2000.
- [1697] Y.H. Lin, J.L. Xu, J. Hu, L.H. Wang, S.L. Ong, J.R. Leadbetter, and L.H. Zhang. Acyl-homoserine lactone acylase from *Ralstonia* strain XJ12B represents a novel and potent class of quorum-quenching enzymes. *Mol. Microbiol.*, 47:849–860, 2003.
- [1698] Z. Lin, X. Su, W. Chen, B. Ci, S. Zhang, and H. Lin. Dph7 catalyzes a previously unknown demethylation step in diphthamide biosynthesis. *J. Am. Chem. Soc.*, 136:6179–6182, 2014.
- [1699] T. Lindahl, J.A. Gally, and G.M. Edelman. Deoxyribonuclease IV: a new exonuclease from mammalian tissues. *Proc. Natl. Acad. Sci. USA*, 62:597–603, 1969.
- [1700] T. Lindahl, J.A. Gally, and G.M. Edelman. Properties of deoxyribonuclease 3 from mammalian tissues. *J. Biol. Chem.*, 244:5014–5019, 1969.
- [1701] H. Lindner, S. Hopfner, M. Tafler-Naumann, M. Miko, L. Konrad, and K.H. Rohm. The distribution of aminoacylase I among mammalian species and localization of the enzyme in porcine kidney. *Biochimie*, 82:129–137, 2000.
- [1702] P.A. Lindquist, K. Fujikawa, and E.W. Davie. Activation of bovine factor IX (Christmas factor) by factor XIa (activated plasma thromboplastin antecent) and a protease from Russell’s viper venom. *J. Biol. Chem.*, 253:1902–1909, 1978.
- [1703] L.L. Lindsay, M.J. Wieduwilt, and J.L. Hedrick. Oviductin, the *Xenopus laevis* oviductal protease that processes egg envelope glycoprotein gp43, increases sperm binding to envelopes, and is translated as part of an unusual mosaic protein composed of two protease and several CUB domains. *Biol. Reprod.*, 60:989–995, 1999.
- [1704] H. Lineweaver and E.F. Jansen. Pectic enzymes. *Adv. Enzymol. Relat. Subj. Biochem.*, 11:267–295, 1951.
- [1705] V. Ling. Partial digestion of  $^{32}\text{P}$ -fd DNA with  $\text{T}^4$  endonuclease IV. *FEBS Lett.*, 19:50–54, 1971.
- [1706] R.P. Link and F.J. Castellino. Kinetic comparison of bovine blood coagulation factor IX $\alpha$  and IX $\beta$  towards bovine factor X. *Biochemistry*, 22:4033–4041, 1983.
- [1707] A. Linker, K. Meyer, and P. Hoffman. The production of hyaluronate oligosaccharides by leech hyaluronidase and alkali. *J. Biol. Chem.*, 235:924–927, 1960.
- [1708] T.C. Linn, J.W. Pelley, F.H. Petit, F. Hucho, D.D. Randall, and L.J. Reed.  $\alpha$ -Keto acid dehydrogenase complexes. XV. Purification and properties of the component enzymes of the pyruvate dehydrogenase complexes from bovine kidney and heart. *Arch. Biochem. Biophys.*, 148:327–342, 1972.
- [1709] M.N. Lioubin, P.A. Algate, S. Tsai, K. Carlberg, A. Aebersold, and L.R. Rohrschneider. p150Ship, a signal transduction molecule with inositol polyphosphate-5-phosphatase activity. *Genes Dev.*, 10:1084–1095, 1996.
- [1710] B. Lipinska, M. Zylicz, and C. Georgopoulos. The HtrA (DegP) protein, essential for *Escherichia coli* survival at high temperatures, is an endopeptidase. *J. Bacteriol.*, 172:1791–1797, 1990.
- [1711] D.L. Lips and P.W. Majerus. The discovery of a 3-phosphomonoesterase that hydrolyzes phosphatidylinositol 3-phosphate in NIH 3T3 cells. *J. Biol. Chem.*, 264:19911–19915, 1989.
- [1712] C. Little and A.-B. Otnass. The metal ion dependence of phospholipase C from *Bacillus cereus*. *Biochim. Biophys. Acta*, 391:326–333, 1975.
- [1713] J.W. Little. An exonuclease induced by bacteriophage lambda. II. Nature of the enzymatic reaction. *J. Biol. Chem.*, 242:679–686, 1967.
- [1714] J.W. Little, B. Kim, K.L. Roland, M.H. Smith, L.-L. Lin, and S.N. Slilaty. Cleavage of LexA repressor. *Methods Enzymol.*, 244:266–284, 1994.
- [1715] B. Liu, I.N. Clarke, and P.R. Lambden. Polyprotein processing in Southampton virus: identification of 3C-like protease cleavage sites by in vitro mutagenesis. *J. Virol.*, 70:2605–2610, 1996.

- [1716] B.L. Liu, G.J. Viljoen, I.N. Clarke, and P.R. Lambden. Identification of further proteolytic cleavage sites in the Southampton calicivirus polyprotein by expression of the viral protease in *E. coli*. *J. Gen. Virol.*, 80:291–296, 1999.
- [1717] C. Liu, D. Li, Y.H. Liang, L.F. Li, and X.D. Su. Ring-opening mechanism revealed by crystal structures of NagB and its ES intermediate complex. *J. Mol. Biol.*, 379:73–81, 2008.
- [1718] D. Liu, B.W. Lepore, G.A. Petsko, P.W. Thomas, E.M. Stone, W. Fast, and D. Ringe. Three-dimensional structure of the quorum-quenching *N*-acyl homoserine lactone hydrolase from *Bacillus thuringiensis*. *Proc. Natl. Acad. Sci. USA*, 102:11882–11887, 2005.
- [1719] F. Liu and M.E. Tanner. PseG of pseudaminic acid biosynthesis: a UDP-sugar hydrolase as a masked glycosyltransferase. *J. Biol. Chem.*, 281:20902–20909, 2006.
- [1720] H. Liu, N.D. Rodgers, X. Jiao, and M. Kiledjian. The scavenger mRNA decapping enzyme DcpS is a member of the HIT family of pyrophosphatases. *EMBO J.*, 21:4699–4708, 2002.
- [1721] J.-Q. Liu, T. Kurihara, A.K.M.Q. Hasan, V. Nardi-Dei, H. Koshikawa, N. Esaki, and K. Soda. Purification and characterization of thermostable and nonthermostable 2-haloacid dehalogenases with different stereospecificities from *Pseudomonas* sp. strain YL. *Appl. Environ. Microbiol.*, 60:2389–2393, 1994.
- [1722] K. Liu, T.T. Liu, and N.Y. Zhou. HbzF catalyzes direct hydrolysis of maleylpyruvate in the gentisate pathway of *Pseudomonas alcaligenes* NCIMB 9867. *Appl. Environ. Microbiol.*, 79:1044–1047, 2013.
- [1723] L. Liu, J.F. Wu, Y.F. Ma, S.Y. Wang, G.P. Zhao, and S.J. Liu. A novel deaminase involved in chloronitrobenzene and nitrobenzene degradation with *Comamonas* sp. strain CNB-1. *J. Bacteriol.*, 189:2677–2682, 2007.
- [1724] Q. Liu, I.A. Kriksunov, R. Graeff, C. Munshi, H.C. Lee, and Q. Hao. Crystal structure of human CD38 extracellular domain. *Structure*, 13:1331–1339, 2005.
- [1725] S.W. Liu, V. Rajagopal, S.S. Patel, and M. Kiledjian. Mechanistic and kinetic analysis of the DcpS scavenger decapping enzyme. *J. Biol. Chem.*, 283:16427–16436, 2008.
- [1726] T.-Y. Liu and S.D. Elliott. Streptococcal proteinase. In P.D. Boyer, editor, *The Enzymes*, pages 609–647. Academic Press, New York, 3rd edition, 1971.
- [1727] Z. Liu and D.A. Bryant. Identification of a gene essential for the first committed step in the biosynthesis of bacteriochlorophyll *c*. *J. Biol. Chem.*, 286:22393–22402, 2011.
- [1728] M.O. Lively, A.L. Newsome, and M. Nusier. Eukaryote microsomal signal peptidases. *Methods Enzymol.*, 244:301–314, 1994.
- [1729] M. Llanillo, N. Perez, and J.A. Cabezas.  $\beta$ -Galactosidase and  $\beta$ -glucosidase activities of the same enzyme from rabbit liver. *Int. J. Biochem.*, 8:557–564, 1977.
- [1730] S.-S. Lo, B.A. Fraser, and T.-Y. Liu. The mixed disulphide in the zymogen of streptococcal proteinase. Characterization and implication for its biosynthesis. *J. Biol. Chem.*, 259:11041–11045, 1984.
- [1731] S.L. Loayza, M. Trikha, F.S. Markland, P. Riquelme, and J. Kuo. Resolution of isoforms of natural and recombinant fibrolase, the fibrinolytic enzyme from *Agkistrodon contortrix contortrix* snake venom, and comparison of their EDTA sensitivities. *J. Chromatogr. B Biomed. Appl.*, 662:227–243, 1994.
- [1732] Y.P. Loh. Kinetic studies on the processing of human  $\beta$ -lipotropin by bovine pituitary intermediate lobe *pro*-opiomelanocortin-converting enzyme. *J. Biol. Chem.*, 261:11949–11955, 1986.
- [1733] Y.P. Loh, D.C. Parish, and R. Tuteja. Purification and characterization of a paired basic residue-specific *pro*-opiomelanocortin converting enzyme from bovine pituitary intermediate lobe secretory vesicles. *J. Biol. Chem.*, 260:7194–7205, 1985.
- [1734] J. London, S.Z. Hausman, and J. Thompson. Characterization of a membrane-regulated sugar phosphate phosphohydrolase from *Lactobacillus casei*. *J. Bacteriol.*, 163:951–956, 1985.

- [1735] N. London, J.D. Farelli, S.D. Brown, C. Liu, H. Huang, M. Korczynska, N.F. Al-Obaidi, P.C. Babbitt, S.C. Almo, K.N. Allen, and B.K. Shoichet. Covalent docking predicts substrates for haloalkanoate dehalogenase superfamily phosphatases. *Biochemistry*, 54:528–537, 2015.
- [1736] H. Lorenz-Meyer, A.L. Blum, H.P. Haemmerli, and G. Semenza. A second enzyme defect in acquired lactase deficiency: lack of small-intestinal phlorizin-hydrolase. *Eur. J. Clin. Invest.*, 2:326–331, 1972.
- [1737] S. Lory and M.S. Strom. Structure-function relationship of type-IV prepilin peptidase of *Pseudomonas aeruginosa* - A review. *Gene*, 192:117–121, 1997.
- [1738] D.J. Loskutoff and R.R. Schleef. Plasminogen activators and their inhibitors. *Methods Enzymol.*, 163:293–302, 1988.
- [1739] T.J. Lott, L.S. Page, P. Boiron, J. Benson, and E. Reiss. Nucleotide sequence of the *Candida albicans* aspartyl proteinase gene. *Nucleic Acids Res.*, 17:1779–1779, 1989.
- [1740] R. Lottenberg, U. Christiansen, C.M. Jackson, and P.L. Coleman. Assay of coagulation proteases using peptide chromogenic and fluorogenic substrates. *Methods Enzymol.*, 80:341–361, 1981.
- [1741] A.L. Lovering, S.S. Lee, Y.W. Kim, S.G. Withers, and N.C. Strynadka. Mechanistic and structural analysis of a family 31  $\alpha$ -glycosidase and its glycosyl-enzyme intermediate. *J. Biol. Chem.*, 280:2105–2115, 2005.
- [1742] M.G. Low and A.R.S. Prasad. A phospholipase D specific for the phosphatidylinositol anchor of cell-surface proteins is abundant in plasma. *Proc. Natl. Acad. Sci. USA*, 85:980–984, 1988.
- [1743] A.Y. Lu, D. Ryan, D.M. Jerina, J.W. Daly, and W. Levin. Liver microsomal epoxide hydrase. Solubilization, purification, and characterization. *J. Biol. Chem.*, 250:8283–8288, 1975.
- [1744] G. Lu, J. Zhang, Y. Li, Z. Li, N. Zhang, X. Xu, T. Wang, Z. Guan, G.F. Gao, and J. Yan. hNUDT16: a universal decapping enzyme for small nucleolar RNA and cytoplasmic mRNA. *Protein Cell*, 2:64–73, 2011.
- [1745] P.-J. Lu, H.-C. Liu, and I.-H. Tsai. The midgut trypsins of shrimp (*Penaeus monodon*). High efficiency toward native protein substrates including collagens. *Biol. Chem. Hoppe-Seyler*, 371:851–859, 1990.
- [1746] W.-P. Lu and D.P. Kelly. Cellular location and partial purification of the ‘thiosulphate-oxidizing enzyme’ and ‘trithionate hydrolase’ from *Thiobacillus tepidarius*. *J. Gen. Microbiol.*, 134:877–885, 1988.
- [1747] Z. Lu, L. Wang, D. Dunaway-Mariano, and K.N. Allen. Structure-function analysis of 2-keto-3-deoxy-D-glycero-D-galactononate-9-phosphate phosphatase defines specificity elements in type C<sub>0</sub> haloalkanoate dehalogenase family members. *J. Biol. Chem.*, 284:1224–1233, 2009.
- [1748] A.L. Luaces and A.J. Barrett. Affinity purification and biochemical characterization of histolysin: the major cysteine proteinase of *Entamoeba histolytica*. *Biochem. J.*, 250:903–909, 1988.
- [1749] W.A. Lubas and R.G. Spiro. Golgi endo- $\alpha$ -D-mannosidase from rat liver, a novel N-linked carbohydrate unit processing enzyme. *J. Biol. Chem.*, 262:3775–3781, 1987.
- [1750] J. Lubkowski, A. Wlodawer, H.L. Ammon, T.D. Copeland, and A.L. Swain. Structural characterization of *Pseudomonas* 7A glutaminase-asparaginase. *Biochemistry*, 33:10257–10265, 1994.
- [1751] C. Lugnier, T. Keravis, A. Le Bec, O. Pauvert, S. Proteau, and E. Rousseau. Characterization of cyclic nucleotide phosphodiesterase isoforms associated to isolated cardiac nuclei. *Biochim. Biophys. Acta*, 1472:431–446, 1999.
- [1752] P. Lund and B. Magasanik. N-Formimino-L-glutamate formiminohydrolase of *Aerobacter aerogenes*. *J. Biol. Chem.*, 240:4316–4319, 1965.
- [1753] R.L. Lundblad, H.S. Kingdon, and K.G. Mann. Thrombin. *Methods Enzymol.*, 45:156–176, 1976.
- [1754] J.E. Lunn, A.R. Ashton, M.D. Hatch, and H.W. Heldt. Purification, molecular cloning, and sequence analysis of sucrose-6<sup>F</sup>-phosphate phosphohydrolase from plants. *Proc. Natl. Acad. Sci. USA*, 97:12914–12919, 2000.
- [1755] J.E. Lunn and E. MacRae. New complexities in the synthesis of sucrose. *Curr. Opin. Plant Biol.*, 6:208–214, 2003.

- [1756] M.J. Lush, Y. Li, D.J. Read, A.C. Willis, and P. Glynn. Neuropathy target esterase and a homologous *Drosophila* neurodegeneration-associated mutant protein contain a novel domain conserved from bacteria to man. *Biochem. J.*, 332:1–4, 1998.
- [1757] W.B. Lushbaugh, A.F. Hofbauer, and F.E. Pittman. *Entamoeba histolytica*: purification of cathepsin B. *Exp. Parasitol.*, 59:328–336, 1985.
- [1758] R. Luthi-Carter, U.V. Berger, A.K. Barczak, M. Enna, and J.T. Coyle. Isolation and expression of a rat brain cDNA encoding glutamate carboxypeptidase II. *Proc. Natl. Acad. Sci. USA*, 95:3215–3220, 1998.
- [1759] K.R. Lynn and N.A. Clevette-Radford. Two proteases from the latex of *Elaeophorbium drupifera*. *Phytochemistry*, 24:2843–2845, 1985.
- [1760] W.S. Lynn and N.C. Perryman. Properties and purification of adipose tissue lipase. *J. Biol. Chem.*, 235:1912–1916, 1960.
- [1761] M., Ikeda Kato, , and K. 1 and K2, isolated from mycelia of *Aspergillus oryzae*. I. Isolation and purification of DNases K1 and K2. *J. Biochem. (Tokyo)*, 64:321–328, 1968.
- [1762] S. Maas, A.P. Gerber, and A. Rich. Identification and characterization of a human tRNA-specific adenosine deaminase related to the ADAR family of pre-mRNA editing enzymes. *Proc. Natl. Acad. Sci. USA*, 96:8895–8900, 1999.
- [1763] M.R. Macbeth, H.L. Schubert, A.P. Vandemark, A.T. Lingam, C.P. Hill, and B.L. Bass. Inositol hexakisphosphate is bound in the ADAR2 core and required for RNA editing. *Science*, 309:1534–1539, 2005.
- [1764] R.T.A. MacGillivray and E.W. Davie. Characterization of bovine prothrombin mRNA and its translation product. *Biochemistry*, 23:1626–1634, 1984.
- [1765] V.L. Mackay, J. Armstrong, C. Yip, S. Welch, K. Walker, S. Osborn, P. Sheppard, and J. Forstrom. Characterization of the bar proteinase, an extracellular enzyme from the yeast *Saccharomyces cerevisiae*. *Adv. Exp. Med. Biol.*, 306:161–172, 1991.
- [1766] V.L. Mackay, S.K. Welch, M.Y. Insley, T.R. Manney, J. Holly, G.C. Saari, and M.L. Parker. The *Saccharomyces cerevisiae* BAR1 gene encodes an exported protein with homology to pepsin. *Proc. Natl. Acad. Sci. USA*, 85:55–59, 1988.
- [1767] C.M. MacKinnon, P.E. Carter, S.J. Smyth, B. Dunbar, and J.E. Fothergill. Molecular cloning of cDNA for human complement component C1s. The complete amino acid sequence. *Eur. J. Biochem.*, 169:547–553, 1987.
- [1768] M.I. Mackness, H.M. Thompson, A.R. Hardy, and C.H. Walker. Distinction between 'A'-esterases and arylesterases. Implications for esterase classification. *Biochem. J.*, 245:293–296, 1987.
- [1769] T.K. MacLachlan and W.S. El-Deiry. Apoptotic threshold is lowered by p53 transactivation of caspase-6. *Proc. Natl. Acad. Sci. USA*, 99:9492–9497, 2002.
- [1770] S. Mafu, M.L. Hillwig, and R.J. Peters. A novel labda-7,13E-dien-15-ol-producing bifunctional diterpene synthase from *Selaginella moellendorffii*. *Chembiochem.*, 12:1984–1987, 2011.
- [1771] J.L. Magnard, A. Rocchia, J.C. Caissard, P. Vergne, P. Sun, R. Hecquet, A. Dubois, L. Hibrand-Saint Oyant, F. Jullien, F. Nicole, O. Raymond, S. Huguet, R. Baltenweck, S. Meyer, P. Claudel, J. Jeauffre, M. Rohmer, F. Foucher, P. Hugueney, M. Bendahmane, and S. Baudino. Plant volatiles. Biosynthesis of monoterpene scent compounds in roses. *Science*, 349:81–83, 2015.
- [1772] S. Magnusson. Bovine prothrombin and thrombin. *Methods Enzymol.*, 19:157–184, 1970.
- [1773] S. Mahadevan and F. Sauer. Carnitine ester hydrolase of rat liver. *J. Biol. Chem.*, 244:4448–4453, 1969.
- [1774] S. Mahadevan and A.L. Tappel.  $\beta$ -Aspartylglucosylamine amido hydrolase of rat liver and kidney. *J. Biol. Chem.*, 242:4568–4576, 1967.
- [1775] D. Mahuran, P. Clements, and J. Hopwood. A rapid four column purification of 2-deoxy-D-glucoside-2-sulphamate sulphohydrolase from human liver. *Biochim. Biophys. Acta*, 757:359–365, 1983.
- [1776] V.P. Maier, S. Hasegawa, and E. Hera. Limonin D-ring-lactone hydrolase. A new enzyme from *Citrus* seeds. *Phytochemistry*, 8:405–407, 1969.

- [1777] A.R. Main. The differentiation of the A-type esterases in sheep serum. *Biochem. J.*, 75:188–195, 1960.
- [1778] C. Mainardi. L. and Hasty, K. A. Secretion and glycosylation of rabbit macrophage type V collagenase. *Matrix*, 10:84–90, 1990.
- [1779] G. Mains, M. Takahashi, J. Sodek, and T. Hofmann. The specificity of penicillopepsin. *Can. J. Biochem.*, 49:1134–1149, 1971.
- [1780] T. Maita, S. Nagata, G. Matsuda, S. Maruta, K. Oda, S. Murao, and D. Tsuru. Complete amino acid sequence of *Scytalidium lignicolum* acid protease B. *J. Biochem. (Tokyo)*, 95:465–473, 1984.
- [1781] G.S. Maitz, E.M. Haas, and P.A. Castric. Purification and properties of the allophanate hydrolase from *Chlamydomonas reinhardtii*. *Biochim. Biophys. Acta*, 714:486–491, 1982.
- [1782] E. Majima, K. Oda, S. Murao, and E. Ichishima. Comparative study on the specificities of several fungal aspartic and acidic proteinases towards the tetradecapeptide of a renin substrate. *Agric. Biol. Chem.*, 52:787–793, 1988.
- [1783] K.K. Makinen and P.-L. Makinen. Purification and properties of an extracellular collagenolytic protease produced by the human oral bacterium *Bacillus cereus* (strain Soc 67). *J. Biol. Chem.*, 262:12488–12495, 1987.
- [1784] P.-L. Mäkinen, D.B. Clewell, F. An, , and K.K. Purification and substrate specificity of a strongly hydrophobic extracellular metalloendopeptidase (“gelatinase”) from *Streptococcus faecalis* (strain 0G1-10). *J. Biol. Chem.*, 264:3325–3334, 1989.
- [1785] M.H. Malamy and B.L. Horecker. Purification and crystallization of the alkaline phosphatase of *Escherichia coli*. *Biochemistry*, 3:1893–1897, 1964.
- [1786] P. Malathi and R.K. Crane. Phlorizin hydrolase: a  $\beta$ -glucosidase of hamster intestinal brush border membrane. *Biochim. Biophys. Acta*, 173:245–256, 1969.
- [1787] B. Malfroy, P.R. Schofield, W.-J. Kuang, P.H. Seeburg, Mason, and W.J. Molecular cloning and amino acid sequence of rat enkephalinase. *Biochem. Biophys. Res. Commun.*, 144:59–66, 1987.
- [1788] O.P. Malhotra and A.M. Kayastha. Chemical inactivation and active site groups of phosphoenolpyruvate-phosphatase from germinating mung beans (*Vigna radiata*). *Plant Sci.*, 65:161–170, 1989.
- [1789] O.P. Malhotra and A.M. Kayastha. Isolation and characterization of phosphoenolpyruvate phosphatase from germinating mung beans (*Vigna radiata*). *Plant Physiol.*, 93:194–200, 1990.
- [1790] O.P. Malhotra and G. Philip. Specificity of goat intestinal esterase. *Biochem. Z.*, 346:386–402, 1966.
- [1791] A.-S. Malik and M.G. Low. Conversion of human placental alkaline phosphatase from a high  $M_r$  form to a low  $M_r$  form during butanol extraction. An investigation of the role of endogenous phosphoinositide-specific phospholipases. *Biochem. J.*, 240:519–527, 1986.
- [1792] P.A. Mallory and J. Travis. Human pancreatic enzymes: purification and characterization of a nonelastolytic enzyme, protease E, resembling elastase. *Biochemistry*, 14:722–729, 1975.
- [1793] H. Malmgren. Enzymatic behavior of polymetaphosphate. V. Purification and specificity of the enzyme. *Acta Chem. Scand.*, 6:16–26, 1952.
- [1794] S.E. Maloney, A. Maule, and A.R. Smith. Purification and preliminary characterization of permethrinase from a pyrethroid-transforming strain of *Bacillus cereus*. *Appl. Environ. Microbiol.*, 59:2007–2013, 1993.
- [1795] N. Malys and J.E. McCarthy. Dcs2, a novel stress-induced modulator of m<sup>7</sup>GpppX pyrophosphatase activity that locates to P bodies. *J. Mol. Biol.*, 363:370–382, 2006.
- [1796] M. Mancini, C.E. Machamer, S. Roy, D.W. Nicholson, N.A. Thornberry, L.A. Casciola-Rosen, and A. Rosen. Caspase-2 is localized at the Golgi complex and cleaves golgin-160 during apoptosis. *J. Cell Biol.*, 149:603–612, 2000.
- [1797] F.J. Mandelbaum, A.P. Reichel, and M.T. Assakura. Isolation and characterization of a proteolytic enzyme from the venom of the snake *Bothrops jararaca* (jararaca). *Toxicon*, 20:955–972, 1982.

- [1798] F.R. Mandelbaum, A.P. Reichl, and M.T. Assakura. Some physical and biochemical characteristics of HF2, one of the hemorrhagic factors in the venom of *Bothrops jararaca*. In A. Ohsaka, K. Hayashi, and Y. Sawai, editors, *Animal, Plant and Microbial Toxins*, pages 111–121. Plenum Press, New York, 1976.
- [1799] K.G. Mann. Prothrombin. *Methods Enzymol.*, 45:123–156, 1976.
- [1800] D.J. Manners. Enzymic synthesis and degradation of starch and glycogen. *Adv. Carbohydr. Chem.*, 17:371–430, 1962.
- [1801] D.J. Manners. Observations on the specificity and nomenclature of starch debranching enzymes. *J. Appl. Glycosci.*, 44:83–85, 1997.
- [1802] D.S. Manning and J.C. Leong. Expression in *Escherichia coli* of the large genomic segment of infectious pancreatic necrosis virus. *Virology*, 179:16–25, 1990.
- [1803] E. Manser, , and D. , Loo,L., Goh, P.Y., Monfries, C., Hall, C. and Lim, L. Human carboxypeptidase E: isolation and characterisation of the cDNA, sequence conservation, expression and processing in vitro. *Biochem. J.*, 267:517–525, 1990.
- [1804] C.A. Mapes and C.C. Sweeley. Substrate specificity of ceramide trihexosidase. *FEBS Lett.*, 25:279–281, 1972.
- [1805] B. Maras, D. Barra, S. Duprè, and G. Pitari. Is pantetheinase the actual identity of mouse and human vanin-1 proteins? *FEBS Lett.*, 461:149–152, 1999.
- [1806] P. Marchant, V.A. Manneh, and J. Blankenship.  $N^1$ -Acetylspermidine is not a substrate for  $N$ -acetylspermidine deacetylase. *Biochim. Biophys. Acta*, 881:297–299, 1986.
- [1807] S. Marcinowski and H. Grisebach. Enzymology of lignification. Cell-wall bound  $\beta$ -glucosidase for coniferin from spruce (*Picea abies*) seedlings. *Eur. J. Biochem.*, 87:37–44, 1978.
- [1808] E.M. Marcotte, A.F. Monzingo, S.R. Ernst, R. Brzezinski, and J.D. Robertus. X-ray structure of an anti-fungal chitosanase from *Streptomyces* N174. *Nat. Struct. Biol.*, 3:155–162, 1996.
- [1809] N. Margolin, S.A. Raybuck, K.P. Wilson, W.Y. Chen, T. Fox, Y. Gu, and D.J. Livingston. Substrate and inhibitor specificity of interleukin-1 $\beta$ -converting enzyme and related caspases. *J. Biol. Chem.*, 272:7223–7228, 1997.
- [1810] E. Margolles-Clark, M. Tenkanen, T. Nakari-Setälä, and M. Penttilä. Cloning of genes encoding  $\alpha$ -L-arabinofuranosidase and  $\beta$ -xylosidase from *Trichoderma reesei* by expression in *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.*, 62:3840–3846, 1996.
- [1811] E. Margolles-Clark, M. Tenkanen, H. Söderlund, , and M. Acetyl xylan esterase from *Trichoderma reesei* contains an active site serine and a cellulose binding domain. *Eur. J. Biochem.*, 237:553–560, 1996.
- [1812] M.J. Mark and S. Kornfeld. Partial purification and characterization of the glucosidases involved in the processing of asparagine-linked oligosaccharides. *Arch. Biochem. Biophys.*, 199:249–258, 1980.
- [1813] F.S. Markland, C. Kettner, S. Schiffmann, E. Shaw, S.S. Bajwa, K.N.N. Reddy, H. Kirakossian, G.B. Patkos, I. Theodor, and H. Pirkle. Kallikrein-like activity of crotalase, a snake venom enzyme that clots fibrinogen. *Proc. Natl. Acad. Sci. USA*, 79:1688–1692, 1982.
- [1814] F.S. Markland and E.L. Smith. Subtilisins: primary structure, chemical and physical properties. In P.D. Boyer, editor, *The Enzymes*, volume 3, pages 561–608. Academic Press, New York, 3rd edition, 1971.
- [1815] F. Marks and I. Raab. The second messenger system of mouse epidermis. IV. Cyclic AMP and cyclic GMP phosphodiesterase. *Biochim. Biophys. Acta*, 334:368–377, 1974.
- [1816] M. Martens, M. Cottenie-Ruysschaert, R. Hanselaer, L. De Cooman, K.V. Castele, and F. Van Sumere. Characteristics and specificity of purified  $N$ -feruloylglycine amidohydrolase from isolated barley embryos. *Phytochemistry*, 27:2465–2475, 1988.
- [1817] M. Martens, M. Cottenie-Ruysschaert, R. Hanselaer, L. De Cooman, K.V. Castele, and F. Van Sumere.  $N$ -Feruloylglycine amidohydrolase from barley seeds and isolated barley embryos. *Phytochemistry*, 27:2457–2463, 1988.



- [1818] F. Martin, F. Malergue, G. Pitari, J.M. Philippe, S. Philips, C. Chabret, S. Granjeaud, M.G. Mattei, A.J. Mungall, P. Naquet, and F. Galland. Vanin genes are clustered (human 6q22-24 and mouse 10A2B1) and encode isoforms of pantetheinase ectoenzymes. *Immunogenetics*, 53:296–306, 2001.
- [1819] P. Martin, P. Trieu-Cuot, J.-C. Collin, and B. Ribadeau Dumas. Purification and characterization of bovine gastricsin. *Eur. J. Biochem.*, 122:31–39, 1982.
- [1820] S.J. Martin, G.P. Amarante-Mendes, L. Shi, T.H. Chuang, C.A. Casiano, G.A. O'Brien, P. Fitzgerald, E.M. Tan, G.M. Bokoch, A.H. Greenberg, and D.R. Green. The cytotoxic cell protease granzyme B initiates apoptosis in a cell-free system by proteolytic processing and activation of the ICE/CED-3 family protease, CPP32, via a novel two-step mechanism. *EMBO J.*, 15:2407–2416, 1996.
- [1821] S.S. Martin and H.E. Senior. Membrane adenosine triphosphatase activities in rat pancreas. *Biochim. Biophys. Acta*, 602:401–418, 1980.
- [1822] S. Martínez-Rodríguez, J.M. Clemente-Jiménez, F. Rodríguez-Vico, and F.J. Las Heras-Vázquez. Molecular cloning and biochemical characterization of L-N-carbamoylase from *Sinorhizobium meliloti* CECT4114. *J. Mol. Microbiol. Biotechnol.*, 9:16–25, 2005.
- [1823] F. Martinon and J. Tschopp. Inflammatory caspases: linking an intracellular innate immune system to autoinflammatory diseases. *Cell*, 117:561–574, 2004.
- [1824] L.M. Martins, B.E. Turk, V. Cowling, A. Borg, E.T. Jarrell, L.C. Cantley, and J. Downward. Binding specificity and regulation of the serine protease and PDZ domains of HtrA2/Omi. *J. Biol. Chem.*, 278:49417–49427, 2003.
- [1825] K. Maruta, T. Nakada, M. Kubota, H. Chaen, T. Sugimoto, M. Kurimoto, , and Y. Formation of trehalose from maltooligosaccharides by a novel enzymatic system. *Biosci. Biotechnol. Biochem.*, 59:1829–1834, 1995.
- [1826] C. Maruyama and Y. Hamano. The biological function of the bacterial isochorismatase-like hydrolase SttH. *Biosci. Biotechnol. Biochem.*, 73:2494–2500, 2009.
- [1827] K. Maruyama. Purification and properties of 2-pyrone-4,6-dicarboxylate hydrolase. *J. Biochem. (Tokyo)*, 93:557–565, 1983.
- [1828] Y. Maruyama, Y. Nakamichi, T. Itoh, B. Mikami, W. Hashimoto, and K. Murata. Substrate specificity of streptococcal unsaturated glucuronyl hydrolases for sulfated glycosaminoglycan. *J. Biol. Chem.*, 284:18059–18069, 2009.
- [1829] S.P. Marx, J. Nösberger, and M. Frehner. Hydrolysis of fructan in grasses: A  $\beta$ -(2-6)-linkage specific fructan- $\beta$ -fructosidase from stubble of *Lolium perenne*. *New Phytol.*, 135:279–290, 1997.
- [1830] T. Masaki, T. Fujihashi, K. Nakamura, and M. Soejima. Studies on a new proteolytic enzyme from *Achromobacter lyticus* M497-1. II. Specificity and inhibition studies of *Achromobacter* protease I. *Biochim. Biophys. Acta*, 660:51–55, 1981.
- [1831] T. Masaki, M. Tanabe, K. Nakamura, and M. Soejima. Studies on a new proteolytic enzyme from *Achromobacter lyticus* M497-1. I. Purification and some enzymatic properties. *Biochim. Biophys. Acta.*, 660:44–50, 1981.
- [1832] Z. Mashhadi, H. Xu, and R.H. White. An Fe<sup>2+</sup>-dependent cyclic phosphodiesterase catalyzes the hydrolysis of 7,8-dihydro-D-neopterin 2',3'-cyclic phosphate in methanopterin biosynthesis. *Biochemistry*, 48:9384–9392, 2009.
- [1833] D. Mason, G.V. Mallo, M.R. Terebiznik, B. Payrastra, B.B. Finlay, J.H. Brumell, L. Rameh, and S. Grinstein. Alteration of epithelial structure and function associated with PtdIns(4,5)P<sub>2</sub> degradation by a bacterial phosphatase. *J. Gen. Physiol.*, 129:267–283, 2007.
- [1834] R. Matalon, J.A. Cifonelli, and A. Dorfman. L-Iduronidase in cultured human fibroblasts and liver. *Biochem. Biophys. Res. Commun.*, 42:340–345, 1971.
- [1835] Z. Mathew, T.M. Knox, and C.G. Miller. *Salmonella enterica* serovar typhimurium peptidase B is a leucyl aminopeptidase with specificity for acidic amino acids. *J. Bacteriol.*, 182:3383–3393, 2000.
- [1836] R. Matsas, I.S. Fulcher, A.J. Kenny, and A.J. Turner. Substance P and [Leu]enkephalin are hydrolyzed by an enzyme in pig caudate synaptic membranes that is identical with the endopeptidase of kidney microvilli. *Proc. Natl Acad. Sci. USA*, 80:3111–3115, 1983.

- [1837] Y. Matsuda. Gametolysin. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 1140–1143. Academic Press, London, 1998.
- [1838] E. Matsumura, T. Shin, S. Murao, M. Sakaguchi, and T. Kawano. A novel enzyme,  $N^\alpha$ -benzyloxycarbonyl amino acid urethane hydrolase IV. *Agric. Biol. Chem.*, 49:3643–3645, 1985.
- [1839] M. Matsushita and T. Fujita. Activation of the classical complement pathway by mannose-binding protein in association with a novel C1s-like serine protease. *J. Exp. Med.*, 176:1497–1502, 1992.
- [1840] S. Matsushita and I.D. Raacke. Purification of nucleoside triphosphatases from pea seedling ribosomes. *Biochim. Biophys. Acta*, 166:707–710, 1968.
- [1841] T. Matsuyama and K. Tanaka. On the enzyme of *Aspergillus fumigatus* producing difructose anhydride I from inulobiose. *Agric. Biol. Chem.*, 53:831–832, 1989.
- [1842] T. Matsuyama, K. Tanaka, M. Mashiko, and M. Kanamoto. Enzymic formation of di-D-fructose 1,2'; 2,1' dianhydride from inulobiose by *Aspergillus fumigatus*. *J. Biochem. (Tokyo)*, 92:1325–1328, 1982.
- [1843] H. Matsuzawa, K. Tokugawa, M. Hamaoki, M. Mizoguchi, H. Taguchi, I. Terada, S.T. Kwon, and T. Ohta. Purification and characterization of aqualysin I (a thermophilic alkaline serine protease) produced by *Thermus aquaticus* YT-1. *Eur. J. Biochem.*, 171:441–447, 1988.
- [1844] H. Mattenheimer. Die Substratspezifität "anorganischer" Poly- und Metaphosphatase. I. Optimale Wirkungsbedingungen für den enzymatischen Abbau von Poly- und Metaphosphaten. *Hoppe-Seyler's Z. Physiol. Chem.*, 303:107–114, 1956.
- [1845] E. Mattern-Dogru, X. Ma, J. Hartmann, H. Decker, and J. Stöckigt. Potential active-site residues in polyneuridine aldehyde esterase, a central enzyme of indole alkaloid biosynthesis, by modelling and site-directed mutagenesis. *Eur. J. Biochem.*, 269:2889–2896, 2002.
- [1846] B. Matthews. W. Structural basis of the action of thermolysin and related zinc peptidases. *Acc. Chem. Res.*, 21:333–340, 1988.
- [1847] M.L. Matthews, G. Periyannan, C. Hajdin, T.K. Sidgel, B. Bennett, and M.W. Crowder. Probing the reaction mechanism of the D-ala-D-ala dipeptidase, VanX, by using stopped-flow kinetic and rapid-freeze quench EPR studies on the Co(II)-substituted enzyme. *J. Am. Chem. Soc.*, 128:13050–13051, 2006.
- [1848] J. Mauger, T. Nagasawa, and H. Yamada. Occurrence of a novel nitrilase, arylacetone nitrilase, in *Alcaligenes faecalis* JM3. *Arch. Microbiol.*, 155:1–6, 1990.
- [1849] W.P. Maughan and S. Shuman. Characterization of 3'-phosphate RNA ligase paralogs RtcB1, RtcB2, and RtcB3 from *Myxococcus xanthus* highlights DNA and RNA 5'-phosphate capping activity of RtcB3. *J. Bacteriol.*, 197:3616–3624, 2015.
- [1850] M.R. Maurizi, W.P. Clark, Y. Katayama, S. Rudikoff, J. Pumphrey, B. Bowers, and S. Gottesman. Sequence and structure of Clp P, the proteolytic component of the ATP-dependent Clp protease of *Escherichia coli*. *J. Biol. Chem.*, 265:12536–12545, 1990.
- [1851] M.R. Maurizi, M.W. Thompson, S.K. Singh, and S.-H. Kim. Endopeptidase Clp: the ATP-dependent Clp protease from *Escherichia coli*. *Methods Enzymol.*, 244:314–331, 1994.
- [1852] D.V. Mavrodi, R.F. Bonsall, S.M. Delaney, M.J. Soule, G. Phillips, and L.S. Thomashow. Functional analysis of genes for biosynthesis of pyocyanin and phenazine-1-carboxamide from *Pseudomonas aeruginosa* PAO1. *J. Bacteriol.*, 183:6454–6465, 2001.
- [1853] M. Mayer, S. Schiffer, and A. Marchfelder. tRNA 3' processing in plants: nuclear and mitochondrial activities differ. *Biochemistry*, 39:2096–2105, 2000.
- [1854] J.T. Maynes, C. Garen, M.M. Cherney, G. Newton, D. Arad, Y. Av-Gay, R.C. Fahey, and M.N. James. The crystal structure of 1-D-myo-inositol 2-acetamido-2-deoxy- $\alpha$ -D-glucopyranoside deacetylase (MshB) from *Mycobacterium tuberculosis* reveals a zinc hydrolase with a lactate dehydrogenase fold. *J. Biol. Chem.*, 278:47166–47170, 2003.

- [1855] D. Mazel, S. Pochet, and P. Marliere. Genetic characterization of polypeptide deformylase, a distinctive enzyme of eubacterial translation. *EMBO J.*, 13:914–923, 1994.
- [1856] M. Mazelis and R.K. Creveling. An adenosine hydrolase from brussels sprouts. *J. Biol. Chem.*, 238:3358–3361, 1963.
- [1857] M. Mazelis, B. Levin, and N. Mallinson. Decomposition of methyl methionine sulfonium salts by a bacterial enzyme. *Biochim. Biophys. Acta*, 105:106–114, 1965.
- [1858] D. Mazumder-Shivakumar and T.C. Bruice. Computational study of IAG-nucleoside hydrolase: determination of the preferred ground state conformation and the role of active site residues. *Biochemistry*, 44:7805–7817, 2005.
- [1859] L. Mazzaferro, L. Pi nuel, M. Minig, and J.D. Breccia. Extracellular monoenzyme deglycosylation system of 7-O-linked flavonoid  $\beta$ -rutinosides and its disaccharide transglycosylation activity from *Stilbella fimetaria*. *Arch. Microbiol.*, 192:383–393, 2010.
- [1860] L. Mazzaferro, L. Pi nuel, M. Minig, and J.D. Breccia. Erratum to: Extracellular monoenzyme deglycosylation system of 7-O-linked flavonoid  $\beta$ -rutinosides and its disaccharide transglycosylation activity from *Stilbella fimetaria*. *Arch. Microbiol.*, 193:461–461, 2011.
- [1861] D.G. McCafferty, I.A. Lessard, and C.T. Walsh. Mutational analysis of potential zinc-binding residues in the active site of the enterococcal D-Ala-D-Ala dipeptidase VanX. *Biochemistry*, 36:10498–10505, 1997.
- [1862] J.D. McCarter, D. Stephens, K. Shoemaker, S. Rosenberg, J.F. Kirsch, and G. Georgiou. Substrate specificity of the *Escherichia coli* outer membrane protease OmpT. *J. Bacteriol.*, 186:5919–5925, 2004.
- [1863] S.J. McClure and P.J. Robinson. Dynamin, endocytosis and intracellular signalling. *Mol. Membr. Biol.*, 13:189–215, 1996.
- [1864] R.M. McCready and C.G. Seegmiller. Action of pectic enzymes on oligogalacturonic acids and some of their derivatives. *Arch. Biochem. Biophys.*, 50:440–450, 1954.
- [1865] J.L. McCulloch, B.A. Chabner, and J.R. Bertino. Purification and properties of carboxypeptidase G<sub>1</sub>. *J. Biol. Chem.*, 246:7207–7213, 1971.
- [1866] J.K. McDonald. Dipeptidyl-peptidase III. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 536–538. Handbook of Proteolytic Enzymes, London, 1998.
- [1867] J.K. McDonald and S. Ellis. On the substrate specificity of cathepsins B1 and B2 including a new fluorogenic substrate for cathepsin B1. *Life Sci.*, 17:1269–1276, 1975.
- [1868] J.K. McDonald, T.J. Reilly, B.B. Zeitman, and S. Ellis. Dipeptidyl arylamidase II of the pituitary. Properties of lysyl-alanyl- $\beta$ -naphthylamide hydrolysis: inhibition by cations, distribution in tissues and subcellular localization. *J. Biol. Chem.*, 243:4143–4150, 1968.
- [1869] J.K. McDonald and C. Schwabe. Intracellular exopeptidases. In A.J. Barrett, editor, *Proteinases in Mammalian Cells and Tissues*, pages 311–391. North-Holland Publishing Co., Amsterdam, 1977.
- [1870] J.K. McDonald, B.B. Zeitman, T.J. Reilly, and S. Ellis. New observations on the substrate specificity of cathepsin C (dipeptidyl aminopeptidase I) including the degradation of  $\beta$ -corticotropin and other peptide hormones. *J. Biol. Chem.*, 244:2693–2709, 1969.
- [1871] J.W. McGrath, G.B. Wisdom, G. McMullan, M.J. Larkin, , and J.P. The purification and properties of phosphonoacetate hydrolase, a novel carbon-phosphorus bond-cleavage enzyme from *Pseudomonas fluorescens* 23F. *Eur. J. Biochem.*, 234:225–230, 1995.
- [1872] M.E. McGrath, J.L. Klaus, M.G. Barnes, and D. Brömme. Crystal structure of human cathepsin K complexed with a potent inhibitor. *Nature Struct. Biol.*, 4:105–109, 1997.
- [1873] J.J. McGuire and J.K. Coward. Pteroylpolyglutamates: biosynthesis, degradation and function. In R.L. Blakley and S.J. Benkovic, editors, *Folates and Pterins*, pages 135–191. John Wiley and Sons, New York, 1984.

- [1874] B.A. McMullen, K. Fujikawa, W. Kisiel, T. Sasagawa, W.N. Howald, E.Y. Kwa, and B. Weinstein. Complete amino acid sequence of the light chain of human blood coagulation factor X: evidence for identification of residue 63 as  $\beta$ -hydroxyaspartic acid. *Biochemistry*, 22:2875–2884, 1983.
- [1875] G.A. McQuibban, S. Saurya, and M. Freeman. Mitochondrial membrane remodelling regulated by a conserved rhomboid protease. *Nature*, 423:537–541, 2003.
- [1876] B.J. McRae, K. Kurachi, R.L. Heimark, K. Fujikawa, E.W. Davie, and J.C. Powers. Mapping the active sites of bovine thrombin, factor IXa, factor Xa, factor XIa, factor XIIa, plasma kallikrein, and trypsin with amino acid and peptide thioesters: development of new sensitive substrates. *Biochemistry*, 20:7196–7206, 1981.
- [1877] E. Mehl and H. Jatzkewitz. A cerebrosidesulfatase from swine kidney. *Hoppe-Seyler's Z. Physiol. Chem.*, 339:260–276, 1964.
- [1878] A.H. Mehler and W.E. Knox. The conversion of tryptophan to kynurenine in liver. II. The enzymatic hydrolysis of formylkynurenine. *J. Biol. Chem.*, 187:431–438, 1950.
- [1879] P. Meighan and R. Pirzad. Mite endopeptidase 1. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 1187–1189. Elsevier, London, 2nd edition, 2004.
- [1880] A. Meister and J.P. Greenstein. Enzymatic hydrolysis of 2,4-diketo acids. *J. Biol. Chem.*, 175:573–588, 1948.
- [1881] A. Meister, L. Levintow, R.E. Greenfield, and P.A. Abendschein. Hydrolysis and transfer reactions catalyzed by  $\omega$ -amidase preparations. *J. Biol. Chem.*, 215:441–460, 1955.
- [1882] A. Meister, A.N. Radhakrishnan, and S.D. Buckley. Enzymatic synthesis of L-pipecolic acid and L-proline. *J. Biol. Chem.*, 229:789–800, 1957.
- [1883] B. Meloun, M. Baudyš, V. Kostka, G. Hausdorf, C. Frömmel, and W.E. Höhne. Complete primary structure of thermolysin from *Thermoactinomyces vulgaris* and its structural features related to the subtilisin-type proteinases. *FEBS Lett.*, 183:195–200, 1985.
- [1884] R. Ménard and A.C. Storer. Papain. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 555–557. Academic Press, London, 1998.
- [1885] V. Mendes, A. Maranha, S. Alarico, M.S. da Costa, and N. Empadinhas. *Mycobacterium tuberculosis* Rv2419c, the missing glucosyl-3-phosphoglycerate phosphatase for the second step in methylglucose lipopolysaccharide biosynthesis. *Sci Rep*, 1:177–177, 2011.
- [1886] J. Mendicino and J.A. Muntz. The activating effect of adenosine triphosphate on brain adenylic deaminase. *J. Biol. Chem.*, 233:178–183, 1958.
- [1887] R. Mentlein, S. Heiland, and E. Heymann. Simultaneous purification and comparative characterization of six serine hydrolases from rat liver microsomes. *Arch. Biochem. Biophys.*, 200:547–559, 1980.
- [1888] R. Mentlein, G. Reuter, and E. Heymann. Specificity of two different purified acylcarnitine hydrolases from rat liver, their identity with other carboxylesterases, and their possible function. *Arch. Biochem. Biophys.*, 240:801–810, 1985.
- [1889] R. Mentlein, M. Schumann, and E. Heymann. Comparative chemical and immunological characterization of five lipolytic enzymes (carboxylesterases) from rat liver microsomes. *Arch. Biochem. Biophys.*, 234:612–621, 1984.
- [1890] J.R. Merkel and J.H. Dreisbach. Purification and characterization of a marine bacterial collagenase. *Biochemistry*, 17:2857–2863, 1978.
- [1891] R.M. Metrione, A.G. Neves, and J.S. Fruton. Purification and properties of dipeptidyl transferase (cathepsin C). *Biochemistry*, 5:1597–1604, 1966.
- [1892] G. Metz and K.-H. Rohm. Yeast aminopeptidase I. Chemical composition and catalytic properties. *Biochim. Biophys. Acta*, 429:933–949, 1976.
- [1893] R. Metz, S. Henning, and W.P. Hammes. LD-Carboxypeptidase activity in *Escherichia coli*. II. Isolation, purification and characterization of the enzyme from *E. coli* K 12. *Arch. Microbiol.*, 144:181–186, 1986.

- [1894] F. Meussdoerffer, P. Tortora, and H. Holzer. Purification and properties of proteinase A from yeast. *J. Biol. Chem.*, 255:12087–12093, 1980.
- [1895] J.J.M. Meyer, N. Grobbelaar, R. Vlegaar, and A.I. Louw. Fluoroacetyl-coenzyme-A hydrolase-like activity in *Dichapetalum cymosum*. *J. Plant Physiol.*, 139:369–372, 1992.
- [1896] K. Meyer, P. Hoffman, and A. Linker. Hyaluronidases. In P.D. Boyer, H. Lardy, and K. Myrback, editors, *The Enzymes*, volume 4, pages 447–460. Academic Press, New York, 2nd edition, 1960.
- [1897] E.C. Meyer-Barton, J.R. Klein, M. Imam, and R. Plapp. Cloning and sequence analysis of the X-prolyl-dipeptidyl-aminopeptidase gene (*pepX*) from *Lactobacillus delbrückii* ssp. *lactis* DSM7290. *Appl. Microbiol. Biotechnol.*, 40:82–89, 1993.
- [1898] O. Meyerhof, R. Shatas, and A. Kaplan. Heat of hydrolysis of trimetaphosphate. *Biochim. Biophys. Acta*, 12:121–127, 1953.
- [1899] G. Meyers, C. Rossi, and H.J. Thiel. Calicivirus endopeptidases. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 1380–1382. Elsevier, London, 2nd edition, 2004.
- [1900] H.Q. Miao, E. Navarro, S. Patel, D. Sargent, H. Koo, H. Wan, A. Plata, Q. Zhou, D. Ludwig, P. Bohlen, and P. Kussie. Cloning, expression, and purification of mouse heparanase. *Protein Expr. Purif.*, 26:425–431, 2002.
- [1901] M.L. Michaels, J. Tchou, A.P. Grollman, and J.H. Miller. A repair system for 8-oxo-7,8-dihydrodeoxyguanine. *Biochemistry*, 31:10964–10968, 1992.
- [1902] G. Michel, T. Barbeyron, D. Flament, T. Vernet, B. Kloareg, and O. Dideberg. Expression, purification, crystallization and preliminary x-ray analysis of the  $\kappa$ -carrageenase from *Pseudoalteromonas carrageenovora*. *Acta Crystallogr. D Biol. Crystallogr.*, 55:918–920, 1999.
- [1903] G. Michel, L. Chantalat, E. Duee, T. Barbeyron, B. Henrissat, B. Kloareg, and O. Dideberg. The  $\kappa$ -carrageenase of *P. carrageenovora* features a tunnel-shaped active site: a novel insight in the evolution of Clan-B glycoside hydrolases. *Structure*, 9:513–525, 2001.
- [1904] G. Michel, L. Chantalat, E. Fanchon, B. Henrissat, B. Kloareg, and O. Dideberg. The  $\iota$ -carrageenase of *Alteromonas fortis*. A  $\beta$ -helix fold-containing enzyme for the degradation of a highly polyanionic polysaccharide. *J. Biol. Chem.*, 276:40202–40209, 2001.
- [1905] G. Michel, W. Helbert, R. Kahn, O. Dideberg, and B. Kloareg. The structural bases of the processive degradation of  $\iota$ -carrageenan, a main cell wall polysaccharide of red algae. *J. Mol. Biol.*, 334:421–433, 2003.
- [1906] E. Miclet, V. Stoven, P.A. Michels, F.R. Opperdoes, J.-Y. Lallemand, and F. Duffieux. NMR spectroscopic analysis of the first two steps of the pentose-phosphate pathway elucidates the role of 6-phosphogluconolactonase. *J. Biol. Chem.*, 276:34840–34846, 2001.
- [1907] C.L. Middleton, J.L. Parker, D.J. Richard, M.F. White, and C.S. Bond. Crystallization and preliminary X-ray diffraction studies of Hje, a Holliday junction resolving enzyme from *Sulfolobus solfataricus*. *Acta Crystallogr. D Biol. Crystallogr.*, 59:171–173, 2003.
- [1908] M. Mihoub, J. Abdallah, B. Gontero, J. Dairou, and G. Richarme. The DJ-1 superfamily member Hsp31 repairs proteins from glycation by methylglyoxal and glyoxal. *Biochem. Biophys. Res. Commun.*, 463:1305–1310, 2015.
- [1909] A.J. Mikulski and M. Laskowski. , Sr. Mung bean nuclease I. 3. Purification procedure and (3')  $\omega$  monophosphatase activity. *J. Biol. Chem.*, 245:5026–5031, 1970.
- [1910] L.C. Miletti, C. Marino, K. Marino, R.M. de Lederkremer, W. Colli, and M.J.M. Alves. Immobilized 4-aminophenyl-1-thio- $\beta$ -D-galactofuranoside as a matrix for affinity purification of an exo- $\beta$ -D-galactofuranosidase. *Carbohydr. Res.*, 320:176–182, 1999.
- [1911] P.J. Mill and R. Tuttobello. The pectic enzymes of *Aspergillus niger*. 2. Endopolygalacturonase. *Biochem. J.*, 79:57–64, 1961.

- [1912] D. Miller, K. O'Brien, H. Xu, and R.H. White. Identification of a 5'-deoxyadenosine deaminase in *Methanocaldococcus jannaschii* and its possible role in recycling the radical S-adenosylmethionine enzyme reaction product 5'-deoxyadenosine. *J. Bacteriol.*, 196:1064–1072, 2014.
- [1913] J.D. Miller, H. Wilhelm, L. Gierasch, R. Gilmore, and P. Walter. GTP binding and hydrolysis by the signal recognition particle during initiation of protein translocation. *Nature*, 366:351–354, 1993.
- [1914] J.J. Miller, D.G. Changaris, and R.S. Levy. Purification, subunit structure and inhibitor profile of cathepsin-A. *J. Chromatogr.*, 627:153–162, 1992.
- [1915] K.D. Miller. Horse prothrombin. *Methods Enzymol.*, 19:140–145, 1970.
- [1916] K.D. Miller and W.H. Copeland. A blood trans- $\alpha$ -glucosylase. *Biochim. Biophys. Acta*, 22:193–194, 1956.
- [1917] J. Millet and R. Acher. Spécificité de la mégatéropeptidase: une amino-endopeptidase à caractère hydrophobe. *Eur. J. Biochem.*, 9:456–462, 1969.
- [1918] G.B. Mills. A biochemical study of *Pseudomonas prunicola* Wormald. I. Pectin esterase. *Biochem. J.*, 44:302–305, 1949.
- [1919] S. Milstien and P. Goldman. Metabolism of guanidinosuccinic acid. I. Characterization of a specific amidino hydrolase from *Pseudomonas chlororaphis*. *J. Biol. Chem.*, 247:6280–6283, 1972.
- [1920] M. Mimura, R. Zallot, T.D. Niehaus, G. Hasnain, S.K. Gidda, T.N. Nguyen, E.M. Anderson, R.T. Mullen, G. Brown, A.F. Yakunin, V. de Crecy-Lagard, J.F. Gregory, McCarty 3rd, Hanson D.R., and A.D. *Arabidopsis* TH<sub>2</sub> encodes the orphan enzyme thiamin monophosphate phosphatase. *Plant Cell*, 28:2683–2696, 2016.
- [1921] A. Minagawa, H. Takaku, M. Takagi, and M. Nashimoto. A novel endonucleolytic mechanism to generate the CCA 3' termini of tRNA molecules in *Thermotoga maritima*. *J. Biol. Chem.*, 279:15688–15697, 2004.
- [1922] M. Minami, S. Ohno, H. Kawasaki, O. Rådmark, B. Samuelsson, H. Jörnvall, T. Shimizu, Y. Seyama, and K. Suzuki. Molecular cloning of a cDNA coding for human leukotriene A<sub>4</sub> hydrolase - complete primary structure of an enzyme involved in eicosanoid synthesis. *J. Biol. Chem.*, 262:13873–13876, 1987.
- [1923] T. Minamikawa, N.P. Jayasankar, B.A. Bohm, I.E. Taylor, and G.H. Towers. An inducible hydrolase from *Aspergillus niger*, acting on carbon-carbon bonds, for phlorrhizin and other C-acylated phenols. *Biochem. J.*, 116:889–897, 1970.
- [1924] D. Ming, L. Ruiz-Avila, and R.F. Margolskee. Characterization and solubilization of bitter-responsive receptors that couple to gustducin. *Proc. Natl. Acad. Sci. USA*, 95:8933–8938, 1998.
- [1925] A.C. Minson and E.H. Creaser. Purification of a trifunctional enzyme, catalysing three steps of the histidine pathway, from *Neurospora crassa*. *Biochem. J.*, 114:49–56, 1969.
- [1926] M. Misago, Y.F. Liao, S. Kudo, S. Eto, M.G. Mattei, K.W. Moremen, and M.N. Fukuda. Molecular cloning and expression of cDNAs encoding human  $\alpha$ -mannosidase II and a previously unrecognized  $\alpha$ -mannosidase IIx isozyme. *Proc. Natl Acad. Sci. USA*, 92:11766–11770, 1995.
- [1927] V. Mishra and D.R. Ronning. Crystal structures of the *Helicobacter pylori* MTAN enzyme reveal specific interactions between S-adenosylhomocysteine and the 5'-alkylthio binding subsite. *Biochemistry*, 51:9763–9772, 2012.
- [1928] K. Misra, A.B. Banerjee, S. Ray, and M. Ray. Glyoxalase III from *Escherichia coli*: a single novel enzyme for the conversion of methylglyoxal into D-lactate without reduced glutathione. *Biochem. J.*, 305:999–1003, 1995.
- [1929] M.Y. Mistou and J.C. Gripon. Catalytic properties of the cysteine aminopeptidase PepC, a bacterial bleomycin hydrolase. *Biochim. Biophys. Acta*, 1383:63–70, 1998.
- [1930] Y. Misumi, Y. Hayashi, F. Arakawa, and Y. Ikehara. Molecular cloning and sequence analysis of human dipeptidyl peptidase IV, a serine proteinase on the cell surface. *Biochim. Biophys. Acta*, 1131:333–336, 1992.
- [1931] W.M. Mitchell. Cleavage at arginine residues by clostripain. *Methods Enzymol.*, 47:165–170, 1977.
- [1932] T. Mitsui, L.M. Riddiford, and G. Bellamy. Metabolism of juvenile hormone by the epidermis of the tobacco hornworm (*Manduca sexta*). *Insect Biochem.*, 9:637–643, 1979.

- [1933] Y. Mitsuishi, M. Kobayashi, and K. Matsuda. Dextran  $\alpha$ -1,2-debranching enzyme from *Flavobacterium* sp. M-73: its production and purification. *Agric. Biol. Chem.*, 43:2283–2290, 1979.
- [1934] Y. Mitsuishi, M. Kobayashi, and K. Matsuda. Dextran  $\alpha$ -(1 $\rightarrow$ 2)-debranching enzyme from *Flavobacterium* sp. M-73. Properties and mode of action. *Carbohydr. Res.*, 83:303–313, 1980.
- [1935] M. Miwa and T. Sugimura. Splitting of the ribose-ribose linkage of poly(adenosine diphosphate-ribose) by a calf thymus extract. *J. Biol. Chem.*, 246:6362–6364, 1971.
- [1936] E. Miyagawa, H. Takahiro, and M. Yoshinobu. Purification and properties of *N*-benzoyl-L-alanine amidohydrolase from *Corynebacterium equii*. *Agric. Biol. Chem.*, 50:1527–1531, 1986.
- [1937] M. Miyake, T. Innami, and Y. Kakimoto. A  $\beta$ -citryl-L-glutamate-hydrolysing enzyme in rat testes. *Biochim. Biophys. Acta*, 760:206–214, 1983.
- [1938] K. Miyazaki, Y. Hattori, F. Umenishi, H. Yasumitsu, and M. Umeda. Purification and characterization of extracellular matrix-degrading metalloproteinase, matrin (pump-1), secreted from human rectal carcinoma cell line. *Cancer Res.*, 50:7758–7764, 1990.
- [1939] S. Miyazawa, S. Furuta, and T. Hashimoto. Induction of a novel long-chain acyl-CoA hydrolase in rat liver by administration of peroxisome proliferators. *Eur. J. Biochem.*, 117:425–430, 1981.
- [1940] K. Mizuno, T. Nakamura, T. Ohshima, S. Tanaka, and H. Matsuo. Yeast KEX2 gene encodes an endopeptidase homologous to subtilisin-like serine proteases. *Biochem. Biophys. Res. Commun.*, 156:246–254, 1988.
- [1941] K. Mizuno, T. Nakamura, T. Ohshima, S. Tanaka, and H. Matsuo. Characterization of KEX2-encoded endopeptidase from yeast *Saccharomyces cerevisiae*. *Biochem. Biophys. Res. Commun.*, 159:305–311, 1989.
- [1942] K. Mizusawa and F. Yoshida. Thermophilic *Streptomyces* alkaline proteinase. *J. Biol. Chem.*, 247:6978–6984, 1972.
- [1943] I. Mochalkin, J.D. Knafels, and S. Lightle. Crystal structure of LpxC from *Pseudomonas aeruginosa* complexed with the potent BB-78485 inhibitor. *Protein Sci.*, 17:450–457, 2008.
- [1944] C.M. Moehle, R. Tizard, S.K. Lemmon, J. Smart, and E.W. Jones. Protease B of the lysosome like vacuole of the yeast *Saccharomyces cerevisiae* is homologous to the subtilisin family of serine proteases. *Mol. Cell. Biol.*, 7:4390–4399, 1987.
- [1945] S. Mogelson and L.G. Lange. Nonoxidative ethanol metabolism in rabbit myocardium: purification to homogeneity of fatty acyl ethyl ester synthase. *Biochemistry*, 23:4075–4081, 1984.
- [1946] G. Moiseyev, Y. Chen, Y. Takahashi, B.X. Wu, and J.X. Ma. RPE65 is the isomerohydrolase in the retinoid visual cycle. *Proc. Natl. Acad. Sci. USA*, 102:12413–12418, 2005.
- [1947] A. Mokhtari, V.S. Blancato, G.D. Repizo, C. Henry, A. Píkis, A. Bourand, M. de Fatima Alvarez, S. Immel, A. Mechakra-Maza, A. Hartke, J. Thompson, C. Magni, and J. Deutscher. *Enterococcus faecalis* utilizes maltose by connecting two incompatible metabolic routes via a novel maltose 6'-phosphate phosphatase (MapP). *Mol. Microbiol.*, 88:234–253, 2013.
- [1948] S.K. Mokkaapati, A.R. Fernandez de Henestrosa, and A.S. Bhagwat. *Escherichia coli* DNA glycosylase Mug: a growth-regulated enzyme required for mutation avoidance in stationary-phase cells. *Mol. Microbiol.*, 41:1101–1111, 2001.
- [1949] L.C. Mokrash and R.N. McGilvery. Purification and properties of fructose-1,6-diphosphatase. *J. Biol. Chem.*, 221:909–917, 1956.
- [1950] A. Molgaard, S. Kauppinen, and S. Larsen. Rhamnogalacturonan acetylerase elucidates the structure and function of a new family of hydrolases. *Structure*, 8:373–383, 2000.
- [1951] J.S. Moncrief, R. Obiso, Barroso Jr., Kling L.A., Wright J.J., Van Tassel R.L., Lysterly R.L., Wilkins D.M., and T.D. The enterotoxin of *Bacteroides fragilis* is a metalloprotease. *Infect. Immun.*, 63:175–181, 1995.
- [1952] V. Monnet, J.P. Ley, and S. Gonzalez. Substrate specificity of the cell envelope-located proteinase of *Lactococcus lactis* subsp. *lactis* NCDO763. *Int. J. Biochem.*, 24:707–718, 1992.

- [1953] J. Monod and M. Cohn. La biosynthèse induite des enzymes (adaptation enzymatique). *Adv. Enzymol. Relat. Subj. Biochem.*, 13:67–119, 1952.
- [1954] C. Montecucco and G. Schiavo. Mechanism of action of tetanus and botulinum neurotoxins. *Mol. Microbiol.*, 8:1–13, 1994.
- [1955] D. Monti, G. Carrea, S. Riva, E. Baldaro, and G. Frare. Characterization of an industrial biocatalyst: immobilized glutaryl-7-ACA acylase. *Biotechnol. Bioeng.*, 70:239–244, 2000.
- [1956] P. Moog and K. Krisch. [The purification and characterization of atropine esterase from rabbit liver microsomes]. *Hoppe-Seyler's Z. Physiol. Chem.*, 355:529–542, 1974.
- [1957] J.H. Moore and D.L. Williams. Some observations on the specificity of phospholipase A. *Biochim. Biophys. Acta*, 84:41–54, 1964.
- [1958] S.J. Moore, A.D. Lawrence, R. Biedendieck, E. Deery, S. Frank, M.J. Howard, S.E. Rigby, and M.J. Warren. Elucidation of the anaerobic pathway for the corrin component of cobalamin (vitamin B<sub>12</sub>). *Proc. Natl Acad. Sci. USA*, 110:14906–14911, 2013.
- [1959] J. Mora, R. Tarrab, J. Martuscelli, and G. Soberón. Characteristics of arginases from ureotelic and non-ureotelic animals. *Biochem. J.*, 96:588–594, 1965.
- [1960] H.M. Mora-Montes, S. Bates, M.G. Netea, D.F. Diaz-Jimenez, E. Lopez-Romero, S. Zinker, P. Ponce-Noyola, B.J. Kullberg, A.J. Brown, F.C. Odds, A. Flores-Carreón, and N.A. Gow. Endoplasmic reticulum  $\alpha$ -glycosidases of *Candida albicans* are required for N glycosylation, cell wall integrity, and normal host-fungus interaction. *Eukaryot Cell*, 6:2184–2193, 2007.
- [1961] M. Moracci, B. Cobucci Ponzano, A. Trincone, S. Fusco, M. De Rosa, J. van Der Oost, C.W. Sensen, R.L. Charlebois, and M. Rossi. Identification and molecular characterization of the first  $\alpha$ -xylosidase from an archaeon. *J. Biol. Chem.*, 275:22082–22089, 2000.
- [1962] A. Morana, A. Esposito, L. Maurelli, G. Ruggiero, E. Ionata, M. Rossi, and F. La Cara. A novel thermoacidophilic cellulase from *Alicyclobacillus acidocaldarius*. *Protein Pept. Lett.*, 15:1017–1021, 2008.
- [1963] R.A. Moreau and A.H.C. Huang. Enzymes of wax ester catabolism in jojoba. *Methods Enzymol.*, 71:804–813, 1981.
- [1964] K.W. Moremen and P.W. Robbins. Isolation, characterization, and expression of cDNAs encoding murine  $\alpha$ -mannosidase II, a Golgi enzyme that controls conversion of high mannose to complex N-glycans. *J. Cell Biol.*, 115:1521–1534, 1991.
- [1965] H. Mori, J.H. Lee, M. Okuyama, M. Nishimoto, M. Ohguchi, D. Kim, A. Kimura, and S. Chiba. Catalytic reaction mechanism based on  $\alpha$ -secondary deuterium isotope effects in hydrolysis of trehalose by European honeybee trehalase. *Biosci. Biotechnol. Biochem.*, 73:2466–2473, 2009.
- [1966] N. Mori, T. Nikai, H. Sugihara, and A.T. Tu. Biochemical characterization of hemorrhagic toxins with fibrinogenase activity isolated from *Crotalus ruber ruber* venom. *Arch. Biochem. Biophys.*, 253:108–121, 1987.
- [1967] M. Moriguchi, K. Sakai, Y. Katsuno, T. Maki, , and M. Purification and characterization of novel N-acyl-D-aspartate amidohydrolase from *Alcaligenes xylosoxydans* subsp. *xylosoxydans* A-6. *Biosci. Biotechnol. Biochem.*, 57:1145–1148, 1993.
- [1968] K. Morihara and T. Oka. Comparative specificity of microbial acid proteinases for synthetic peptides. III. Relationship with their trypsinogen activating ability. *Arch. Biochem. Biophys.*, 157:561–572, 1973.
- [1969] K. Morihara, T. Oka, and H. Tsuzuki. Comparative study of various serine alkaline proteinases from microorganisms. Esterase activity against N-acylated peptide ester substrates. *Arch. Biochem. Biophys.*, 165:72–79, 1974.
- [1970] K. Morihara and H. Tsuzuki. Comparative study of various neutral proteinases from microorganisms: specificity with oligopeptides. *Arch. Biochem. Biophys.*, 146:291–296, 1971.
- [1971] K. Morihara and H. Tsuzuki. *Pseudomonas aeruginosa* elastase: affinity chromatography and some properties as a metallo-neutral proteinase. *Agric. Biol. Chem.*, 39:1123–1128, 1975.



- [1972] K. Morihara and H. Tsuzuki. Specificity of proteinase K from *Tritirachium album* Limber for synthetic peptides. *Agric. Biol. Chem.*, 39:1489–1492, 1975.
- [1973] K. Morihara, H. Tsuzuki, and T. Oka. Comparison of the specificities of various neutral proteinases from microorganisms. *Arch. Biochem. Biophys.*, 123:572–588, 1968.
- [1974] K. Morihara, H. Tsuzuki, and T. Oka. On the specificity of *Pseudomonas aeruginosa* alkaline proteinase with synthetic peptides. *Biochim. Biophys. Acta*, 309:414–429, 1973.
- [1975] N. Morimoto, W. Fukuda, N. Nakajima, T. Masuda, Y. Terui, T. Kanai, T. Oshima, T. Imanaka, and S. Fujiwara. Dual biosynthesis pathway for longer-chain polyamines in the hyperthermophilic archaeon *Thermococcus kodakarensis*. *J. Bacteriol.*, 192:4991–5001, 2010.
- [1976] S. Morimoto, T. Harioka, and Y. Shoyama. Purification and characterization of flavone-specific  $\beta$ -glucuronidase from callus cultures of *Scutellaria baicalensis* Georgi. *Planta*, 195:535–540, 1995.
- [1977] C. Morisseau and B.D. Hammock. Epoxide hydrolases: mechanisms, inhibitor designs, and biological roles. *Annu. Rev. Pharmacol. Toxicol.*, 45:311–333, 2005.
- [1978] A. Moriyama, M. Nakanishi, and M. Sasaki. Porcine muscle prolyl endopeptidase and its endogenous substrates. *J. Biochem. (Tokyo)*, 104:112–117, 1988.
- [1979] M. Mörl and A. Marchfelder. The final cut. The importance of tRNA 3'-processing. *EMBO Rep.*, 2:17–20, 2001.
- [1980] B.J. Morley and R.D. Campbell. Internal homologies of the Ba fragment from human complement component Factor B, as a class III MHC antigen. *EMBO J.*, 3:153–157, 1984.
- [1981] N. Morley and A. Kuksis. Positional specificity of lipoprotein lipase. *J. Biol. Chem.*, 247:6389–6393, 1972.
- [1982] O.V. Moroz, A.G. Murzin, K.S. Makarova, E.V. Koonin, K.S. Wilson, and M.Y. Galperin. Dimeric dUTPases, HisE, and MazG belong to a new superfamily of all- $\alpha$  NTP pyrophosphohydrolases with potential "house-cleaning" functions. *J. Mol. Biol.*, 347:243–255, 2005.
- [1983] F.-M. Mörsberger, R. Müller, M.K. Otto, F. Lingens, and K.D. Kulbe. Purification and characterization of 2-halocarboxylic acid dehalogenase II from *Pseudomonas* spec. CBS 3. *Biol. Chem. Hoppe-Seyler*, 372:915–922, 1991.
- [1984] R.K. Morton. Alkaline phosphatase of milk. 2. Purification of the enzyme. *Biochem. J.*, 55:795–800, 1953.
- [1985] J. Moss, M.K. Jacobson, and S.J. Stanley. Reversibility of arginine-specific mono(ADP-ribosyl)ation: identification in erythrocytes of an ADP-ribose-L-arginine cleavage enzyme. *Proc. Natl. Acad. Sci. USA*, 82:5603–5607, 1985.
- [1986] J. Moss, S.J. Stanley, M.S. Nightingale, J.J. Murtagh, Monaco Jr., Mishima L., Chen K., Williamson H.C., Tsai K.C., and S.C. Molecular and immunological characterization of ADP-ribosylarginine hydrolases. *J. Biol. Chem.*, 267:10481–10488, 1992.
- [1987] K. Motosugi, N. Esahi, and K. Soda. Bacterial assimilation of D- and L-2-chloropropionates and occurrence of a new dehalogenase. *Arch. Microbiol.*, 131:179–183, 1982.
- [1988] K. Motosugi, N. Esahi, and K. Soda. Enzymatic preparation of D- and L-lactic acid from racemic 2-chloropropionic acid. *Biotechnol. Bioeng.*, 26:805–806, 1984.
- [1989] K. Motosugi, N., Soda Esahi, properties of a new enzyme K. Purification, and DL. 2-haloacid dehalogenase, from *Pseudomonas* sp. *J. Bacteriol.*, 150:522–527, 1982.
- [1990] M. Motosugi, N. Esaki, and K. Soda. Preparation and properties of 2-halo acid dehalogenase from *Pseudomonas putida*. *Agric. Biol. Chem.*, 46:837–838, 1982.
- [1991] L.A. Mounter. Enzymic hydrolysis of organophosphorus compounds. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 4, pages 541–550. Academic Press, New York, 2nd edition, 1960.
- [1992] D. Mukhopadhyay and M. Dasso. Modification in reverse: the SUMO proteases. *Trends Biochem. Sci.*, 32:286–295, 2007.

- [1993] A.M. Mulichak, M.J. Theisen, B. Essigmann, C. Benning, and R.M. Garavito. Crystal structure of SQD1, an enzyme involved in the biosynthesis of the plant sulfolipid headgroup donor UDP-sulfoquinovose. *Proc. Natl. Acad. Sci. USA*, 96:13097–13102, 1999.
- [1994] D. Muller, B. Quantin, M.-C. Gesnel, R. Millon-Collard, J. Abecassis, and R. Breathnach. The collagenase gene family in humans consists of at least four members. *Biochem. J.*, 253:187–192, 1988.
- [1995] E.G. Müller. Endoribonuclease IV. A poly(A)-specific ribonuclease from chick oviduct. 1. Purification of the enzyme. *Eur. J. Biochem.*, 70:241–248, 1976.
- [1996] E.G. Müller, G. Seibert, R. Steffen, and R.K. Zahn. Endoribonuclease IV. 2. Further investigation on the specificity. *Eur. J. Biochem.*, 70:249–258, 1976.
- [1997] I.B. Muller, B. Bergmann, M.R. Groves, I. Couto, L. Amaral, T.P. Begley, R.D. Walter, and C. Wrenger. The vitamin B<sub>1</sub> metabolism of *Staphylococcus aureus* is controlled at enzymatic and transcriptional levels. *PLoS One*, 4:e7656–e7656, 2009.
- [1998] R. Müller, J. Thiele, U. Klages, and F. Lingens. Incorporation of [<sup>18</sup>O]water into 4-hydroxybenzoic acid in the reaction of 4-chlorobenzoate dehalogenase from *Pseudomonas spec. CBS 3*. *Biochem. Biophys. Res. Commun.*, 124:178–182, 1984.
- [1999] W. Muller and E. Leistner. 1,4-Naphthoquinone, an intermediate in juglone (5-hydroxy-1,4-naphthoquinone) biosynthesis. *Phytochemistry*, 15:407–410, 1976.
- [2000] H.J. Müller-Eberhard. Molecular organization and function of the complement system. *Annu. Rev. Biochem.*, 57:321–347, 1988.
- [2001] W. Müller-Esterl and H. Fritz. Sperm acrosin. *Methods Enzymol.*, 80:621–632, 1981.
- [2002] R.S. Munford and J.P. Hunter. Acyloxyacyl hydrolase, a leukocyte enzyme that deacylates bacterial lipopolysaccharides, has phospholipase, lysophospholipase, diacylglycerollipase, and acyltransferase activities in vitro. *J. Biol. Chem.*, 267:10116–10121, 1992.
- [2003] K. Murakami-Murofushi, A. Hiratsuka, and J. Ohta. A novel acid protease from haploid amoebae of *Physarum polycephalum*, and its changes during mating and subsequent differentiation into diploid plasmodia. *Cell Struct. Funct.*, 9:311–315, 1984.
- [2004] T. Muraki, M. Taki, Y. Hasegawa, H. Iwaki, and P.C. Lau. Prokaryotic homologs of the eukaryotic 3-hydroxyanthranilate 3,4-dioxygenase and 2-amino-3-carboxymuconate-6-semialdehyde decarboxylase in the 2-nitrobenzoate degradation pathway of *Pseudomonas fluorescens* strain KU-7. *Appl. Environ. Microbiol.*, 69:1564–1572, 2003.
- [2005] S. Murao, E., Kawano Matsumura, characterization of a novel enzyme T. Isolation, and N.  $\alpha$ -benzyloxycarbonyl amino acid urethane hydrolase, from *Streptococcus faecalis* R ATCC 8043. *Agric. Biol. Chem.*, 49:967–972, 1985.
- [2006] S. Murao, S. Funakoshi, and K. Oda. Purification, crystallization and some enzymatic properties of acid protease of *Cladosporium sp. No. 45-2*. *Agric. Biol. Chem.*, 36:1327–1333, 1972.
- [2007] A.S. Murkin, W.K. Chou, W.W. Wakarchuk, and M.E. Tanner. Identification and mechanism of a bacterial hydrolyzing UDP-N-acetylglucosamine 2-epimerase. *Biochemistry*, 43:14290–14298, 2004.
- [2008] T. Muro, T. Kuramoto, K. Imoto, and S. Okada. Purification and some properties of glycyrrhizic acid hydrolase from *Aspergillus niger* GRM3. *Agric. Biol. Chem.*, 50:687–692, 1986.
- [2009] G. Murphy, C.G. McAlpine, C.T. Poll, and J.J. Reynolds. Purification and characterization of a bone metalloproteinase that degrades gelatin and types IV and V collagen. *Biochim. Biophys. Acta*, 831:49–58, 1985.
- [2010] T. Muta, R. Hashimoto, T. Miyata, H. Nishimura, Y. Toh, and S. Iwanaga. Proclotting enzyme from horseshoe crab hemocytes. cDNA cloning, disulfide locations, and subcellular localization. *J. Biol. Chem.*, 265:22426–22433, 1990.
- [2011] T. Muta, T. Miyata, Y. Misumi, F. Tokunaga, T. Nakamura, Y. Toh, Y. Ikehara, and S. Iwanaga. Limulus factor C. An endotoxin-sensitive serine protease zymogen with a mosaic structure of complement-like, epidermal growth factor-like, and lectin-like domains. *J. Biol. Chem.*, 266:6554–6561, 1991.

- [2012] M. Mutter, G. Beldman, S.M. Pitson, H.A. Schols, and A.G. Voragen. Rhamnogalacturonan  $\alpha$ -D-galactopyranosyluronohydrolase. An enzyme that specifically removes the terminal nonreducing galacturonosyl residue in rhamnogalacturonan regions of pectin. *Plant Physiol.*, 117:153–163, 1998.
- [2013] M. Mutter, G. Beldman, H.A. Schols, and A.G. Voragen. Rhamnogalacturonan  $\alpha$ -L-rhamnopyranohydrolase. A novel enzyme specific for the terminal nonreducing rhamnosyl unit in rhamnogalacturonan regions of pectin. *Plant Physiol.*, 106:241–250, 1994.
- [2014] M. Muzio, A.M. Chinnaiyan, F.C. Kischkel, K. O'Rourke, A. Shevchenko, J. Ni, C. Scaffidi, J.D. Bretz, M. Zhang, R. Gentz, M. Mann, P.H. Kramer, M.E. Peter, and V.M. Dixit. FLICE, a novel FADD-homologous ICE/CED-3-like protease, is recruited to the CD95 (Fas/APO-1) death-inducing signaling complex. *Cell*, 85:817–827, 1996.
- [2015] F.L. Myers and D.H. Northcote. Partial purification and some properties of a cellulase from *Helix pomatia*. *Biochem. J.*, 71:749–756, 1959.
- [2016] R.W. Myers, J.W. Wray, S. Fish, and R.H. Abeles. Purification and characterization of an enzyme involved in oxidative carbon-carbon bond cleavage reactions in the methionine salvage pathway of *Klebsiella pneumoniae*. *J. Biol. Chem.*, 268:24785–24791, 1993.
- [2017] K. Myrback. Invertases. In P.D. Boyer, H. Lardy, and K. Myrback, editors, *The Enzymes*, volume 4, pages 379–396. Academic Press, New York, 2nd edition, 1960.
- [2018] K. Myrback and B. Ortenblad. Trehalose und Hefe. II. Trehalosewirkung von Hefepräparaten. *Biochem. Z.*, 291:61–69, 1937.
- [2019] C.E. Argara na, H.S. Barra, and R. Caputto. Tubuliny-tyrosine carboxypeptidase from chicken brain: properties and partial purification. *J. Neurochem.*, 34:114–118, 1980.
- [2020] A. Naamati, N. Regev-Rudzki, S. Galperin, R. Lill, and O. Pines. Dual targeting of Nfs1 and discovery of its novel processing enzyme, Icp55. *J. Biol. Chem.*, 284:30200–30208, 2009.
- [2021] D. Nachmansohn and I.B. Wilson. The enzymic hydrolysis and synthesis of acetylcholine. *Adv. Enzymol. Relat. Subj. Biochem.*, 12:259–339, 1951.
- [2022] Y. Nadai. Arginase. II. Distribution and properties of D-arginase. *J. Biochem. (Tokyo)*, 45:1011–1020, 1958.
- [2023] S. Nagasawa, C. Ichihara, and R.M. Stroud. Cleavage of C4b by C3b inactivator: production of a nicked form from C4b, C4b', as an intermediate cleavage product of C4b by C3b inactivator. *J. Immunol.*, 125:578–582, 1980.
- [2024] T. Nagasawa, J. Mauger, and H. Yamada. A novel nitrilase, arylacetone nitrilase, of *Alcaligenes faecalis* JM3. Purification and characterization. *Eur. J. Biochem.*, 194:765–772, 1990.
- [2025] K. Nagata, N. Yoshida, F. Ogata, M. Araki, and K. Noda. Subsite mapping of an acidic amino acid-specific endopeptidase from *Streptomyces griseus*, GluSGP, and protease V8. *J. Biochem. (Tokyo)*, 110:859–862, 1991.
- [2026] D.K. Nägler and R. Ménard. Human cathepsin X: A novel cysteine protease of the papain family with a very short proregion and unique insertions. *FEBS Lett.*, 434:135–139, 1998.
- [2027] D.K. Nägler, R. Zhang, W. Tam, T. Sulea, E.O. Purisima, and R. Ménard. Human cathepsin X: A cysteine protease with unique carboxypeptidase activity. *Biochemistry*, 38:12648–12654, 1999.
- [2028] D.K. Sulea Nägler, Ménard T., and R. Full length cDNA of human cathepsin F predicts the presence of a cystatin domain at the N-terminus of the cysteine protease zymogen. *Biochem. Biophys. Res. Commun.*, 257:313–318, 1999.
- [2029] K.G. Nair. Purification and properties of 3',5'-cyclic nucleotide phosphodiesterase from dog heart. *Biochemistry*, 5:150–157, 1966.
- [2030] P.P. Nair, M. Gordon, and J. Reback. The enzymatic cleavage of the carbon-nitrogen bond in 3 $\alpha$ ,7 $\alpha$ ,12 $\alpha$ -trihydroxy-5- $\beta$ -cholan-24-oylglycine. *J. Biol. Chem.*, 242:7–11, 1967.
- [2031] T. Nakada, S. Ikegami, H. Chaen, M. Kubota, S. Fukuda, T. Sugimoto, M. Kurimoto, , and Y. Purification and characterization of thermostable maltoooligosyl trehalose trehalohydrolase from the thermoacidophilic archaeobacterium *Sulfolobus acidocaldarius*. *Biosci. Biotechnol. Biochem.*, 60:267–270, 1996.

- [2032] T. Nakada, K. Maruta, H. Mitsuzumi, M. Kubota, H. Chaen, , and T. , Kurimoto M., Tsujisaka, Y. Purification and characterization of a novel enzyme, maltooligosyl trehalose trehalohydrolase, from *Arthrobacter* sp. Q36. *Biosci. Biotechnol. Biochem.*, 59:2215–2218, 1995.
- [2033] T. Nakadai, S. Nasuno, and N. Iguchi. Purification and properties of neutral proteinase II from *Aspergillus oryzae*. *Agric. Biol. Chem.*, 37:2703–2708, 1973.
- [2034] T. Nakagawa, H. Flores-Rozas, and R.D. Kolodner. The MER3 helicase involved in meiotic crossing over is stimulated by single-stranded DNA-binding proteins and unwinds DNA in the 3' to 5' direction. *J. Biol. Chem.*, 276:31487–31493, 2001.
- [2035] Y. Nakagawa. Alkaline proteinases from *Aspergillus*. *Methods Enzymol.*, 19:581–591, 1970.
- [2036] Y.S. Nakagawa, V.G. Eijnsink, K. Totani, and G. Vaaje-Kolstad. Conversion of  $\alpha$ -chitin substrates with varying particle size and crystallinity reveals substrate preferences of the chitinases and lytic polysaccharide monoxygenase of *Serratia marcescens*. *J. Agric. Food Chem.*, 61:11061–11066, 2013.
- [2037] K. Nakahama, K. Yoshimura, R. Marumoto, M. Kikuchi, I.S. Lee, T. Hase, and H. Matsubara. Cloning and sequencing of *Serratia* protease gene. *Nucleic Acids Res.*, 14:5843–5856, 1986.
- [2038] M. Nakajima, K. Mizusawa, and F. Yoshida. Purification and properties of an extracellular proteinase of psychrophilic *Escherichia freundii*. *Eur. J. Biochem.*, 44:87–96, 1974.
- [2039] M. Nakajima, Y. Shirokane, and K. Mizusawa. A new amidinohydrolase, methylguanidine amidinohydrolase from *Alcaligenes* sp. N-42. *FEBS Lett.*, 110:43–46, 1980.
- [2040] T. Nakajima and C.E. Ballou. Structure of the linkage region between the polysaccharide and protein parts of *Saccharomyces cerevisiae* mannan. *J. Biol. Chem.*, 249:7685–7694, 1974.
- [2041] T. Nakajima, S.K. Maitra, and C.E. Ballou. An endo- $\alpha$ -1 $\rightarrow$ 6-D-mannanase from a soil bacterium. Purification, properties, and mode of action. *J. Biol. Chem.*, 251:174–181, 1976.
- [2042] Y. Nakajima, N. Fukunaga, S. Sasaki, and S. Usami. Preparation and properties of NADP pyrophosphatase from *Proteus vulgaris*. *Biochim. Biophys. Acta*, 293:242–255, 1973.
- [2043] T. Nakakuki, K. Azuma, and K. Kainuma. Action patterns of various exo-amylases and the anomeric configurations of their products. *Carbohydr. Res.*, 128:297–310, 1984.
- [2044] Y. Nakamichi, Y. Maruyama, B. Mikami, W. Hashimoto, and K. Murata. Structural determinants in streptococcal unsaturated glucuronyl hydrolase for recognition of glycosaminoglycan sulfate groups. *J. Biol. Chem.*, 286:6262–6271, 2011.
- [2045] N. Nakamura, N. Inoue, R. Watanabe, M. Takahashi, J. Takeda, V.L. Stevens, and T. Kinoshita. Expression cloning of PIG-L, a candidate *N*-acetylglucosaminyl-phosphatidylinositol deacetylase. *J. Biol. Chem.*, 272:15834–15840, 1997.
- [2046] T. Nakamura, T. Horiuchi, T. Morita, and S. Iwanaga. Purification and properties of intracellular clotting factor, factor B, from horseshoe crab (*Tachypleus tridentatus*) hemocytes. *J. Biochem. (Tokyo)*, 99:847–857, 1986.
- [2047] T. Nakamura, S. Meshitsuka, S. Kitagawa, N. Abe, J. Yamada, T. Ishino, H. Nakano, T. Tsuzuki, T. Doi, Y. Kobayashi, S. Fujii, M. Sekiguchi, and Y. Yamagata. Structural and dynamic features of the MutT protein in the recognition of nucleotides with the mutagenic 8-oxoguanine base. *J. Biol. Chem.*, 285:444–452, 2010.
- [2048] T. Nakamura, T. Morita, and S. Iwanaga. Lipopolysaccharide-sensitive serine-protease zymogen (factor C) found in *Limulus* hemocytes. Isolation and characterization. *Eur. J. Biochem.*, 154:511–521, 1986.
- [2049] C. Nakano, T. Ootsuka, K. Takayama, T. Mitsui, T. Sato, and T. Hoshino. Characterization of the Rv3378c gene product, a new diterpene synthase for producing tuberculosinol and (13*R*,*S*)-isotuberculosinol (nosyberkol), from the *Mycobacterium tuberculosis* H37Rv genome. *Biosci. Biotechnol. Biochem.*, 75:75–81, 2011.
- [2050] C. Nakano, M. Oshima, N. Kurashima, and T. Hoshino. Identification of a new diterpene biosynthetic gene cluster that produces *O*-methylkolavelool in *Herpetosiphon aurantiacus*. *Chembiochem*, 16:772–781, 2015.

- [2051] K. Nakayama, H. Honda, Y. Ogawa, T., Ota Ozawa, , and L. carnitinamide hydrolase and method for producing same, 1988.
- [2052] K. Nakayama, W.S. Kim, T. Nakagawa, M. Nagahama, and K. Murakami. Substrate specificity of prorenin converting enzyme of mouse submandibular gland. Analysis using site-directed mutagenesis. *J. Biol. Chem.*, 265:21027–21031, 1990.
- [2053] N. Nakayama, T. Matsubara, T. Ohshiro, Y. Moroto, Y. Kawata, K. Koizumi, Y. Hirakawa, M. Suzuki, K. Maruhashi, Y. Izumi, and R. Kurane. A novel enzyme, 2'-hydroxybiphenyl-2-sulfinate desulfinate (DszB), from a dibenzothiophene-desulfurizing bacterium *Rhodococcus erythropolis* KA2-5-1: gene overexpression and enzyme characterization. *Biochim. Biophys. Acta*, 1598:122–130, 2002.
- [2054] K. Nakazawa and S. Suzuki. Purification of keratan sulfate-endogalactosidase and its action on keratan sulfates of different origin. *J. Biol. Chem.*, 250:912–917, 1975.
- [2055] F. Nanjo, R. Katsumi, and K. Sakai. Purification and characterization of an exo- $\beta$ -D-glucosaminidase, a novel type of enzyme, from *Nocardia orientalis*. *J. Biol. Chem.*, 265:10088–10094, 1990.
- [2056] A. Naqvi, E. Tinsley, and S.A. Khan. Purification and characterization of the PcrA helicase of *Bacillus anthracis*. *J. Bacteriol.*, 185:6633–6639, 2003.
- [2057] N.J. Nardacci, S. Mukhopadhyay, and B.J. Campbell. Partial purification and characterization of the antidiuretic hormone in toad bladder. *Biochim. Biophys. Acta*, 377:146–157, 1975.
- [2058] A.V. Naumov, Y.A. Shabalin, V.M. Vagabov, and I.S. Kulaev. Two pathways of dephosphorylation of dolichyl diphosphate in yeasts. *Biochemistry (Mosc)*, 50:551–556, 1985.
- [2059] J.M. La Nauze, J.R. Coggins, and H.B.F. Dixon. Aldolase-like imine formation in the mechanism of action of phosphonoacetaldehyde hydrolase. *Biochem. J.*, 165:409–411, 1977.
- [2060] J.M. La Nauze and H. Rosenberg. The identification of 2-phosphonoacetaldehyde as an intermediate in the degradation of 2-aminoethylphosphonate by *Bacillus cereus*. *Biochim. Biophys. Acta*, 165:438–447, 1968.
- [2061] J.M. La Nauze, H. Rosenberg, and D.C. Shaw. The enzymic cleavage of the carbon-phosphorus bond: purification and properties of phosphonate. *Biochim. Biophys. Acta*, 212:332–350, 1970.
- [2062] P. Neddermann and J. Jiricny. The purification of a mismatch-specific thymine-DNA glycosylase from HeLa cells. *J. Biol. Chem.*, 268:21218–21224, 1993.
- [2063] P. Nedkov, W. Oberthür, and G. Braunitzer. Determination of the complete amino acid sequence of subtilisin DY and its comparison with the primary structures of the subtilisins BPN, Carlsberg and amylosacchariticus. *Biol. Chem. Hoppe-Seyler*, 366:421–430, 1985.
- [2064] E.J. Neer. Heterotrimeric G proteins: organizers of transmembrane signals. *Cell*, 80:249–259, 1995.
- [2065] M. Negi, R. Tsuboi, T. Matsui, and H. Ogawa. Isolation and characterization of proteinase from *Candida albicans*: substrate specificity. *J. Invest. Dermatol.*, 83:32–36, 1984.
- [2066] O. Negishi, T. Ozawa, and H. Imagawa. N-Methyl nucleosidase from tea leaves. *Agric. Biol. Chem.*, 52:169–175, 1988.
- [2067] S. Negoro, N. Shibata, Y. Tanaka, K. Yasuhira, H. Shibata, H. Hashimoto, Y.H. Lee, S. Oshima, R. Santa, S. Oshima, K. Mochiji, Y. Goto, T. Ikegami, K. Nagai, D. Kato, M. Takeo, and Y. Higuchi. Three-dimensional structure of nylon hydrolase and mechanism of nylon-6 hydrolysis. *J. Biol. Chem.*, 287:5079–5090, 2012.
- [2068] T.E. Nelson, E. Kolb, and J. Larner. Purification and properties of rabbit muscle amylo-1,6-glucosidase-oligo-1,4-1,4-transferase. *Biochemistry*, 8:1419–1428, 1969.
- [2069] Y. Nemerson and M.P. Esnouf. Activation of a proteolytic system by a membrane lipoprotein: mechanism of action of tissue factor. *Proc. Natl. Acad. Sci. USA*, 70:310–314, 1973.
- [2070] F.C. Neuhaus and W.L. Byrne. Metabolism of phosphoserine. II. Purification and properties of O-phosphoserine phosphatase. *J. Biol. Chem.*, 234:113–121, 1959.

- [2071] N.P. Neumann and J.O. Lampen. Purification and properties of yeast invertase. *Biochemistry*, 6:468–475, 1967.
- [2072] D. Newman, N. Abuladze, K. Scholz, W. Dekant, V. Tsuprun, S. Ryazantsev, G. Bondar, P. Sassani, I. Kurtz, and A. Pushkin. Specificity of aminoacylase III-mediated deacetylation of mercapturic acids. *Drug Metab. Dispos.*, 35:43–50, 2007.
- [2073] J.W. Newman, C. Morisseau, and B.D. Hammock. Epoxide hydrolases: their roles and interactions with lipid metabolism. *Prog. Lipid Res.*, 44:1–51, 2005.
- [2074] J.W. Newman, C. Morisseau, T.R. Harris, and B.D. Hammock. The soluble epoxide hydrolase encoded by EPXH<sub>2</sub> is a bifunctional enzyme with novel lipid phosphate phosphatase activity. *Proc. Natl. Acad. Sci. USA*, 100:1558–1563, 2003.
- [2075] G.L. Newton, Y. Av-Gay, and R.C. Fahey. A novel mycothiol-dependent detoxification pathway in mycobacteria involving mycothiol S-conjugate amidase. *Biochemistry*, 39:10739–10746, 2000.
- [2076] G.L. Newton, Y. Av-Gay, and R.C. Fahey. N-Acetyl-1-D-myo-inosityl-2-amino-2-deoxy- $\alpha$ -D-glucopyranoside deacetylase (MshB) is a key enzyme in mycothiol biosynthesis. *J. Bacteriol.*, 182:6958–6963, 2000.
- [2077] S.Y. Ng, D.J. VanDyke, B. Chaban, J. Wu, Y. Nosaka, S. Aizawa, and K.F. Jarrell. Different minimal signal peptide lengths recognized by the archaeal prepilin-like peptidases FlaK and PibD. *J. Bacteriol.*, 191:6732–6740, 2009.
- [2078] P.K. Ngai and M.P. Walsh. Inhibition of smooth muscle actin-activated myosin Mg<sup>2+</sup>-ATPase activity by caldesmon. *J. Biol. Chem.*, 259:13656–13659, 1984.
- [2079] D. Nicholson and N.A. Thornberry. Caspase-3 and caspase-7. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 1298–1302. Elsevier, London, 2nd edition, 2004.
- [2080] R. Nicholson, G. Murphy, and R. Breathnach. Human and rat malignant-tumor-associated mRNAs encode stromelysin-like metalloproteinases. *Biochemistry*, 28:5195–5203, 1989.
- [2081] M.J.H. Nicklin, K.S. Harris, P.V. Pallai, and E. Wimmer. Poliovirus proteinase 3C: large-scale expression, purification, and specific cleavage activity on natural and synthetic substrates in vitro. *J. Virol.*, 62:4586–4593, 1988.
- [2082] K. Niebuhr, S. Giuriato, T. Pedron, D.J. Philpott, F. Gaits, J. Sable, M.P. Sheetz, C. Parsot, P.J. Sansonetti, and B. Payrasstre. Conversion of PtdIns(4,5)P<sub>2</sub> into PtdIns(5)P by the *S. flexneri* effector IpgD reorganizes host cell morphology. *EMBO J.*, 21:5069–5078, 2002.
- [2083] T. Nihira, K. Chiku, E. Suzuki, M. Nishimoto, S. Fushinobu, M. Kitaoka, K. Ohtsubo, and H. Nakai. An inverting  $\beta$ -1,2-mannosidase belonging to glycoside hydrolase family 130 from *Dyadobacter fermentans*. *FEBS Lett.*, 589:3604–3610, 2015.
- [2084] T. Niimura, T. Tokieda, and T. Yamaha. Partial purification and some properties of cyclamate sulfamatase. *J. Biochem. (Tokyo)*, 75:407–417, 1974.
- [2085] T. Nikai, N. Mori, M. Kishida, H. Sugihara, and A.T. Tu. Isolation and biochemical characterization of hemorrhagic toxin f from the venom of *Crotalus atrox* (western diamondback rattlesnake). *Arch. Biochem. Biophys.*, 231:309–319, 1984.
- [2086] O. Nikolaeva, Y. Takahashi, G. Moiseyev, and J.X. Ma. Purified RPE65 shows isomerohydrolase activity after reassociation with a phospholipid membrane. *FEBS J.*, 276:3020–3030, 2009.
- [2087] J. Nikus, A. Esen, and L.M.V. Jonsson. Cloning of a plastidic rye (*Secale cereale*)  $\beta$ -glucosidase cDNA and its expression in *Escherichia coli*. *Physiol. Plantarum*, 118:337–348, 2003.
- [2088] A. Nilsson. The presence of spingomyelin- and ceramide-cleaving enzymes in the small intestinal tract. *Biochim. Biophys. Acta*, 176:339–347, 1969.
- [2089] P. Nilsson-Ehle, P. Belfrage, and B. Borgström. Purified human lipoprotein lipase: positional specificity. *Biochim. Biophys. Acta*, 248:114–120, 1971.
- [2090] R.H. Nimmo-Smith. Aromatic N-deacylation by chick-kidney mitochondria. *Biochem. J.*, 75:284–293, 1960.
- [2091] V. Ninjoor, S.L. Taylor, and A.L. Tappel. Purification and characterization of rat liver lysosomal cathepsin B2. *Biochim. Biophys. Acta*, 370:308–321, 1974.

- [2092] W. Nishii and K. Takahashi. Determination of the cleavage sites in Sula, a cell division inhibitor, by the ATP-dependent HslVU protease from *Escherichia coli*. *FEBS Lett.*, 553:351–354, 2003.
- [2093] W. Nishii, T. Ueki, R. Miyashita, M. Kojima, Y.T. Kim, N. Sasaki, K. Murakami-Murofushi, and K. Takahashi. Structural and enzymatic characterization of physarolisin (formerly physaropepsin) proves that it is a unique serine-carboxyl proteinase. *Biochem. Biophys. Res. Commun.*, 301:1023–1029, 2003.
- [2094] N. Nishino and J.C. Powers. *Pseudomonas aeruginosa* elastase. Development of a new substrate, inhibitors, and an affinity ligand. *J. Biol. Chem.*, 255:3482–3486, 1980.
- [2095] K. Nishitani and D.J. Nevins. Enzymic analysis of feruloylated arabinoxylans (Feraxan) derived from *Zea mays* cell walls. I. Purification of novel enzymes capable of dissociating Feraxan fragments from *Zea mays* coleoptile cell wall. *Plant Physiol.*, 87:883–890, 1988.
- [2096] K. Nishizawa and Y. Hashimoto. Cellulose splitting enzymes. VI. Difference in the specificities of cellulase and  $\beta$ -glucosidase from *Irpex lacteus*. *Arch. Biochem. Biophys.*, 81:211–222, 1959.
- [2097] M. Nishizawa, T. Yasuhara, T. Nakai, Y. Fujiki, and A. Ohashi. Molecular cloning of the aminopeptidase Y gene of *Saccharomyces cerevisiae*. Sequence analysis and gene disruption of a new aminopeptidase. *J. Biol. Chem.*, 269:13651–13655, 1994.
- [2098] Y. Nishizawa, T. Kurihara, and Y. Takahashi. Spectrophotometric assay, solubilization and purification of brain 2':3'-cyclic nucleotide 3'-phosphodiesterase. *Biochem. J.*, 191:71–82, 1980.
- [2099] H. Nitta, T. Imamura, Y. Wada, A. Irie, H. Kobayashi, K. Okamoto, and H. Baba. Production of C5a by ASP, a serine protease released from *Aeromonas sobria*. *J. Immunol.*, 181:3602–3608, 2008.
- [2100] H. Nitta, H. Kobayashi, A. Irie, H. Baba, K. Okamoto, and T. Imamura. Activation of prothrombin by ASP, a serine protease released from *Aeromonas sobria*. *FEBS Lett.*, 581:5935–5939, 2007.
- [2101] D. Njus, J. Knoth, and M. Zallakian. Proton-linked transport in chromaffin granules. *Curr. Top. Bioenerg.*, 11:107–147, 1981.
- [2102] A. Nobre, N. Empadinhas, M.F. Nobre, E.C. Lourenco, C. Maycock, M.R. Ventura, A. Mingote, and M.S. da Costa. The plant *Selaginella moellendorffii* possesses enzymes for synthesis and hydrolysis of the compatible solutes mannosylglycerate and glucosylglycerate. *Planta*, 237:891–901, 2013.
- [2103] J. Nogales, A. Canales, J. Jiménez-Barbero, Pingarrón Serra B., García J. M., Díaz J. L., and E. Unravelling the gallic acid degradation pathway in bacteria: the *gal* cluster from *Pseudomonas putida*. *Mol. Microbiol.*, 79:359–374, 2011.
- [2104] M. Nogawa, H. Takahashi, A. Kashiwagi, K. Ohshima, H. Okada, and Y. Morikawa. Purification and characterization of exo- $\beta$ -D-glucosaminidase from a cellulolytic fungus, *Trichoderma reesei* PC-3-7. *Appl. Environ. Microbiol.*, 64:890–895, 1998.
- [2105] H. Nojiri, H. Taira, K. Iwata, K. Morii, J.W. Nam, T. Yoshida, H. Habe, S. Nakamura, K. Shimizu, H. Yamane, and T. Omori. Purification and characterization of meta-cleavage compound hydrolase from a carbazole degrader *Pseudomonas resinovorans* strain CA10. *Biosci. Biotechnol. Biochem.*, 67:36–45, 2003.
- [2106] C. Nolan, L.S. Hall, and G.H. Barlow. Ancrod, the coagulating enzyme from Malayan pit viper (*Agkistrodon rhodostoma*) venom. *Methods Enzymol.*, 45:205–213, 1976.
- [2107] K. Nomura. Specificity of prolyl endopeptidase. *FEBS Lett.*, 209:235–237, 1986.
- [2108] K. Nomura, H. Tanaka, Y. Kikkawa, M. Yamaguchi, and N. Suzuki. The specificity of sea urchin hatching enzyme (envelysin) places it in the mammalian matrix metalloproteinase family. *Biochemistry*, 30:6115–6123, 1991.
- [2109] M. Nonaka, D. Tsuchimoto, K. Sakumi, and Y. Nakabeppu. Mouse RS21-C<sub>6</sub> is a mammalian 2'-deoxycytidine 5'-triphosphate pyrophosphohydrolase that prefers 5-iodocytosine. *FEBS J.*, 276:1654–1666, 2009.
- [2110] R.C. Nordlie. Glucose-6-phosphatase, hydrolytic and synthetic activities. In P.D. Boyer, editor, *The Enzymes*, volume 4, pages 543–610. Academic Press, New York, 3rd edition, 1971.
- [2111] R.C. Nordlie. Metabolic regulation by multifunctional glucose-6-phosphatase. *Curr. Top. Cell. Regul.*, 8:33–33, 1974.

- [2112] A. Nordwig and H. Mayer. The cleavage of prolyl peptidases by kidney peptidases. *Hoppe-Seyler's Z. Physiol. Chem.*, 354:380–383, 1973.
- [2113] F.A. Norris, R.C. Atkins, and P.W. Majerus. The cDNA cloning and characterization of inositol polyphosphate 4-phosphatase type II. Evidence for conserved alternative splicing in the 4-phosphatase family. *J. Biol. Chem.*, 272:23859–23864, 1997.
- [2114] F.A. Norris, V. Auethavekiat, and P.W. Majerus. The isolation and characterization of cDNA encoding human and rat brain inositol polyphosphate 4-phosphatase. *J. Biol. Chem.*, 270:16128–16133, 1995.
- [2115] M.J. North and A. Whyte. Purification and characterization of two acid proteinases from *Dictyostelium discoideum*. *J. Gen. Microbiol.*, 130:123–134, 1984.
- [2116] J. Norton and J.S. Roth. A ribonuclease specific for 2'-O-methylated ribonucleic acid. *J. Biol. Chem.*, 242:2029–2034, 1967.
- [2117] N.G. Nossal and M. Singer. The processive degradation of individual polyribonucleotide chains. I. *Escherichia coli* ribonuclease II. *J. Biol. Chem.*, 243:913–922, 1968.
- [2118] P. Novak and I.K. Dev. Degradation of a signal peptide by protease IV and oligopeptidase A. *J. Bacteriol.*, 170:5067–5075, 1988.
- [2119] N. Nualkaew, W. De-Eknamkul, T.M. Kutchan, and M.H. Zenk. Membrane-bound geranylgeranyl diphosphate phosphatases: purification and characterization from *Croton stellatopilosus* leaves. *Phytochemistry*, 67:1613–1620, 2006.
- [2120] J. Nunnari, T.D. Fox, and P. Walter. A mitochondrial protease with two catalytic subunits of nonoverlapping specificities. *Science*, 262:1997–2004, 1993.
- [2121] G. Nurmman and D. Strack. Sinapine esterase. 1. Characterization of sinapine esterase from cotyledons of *Raphanus sativus*. *Z. Naturforsch. C: Biosci.*, 34:715–720, 1979.
- [2122] V. Nurmikko, E. Salo, H. Hakola, K. Makinen, and E.E. Snell. The bacterial degradation of pantothenic acid. II. Pantothenate hydrolase. *Biochemistry*, 5:399–402, 1966.
- [2123] E.J. Nyns, D. Zach, and E.E. Snell. The bacterial oxidation of vitamin B<sub>6</sub>. 8. Enzymatic breakdown of  $\alpha$ -(N-acetylamino)methylene succinic acid. *J. Biol. Chem.*, 244:2601–2605, 1969.
- [2124] R.J. Obiso, Lyerly Jr., Van Tassell D.M., Wilkins R.L., and T.D. Proteolytic activity of the *Bacteroides fragilis* enterotoxin causes fluid secretion and intestinal damage in vivo. *Infect. Immun.*, 63:3820–3826, 1995.
- [2125] M. Obst, A. Krug, H. Luftmann, and A. Steinbüchel. Degradation of cyanophycin by *Sedimentibacter hongkongensis* strain KI and *Citrobacter amalonaticus* strain G isolated from an anaerobic bacterial consortium. *Appl. Environ. Microbiol.*, 71:3642–3652, 2005.
- [2126] M. Obst, F.B. Oppermann-Sanio, H. Luftmann, and A. Steinbüchel. Isolation of cyanophycin-degrading bacteria, cloning and characterization of an extracellular cyanophycinase gene (*cphE*) from *Pseudomonas anguilliseptica* strain BI. The *cphE* gene from *P. anguilliseptica* BI encodes a cyanophycin-hydrolyzing enzyme. *J. Biol. Chem.*, 277:25096–25105, 2002.
- [2127] M.A. O'Connell, A. Gerber, and L.P. Keegan. Purification of native and recombinant double-stranded RNA-specific adenosine deaminases. *Methods*, 15:51–62, 1998.
- [2128] B. O'Connor and G. O'Cuinn. Purification of and kinetic studies on a narrow specificity synaptosomal membrane pyroglutamate aminopeptidase from guinea-pig brain. *Eur. J. Biochem.*, 150:47–52, 1985.
- [2129] K. Oda, S. Funakoshi, and S. Murao. Some physicochemical properties and substrate specificity of acid protease isolated from *Cladosporium* sp. No. 45-2. *Agric. Biol. Chem.*, 37:1723–1729, 1973.
- [2130] K. Oda, M. Kamada, and S. Murao. Some physicochemical properties and substrate specificity of acid protease of *Rhodotorula glutinis* K-24. *Agric. Biol. Chem.*, 36:1103–1108, 1972.
- [2131] K. Oda and S. Murao. Purification and some enzymatic properties of acid protease A and B of *Scytalidium lignicolum* ATCC 24568. *Agric. Biol. Chem.*, 38:2435–2444, 1974.



- [2132] K. Oda and S. Murao. Action of *Scytalidium lignicolum* acid proteases on insulin B-chain. *Agric. Biol. Chem.*, 40:1221–1225, 1976.
- [2133] K. Oda, H. Nakatani, and B.M. Dunn. Substrate specificity and kinetic properties of pepstatin-insensitive carboxyl proteinase from *Pseudomonas* sp. No. 101. *Biochim. Biophys. Acta*, 1120:208–214, 1992.
- [2134] K. Oda, T. Nakazima, T. Terashita, K. Suzuki, and S. Murao. Purification and properties of an S-PI(pepstatin Ac)-insensitive carboxyl proteinase from a *Xanthomonas* sp. bacterium. *Agric. Biol. Chem.*, 51:3073–3080, 1987.
- [2135] K. Oda, M. Sugitani, K. Fukuhara, and S. Murao. Purification and properties of a pepstatin-insensitive carboxyl proteinase from a Gram-negative bacterium. *Biochim. Biophys. Acta*, 923:463–469, 1987.
- [2136] K. Oda, H. Torishima, and S. Murao. Purification and characterization of acid proteinase C of *Scytalidium lignicolum* ATCC 24568. *Agric. Biol. Chem.*, 50:651–658, 1986.
- [2137] M.N. Oda, S.V. Scott, A. Hefner-Gravink, A.D. Caffarelli, and D.J. Klionsky. Identification of a cytoplasm to vacuole targeting determinant in aminopeptidase I. *J. Cell Biol.*, 132:999–1010, 1996.
- [2138] S. Odake, C.-M. Kam, L. Narasimhan, M. Poe, J.T. Blake, O. Krahenbuhl, J. Tschopp, and J.C. Powers. Human and murine cytotoxic T lymphocyte serine proteases: subsite mapping with peptide thioester substrates and inhibition of enzyme activity and cytolysis by isocoumarins. *Biochemistry*, 30:2217–2227, 1991.
- [2139] C.E. Ody and E.G. Erdős. Human prolylcarboxypeptidase. *Methods Enzymol.*, 80:460–466, 1981.
- [2140] F. Oesch. Mammalian epoxide hydrolases: inducible enzymes catalysing the inactivation of carcinogenic and cytotoxic metabolites derived from aromatic and olefinic compounds. *Xenobiotica*, 3:305–340, 1973.
- [2141] F. Oesch. Purification and specificity of a human microsomal epoxide hydratase. *Biochem. J.*, 139:77–88, 1974.
- [2142] F. Oesch and J. Daly. Solubilization, purification, and properties of a hepatic epoxide hydrolase. *Biochim. Biophys. Acta*, 227:692–697, 1971.
- [2143] F. Oesch, C.W. Timms, C.H. Walker, T.M. Guenther, A. Sparrow, T. Watabe, and C.R. Wolf. Existence of multiple forms of microsomal epoxide hydrolases with radically different substrate specificities. *Carcinogenesis*, 5:7–9, 1984.
- [2144] M. Ogata-Arakawa, T. Muramatsu, and A. Kobata.  $\alpha$ -L-Fucosidases from almond emulsin: characterization of the two enzymes with different specificities. *Arch. Biochem. Biophys.*, 181:353–358, 1977.
- [2145] J. Ogawa, H. Miyake, and S. Shimizu. Purification and characterization of N-carbamoyl-L-amino acid amidohydrolase with broad substrate specificity from *Alcaligenes xylosoxidans*. *Appl. Microbiol. Biotechnol.*, 43:1039–1043, 1995.
- [2146] J. Ogawa, S. Shimizu, and H. N-Carbamoyl-D-amino acid amidohydrolase from *Comamonas* sp. E222c; purification and characterization. *Eur. J. Biochem.*, 212:685–691, 1993.
- [2147] J. Ogawa, C.L. Soong, M. Honda, and S. Shimizu. Imidase, a new dihydropyrimidinase-like enzyme involved in the metabolism of cyclic imides. *Eur. J. Biochem.*, 243:322–327, 1997.
- [2148] J. Ogawa, S. Takeda, S.X. Xie, H. Hatanaka, T. Ashikari, T. Amachi, and S. Shimizu. Purification, characterization, and gene cloning of purine nucleosidase from *Ochrobactrum anthropi*. *Appl. Environ. Microbiol.*, 67:1783–1787, 2001.
- [2149] K. Ogawa, Y. Ijima, W. Guo, N. Watanabe, T. Usui, S. Dong, Q. Tong, and K. Sakata. Purification of a  $\beta$ -primeverosidase concerned with alcoholic aroma formation in tea leaves (cv. Shuxian) to be processed to oolong tea. *J. Agric. Food Chem.*, 45:877–882, 1997.
- [2150] T. Ogawa, M. Kimoto, and K. Sasaoka. Purification and properties of a new enzyme,  $N^G, N^G$ -dimethylarginine dimethylaminohydrolase, from rat kidney. *J. Biol. Chem.*, 264:10205–10209, 1989.
- [2151] T. Ogawa, K. Noguchi, M. Saito, Y. Nagahata, H. Kato, A. Ohtaki, H. Nakayama, N. Dohmae, Y. Matsushita, M. Odaka, M. Yohda, H. Nyunoya, and Y. Katayama. Carbonyl sulfide hydrolase from *Thiobacillus thioparus* strain TH115 is one of the  $\beta$ -carbonic anhydrase family enzymes. *J. Am. Chem. Soc.*, 135:3818–3825, 2013.
- [2152] E.L. Oginsky and R.F. Gehrig. The arginine dihydrolase system of *Streptococcus faecalis*. II. Properties of arginine dihydrolase. *J. Biol. Chem.*, 198:799–805, 1952.

- [2153] E. Ogris, X. Du, K.C. Nelson, E.K. Mak, X.X. Yu, W.S. Lane, and D.C. Pallas. A protein phosphatase methyltransferase (PME-1) is one of several novel proteins stably associating with two inactive mutants of protein phosphatase 2A. *J. Biol. Chem.*, 274:14382–14391, 1999.
- [2154] P. Oh, D.P. McIntosh, and J.E. Schnitzer. Dynamin at the neck of caveolae mediates their budding to form transport vesicles by GTP-driven fission from the plasma membrane of endothelium. *J. Cell Biol.*, 141:101–114, 1998.
- [2155] S.F. O’Handley, C.A. Dunn, and M.J. Bessman. Orf135 from *Escherichia coli* is a Nudix hydrolase specific for CTP, dCTP, and 5-methyl-dCTP. *J. Biol. Chem.*, 276:5421–5426, 2001.
- [2156] S.F. O’Handley, D.N. Frick, L.C. Bullions, A.S. Mildvan, and M.J. Bessman. *Escherichia coli* orf17 codes for a nucleoside triphosphate pyrophosphohydrolase member of the MutT family of proteins. Cloning, purification, and characterization of the enzyme. *J. Biol. Chem.*, 271:24649–24654, 1996.
- [2157] T. Ohara, K. Makino, H. Shinagawa, A. Nakata, S. Norioka, and F. Sakiyama. Cloning, nucleotide sequence, and expression of *Achromobacter* protease I gene. *J. Biol. Chem.*, 264:20625–2063, 1989.
- [2158] T. Ohhashi, T. Katsu, and M. Ikeda. Improvement of reconstitution of the Cl<sup>-</sup>-translocating ATPase isolated from *Acetabularia acetabulum* into liposomes and several anion pump characteristics. *Biochim. Biophys. Acta*, 1106:165–170, 1992.
- [2159] K. Ohishi, K. Murase, T. Ohta, and H. Etoh. Cloning and sequencing of the deacetylase gene from *Vibrio alginolyticus* H-8. *J. Biosci. Bioeng.*, 90:561–563, 2000.
- [2160] K. Ohishi, M. Yamagishi, T. Ohta, M. Motosugi, H. Izumida, H. Sano, K. Adachi, , and T. Purification and properties of two deacetylases produced by *Vibrio alginolyticus* H-8. *Biosci. Biotechnol. Biochem.*, 61:1113–1117, 1997.
- [2161] N. Ohishi, T. Izumi, M. Minami, S. Kitamura, Y. Seyama, S. Ohkawa, S. Terao, H. Yotsumoto, F. Takaku, and T. Shimizu. Leukotriene A<sub>4</sub> hydrolase in the human lung. Inactivation of the enzyme with leukotriene A<sub>4</sub> isomers. *J. Biol. Chem.*, 262:10200–10205, 1987.
- [2162] J.B. Ohlrogge, W.E. Shine, and P.K. Stumpf. Fat metabolism in higher plants. Characterization of plant acyl-ACP and acyl-CoA hydrolases. *Arch. Biochem. Biophys.*, 189:382–391, 1978.
- [2163] E. Ohmura and O. Hayaishi. Enzymatic conversion of formylaspartic acid to aspartic acid. *J. Biol. Chem.*, 227:181–190, 1957.
- [2164] T. Ohno, M. Tsuchiya, H. Osago, N. Hara, J. Jidoi, and M. Shimoyama. Detection of arginine-ADP-ribosylated protein using recombinant ADP-ribosylarginine hydrolase. *Anal. Biochem.*, 10:115–122, 1995.
- [2165] M. Ohnuma, Y. Terui, M. Tamakoshi, H. Mitome, M. Niitsu, K. Samejima, E. Kawashima, and T. Oshima. N<sup>1</sup>-aminopropylagmatine, a new polyamine produced as a key intermediate in polyamine biosynthesis of an extreme thermophile, *Thermus thermophilus*. *J. Biol. Chem.*, 280:30073–30082, 2005.
- [2166] T. Ohnuma, T. Numata, T. Osawa, M. Mizuhara, O. Lampela, A.H. Juffer, K. Skriver, and T. Fukamizo. A class V chitinase from *Arabidopsis thaliana*: gene responses, enzymatic properties, and crystallographic analysis. *Planta*, 234:123–137, 2011.
- [2167] H. Ohshima, N. Hirano, and H. Takahashi. A hexanucleotide sequence (dC1-dC6 tract) restricts the dC-specific cleavage of single-stranded DNA by endonuclease IV of bacteriophage T<sub>4</sub>. *Nucleic Acids Res.*, 35:6681–6689, 2007.
- [2168] M. Ohsugi, J. Khan, C. Hensley, S. Chew, and H.A. Barker. Metabolism of L-β-lysine by a *Pseudomonas*. Purification and properties of a deacetylase-thiolesterase utilizing 4-acetamidobutyryl CoA and related compounds. *J. Biol. Chem.*, 256:7642–7651, 1981.
- [2169] Y. Ohta and Y. Hatada. A novel enzyme, λ-carrageenase, isolated from a deep-sea bacterium. *J. Biochem. (Tokyo)*, 140:475–481, 2006.
- [2170] Y. Ohta, Y. Hatada, M. Miyazaki, Y. Nogi, S. Ito, and K. Horikoshi. Purification and characterization of a novel α-agarase from a *Thalassomonas* sp. *Curr. Microbiol.*, 50:212–216, 2005.

- [2171] Y. Ohta, Y. Hatada, Y. Nogi, M. Miyazaki, Z. Li, M. Akita, Y. Hidaka, S. Goda, S. Ito, and K. Horikoshi. Enzymatic properties and nucleotide and amino acid sequences of a thermostable  $\beta$ -agarase from a novel species of deep-sea *Microbulbifer*. *Appl. Microbiol. Biotechnol.*, 64:505–514, 2004.
- [2172] Y. Ohta, Y. Nogi, M. Miyazaki, Z. Li, Y. Hatada, S. Ito, and K. Horikoshi. Enzymatic properties and nucleotide and amino acid sequences of a thermostable  $\beta$ -agarase from the novel marine isolate, JAMB-A94. *Biosci. Biotechnol. Biochem.*, 68:1073–1081, 2004.
- [2173] Y. Ogura Ohta, Wada Y., and A. Thermostable protease from thermophilic bacteria. I. Thermostability, physicochemical properties, and amino acid composition. *J. Biol. Chem.*, 241:5919–5925, 1966.
- [2174] K. Ohtsuki, K. Taguchi, K. Sato, and M. Kawabata. Purification of ginger proteases by DEAE-Sepharose and isoelectric focusing. *Biochim. Biophys. Acta*, 1243:181–184, 1995.
- [2175] M. Ohtsuru, J. Tang, and R. Delaney. Purification and characterization of rhizopuspesin isozymes from a liquid culture of *Rhizopus chinensis*. *Int. J. Biochem.*, 14:925–932, 1982.
- [2176] T. Ohya, T. Sawai, S. Uemura, and K. Abe. Some catalytic properties of an exo-1,6- $\alpha$ -glucosidase (glucodextranase) from *Arthrobacter globiformis* I42. *Agric. Biol. Chem.*, 42:571–577, 1978.
- [2177] M. Oishi. An ATP-dependent deoxyribonuclease from *Escherichia coli* with a possible role in genetic recombination. *Proc. Natl. Acad. Sci. USA*, 64:1292–1299, 1969.
- [2178] T. Oka, K. Ishino, H. Tsuzuki, K. Morihara, and K. Arima. On the specificity of a rennin-like enzyme from *Mucor pusillus*. *Agric. Biol. Chem.*, 37:1177–1184, 1973.
- [2179] T. Okada, H. Suzuki, K. Wada, H. Kumagai, and K. Fukuyama. Crystal structures of  $\gamma$ -glutamyltranspeptidase from *Escherichia coli*, a key enzyme in glutathione metabolism, and its reaction intermediate. *Proc. Natl. Acad. Sci. USA*, 103:6471–6476, 2006.
- [2180] T. Okada, H. Suzuki, K. Wada, H. Kumagai, and K. Fukuyama. Crystal structure of the  $\gamma$ -glutamyltranspeptidase precursor protein from *Escherichia coli*. Structural changes upon autocatalytic processing and implications for the maturation mechanism. *J. Biol. Chem.*, 282:2433–2439, 2007.
- [2181] Y. Okada, H. Harris Nagase, , and Jr. A metalloproteinase from human rheumatoid synovial fibroblasts that digests connective tissue matrix components. Purification and characterization. *J. Biol. Chem.*, 261:14245–14255, 1986.
- [2182] Y. Okada, T. Morodomi, J.J. Enghild, K. Suzuki, A. Yasui, I. Nakanishi, G. Salvesen, and H. Nagase. Matrix metalloproteinase 2 from human rheumatoid synovial fibroblasts-purification and activation of the precursor and enzymic properties. *Eur. J. Biochem.*, 194:721–730, 1990.
- [2183] Y. Okada, S. Yamada, M. Toyoshima, J. Dong, M. Nakajima, and K. Sugahara. Structural recognition by recombinant human heparanase that plays critical roles in tumor metastasis. Hierarchical sulfate groups with different effects and the essential target disulfated trisaccharide sequence. *J. Biol. Chem.*, 277:42488–42495, 2002.
- [2184] Y. Okamoto, J. Morishita, K. Tsuboi, T. Tonai, and N. Ueda. Molecular characterization of a phospholipase D generating anandamide and its congeners. *J. Biol. Chem.*, 279:5298–5305, 2004.
- [2185] Y. Okawa and T. Yamaguchi. Studies on sterol-ester hydrolase from *Fusarium oxysporum*. I. Partial purification and properties. *J. Biochem. (Tokyo)*, 81:1209–1215, 1977.
- [2186] F. Okazaki, K. Shiraki, Y. Tamaru, T. Araki, and M. Takagi. The first thermodynamic characterization of  $\beta$ -1,3-xylanase from a marine bacterium. *Protein J.*, 24:413–421, 2005.
- [2187] A. Okazawa, L. Tango, Y. Itoh, E. Fukusaki, and A. Kobayashi. Characterization and subcellular localization of chlorophyllase from *Ginkgo biloba*. *Z. Naturforsch. [C]*, 61:111–117, 2006.
- [2188] K. Okemoto, T. Uekita, Y. Tsumuraya, Y. Hashimoto, and T. Kasama. Purification and characterization of an endo- $\beta$ -(1 $\rightarrow$ 6)-galactanase from *Trichoderma viride*. *Carbohydr. Res.*, 338:219–230, 2003.
- [2189] K. Okuda, K. Morihara, Y. Atsumi, H. Takeuchi, S. Kawamoto, H. Kawasaki, K. Suzuki, and J. Fukushima. Complete nucleotide sequence of the structural gene for alkaline proteinase from *Pseudomonas aeruginosa* IFO 3455. *Infect. Immun.*, 58:4083–4088, 1990.

- [2190] M. Okuyama, A. Kaneko, H. Mori, S. Chiba, and A. Kimura. Structural elements to convert *Escherichia coli*  $\alpha$ -xylosidase (YicI) into  $\alpha$ -glucosidase. *FEBS Lett.*, 580:2707–2711, 2006.
- [2191] R.W. Olafson, G.I. Drummond, and J.F. Lee. Studies on 2',3'-cyclic nucleotide-3'-phosphohydrolase from brain. *Can. J. Biochem.*, 47:961–966, 1969.
- [2192] C. Oldfield, O. Pogrebinsky, J. Simmonds, E.S. Öolson, and C.F. Kulpa. Elucidation of the metabolic pathway for dibenzothioophene desulfurization by *Rhodococcus* sp. strain IGTS8 (ATCC 53968). *Microbiology*, 143:2961–2973, 1997.
- [2193] D.B. Olsen, T.W. Hepburn, M. Moos, P.S. Mariano, and D. Dunaway-Mariano. Substrate binding and catalytic groups of the P-C bond cleaving enzyme, phosphonoacetaldehyde hydrolase. *Biochemistry*, 27:2229–2234, 1988.
- [2194] V. Olsen, K. Guruprasad, N.X. Cawley, H.C. Chen, T.L. Blundell, and Y.P. Loh. Cleavage efficiency of the novel aspartic protease yapsin 1 (Yap3p) enhanced for substrates with arginine residues flanking the P1 site: correlation with electronegative active-site pockets predicted by molecular modeling. *Biochemistry*, 37:2768–2777, 1998.
- [2195] M.O.J. Olson, N. Nagabushan, M. Dzwiniel, L.B. Smillie, and D.R. Whitaker. Primary structure of  $\alpha$ -lytic protease: a bacterial homologue of the pancreatic serine proteases. *Nature*, 228:438–442, 1970.
- [2196] T. Omori, K. Sugimura, H. Ishigooka, and Y. Minoda. Purification and some properties of a 2-hydroxy-6-oxo-6-phenylhexa-2,4-dienoic acid hydrolyzing enzyme from *Pseudomonas cruciviae* S93 B1 involved in the degradation of biphenyl. *Agric. Biol. Chem.*, 50:931–937, 1986.
- [2197] T. Omori-Satoh and S. Sadahiro. Resolution of the major hemorrhagic component of *Trimeresurus flavoviridis* venom into two parts. *Biochim. Biophys. Acta*, 580:392–404, 1979.
- [2198] Y. Ono, F. Torii, K. Ojima, N. Doi, K. Yoshioka, Y. Kawabata, D. Labeit, S. Labeit, K. Suzuki, K. Abe, T. Maeda, and H. Sorimachi. Suppressed disassembly of autolyzing p94/CAPN3 by N2A connectin/titin in a genetic reporter system. *J. Biol. Chem.*, 281:18519–18531, 2006.
- [2199] A.T. Orawski, J.P. Susz, and W.H. Simmons. Aminopeptidase-P from bovine lung - solubilization, properties, potential role in bradykinin degradation. *Mol. Cell. Biochem.*, 75:123–132, 1987.
- [2200] M. Orłowski, S. Reznik, J. Ayala, and A.R. Pierotti. Endopeptidase 24.15 from rat testes. Isolation of the enzyme and its specificity toward synthetic and natural peptides, including enkephalin-containing peptides. *Biochem. J.*, 261:951–958, 1989.
- [2201] L. Orning, J.K. Gierse, and F.A. Fitzpatrick. The bifunctional enzyme leukotriene- $A_4$  hydrolase is an arginine aminopeptidase of high efficiency and specificity. *J. Biol. Chem.*, 269:11269–11267, 1994.
- [2202] L.N. Ornston. The conversion of catechol and protocatechuate to  $\beta$ -keto adipate by *Pseudomonas putida*. II. Enzymes of the protocatechuate pathway. *J. Biol. Chem.*, 241:3787–3794, 1966.
- [2203] L.N. Ornston. Conversion of catechol and protocatechuate to  $\beta$ -keto adipate (*Pseudomonas putida*). *Methods Enzymol.*, 17A:529–549, 1970.
- [2204] G. Orsomando, R.D. de la Garza, B.J. Green, M. Peng, P.A. Rea, T.J. Ryan, J.F. Gregory, Hanson 3rd, and A.D. Plant  $\gamma$ -glutamyl hydrolases and folate polyglutamates: characterization, compartmentation, and co-occurrence in vacuoles. *J. Biol. Chem.*, 280:28877–28884, 2005.
- [2205] E. Ortlund, M.W. Lacount, K. Lewinski, and L. Lebioda. Reactions of *Pseudomonas* 7A glutaminase-asparaginase with diazo analogues of glutamine and asparagine result in unexpected covalent inhibitions and suggests an unusual catalytic triad Thr-Tyr-Glu. *Biochemistry*, 39:1199–1204, 2000.
- [2206] J. Oshiro, G.S. Han, and G.M. Carman. Diacylglycerol pyrophosphate phosphatase in *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta*, 1635:1–9, 2003.
- [2207] T. Østerlund, B. Danielsson, E. Degerman, J.A. Contreras, G. Edgren, R.C. Davis, M.C. Schotz, and C. Holm. Domain-structure analysis of recombinant rat hormone-sensitive lipase. *Biochem. J.*, 319 ( Pt 2):411–420, 1996.
- [2208] A.L. Osterman, V.M. Stepanov, G.N. Rudenskaya, O.M. Khodova, I.A. Tsaplina, M.B. Yakovleva, and L.G. Loginova. Carboxypeptidase T - an extracellular carboxypeptidase of thermophilic actinomycetes - a remote analog of animal carboxypeptidases. *Biochemistry (USSR)*, 49:292–301, 1984.

- [2209] V.I. Ostoslavskaya, E.K. Kotlova, V.M. Stepanov, G.H. Rudenskaya, L.A. Baratova, and L.P. Belyanova. Aspergillopepsin F-A carboxylic proteinase from *Aspergillus foetidus*. *Bioorg. Khim.*, 5:595–603, 1976.
- [2210] V.I. Ostoslavskaya, L.P. Revina, E.K. Kotlova, I.A. Surova, E.D. Levin, E.A. Timokhima, and V.M. Stepanov. The primary structure of aspergillopepsin A, aspartic proteinase from *Aspergillus awamori*. IV. Amino acid sequence of the enzyme. *Bioorg. Khim.*, 12:1030–1047, 1986.
- [2211] S. Ota, E. Muta, Y. Katanita, and Y. Okamoto. Reinvestigation of fractionation and some properties of the proteolytically active components of stem and fruit bromelains. *J. Biochem. (Tokyo)*, 98:219–228, 1985.
- [2212] Y. Otaka, T. Uchida, and T. Sakai. Purification and properties of ribonuclease from yeast. *J. Biochem. (Tokyo)*, 54:322–327, 1963.
- [2213] G.A. O'Toole, J.R. Trzebiatowski, and J.C. Escalante-Semerena. The cobC gene of *Salmonella typhimurium* codes for a novel phosphatase involved in the assembly of the nucleotide loop of cobalamin. *J. Biol. Chem.*, 269:26503–26511, 1994.
- [2214] M. Ottesen and W. Rickert. The acid protease of *Mucor miehei*. *Methods Enzymol.*, 19:459–460, 1970.
- [2215] M. Ottesen and I. Svendsen. The subtilisins. *Methods Enzymol.*, 19:199–215, 1970.
- [2216] K. Otto and H. Riesenköning. Improved purification of cathepsin B1 and cathepsin B2. *Biochim. Biophys. Acta*, 379:462–475, 1975.
- [2217] M. Ouaisi and A. Ouaisi. Histone deacetylase enzymes as potential drug targets in cancer and parasitic diseases. *J. Biomed. Biotechnol.*, 2006:13474–13474, 2006.
- [2218] H. Outtrup and B.E. Norman. Properties and application of a thermostable maltogenic amylase produced by a strain of *Bacillus* modified by recombinant-DNA techniques. *Stärke*, 36:405–411, 1984.
- [2219] A.Z. Ozsoy, J.J. Sekelsky, and S.W. Matson. Biochemical characterization of the small isoform of *Drosophila melanogaster* RECQ5 helicase. *Nucleic Acids Res.*, 29:2986–2993, 2001.
- [2220] H.C. Pace and C. Brenner. The nitrilase superfamily: classification, structure and function. *Genome Biol.*, 2:0001.–001.9, 2001.
- [2221] C.R. Pace-Asciak. Formation and metabolism of hepoxilin A<sub>3</sub> by the rat brain. *Biochem. Biophys. Res. Commun.*, 151:493–498, 1988.
- [2222] C.R. Pace-Asciak and W.-S. Lee. Purification of hepoxilin epoxide hydrolase from rat liver. *J. Biol. Chem.*, 264:9310–9313, 1989.
- [2223] W.K. Paik, L. Bloch-Frankenthal, S.M. Birnbaum, M. Winitz, and J.P. Greenstein. ε-Lysine acylase. *Arch. Biochem. Biophys.*, 69:56–66, 1957.
- [2224] M.J.I. Paine, H.P. Desmond, R.D.G. Theakston, and J.M. Crampton. Purification, cloning, and molecular characterization of a high molecular weight hemorrhagic metalloprotease, jararhagin, from *Bothrops jararaca* venom. Insights into the disintegrin gene family. *J. Biol. Chem.*, 267:22869–22876, 1992.
- [2225] C. Palaniappan, G.M. Fuentes, L. Rodriguez-Rodriguez, P.J. Fay, and R.A. Bambara. Helix structure and ends of RNA/DNA hybrids direct the cleavage specificity of HIV-1 reverse transcriptase RNase H. *J. Biol. Chem.*, 271:2063–2070, 1996.
- [2226] M.J. Pallen and B.W. Wren. The HtrA family of serine proteases. *Mol. Microbiol.*, 26:209–221, 1997.
- [2227] N.J. Palleroni and M. Doudoroff. Metabolism of carbohydrates by *Pseudomonas saccharophilla*. III. Oxidation of D-arabinose. *J. Bacteriol.*, 74:180–185, 1957.
- [2228] G.J. Palm and K.H. Rohm. Aminoacylase I from porcine kidney: identification and characterization of two major protein domains. *J. Protein Chem.*, 14:233–240, 1995.
- [2229] J.L. Palmer and R.H. Abeles. The mechanism of action of S-adenosylhomocysteinase. *J. Biol. Chem.*, 254:1217–1226, 1979.

- [2230] R.E. Palmer and R.L. Anderson. Cellobiose metabolism in *Aerobacter aerogenes*. 3. Cleavage of cellobiose monophosphate by a phospho- $\beta$ -glucosidase. *J. Biol. Chem.*, 247:3420–3423, 1972.
- [2231] M. Panneerselvam and S.C. Dhar. Studies on the peptide bond specificity and the essential groups of an acid proteinase from *Aspergillus fumigatus*. *Ital. J. Biochem.*, 30:207–216, 1981.
- [2232] A.D. Pannifer, T.Y. Wong, R. Schwarzenbacher, M. Renatus, C. Petosa, J. Bienkowska, D.B. Lacy, R.J. Collier, S. Park, S.H. Leppla, P. Hanna, and R.C. Liddington. Crystal structure of the anthrax lethal factor. *Nature*, 414:229–233, 2001.
- [2233] K. Pari, G.A. Mueller, E.F. DeRose, T.W. Kirby, and R.E. London. Solution structure of the RNase H domain of the HIV-1 reverse transcriptase in the presence of magnesium. *Biochemistry*, 42:639–650, 2003.
- [2234] S.S. Parikh, C.D. Putnam, and J.A. Tainer. Lessons learned from structural results on uracil-DNA glycosylase. *Mutat. Res.*, 460:183–199, 2000.
- [2235] S.J. Park and S.Y. Lee. Identification and characterization of a new enoyl coenzyme A hydratase involved in biosynthesis of medium-chain-length polyhydroxyalkanoates in recombinant *Escherichia coli*. *J. Bacteriol.*, 185:5391–5397, 2003.
- [2236] S.Y. Park, S.J. Lee, T.K. Oh, J.W. Oh, B.T. Koo, D.Y. Yum, and J.K. Lee. AhlD, an *N*-acylhomoserine lactonase in *Arthrobacter sp.*, and predicted homologues in other bacteria. *Microbiology*, 149:1541–1550, 2003.
- [2237] D.W. Parkin. Purine-specific nucleoside *N*-ribohydrolase from *Trypanosoma brucei brucei*. Purification, specificity, and kinetic mechanism. *J. Biol. Chem.*, 271:21713–21719, 1996.
- [2238] S. Parrish and B. Moss. Characterization of a second vaccinia virus mRNA-decapping enzyme conserved in poxviruses. *J. Virol.*, 81:12973–12978, 2007.
- [2239] S. Parrish, W. Resch, and B. Moss. Vaccinia virus D10 protein has mRNA decapping activity, providing a mechanism for control of host and viral gene expression. *Proc. Natl. Acad. Sci. USA*, 104:2139–2144, 2007.
- [2240] J.F. Parsons, K. Calabrese, E. Eisenstein, and J.E. Ladner. Structure and mechanism of *Pseudomonas aeruginosa* PhzD, an isochorismatase from the phenazine biosynthetic pathway. *Biochemistry*, 42:5684–5693, 2003.
- [2241] R. Parthasarathy, L. Parthasarathy, and R. Vadnal. Brain inositol monophosphatase identified as a galactose 1-phosphatase. *Brain Res.*, 778:99–106, 1997.
- [2242] R. Parvin and R.A. Smith. Phosphoramidates. V. Probable identity of rat liver microsomal glucose 6-phosphatase, phosphoramidase, and phosphoramidate-hexose phosphotransferase. *Biochemistry*, 8:1748–1748, 1969.
- [2243] M.D. Pato and R.S. Adelstein. Purification and characterization of a multisubunit phosphatase from turkey gizzard smooth muscle. The effect of calmodulin binding to myosin light chain kinase on dephosphorylation. *J. Biol. Chem.*, 258:7047–7054, 1983.
- [2244] M.P. Patricelli and B.F. Cravatt. Characterization and manipulation of the acyl chain selectivity of fatty acid amide hydrolase. *Biochemistry*, 40:6107–6115, 2001.
- [2245] M.P. Patricelli, H.A. Lashuel, D.K. Giang, J.W. Kelly, and B.F. Cravatt. Comparative characterization of a wild type and transmembrane domain-deleted fatty acid amide hydrolase: identification of the transmembrane domain as a site for oligomerization. *Biochemistry*, 37:15177–15187, 1998.
- [2246] T.N. Pattabiraman, , and B.K. Purification of glucosamine 6-phosphate deaminase from human brain. *Biochim. Biophys. Acta*, 54:273–283, 1961.
- [2247] J.M. Patti, A. Schneider, N. Garza, and J.O. Boles. Isolation and characterization of pcp, a gene encoding a pyrrolidone carboxyl peptidase in *Staphylococcus aureus*. *Gene*, 166:95–99, 1995.
- [2248] M. Pauly, L.N. Andersen, S. Kaupinnen, L.V. Kofod, W.S. York, P. Albersheim, and A. Darvill. A xyloglucan specific endo- $\beta$ -1,4-glucanase from *Aspergillus aculeatus*: expression cloning in yeast, purification and characterization of the recombinant enzyme. *Glycobiology*, 9:93–100, 1999.
- [2249] R.A. Pauptit, R. Karlson, D. Picot, J.A. Jenkins, A.-S. Niklaus-Reimer, and J.N. Jansonius. Crystal structure of neutral protease from *Bacillus cereus* refined at 3.0 Å resolution and comparison with the homologous but more thermostable enzyme thermolysin. *J. Mol. Biol.*, 199:525–537, 1988.

- [2250] N. Pavloff, J. Potempa, R.N. Pike, V. Prochazka, M.C. Kiefer, J. Travis, and P.J. Barr. Molecular cloning and structural characterization of the Arg-gingipain proteinase of *Porphyromonas gingivalis*. Biosynthesis as a proteinase-adhesin polyprotein. *J. Biol. Chem.*, 270:1007–1010, 1995.
- [2251] R.J. Peanasky, D. Gratecos, J. Baratti, and M. Rovey. Mode of activation and N-terminal sequence of subunit II in bovine procarboxypeptidase A and of porcine chymotrypsinogen C. *Biochim. Biophys. Acta*, 181:82–92, 1969.
- [2252] D. Pei. Peptide deformylase: a target for novel antibiotics? *Emerging Therapeutic Targets*, 5:23–40, 2001.
- [2253] S. Pelkonen, J. Pelkonen, and J. Finne. Common cleavage pattern of polysialic acid by bacteriophage endosialidases of different properties and origins. *J. Virol.*, 65:4409–4416, 1989.
- [2254] P. Pellerin and J.M. Brillouet. Purification and properties of an exo-(1→3)- $\beta$ -D-galactanase from *Aspergillus niger*. *Carbohydr. Res.*, 264:281–291, 1994.
- [2255] D. Pennica, W.E. Holmes, W.J. Kohr, R.N. Harkins, G.A. Vehar, C.A. Ward, W.F. Bennett, E. Yelverton, P.H. Seeburg, H.L. Heyneker, D.V. Goeddel, and D. Collen. Cloning and expression of human tissue-type plasminogen activator cDNA in *E. coli*. *Nature*, 301:214–221, 1983.
- [2256] R. Pensl and R. Suetfeld. Occurrence of 3,4-diacetylbutinylbithiophene in *Tagetes patula* and its enzymatic conversion. *Z. Naturforsch. C: Biosci.*, 40:3–7, 1985.
- [2257] A. Peracchi, M. Veiga da Cunha, T. Kuhara, K.W. Ellens, N. Paczia, V. Stroobant, A.K. Seliga, S. Marlaire, S. Jaisson, G.T. Bommer, J. Sun, K. Huebner, C.L. Linster, A.J.L. Cooper, and E. Van Schaftingen. Nit1 is a metabolite repair enzyme that hydrolyzes deaminated glutathione. *Proc. Natl Acad. Sci. USA*, 114:E3233–E3242, 2017.
- [2258] X. Pesesse, C. Moreau, A.L. Drayer, R. Woscholski, P. Parker, and C. Erneux. The SH<sub>2</sub> domain containing inositol 5-phosphatase SHIP2 displays phosphatidylinositol 3,4,5-trisphosphate and inositol 1,3,4,5-tetrakisphosphate 5-phosphatase activity. *FEBS Lett.*, 437:301–303, 1998.
- [2259] J.L. Pesquero, P. Boschcov, M.C.F. Oliveira, and A.C.M. Paiva. Effect of substrate size on tonin activity. *Biochem. Biophys. Res. Commun.*, 108:1441–1446, 1982.
- [2260] C. Petersen and L.B. Møller. The RihA, RihB, and RihC ribonucleoside hydrolases of *Escherichia coli*. Substrate specificity, gene expression, and regulation. *J. Biol. Chem.*, 276:884–894, 2001.
- [2261] L.C. Petersen, M. Johannessen, D. Forster, A. Kumar, and E. Mulvihill. The effect of polymerised fibrin on the catalytic activities of one-chain tissue-type plasminogen activator as revealed by an analogue resistant to plasmin cleavage. *Biochim. Biophys. Acta*, 952:245–254, 1988.
- [2262] L.N. Petersen, S. Marineo, S. Mandala, F. Davids, B.T. Sewell, and R.A. Ingle. The missing link in plant histidine biosynthesis: *Arabidopsis* myoinositol monophosphatase-like2 encodes a functional histidinol-phosphate phosphatase. *Plant Physiol.*, 152:1186–1196, 2010.
- [2263] T.N. Petersen, S. Christgau, L.V. Kofod, S. Kauppinen, A.H. Johnson, and S. Larsen. Crystallization and preliminary X-ray studies of rhamnogalacturonase A from *Aspergillus aculeatus*. *Acta Crystallogr. D Biol. Crystallogr.*, 53:105–107, 1997.
- [2264] T.N. Petersen, S. Kauppinen, and S. Larsen. The crystal structure of rhamnogalacturonase A from *Aspergillus aculeatus*: a right-handed parallel  $\beta$  helix. *Structure*, 5:533–544, 1997.
- [2265] S.B. Peterson and J. Liu. Unraveling the specificity of heparanase utilizing synthetic substrates. *J. Biol. Chem.*, 285:14504–14513, 2010.
- [2266] S. Petit, N. Lejal, J.C. Huet, and B. Delmas. Active residues and viral substrate cleavage sites of the protease of the birnavirus infectious pancreatic necrosis virus. *J. Virol.*, 74:2057–2066, 2000.
- [2267] P.H. Petra. Bovine procarboxypeptidase and carboxypeptidase A. *Methods Enzymol.*, 19:460–503, 1970.
- [2268] B. Petrack, P. Greengard, A. Craston, and F. Sheppy. Nicotinamide deamidase from mammalian liver. *J. Biol. Chem.*, 240:1725–1730, 1965.

- [2269] B. Petrack, L. Sullivan, and S. Ratner. Behavior of purified arginine desiminase from *S. faecalis*. *Arch. Biochem. Biophys.*, 69:186–197, 1957.
- [2270] A. Pfitzner and J. Stöckigt. Characterization of polyneuridine aldehyde esterase, a key enzyme in the biosynthesis of sarpagine ajmaline type alkaloids. *Planta Med.*, 48:221–227, 1983.
- [2271] A. Pfitzner and J. Stöckigt. Polyneuridine aldehyde esterase: an unusual specific enzyme involved in the biosynthesis of sarpagine type alkaloids. *J. Chem. Soc. Chem. Commun.*, pages 459–460, 1983.
- [2272] H.J. Phaff and A.L. Demain. The unienzymatic nature of yeast polygalacturonase. *J. Biol. Chem.*, 218:875–884, 1956.
- [2273] T.N. Phan, N.Z. Ehtesham, R. Tuteja, and N. Tuteja. A novel nuclear DNA helicase with high specific activity from *Pisum sativum* catalytically translocates in the 3'→5' direction. *Eur. J. Biochem.*, 270:1735–1745, 2003.
- [2274] C.J. Phiel, F. Zhang, E.Y. Huang, M.G. Guenther, M.A. Lazar, and P.S. Klein. Histone deacetylase is a direct target of valproic acid, a potent anticonvulsant, mood stabilizer, and teratogen. *J. Biol. Chem.*, 276:36734–36741, 2001.
- [2275] M. Philipp and M.L. Bender. Kinetics of subtilisin and thiolsubtilisin. *Mol. Cell. Biochem.*, 51:5–32, 1983.
- [2276] M.E. Piccone, M. Zellner, T.F. Kumosinski, P.W. Mason, and M.J. Grubman. Identification of the active-site residues of the L proteinase of foot-and-mouth disease virus. *J. Virol.*, 69:4950–4956, 1995.
- [2277] R.J. Pierce, G. Spik, and J. Montreuil. Cytosolic location of an endo-*N*-acetyl- $\beta$ -D-glucosaminidase activity in rat liver and kidney. *Biochem. J.*, 180:673–673, 1979.
- [2278] R.J. Pierce, G. Spik, and J. Montreuil. Demonstration and cytosolic location of an endo-*N*-acetyl- $\beta$ -D-glucosaminidase activity towards an asialo-*N*-acetyl-lactosaminic-type substrate in rat liver. *Biochem. J.*, 185:261–264, 1980.
- [2279] A. Pierotti, K.W. Dong, M.J. Glucksman, M. Orłowski, and J.L. Roberts. Molecular cloning and primary structure of rat testes metalloendopeptidase EC 3.4.24.15. *Biochemistry*, 29:10323–10329, 1990.
- [2280] A.R. Pierotti, A. Prat, V. Chesneau, F. Gaudoux, A.-M. Leseney, T. Foulon, and P. Cohen. *N*-Arginine dibasic convertase, a metalloendopeptidase as a prototype of a class of processing enzymes. *Proc. Natl. Acad. Sci. USA*, 91:6078–6082, 1994.
- [2281] W.W. Pigman. Action of almond emulsin on the phenyl glucosides of synthetic sugars and on  $\beta$ -thiophenyl *d*-glucoside. *J. Res. Nat. Bur. Stand.*, 26:197–204, 1941.
- [2282] D.S. Pikas, J.P. Li, I. Vlodavsky, and U. Lindahl. Substrate specificity of heparanases from human hepatoma and platelets. *J. Biol. Chem.*, 273:18770–18777, 1998.
- [2283] A.C. Pike, B. Shrestha, V. Popuri, N. Burgess-Brown, L. Muzzolini, S. Costantini, A. Vindigni, and O. Gileadi. Structure of the human RECQ1 helicase reveals a putative strand-separation pin. *Proc. Natl. Acad. Sci. USA*, 106:1039–1044, 2009.
- [2284] R. Pike, W. McGraw, J. Potempa, and J. Travis. Lysine- and arginine-specific proteinases from *Porphyromonas gingivalis*. Isolation, characterization, and evidence for the existence of complexes with hemagglutinins. *J. Biol. Chem.*, 269:406–411, 1994.
- [2285] E.V. Pindel, N.Y. Kedishvili, T.L. Abraham, M.R. Brzezinski, J. Zhang, R.A. Dean, and W.F. Bosron. Purification and cloning of a broad substrate specificity human liver carboxylesterase that catalyzes the hydrolysis of cocaine and heroin. *J. Biol. Chem.*, 272:14769–14775, 1997.
- [2286] H. Pirkle, I. Theodor, D. Miyada, and G. Simmons. Thrombin-like enzyme from the venom of *Bitis gabonica*. Purification, properties and coagulant actions. *J. Biol. Chem.*, 261:8830–8835, 1986.
- [2287] G. Pitari, F. Malergue, F. Martin, J.M. Philippe, M.T. Massucci, C. Chabret, B. Maras, S. Duprè, P. Naquet, and F. Galland. Pantetheinase activity of membrane-bound Vanin-1: lack of free cysteamine in tissues of Vanin-1 deficient mice. *FEBS Lett.*, 483:149–154, 2000.
- [2288] H.C. Pitot and E. Gohda. Cathepsin T. *Methods Enzymol.*, 142:279–289, 1987.
- [2289] S.M. Pitson, M. Mutter, L.A. van den Broek, A.G. Voragen, and G. Beldman. Stereochemical course of hydrolysis catalysed by  $\alpha$ -L-rhamnosyl and  $\alpha$ -D-galacturonosyl hydrolases from *Aspergillus aculeatus*. *Biochem. Biophys. Res. Commun.*, 242:552–559, 1998.



- [2290] W.J. Placzek, M.S. Almeida, and K. Wuthrich. NMR structure and functional characterization of a human cancer-related nucleoside triphosphatase. *J. Mol. Biol.*, 367:788–801, 2007.
- [2291] R.J. Planta, J. Gorter, and M. Gruber. The catalytic properties of cathepsin C. *Biochim. Biophys. Acta*, 89:511–519, 1964.
- [2292] A.G. Plaut. The IgA1 proteases of pathogenic bacteria. *Annu. Rev. Microbiol.*, 37:603–622, 1983.
- [2293] I Plevrakis, C. Créminon, C. Clamagirand, N. Brakch, M. Rholam, and P. Cohen. Proocytocin/neurophysin convertase from bovine neurohypophysis and corpus luteum secretory granules: complete purification, structure-function relationships, and competitive inhibitor. *Biochemistry*, 28:2705–2710, 1989.
- [2294] T.H. Plummer, Erdös Jr., and E.G. Human plasma carboxypeptidase N. *Methods Enzymol.*, 80:442–449, 1981.
- [2295] T.H. Plummer, Tarentino Jr., and A.L. Facile cleavage of complex oligosaccharides from glycopeptides by almond emulsin peptide: *N*-glycosidase. *J. Biol. Chem.*, 256:10243–10246, 1981.
- [2296] K. Podzelinska, S.M. He, M. Wathier, A. Yakunin, M. Proudfoot, B. Hove-Jensen, D.L. Zechel, and Z. Jia. Structure of PhnP, a phosphodiesterase of the carbon-phosphorus lyase pathway for phosphonate degradation. *J. Biol. Chem.*, 284:17216–17226, 2009.
- [2297] M. Poe, J.T. Blake, D.A. Boulton, M. Gammon, N.H. Sigal, J.K. Wu, and H.J. Zweerink. Human cytotoxic lymphocyte granzyme B. Its purification from granules and the characterization of substrate and inhibitor specificity. *J. Biol. Chem.*, 266:98–103, 1991.
- [2298] M. Poe, J.K. Wu, J.R. Florance, J.A. Rodkey, C.D. Bennett, and K. Hoogsteen. Purification and properties of renin and  $\gamma$ -renin from the mouse submaxillary gland. *J. Biol. Chem.*, 258:2209–2216, 1983.
- [2299] M. Pogorevc and K. Faber. Purification and characterization of an inverting stereo- and enantioselective sec-alkylsulfatase from the gram-positive bacterium *Rhodococcus ruber* DSM 44541. *Appl. Environ. Microbiol.*, 69:2810–2815, 2003.
- [2300] J. Pohl and B.M. Dunn. Secondary enzyme-substrate interactions: kinetic evidence for ionic interactions between substrate side chains and the pepsin active site. *Biochemistry*, 27:4827–4834, 1988.
- [2301] L. Polgár. Isolation of highly active papaya peptidases A and B from commercial chymopapain. *Biochim. Biophys. Acta*, 658:262–269, 1981.
- [2302] L. Polgár. Problems of classification of papaya latex proteinases. *Biochem. J.*, 221:555–556, 1984.
- [2303] L. Polgár. Structure and function of serine proteases. In A. Neuberger and K. Brocklehurst, editors, *New Comprehensive Biochemistry: Hydrolytic Enzymes*, volume 16, pages 159–200. Elsevier, Amsterdam, 1987.
- [2304] L. Polgár and C. Csoma. Dissociation of ionizing groups in the binding cleft inversely controls the endo- and exopeptidase activities of cathepsin B. *J. Biol. Chem.*, 262:14448–14453, 1987.
- [2305] M.R. Pollard, L. Anderson, C. Fan, D.J. Hawkins, , and H.M. A specific acyl-ACP thioesterase implicated in medium-chain fatty acid production in immature cotyledons of *Umbellularia californica*. *Arch. Biochem. Biophys.*, 284:306–312, 1991.
- [2306] M.R. Pollock. Penicillinase. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 4, pages 269–278. Academic Press, New York, 2nd edition, 1960.
- [2307] M.R. Pollock, A.-M. Torriani, and E.G. Tridgell. Crystalline bacterial penicillinase. *Biochem. J.*, 62:387–391, 1956.
- [2308] L. Polz, H. Schübel, and J. Stöckigt. Characterization of 2 $\beta$ (*R*)-17-*O*-acetyljmalan:acetylerase—a specific enzyme involved in the biosynthesis of the *Rauwolfia* alkaloid ajmaline. *Z. Naturforsch. [C]*, 42:333–342, 1987.
- [2309] K. Ponnuraj, S. Rowland, C. Nessi, P. Setlow, and M.J. Jedrzejewski. Crystal structure of a novel germination protease from spores of *Bacillus megaterium*: Structural arrangement and zymogen activation. *J. Mol. Biol.*, 300:1–10, 2000.
- [2310] S. Pontremoli, S. Traniello, B. Luppis, and W.A. Wood. Fructose diphosphatase from rabbit liver. I. Purification and properties. *J. Biol. Chem.*, 240:3459–3463, 1965.

- [2311] J.L. Pope, J.C. McPherson, and H.C. Tidwell. A study of a monoglyceride-hydrolyzing enzyme of intestinal mucosa. *J. Biol. Chem.*, 241:2306–2310, 1966.
- [2312] M. Popowska, M. Kusio, P. Szymanska, and Z. Markiewicz. Inactivation of the wall-associated de-*N*-acetylase (PgDA) of *Listeria monocytogenes* results in greater susceptibility of the cells to induced autolysis. *J. Microbiol. Biotechnol.*, 19:932–945, 2009.
- [2313] M.A. Porzio and A.M. Pearson. Isolation of an extracellular neutral proteinase from *Pseudomonas fragi*. *Biochim. Biophys. Acta*, 384:235–241, 1975.
- [2314] J. Potempa, A. Dubin, and J. Travis. Staphylopain. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 669–671. Handbook of Proteolytic Enzymes, London, 1998.
- [2315] J. Potempa, Z. Porwit-Bohr, and J. Travis. Stabilization vs. degradation of *Staphylococcus aureus* metalloproteinase. *Biochim. Biophys. Acta*, 993:301–304, 1989.
- [2316] P. Potin, C. Richard, T. Barbeyron, B. Henrissat, C. Gey, Y. Petillot, E. Forest, O. Dideberg, C. Rochas, and B. Kloareg. Processing and hydrolytic mechanism of the *cgkA*-encoded  $\kappa$ -carrageenase of *Alteromonas carrageenovora*. *Eur. J. Biochem.*, 228:971–975, 1995.
- [2317] P. Potin, C. Richard, C. Rochas, and B. Kloareg. Purification and characterization of the  $\alpha$ -agarase from *Alteromonas agarlyticus* (Cataldi) comb. nov., strain GJ1B. *Eur. J. Biochem.*, 214:599–607, 1993.
- [2318] P. Potin, A. Sanseau, Y. Le Gall, C. Rochas, and B. Kloareg. Purification and characterization of a new  $\kappa$ -carrageenase from a marine *Cytophaga*-like bacterium. *Eur. J. Biochem.*, 201:241–247, 1991.
- [2319] K. Poutanen, M. Sundberg, H. Korte, , and J. Deacetylation of xylans by acetyl esterases of *Trichoderma reesei*. *Appl. Microbiol. Biotechnol.*, 33:506–510, 1990.
- [2320] J.F. Powell and J.R. Hunter. Adenosine deaminase and ribosidase in spores of *Bacillus cereus*. *Biochem. J.*, 62:381–387, 1956.
- [2321] J.C. Powers, T. Tanaka, J.W. Harper, Y. Minematsu, L. Barker, D. Lincoln, K.V. Crumley, J.E. Fraki, N.M. Schechter, G.G. Lazarus, K. Nakajima, K. Nakashino, H. Neurath, and R.G. Woodbury. Mammalian chymotrypsin-like enzymes. Comparative reactivities of rat mast cell proteases, human and dog skin chymases, and human cathepsin G with peptide 4-nitroanilide substrates and with peptide chloromethyl ketone and sulfonyl fluoride inhibitors. *Biochemistry*, 24:2048–2058, 1985.
- [2322] A.G. Pratt, E.J. Crawford, and M. Friedkin. The hydrolysis of mono-, di-, and triglutamate derivatives of folic acid with bacterial enzymes. *J. Biol. Chem.*, 243:6367–6372, 1968.
- [2323] J.M. Prescott and S.H. Wilkes. *Aeromonas* aminopeptidase: purification and some general properties. *Arch. Biochem. Biophys.*, 117:328–336, 1966.
- [2324] A. Preumont, R. Rzem, D. Vertommen, and E. Van Schaftingen. HDHD1, which is often deleted in X-linked ichthyosis, encodes a pseudouridine-5'-phosphatase. *Biochem. J.*, 431:237–244, 2010.
- [2325] J.M. Prien, D.J. Ashline, A.J. Lapadula, H. Zhang, and V.N. Reinhold. The high mannose glycans from bovine ribonuclease B isomer characterization by ion trap MS. *J Am Soc Mass Spectrom*, 20:539–556, 2009.
- [2326] G.G. Pritchard and T. Coolbear. The physiology and biochemistry of the proteolytic system in lactic acid bacteria. *FEMS Microbiol. Rev.*, 12:179–206, 1993.
- [2327] M. Proudfoot, E. Kuznetsova, G. Brown, N.N. Rao, M. Kitagawa, H. Mori, A. Savchenko, and A.F. Yakunin. General enzymatic screens identify three new nucleotidases in *Escherichia coli*. Biochemical characterization of SurE, YfbR, and YjjG. *J. Biol. Chem.*, 279:54687–54694, 2004.
- [2328] E. Psylinakis, I.G. Boneca, K. Mavromatis, A. Deli, E. Hayhurst, S.J. Foster, K.M. Varum, and V. Bouriotis. Peptidoglycan *N*-acetylglucosamine deacetylases from *Bacillus cereus*, highly conserved proteins in *Bacillus anthracis*. *J. Biol. Chem.*, 280:30856–30863, 2005.
- [2329] J. Puls.  $\alpha$ -Glucuronidases in the hydrolysis of wood xylans. In J. Visser, M.A. Kusters van Someren, G. Beldman, and A.G.J. Voragen, editors, *Xylans and Xylanases*, pages 213–224. Elsevier, Amsterdam, 1992.

- [2330] R.E. Purdy and P.E. Kolattukudy. Hydrolysis of plant cuticle by plant pathogens. Properties of cutinase I, cutinase II, and a nonspecific esterase isolated from *Fusarium solani pisi*. *Biochemistry*, 14:2832–2840, 1975.
- [2331] R.E. Purdy and P.E. Kolattukudy. Hydrolysis of plant cuticle by plant pathogens. Purification, amino acid composition, and molecular weight of two isoenzymes of cutinase and a nonspecific esterase from *Fusarium solani* f. *pisii*. *Biochemistry*, 14:2824–2831, 1975.
- [2332] C. Purwin, M. Laux, and H. Holzer. Fructose 2-phosphate, an intermediate of the dephosphorylation of fructose 2,6-bisphosphate with purified yeast enzyme. *Eur. J. Biochem.*, 164:27–30, 1986.
- [2333] C. Purwin, M. Laux, and H. Holzer. Fructofuranose 2-phosphate is the product of dephosphorylation of fructose 2,6-bisphosphate. *Eur. J. Biochem.*, 165:543–545, 1987.
- [2334] A. Pushkin, G. Carpenito, N. Abuladze, D. Newman, V. Tsuprun, S. Ryazantsev, S. Motemoturu, P. Sassani, N. Solovieva, R. Dukkupati, and I. Kurtz. Structural characterization, tissue distribution, and functional expression of murine aminocyclase III. *Am. J. Physiol. Cell Physiol.*, 286:C848–C856, 2004.
- [2335] A. Qamar and D. Golemi-Kotra. Dual roles of FmtA in *Staphylococcus aureus* cell wall biosynthesis and autolysis. *Antimicrob. Agents Chemother.*, 56:3797–3805, 2012.
- [2336] S. Qian, H. Yu, C. Zhang, M. Lu, H. Wang, and F. Jin. Purification and characterization of dioscin- $\alpha$ -L-rhamnosidase from pig liver. *Chem Pharm Bull (Tokyo)*, 53:911–914, 2005.
- [2337] Y.M. Qian, O. Varlamov, and L.D. Fricker. Glu<sup>300</sup> of rat carboxypeptidase E is essential for enzymatic activity but not substrate binding or routing to the regulated secretory pathway. *J. Biol. Chem.*, 274:11582–11586, 1999.
- [2338] Y. Qu and J.C. Spain. Biodegradation of 5-nitroanthranilic acid by *Bradyrhizobium* sp. strain JS329. *Appl. Environ. Microbiol.*, 76:1417–1422, 2010.
- [2339] Y. Qu and J.C. Spain. Catabolic pathway for 2-nitroimidazole involves a novel nitrohydrolase that also confers drug resistance. *Environ Microbiol*, 13:1010–1017, 2011.
- [2340] Y. Qu and J.C. Spain. Molecular and biochemical characterization of the 5-nitroanthranilic acid degradation pathway in *Bradyrhizobium* sp. strain JS329. *J. Bacteriol.*, 193:3057–3063, 2011.
- [2341] B. Quantin, G. Murphy, and R. Breathnach. Pump-1 cDNA codes for a protein with characteristics similar to those of classical collagenase family members. *Biochemistry*, 28:5327–5334, 1989.
- [2342] B.G. Que and E.W. Davie. Characterization of a cDNA coding for human factor XII (Hageman factor). *Biochemistry*, 25:1525–1528, 1986.
- [2343] A. Quentmeier and C.G. Friedrich. The cysteine residue of the SoxY protein as the active site of protein-bound sulfur oxidation of *Paracoccus pantotrophus* GB17. *FEBS Lett.*, 503:168–172, 2001.
- [2344] A. Quentmeier, P. Hellwig, F. Bardischewsky, G. Grelle, R. Kraft, and C.G. Friedrich. Sulfur oxidation in *Paracoccus pantotrophus*: interaction of the sulfur-binding protein SoxYZ with the dimanganese SoxB protein. *Biochem. Biophys. Res. Commun.*, 312:1011–1018, 2003.
- [2345] G.B. Quistad, C. Barlow, C.J. Winrow, S.E. Sparks, and J.E. Casida. Evidence that mouse brain neuropathy target esterase is a lysophospholipase. *Proc. Natl. Acad. Sci. USA*, 100:7983–7987, 2003.
- [2346] U. Rabausch, N. Ilmberger, and W.R. Streit. The metagenome-derived enzyme RhaB opens a new subclass of bacterial B type  $\alpha$ -L-rhamnosidases. *J. Biotechnol.*, 191:38–45, 2014.
- [2347] D. Rabier and M.J. Desmazeaud. Inventaire des différentes activités peptidasiques intracellulaires de *Streptococcus thermophilus*. Purification et propriétés d'une dipeptide-hydrolase et d'une aminopeptidase. *Biochimie*, 55:389–404, 1973.
- [2348] J.C. Rabinowitz and H.A. Barker. Purine fermentation by *Clostridium cylindrosporium*. II. Purine transformations. *J. Biol. Chem.*, 218:161–173, 1956.
- [2349] J.C. Rabinowitz and W.E. Pricer. Purine fermentation by *Clostridium cylindrosporium*. V. Formiminoglycine. *J. Biol. Chem.*, 222:537–554, 1956.

- [2350] J.C. Rabinowitz and W.E. Pricer. The enzymatic synthesis of  $N^{10}$ -formyltetrahydrofolic acid and its role in ATP formation during formiminoglycine degradation. *J. Am. Chem. Soc.*, 78:4176–4178, 1956.
- [2351] P.R. Race, M.L. Bentley, J.A. Melvin, A. Crow, R.K. Hughes, W.D. Smith, R.B. Sessions, M.A. Kehoe, D.G. McCafferty, and M.J. Banfield. Crystal structure of *Streptococcus pyogenes* sortase A: implications for sortase mechanism. *J. Biol. Chem.*, 284:6924–6933, 2009.
- [2352] E. Racker. Spectrophotometric measurements of the metabolic formation and degradation of thiol esters and enediol compounds. *Biochim. Biophys. Acta*, 9:577–579, 1952.
- [2353] E. Racker. Sedoheptulose-1,7-diphosphatase from yeast. *Methods Enzymol.*, 5:270–272, 1962.
- [2354] C.R.H. Raetz, B. Hirschberg, W. Dowhan, W.T. Wickner, and E.P. Kennedy. A membrane-bound pyrophosphatase in *Escherichia coli* catalyzing the hydrolysis of cytidine diphosphate-diglyceride. *J. Biol. Chem.*, 247:2245–2247, 1972.
- [2355] G.W. Rafter. Pyrophosphate metabolism in liver mitochondria. *J. Biol. Chem.*, 235:2475–2477, 1960.
- [2356] S. Ragasu, P. Mouchet, C. Lazennec, V. Dive, and T. Meinnel. Substrate recognition and selectivity of peptide deformylase. Similarities and differences with metzincins and thermolysin. *J. Mol. Biol.*, 289:1445–1457, 1999.
- [2357] F.L. Ragone, J.L. Spears, J.M. Wohlgamuth-Benedum, N. Kreel, F.N. Papavasiliou, and J.D. Alfonzo. The C-terminal end of the *Trypanosoma brucei* editing deaminase plays a critical role in tRNA binding. *RNA*, 17:1296–1306, 2011.
- [2358] M.M. Rahman, H.N. Hunter, S. Prova, V. Verma, A. Qamar, and D. Golemi-Kotra. The *Staphylococcus aureus* methicillin resistance factor FmtA is a D-amino esterase that acts on teichoic acids. *MBio*, 7:e02070–e02015, 2016.
- [2359] L. Rajijman, S. Grisolia, and H. Edelhofer. Further purification and properties of brain acyl phosphatase. *J. Biol. Chem.*, 235:2340–2342, 1960.
- [2360] P.T.R. Rajagopalan, S. Grimme, and D. Pei. Characterization of cobalt(II)-substituted peptide deformylase: function of the metal ion and the catalytic residue Glu-133. *Biochemistry*, 39:779–790, 2000.
- [2361] P.T.R. Rajagopalan, X.C. Yu, and D. Pei. Peptide deformylase: a new type of mononuclear iron protein. *J. Am. Chem. Soc.*, 119:12418–12419, 1997.
- [2362] S. Rajapakse, K. Ogiwara, N. Takano, A. Moriyama, and T. Takahashi. Biochemical characterization of human kallikrein 8 and its possible involvement in the degradation of extracellular matrix proteins. *FEBS Lett.*, 579:6879–6884, 2005.
- [2363] T.W. Rall, W.D. Wosilait, and E.W. Sutherland. The interconversion of phosphorylase *a* and phosphorylase *b* from dog heart muscle. *Biochim. Biophys. Acta*, 20:69–76, 1956.
- [2364] R. Ramachandran, C. Hartmann, H.K. Song, R. Huber, and M. Bochtler. Functional interactions of HslV (ClpQ) with the ATPase HslU (ClpY). *Proc. Natl. Acad. Sci. USA*, 99:7396–7401, 2002.
- [2365] S. Ramaswamy and A.N. Radhakrishnan. Lactase-phlorizin hydrolase complex from monkey small intestine. Purification, properties and evidence for two catalytic sites. *Biochim. Biophys. Acta*, 403:446–455, 1975.
- [2366] S.G. Ramaswamy and W.B. Jakoby. (2')3',5'-Bisphosphate nucleotidase. *J. Biol. Chem.*, 262:10044–10047, 1987.
- [2367] G. Ramponi, A. Gueritore, C. Treves, P. Nassi, and V. Baccari. Horse muscle acyl phosphatase: purification and some properties. *Arch. Biochem. Biophys.*, 130:362–369, 1969.
- [2368] G. Ramponi, P. Nassi, G. Cappugi, C. Treves, and G. Manao. Purification and some molecular properties of horse liver acyl phosphatase. *Biochim. Biophys. Acta*, 284:485–496, 1972.
- [2369] D.D. Randall and N.E. Tolbert. 3-Phosphoglycerate phosphatase in plants. I. Isolation and characterization from sugarcane leaves. *J. Biol. Chem.*, 246:5510–5517, 1971.
- [2370] L. Randau, B.J. Stanley, A. Kohlway, S. Mechta, Y. Xiong, and D. Söll. A cytidine deaminase edits C to U in transfer RNAs in Archaea. *Science*, 324:657–659, 2009.
- [2371] F. Randez-Gil, A. Blasco, J.A. Prieto, , and P. DOGR1 and DOGR2: two genes from *Saccharomyces cerevisiae* that confer 2-deoxyglucose resistance when overexpressed. *Yeast*, 11:1233–1240, 1995.

- [2372] A. Randolph, S.H. Chamberlain, H.L.C. Chu, A.D. Retzios, , and F.S. Jr. and Masiarz, F.R. Amino acid sequence of fibrolase, a direct-acting fibrinolytic enzyme from *Agkistrodon contortrix* venom. *Protein Sci.*, 1:590–600, 1992.
- [2373] D.R. Rao and D.M. Greenberg. Studies on the enzymic decomposition of urocanic acid. IV. Purification and properties of 4(5)-imidazolone-5(4)-propionic acid hydrolase. *J. Biol. Chem.*, 236:1758–1763, 1961.
- [2374] N.V. Rao, N.G. Wehner, B.C. Marshall, W.R. Gray, B.H. Gray, and J.R. Hoidal. Characterization of proteinase-3 (PR-3), a neutrophil serine proteinase. Structural and functional properties. *J. Biol. Chem.*, 266:9540–9548, 1991.
- [2375] M.M. Rapport, K. Myer, and A. Linker. Analysis of the products formed on hydrolysis of hyaluronic acid by testicular hyaluronidase. *J. Am. Chem. Soc.*, 73:2416–2420, 1951.
- [2376] A. Rascón, S.H. Soderling, J.B. Schaefer, and J.A. Beavo. Cloning and characterization of a cAMP-specific phosphodiesterase (TbPDE2B) from *Trypanosoma brucei*. *Proc. Natl. Acad. Sci. USA*, 99:4714–4719, 2002.
- [2377] S. Ratner. Urea synthesis and metabolism of arginine and citrulline. *Adv. Enzymol. Relat. Subj. Biochem.*, 15:319–387, 1954.
- [2378] J.W. Rausch, D. Lener, J.T. Miller, J.G. Julias, S.H. Hughes, and S.F. Le Grice. Altering the RNase H primer grip of human immunodeficiency virus reverse transcriptase modifies cleavage specificity. *Biochemistry*, 41:4856–4865, 2002.
- [2379] S.D. Ravindranath and N. Appaji Rao. Nucleotidases in plants. 3. Effect of metabolites on the enzyme hydrolyzing flavine adenine dinucleotide (FAD) from *Phaseolus radiatus*. *Arch. Biochem. Biophys.*, 133:54–59, 1969.
- [2380] M. Rawat, S. Kovacevic, H. Billman-Jacobe, and Y. Av-Gay. Inactivation of *mshB*, a key gene in the mycothiol biosynthesis pathway in *Mycobacterium smegmatis*. *Microbiology*, 149:1341–1349, 2003.
- [2381] R. Rawat, F.J. Sandoval, Z. Wei, R. Winkler, and S. Roje. An FMN hydrolase of the haloacid dehalogenase superfamily is active in plant chloroplasts. *J. Biol. Chem.*, 286:42091–42098, 2011.
- [2382] N.D. Rawlings and A.J. Barrett. Homologues of insulinase, a new superfamily of metalloendopeptidases. *Biochem. J.*, 275:389–391, 1991.
- [2383] N.D. Rawlings and A.J. Barrett. Structure of membrane glutamate carboxypeptidase. *Biochim. Biophys. Acta*, 1339:247–252, 1997.
- [2384] N.D. Rawlings and A.J. Barrett. Tripeptidyl-peptidase I is apparently the CLN2 protein absent in classical late-infantile neuronal ceroid lipofuscinosis. *Biochim. Biophys. Acta*, 1429:496–500, 1999.
- [2385] P.H. Ray and C.D. Benedict. Purification and characterization of specific 3-deoxy-D-manno-octulosonate 8-phosphate phosphatase from *Escherichia coli* B. *J. Bacteriol.*, 142:60–68, 1980.
- [2386] A. Raychaudhuri and P.A. Tipton. Cloning and expression of the gene for soybean hydroxyisourate hydrolase. Localization and implications for function and mechanism. *Plant Physiol.*, 130:2061–2068, 2002.
- [2387] A. Raychaudhuri and P.A. Tipton. A familiar motif in a new context: the catalytic mechanism of hydroxyisourate hydrolase. *Biochemistry*, 42:6848–6852, 2003.
- [2388] P. Raychaudhuri, S. Ghosh, and U. Maitra. Purification and characterization of a guanosine diphosphatase activity from calf liver microsomal salt wash proteins. *J. Biol. Chem.*, 260:8306–8311, 1985.
- [2389] R.J. Read, G.D. Brayer, L. Jurásek, and M.N.G. James. Critical evaluation of comparative model building of *Streptomyces griseus* trypsin. *Biochemistry*, 23:6570–6575, 1984.
- [2390] R.J. Read, M. Fujinaga, A.R. Sielecki, and M.N.G. James. Structure of the complex of *Streptomyces griseus* protease B and the third domain of turkey ovomucoid inhibitor at 1.8-Å resolution. *Biochemistry*, 22:4420–4433, 1983.
- [2391] J. Rech, G. Cathala, and P. Jeanteur. Isolation and characterization of a ribonuclease activity specific for double-stranded RNA (RNase D) from Krebs II ascites cells. *J. Biol. Chem.*, 255:6700–6706, 1980.
- [2392] P.A. Recsei, A.D. Gruss, and R.P. Novick. Cloning, sequence, and expression of the lysostaphin gene from *Staphylococcus simulans*. *Proc. Natl. Acad. Sci. USA*, 84:1127–1131, 1987.

- [2393] K.K. Reddi. Micrococcal nuclease. *Methods Enzymol.*, 12A:257–262, 1967.
- [2394] G.R. Reeck, K.A. Walsh, and H. Neurath. Isolation and characterization of carboxypeptidases A and B from activated pancreatic juice. *Biochemistry*, 10:4690–4698, 1971.
- [2395] L.J. Reed, Z. Damuni, and M.L. Merryfield. Regulation of mammalian pyruvate and branched-chain  $\alpha$ -keto acid dehydrogenase complexes by phosphorylation-dephosphorylation. *Curr. Top. Cell. Regul.*, 27:41–49, 1985.
- [2396] E.T. Reese.  $\beta$ -Mannanases of fungi. *Can. J. Microbiol.*, 11:167–183, 1965.
- [2397] E.T. Reese and M. Mandels.  $\beta$ -D-1,3-Glucanases in fungi. *Can. J. Microbiol.*, 5:173–185, 1959.
- [2398] E.T. Reese, F.W. Parrish, and M. Mandels.  $\beta$ -D-1,2-Glucanases in fungi. *Can. J. Microbiol.*, 7:309–317, 1961.
- [2399] E.T. Reese, F.W. Parrish, and M. Mandels.  $\beta$ -D-1,6-Glucanases in fungi. *Can. J. Microbiol.*, 8:327–334, 1962.
- [2400] A. Reglero and J.A. Cabezas. Glycosidases of molluscs. Purification and properties of  $\alpha$ -L-fucosidase from *Chamelea gallina* L. *Eur. J. Biochem.*, 66:379–387, 1976.
- [2401] K.B.M. Reid, D.M.A. Johnson, J. Gagnon, and R. Prohaska. Preparation of human factor  $\bar{d}$  of the alternative pathway of complement. *Methods Enzymol.*, 80:134–143, 1981.
- [2402] W.A. Reid, L. Vongsorasak, J. Svasti, M.J. Valler, and J. Kay. Identification of the acid proteinase in human seminal fluid as a gastricsin originating in the prostate. *Cell Tissue Res.*, 236:597–600, 1984.
- [2403] J.L. Reissig, W. Lau, and J.E. Glasgow. An endogalactosaminidase from *Streptomyces griseus*. *Can. J. Biochem.*, 53:1237–1249, 1975.
- [2404] M.S. Reitz and V.W. Rodwell.  $\delta$ -Aminovaleramidase of *Pseudomonas putida*. *J. Biol. Chem.*, 245:3091–3096, 1970.
- [2405] H. Rembold and F. Simmersbach. Catabolism of pteridine cofactors. II. A specific pterin deaminase in rat liver. *Biochim. Biophys. Acta*, 184:589–596, 1969.
- [2406] H. Remold, H. Fasold, and F. Staib. Purification and characterization of a proteolytic enzyme from *Candida albicans*. *Biochim. Biophys. Acta*, 167:399–406, 1968.
- [2407] B.S. Remus, A. Jacewicz, and S. Shuman. Structure and mechanism of *E. coli* RNA 2',3'-cyclic phosphodiesterase. *RNA*, 20:1697–1705, 2014.
- [2408] G. Rendina and M.J. Coon. Enzymatic hydrolysis of the coenzyme A thiol esters of  $\beta$ -hydroxypropionic and  $\beta$ -hydroxyisobutyric acids. *J. Biol. Chem.*, 225:523–534, 1957.
- [2409] D. Rennex, B.A. Hemmings, J. Hofsteenge, and S.R. Stone. cDNA cloning of porcine brain prolyl endopeptidase and identification of the active-site seryl residue. *Biochemistry*, 30:2195–2203, 1991.
- [2410] C.E. Requena, G. Perez-Moreno, L.M. Ruiz-Perez, A.E. Vidal, and D. Gonzalez-Pacanowska. The NTP pyrophosphatase DCTPP1 contributes to the homeostasis and cleansing of the dNTP pool in human cells. *Biochem. J.*, 459:171–180, 2014.
- [2411] A.D. Retzios and F.S. Markland. Fibrinolytic enzymes from the venoms of *Agkistrodon contortrix contortrix* and *Crotalus basiliscus basiliscus*: cleavage site specificity towards the  $\alpha$ -chain of fibrin. *Thromb. Res.*, 74:355–367, 1994.
- [2412] M.A. Rey and P.L. Davies. The protease core of the muscle-specific calpain, p94, undergoes  $\text{Ca}^{2+}$ -dependent intramolecular autolysis. *FEBS Lett.*, 532:401–406, 2002.
- [2413] P.E. Reynolds, F. Depardieu, S. Dutka-Malen, M. Arthur, and P. Courvalin. Glycopeptide resistance mediated by enterococcal transposon Tn1546 requires production of VanX for hydrolysis of D-alanyl-D-alanine. *Mol. Microbiol.*, 13:1065–1070, 1994.
- [2414] S.G. Rhee and Y.S. Bae. Regulation of phosphoinositide-specific phospholipase C isozymes. *J. Biol. Chem.*, 272:15045–15048, 1997.
- [2415] S.K. Rhee and G. Fuchs. Phenylacetyl-CoA:acceptor oxidoreductase, a membrane-bound molybdenum-iron-sulfur enzyme involved in anaerobic metabolism of phenylalanine in the denitrifying bacterium *Thauera aromatica*. *Eur. J. Biochem.*, 262:507–515, 1999.

- [2416] S. Riazuddin and L. Grossman. *Micrococcus luteus* correndonucleases. II. Mechanism of action of two endonucleases specific for DNA containing pyrimidine dimers. *J. Biol. Chem.*, 252:6287–6293, 1977.
- [2417] S. Riazuddin and T. Lindahl. Properties of 3-methyladenine-DNA glycosylase from *Escherichia coli*. *Biochemistry*, 17:2110–2118, 1978.
- [2418] C.M. Rice. Hepatitis C virus polyprotein peptidase. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 272–277. Academic Press, London, 1998.
- [2419] C.C. Richardson and A. Kornberg. A deoxyribonucleic acid phosphatase-exonuclease from *Escherichia coli*. I. Purification of the enzyme and characterization of the phosphatase activity. *J. Biol. Chem.*, 239:242–250, 1964.
- [2420] C.C. Richardson, I.R. Lehman, and A. Kornberg. A deoxyribonucleic acid phosphatase-exonuclease from *Escherichia coli*. II. Characterization of the exonuclease activity. *J. Biol. Chem.*, 239:251–258, 1964.
- [2421] G. Richarme, M. Mihoub, J. Dairou, L.C. Bui, T. Leger, and A. Lamouri. Parkinsonism-associated protein DJ-1/Park7 is a major protein deglycase that repairs methylglyoxal- and glyoxal-glycated cysteine, arginine, and lysine residues. *J. Biol. Chem.*, 290:1885–1897, 2015.
- [2422] D. Richter, S. Fehr, and R. Harder. The guanosine 3',5'-bis(diphosphate) (ppGpp) cycle. Comparison of synthesis and degradation of guanosine 3',5'-bis(diphosphate) in various bacterial systems. *Eur. J. Biochem.*, 99:57–64, 1979.
- [2423] R. Richter, M. Hejazi, R. Kraft, K. Ziegler, and W. Lockau. Cyanophycinase, a peptidase degrading the cyanobacterial reserve material multi-L-arginyl-poly-L-aspartic acid (cyanophycin): molecular cloning of the gene of *Synechocystis* sp. PCC 6803, expression in *Escherichia coli*, and biochemical characterization of the purified enzyme. *Eur. J. Biochem.*, 263:163–169, 1999.
- [2424] R.R. Riddle, P.R. Gibbs, R.C. Willson, and M.J. Benedik. Purification and properties of 2-hydroxy-6-oxo-6-(2'-aminophenyl)hexa-2,4-dienoic acid hydrolase involved in microbial degradation of carbazole. *Protein Expr. Purif.*, 28:182–189, 2003.
- [2425] M. Rietschel-Berst, N.H. Jentoft, P.D. Rick, C. Pletcher, F. Fang, and J.E. Gander. Extracellular exo- $\beta$ -galactofuranosidase from *Penicillium charlesii*: isolation, purification, and properties. *J. Biol. Chem.*, 252:3219–3226, 1977.
- [2426] L.U. Rigo, L.R. Maréchal, M.M. Vieira, and L.A. Veiga. Oxidative pathway for L-rhamnose degradation in *Pallularia pullulans*. *Can. J. Microbiol.*, 31:817–822, 1985.
- [2427] M.V. Riley and M.I. Peters. The localization of the anion-sensitive ATPase activity in corneal endothelium. *Biochim. Biophys. Acta*, 644:251–256, 1981.
- [2428] L.A. Del Rio and R.C.W. Berkeley. Exo- $\beta$ -N-acetylmuramidase - a novel hexosaminidase. Production by *Bacillus subtilis* B, purification and characterization. *Eur. J. Biochem.*, 65:3–12, 1976.
- [2429] J.W. Rip, C.A. Rugar, N. Chaudhary, and K.K. Carroll. Localization of a dolichyl phosphate phosphatase in plasma membranes of rat liver. *J. Biol. Chem.*, 256:1929–1934, 1981.
- [2430] A. Ritonja, D.J. Buttle, N.D. Rawlings, V. Turk, and A.J. Barrett. Papaya proteinase IV amino acid sequence. *FEBS Lett.*, 258:109–112, 1989.
- [2431] A. Ritonja, A.B. Rowan, D.J. Buttle, N.D. Rawlings, V. Turk, and A.J. Barrett. Stem bromelain: amino acid sequence and implications for weak binding to cystatin. *FEBS Lett.*, 247:419–429, 1989.
- [2432] P.D. Roach and F.B.S. Palmer. Human erythrocyte cytosol phosphatidyl-inositol-bisphosphate phosphatase. *Biochim. Biophys. Acta*, 661:323–333, 1981.
- [2433] K.C. Robbins, L. Summaria, and R.C. Wohl. Human plasmin. *Methods Enzymol.*, 80:379–387, 1981.
- [2434] D.W.A. Roberts. The wheat leaf phosphatases. II. Pathway of hydrolysis of some nucleotides at pH 5.5. *J. Biol. Chem.*, 222:259–270, 1956.
- [2435] E. Roberts. Glutaminase. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 4, pages 285–300. Academic Press, New York, 2nd edition, 1960.

- [2436] J. Roberts, J.S. Holcenberg, and W.C. Dolowy. Isolation, crystallization, and properties of *Achromobacteraceae* glutaminase-asparaginase with antitumor activity. *J. Biol. Chem.*, 247:84–90, 1972.
- [2437] R.J. Roberts. Restriction enzymes and their isoschizomers. *Nucleic Acids Res.*, 18:2331–2365, 1990.
- [2438] S.J. Roberts, A.J. Stewart, P.J. Sadler, and C. Farquharson. Human PHOSPHO1 displays high specific phospho-ethanolamine and phosphocholine phosphatase activities. *Biochem. J.*, 382:59–65, 2004.
- [2439] H.D. Robertson, S. Altman, and J.D. Smith. Purification and properties of a specific *Escherichia coli* ribonuclease which cleaves a tyrosine transfer ribonucleic acid precursor. *J. Biol. Chem.*, 247:5243–5251, 1972.
- [2440] H.D. Robertson, R.E. Webster, and N.D. Zinder. Purification and properties of ribonuclease III from *Escherichia coli*. *J. Biol. Chem.*, 243:82–91, 1968.
- [2441] G.W. Robinson. Isolation and characterization of papaya peptidase A from commercial chymopapain. *Biochemistry*, 16:3695–3700, 1975.
- [2442] W.G. Robinson and R.H. Hook. Ricinine nitrilase. I. Reaction product and substrate specificity. *J. Biol. Chem.*, 239:4257–4262, 1964.
- [2443] J.F. Robyt and R.J. Ackerman. Isolation, purification, and characterization of a maltotetraose-producing amylase from *Pseudomonas stutzeri*. *Arch. Biochem. Biophys.*, 145:105–114, 1971.
- [2444] J. Roche, G. Lacombe, and H. Girard. Sur la spécificité de certaines déguanidases bactériennes génératrices d'urée et sur l'argininedihydrolase. *Biochim. Biophys. Acta*, 6:210–216, 1950.
- [2445] B. Rodamilans and G. Montoya. Expression, purification, crystallization and preliminary X-ray diffraction analysis of the DDX3 RNA helicase domain. *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.*, 63:283–286, 2007.
- [2446] S.L. Roderick and B.W. Matthews. Crystallization of methionine aminopeptidase from *Escherichia coli*. *J. Biol. Chem.*, 263:16531–16531, 1988.
- [2447] M.V. Rodnina, A. Savelsberg, V.I. Katunin, and W. Wintermeyer. Hydrolysis of GTP by elongation factor G drives tRNA movement on the ribosome. *Nature*, 385:37–41, 1997.
- [2448] J.R. Rodrigues, A. Fernandez, J. Canales, A. Cabezas, J.M. Ribeiro, M.J. Costas, and J.C. Cameselle. Characterization of *Danio rerio* Mn<sup>2+</sup>-dependent ADP-ribose/CDP-alcohol diphosphatase, the structural prototype of the ADPRibase-Mn-like protein family. *PLoS One*, 7:e42249–e42249, 2012.
- [2449] I.R. Rodriguez, F.R. Taravel, and W.J. Whelan. Characterization and function of pig intestinal sucrase-isomaltase and its separate subunits. *Eur. J. Biochem.*, 143:575–582, 1984.
- [2450] J.A. Rodriguez, J.A. Cabezas, and P. Calvo.  $\beta$ -Fucosidase,  $\beta$ -glucosidase and  $\beta$ -galactosidase activities associated in bovine liver. *Int. J. Biochem.*, 14:695–698, 1982.
- [2451] P. Rodriguez, S.T. Bass, and R.G. Hansen. A pyrophosphatase from mammalian tissues specific for derivatives of ADP. *Biochim. Biophys. Acta*, 167:199–201, 1968.
- [2452] M. Röhr. Hippurathydrolase, ein neues Enzym aus Mikroorganismen. I Mitt. Induzierte Biosynthese in *Fusarium semitectum*. Darstellung einer gereinigten Enzympräparation und Untersuchungen zur Substratspezifität. *Monatshfte Chem.*, 99:2255–2277, 1968.
- [2453] M. Röhr. Hippurathydrolase, ein neues Enzym aus Mikroorganismen. II Mitt. Enzymeigenschaften. *Monatshfte Chem.*, 99:2278–2290, 1968.
- [2454] L.H. Rome, A.J. Garvin, and E.F. Neufeld. Human kidney  $\alpha$ -L-iduronidase: purification and characterization. *Arch. Biochem. Biophys.*, 189:344–353, 1978.
- [2455] C. Romero-Segura, C. Sanz, and A.G. Perez. Purification and characterization of an olive fruit  $\beta$ -glucosidase involved in the biosynthesis of virgin olive oil phenolics. *J. Agric. Food Chem.*, 57:7983–7988, 2009.
- [2456] R.J. Roon and B. Levenberg. Urea amidolyase. I. Properties of the enzyme from *Candida utilis*. *J. Biol. Chem.*, 247:4107–4113, 1972.



- [2457] J. De Roover, A. Van Laere, M. De Winter, J.W. Timmermans, and W. Van den Ende. Purification and properties of a second fructan exohydrolase from the roots of *Cichorium intybus*. *Physiol. Plant.*, 106:28–34, 1999.
- [2458] D.R. Rose. Structure, mechanism and inhibition of Golgi  $\alpha$ -mannosidase II. *Curr. Opin. Struct. Biol.*, 22:558–562, 2012.
- [2459] S. Roseman. Glucosamine metabolism. I. *N*-Acetylglucosamine deacetylase. *J. Biol. Chem.*, 226:115–123, 1957.
- [2460] B.P. Rosen, U. Weigel, R.A. Monticello, and B.P. Edwards. Molecular analysis of an anion pump: purification of the ArsC protein. *Arch. Biochem. Biophys.*, 284:381–385, 1991.
- [2461] E. Rosenfeld and G. Wiederschein. The metabolism of L-rhamnose in animal tissues. *Bull. Soc. Chim. Biol.*, 47:1433–1440, 1965.
- [2462] G.W. Ross and M.G. Boulton. Purification of  $\beta$ -lactamases on QAE-sephadex. *Biochim. Biophys. Acta*, 309:430–439, 1973.
- [2463] T.S. Ross and P.W. Majerus. Inositol-1,2-cyclic-phosphate 2-inositolphosphohydrolase. Substrate specificity and regulation of activity by phospholipids, metal ion chelators, and inositol 2-phosphate. *J. Biol. Chem.*, 266:851–856, 1991.
- [2464] V. Rossi, S. Cseh, I. Bally, N.M. Thielens, J.C. Jensenius, and G.J. Arlaud. Substrate specificities of recombinant mannan-binding lectin-associated serine proteases-1 and -2. *J. Biol. Chem.*, 276:40880–40887, 2001.
- [2465] S. Rottloff, R. Stieber, H. Maischak, F.G. Turini, G. Heubl, and A. Mithofer. Functional characterization of a class III acid endochitinase from the traps of the carnivorous pitcher plant genus, *Nepenthes*. *J. Exp. Bot.*, 62:4639–4647, 2011.
- [2466] Y. Rouillé, G. Westermark, S.K. Martin, and D.F. Steiner. Proglucagon is processed to glucagon by prohormone convertase PC2 in alphaTC1-6 cells. *Proc. Natl. Acad. Sci. USA*, 91:3242–3246, 1994.
- [2467] A. Rousset, M. Nguyen-Disteche, R. Minck, and J.-M. Ghuysen. Penicillin-binding proteins and carboxypeptidase/transpeptidase activities in *Proteus vulgaris* P18 and its penicillin-induced stable L-forms. *J. Bacteriol.*, 152:1042–1048, 1982.
- [2468] A.D. Rowan, D.J. Buttle, and A.J. Barrett. Ananain: a novel cysteine proteinase found in pineapple stem. *Arch. Biochem. Biophys.*, 267:262–270, 1988.
- [2469] A.D. Rowan, D.J. Buttle, and A.J. Barrett. The cysteine proteinases of the pineapple plant. *Biochem. J.*, 266:869–875, 1990.
- [2470] A.B. Roy. The steroid sulphatase of *Patella vulgata*. *Biochim. Biophys. Acta*, 15:300–301, 1954.
- [2471] A.B. Roy. The synthesis and hydrolysis of sulfate esters. *Adv. Enzymol. Relat. Subj. Biochem.*, 22:205–235, 1960.
- [2472] A.B. Roy. Sulphatases, lysosomes and disease. *Aust. J. Exp. Biol. Med. Sci.*, 54:111–135, 1976.
- [2473] S. Roychowdhury, D. Panda, L. Wilson, and M.M. Rasenick. G protein  $\alpha$  subunits activate tubulin GTPase and modulate microtubule polymerization dynamics. *J. Biol. Chem.*, 274:13485–13490, 1999.
- [2474] E.L. Rozenfel'd and I.S. Lukomsкая. [The hydrolysis of 1:6 bonds of dextran by animal tissues.]. *Biokhimiya*, 21:412–415, 1956.
- [2475] R. Röchel. Properties of a purified proteinase from the yeast *Candida albicans*. *Biochim. Biophys. Acta*, 659:99–113, 1981.
- [2476] J.A. Ruiz-Maso, S.P. Anand, M. Espinosa, S.A. Khan, and G. del Solar. Genetic and biochemical characterization of the *Streptococcus pneumoniae* PcrA helicase and its role in plasmid rolling circle replication. *J. Bacteriol.*, 188:7416–7425, 2006.
- [2477] M.E. Rumpho, G.E. Edwards, and W.H. Loescher. A pathway for photosynthetic carbon flow to mannitol in celery leaves. Activity and localization of key enzymes. *Plant Physiol.*, 73:869–873, 1983.
- [2478] P. Runeberg-Roos, K. Törmäkangas, and A. Östman. Primary structure of a barley-grain aspartic proteinase. A plant aspartic proteinase resembling mammalian cathepsin D. *Eur. J. Biochem.*, 202:1021–1027, 1991.

- [2479] M.T.C. Runnegar, K. Scott, E.C. Webb, and B. Zerner. Carboxylesterases (EC 3.1.1). Purification and titration of ox liver carboxylesterase. *Biochemistry*, 8:2013–2018, 1969.
- [2480] M. Ruppert, J. Woll, A. Giritch, E. Genady, X. Ma, and J. Stöckigt. Functional expression of an ajmaline pathway-specific esterase from *Rauvolfia* in a novel plant-virus expression system. *Planta*, 222:888–898, 2005.
- [2481] D.W. Russell. The enzymes, regulation, and genetics of bile acid synthesis. *Annu. Rev. Biochem.*, 72:137–174, 2003.
- [2482] A.P. Ryle. The porcine pepsins and pepsinogens. *Methods Enzymol.*, 19:316–336, 1970.
- [2483] Doonan HJ Doonan S, Vernon CA Hanford R, da Airoid LP Walker JM, Barra D Bossa F, Carloni M, and Riva F. The primary structure of aspartate aminotransferase from pig heart muscle. Digestion with a proteinase having specificity for lysine residues. *Biochem. J.*, 149:497–506, 1975.
- [2484] E.J. 's Gravenmade, G.D. Vogels, and C. Van der Drift. Hydrolysis, racemization and absolute configuration of ureidoglycolate, a substrate of allantoicase. *Biochim. Biophys. Acta*, 198:569–582, 1970.
- [2485] P. Sachelaru, E. Schiltz, G.L. Igloi, and R. Brandsch. An  $\alpha/\beta$ -fold C—C bond hydrolase is involved in a central step of nicotine catabolism by *Arthrobacter nicotinovorans*. *J. Bacteriol.*, 187:8516–8519, 2005.
- [2486] L. Sachs and N. Marks. A highly specific aminotripeptidase of rat brain cytosol. Substrate specificity and effects of inhibitors. *Biochim. Biophys. Acta*, 706:229–238, 1982.
- [2487] S. Sadis and L.E. Hightower. Unfolded proteins stimulate molecular chaperone Hsc70 ATPase by accelerating ADP/ATP exchange. *Biochemistry*, 31:9406–9412, 1992.
- [2488] P.D. Sadowski and I. Bakytá. T<sub>4</sub> endonuclease IV. Improved purification procedure and resolution from T<sub>4</sub> endonuclease 3. *J. Biol. Chem.*, 247:405–412, 1972.
- [2489] P.D. Sadowski and J. Hurwitz. Enzymatic breakage of deoxyribonucleic acid. II. Purification and properties of endonuclease IV from T4 phage-infected *Escherichia coli*. *J. Biol. Chem.*, 244:6192–6198, 1969.
- [2490] M.J. Sadowsky, Z. Tong, M. de Souza, and L.P. Wackett. AtzC is a new member of the amidohydrolase protein superfamily and is homologous to other atrazine-metabolizing enzymes. *J. Bacteriol.*, 180:152–158, 1998.
- [2491] S.T. Safrany, J.J. Caffrey, X. Yang, M.E. Bembenek, M.B. Moyer, W.A. Burkhardt, and S.B. Shears. A novel context for the 'MutT' module, a guardian of cell integrity, in a diphosphoinositol polyphosphate phosphohydrolase. *EMBO J.*, 17:6599–6607, 1998.
- [2492] S.T. Safrany, S.W. Ingram, J.L. Cartwright, J.R. Falck, A.G. McLennan, L.D. Barnes, and S.B. Shears. The diadenosine hexaphosphate hydrolases from *Schizosaccharomyces pombe* and *Saccharomyces cerevisiae* are homologues of the human diphosphoinositol polyphosphate phosphohydrolase. Overlapping substrate specificities in a MutT-type protein. *J. Biol. Chem.*, 274:21735–21740, 1999.
- [2493] M. Saharay, H. Guo, and J.C. Smith. Catalytic mechanism of cellulose degradation by a cellobiohydrolase, CelS. *PLoS One*, 5:e1294–e1294, 2010.
- [2494] S.A. Saheb. Purification et caractérisation d'une protéase extracellulaire de *Staphylococcus aureus* inhibée par l'E.D.T.A. *Biochimie*, 58:793–804, 1976.
- [2495] Saier and Jr. Molecular phylogeny as a basis for the classification of transport proteins from bacteria, archaea and eukarya. *Adv. Microb. Physiol.*, 40:81–136, 1998.
- [2496] J.-I. Saito, A. Kita, Y. Higuchi, Y. Nagata, A. Ando, and K. Miki. Crystal structure of chitosanase from *Bacillus circulans* MH-K1 at 1.6-Å resolution and its substrate recognition mechanism. *J. Biol. Chem.*, 274:30818–30825, 1999.
- [2497] K. Saito and D.J. Hanahan. A study of the purification and properties of the phospholipase A of *Crotalus adamanteus* venom. *Biochemistry*, 1:521–532, 1962.
- [2498] K. Saito, K. Kondo, I. Kojima, A. Yokota, and F. Tomita. Purification and characterization of 2,6- $\beta$ -D-fructan 6-levanbiohydrolase from *Streptomyces exfoliatus* F3-2. *Appl. Environ. Microbiol.*, 66:252–256, 2000.
- [2499] K. Saito, Y. Oda, F. Tomita, and A. Yokota. Molecular cloning of the gene for 2,6- $\beta$ -D-fructan 6-levanbiohydrolase from *Streptomyces exfoliatus* F3-2. *FEMS Microbiol. Lett.*, 218:265–270, 2003.

- [2500] K. Sakaguchi and S. Murao. A preliminary report on a new enzyme, "penicillin-amidase". *J. Agric. Chem. Soc. Jpn.*, 23:411–411, 1950.
- [2501] K. Sakai, N. Hamada, and Y. Watanabe. Separation of secondary alcohol oxidase and oxidized poly(vinyl alcohol) hydrolase by hydrophobic and dye-ligand chromatographies. *Agric. Biol. Chem.*, 47:153–155, 1983.
- [2502] K. Sakai, N. Hamada, and Y. Watanabe. A new enzyme,  $\beta$ -diketone hydrolase: a component of a poly(vinyl alcohol)-degrading enzyme preparation. *Agric. Biol. Chem.*, 49:1901–1902, 1985.
- [2503] Y. Sakai, M. Furuichi, M. Takahashi, M. Mishima, S. Iwai, M. Shirakawa, and Y. Nakabeppu. A molecular basis for the selective recognition of 2-hydroxy-dATP and 8-oxo-dGTP by human MTH1. *J. Biol. Chem.*, 277:8579–8587, 2002.
- [2504] Y. Sakano, N. Masuda, and T. Kobayashi. Hydrolysis of pullulan by a novel enzyme from *Aspergillus niger*. *Agric. Biol. Chem.*, 35:971–973, 1971.
- [2505] O. Saksela and D.B. Rifkin. Cell-associated plasminogen activation: regulation and physiological functions. *Annu. Rev. Cell Biol.*, 4:93–126, 1988.
- [2506] K. Sakumi, M. Furuichi, T. Tsuzuki, T. Kakuma, S. Kawabata, H. Maki, and M. Sekiguchi. Cloning and expression of cDNA for a human enzyme that hydrolyzes 8-oxo-dGTP, a mutagenic substrate for DNA synthesis. *J. Biol. Chem.*, 268:23524–23530, 1993.
- [2507] H. Sakura, T.Y. Lin, M. Doi, S. Mizutani, and Y. Kawashima. Purification and properties of oxytocinase, a metalloenzyme. *Biochem. Int.*, 2:173–179, 1981.
- [2508] J.M. Sala-Trepat and W.C. Evans. The meta cleavage of catechol by *Azotobacter* species. 4-Oxalocrotonate pathway. *Eur. J. Biochem.*, 20:400–413, 1971.
- [2509] F. Saleh and P.J. White. Metabolism of DD-2,6-diaminopimelic acid by a diaminopimelate-requiring mutant of *Bacillus megaterium*. *J. Gen. Microbiol.*, 115:95–100, 1979.
- [2510] S.P. Salowe, W.J. Krol, D. Iwatareyul, and C.A. Townsend. Elucidation of the order of oxidations and identification of an intermediate in the multistep clavaminic synthase reaction. *Biochemistry*, 30:2281–2292, 1991.
- [2511] G.S. Salvesen and K.M. Boatright. Caspase-8. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 1293–1296. Elsevier, London, 2nd edition, 2004.
- [2512] G.S. Salvesen and K.M. Boatright. Caspase-9. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 1296–1298. Elsevier, London, 2nd edition, 2004.
- [2513] M.E. Salvucci and G.P. Holbrook. Purification and properties of 2-carboxy-D-arabinitol 1-phosphatase. *Plant Physiol.*, 90:679–685, 1989.
- [2514] J. Sampedro, C. Sieiro, G. Revilla, T. Gonzalez-Villa, and I. Zarra. Cloning and expression pattern of a gene encoding an  $\alpha$ -xylosidase active against xyloglucan oligosaccharides from *Arabidopsis*. *Plant Physiol.*, 126:910–920, 2001.
- [2515] S. Sanda, T. Leustek, M. Theisen, R.M. Garavito, and C. Benning. Recombinant *Arabidopsis* SQD1 converts UDP-glucose and sulfite to the sulfolipid head group precursor UDP-sulfoquinovose in vitro. *J. Biol. Chem.*, 276:3941–3946, 2001.
- [2516] F.J. Sandoval and S. Roje. An FMN hydrolase is fused to a riboflavin kinase homolog in plants. *J. Biol. Chem.*, 280:38337–38345, 2005.
- [2517] H. Sanemori, Y. Egi, and T. Kawasaki. Pathway of thiamine pyrophosphate synthesis in *Micrococcus denitrificans*. *J. Bacteriol.*, 126:1030–1036, 1976.
- [2518] R. Sanishvili, A.F. Yakunin, R.A. Laskowski, T. Skarina, E. Evdokimova, A. Doherty-Kirby, G.A. Lajoie, J.M. Thornton, C.H. Arrowsmith, A. Savchenko, A. Joachimiak, and A.M. Edwards. Integrating structure, bioinformatics, and enzymology to discover function: BioH, a new carboxylesterase from *Escherichia coli*. *J. Biol. Chem.*, 278:26039–26045, 2003.
- [2519] K. Sankaran and H.C. Wu. Bacterial prolipoprotein signal peptidase. *Methods Enzymol.*, 248:169–180, 1995.

- [2520] K. Sano, A. Amemura, and T. Harada. Purification and properties of a  $\beta$ -1,6-glucosidase from *Flavobacterium*. *Biochim. Biophys. Acta*, 377:410–420, 1975.
- [2521] M. Sano, K. Hayakawa, , and I. An enzyme releasing lacto-*N*-biose from oligosaccharides. *Proc. Natl. Acad. Sci. USA*, 89:8512–8516, 1992.
- [2522] M. Sano, K. Hayakawa, , and I. Purification and characterization of an enzyme releasing lacto-*N*-biose from oligosaccharides with type 1 chain. *J. Biol. Chem.*, 268:18560–18566, 1993.
- [2523] I. Santamaría, A.M. Pendás, and C. López-Otín. Genomic structure and chromosomal localization of a human cathepsin O gene (CTSO). *Genomics*, 53:231–234, 1998.
- [2524] I. Santamaría, G. Velasco, M. Cazorla, A. Fueyo, E. Campo, and C. López-Otín. Cathepsin L2, a novel human cysteine proteinase produced by breast and colorectal carcinomas. *Cancer Res.*, 58:1624–1630, 1998.
- [2525] I. Santamaría, G. Velasco, A.M. Pendás, A. Paz, and C. López-Otín. Molecular cloning and structural and functional characterization of cathepsin F, a new cysteine proteinase of the papain family with a long propeptide domain. *J. Biol. Chem.*, 274:13800–13809, 1999.
- [2526] I. Velasco Santamaría, Pendás G., Fueyo A.M., López-Otín A., and C. Cathepsin Z, a novel human cysteine proteinase with a short propeptide domain and a unique chromosomal location. *J. Biol. Chem.*, 273:16816–16823, 1998.
- [2527] S.G. Sarafianos, K. Das, C. Tantillo, A.D. Clark, Ding Jr., Whitcomb J., Boyer J.M., Hughes P.L., Arnold S.H., and E. Crystal structure of HIV-1 reverse transcriptase in complex with a polypurine tract RNA:DNA. *EMBO J.*, 20:1449–1461, 2001.
- [2528] L. Sarda and P. Desnuelle. Action de la lipase pancréatique sur les esters en émulsion. *Biochim. Biophys. Acta*, 30:513–521, 1958.
- [2529] S. Sarge, I. Haase, B. Illarionov, D. Laudert, H.P. Hohmann, A. Bacher, and M. Fischer. Catalysis of an essential step in vitamin B<sub>2</sub> biosynthesis by a consortium of broad spectrum hydrolases. *ChemBioChem*, 16:2466–2469, 2015.
- [2530] S. Sarid, A. Berger, and E. Katchalski. Proline iminopeptidase. II. Purification and comparison with iminopeptidase (prolinase). *J. Biol. Chem.*, 237:2207–2212, 1962.
- [2531] A.D. Sarma, P. Serfozo, K. Kahn, and P.A. Tipton. Identification and purification of hydroxyisourate hydrolase, a novel ureide-metabolizing enzyme. *J. Biol. Chem.*, 274:33863–33865, 1999.
- [2532] D.S.R. Sarma, S. Rajalakshmi, and S. Sarma. Studies on the enzymes involved in nicotinamide adenine dinucleotide metabolism in *Aspergillus niger*. *Biochim. Biophys. Acta*, 81:311–322, 1964.
- [2533] A. Sasaki, M. Yamagishi, T. Mega, S. Norioka, S. Natsuka, and S. Hase. Partial purification and characterization of a novel endo- $\beta$ -mannosidase acting on N-linked sugar chains from *Lilium longiflorum* thumb. *J. Biochem. (Tokyo)*, 125:363–367, 1999.
- [2534] K. Sasaki, F. Taura, Y. Shoyama, and S. Morimoto. Molecular characterization of a novel  $\beta$ -glucuronidase from *Scutellaria baicalensis* Georgi. *J. Biol. Chem.*, 275:27466–27472, 2000.
- [2535] M. Sasaki, T. Kato, and S. Iida. Antigenic determinant common to four kinds of thiol proteases of plant origin. *J. Biochem. (Tokyo)*, 74:635–637, 1973.
- [2536] O. Le Saux, T. Gonzalès, and J. Robert-Baudouy. Mutational analysis of the active site of *Pseudomonas fluorescens* pyrrolidone carboxyl peptidase. *J. Bacteriol.*, 178:3308–3313, 1996.
- [2537] A. Savchenko, M. Proudfoot, T. Skarina, A. Singer, O. Litvinova, R. Sanishvili, G. Brown, N. Chirgadze, and A.F. Yakunin. Molecular basis of the antimutagenic activity of the house-cleaning inosine triphosphate pyrophosphatase RdgB from *Escherichia coli*. *J. Mol. Biol.*, 374:1091–1103, 2007.
- [2538] J.W. Savopoulos, P.S. Carter, S. Turconi, G.R. Pettman, E.H. Karran, C.W. Gray, R.V. Ward, O. Jenkins, and C.L. Creasy. Expression, purification, and functional analysis of the human serine protease HtrA2. *Protein. Expr. Purif.*, 19:227–234, 2000.

- [2539] H. Sawada, K. Iwasaki, F. Kihara-Negishi, H. Ariga, and H. Yokosawa. Localization, expression, and the role in fertilization of spermosin, an ascidian sperm trypsin-like protease. *Biochem. Biophys. Res. Commun.*, 222:499–504, 1996.
- [2540] H. Sawada and T. Someno. Substrate specificity of ascidian sperm trypsin-like proteases, spermosin and acrosin. *Mol. Reprod. Dev.*, 45:240–243, 1996.
- [2541] H. Sawada, H. Yokosawa, and S. Ishii. Purification and characterization of two types of trypsin-like enzymes from sperm of the ascidian (Prochordata) *Halocynthia roretzi*. Evidence for the presence of spermosin, a novel acrosin-like enzyme. *J. Biol. Chem.*, 259:2900–2904, 1984.
- [2542] H. Sawada, H. Yokosawa, T. Someno, T. Saino, and S. Ishii. Evidence for the participation of two sperm proteases, spermosin and acrosin, in fertilization of the ascidian, *Halocynthia roretzi*: inhibitory effects of leupeptin analogs on enzyme activities and fertilization. *Dev. Biol.*, 105:246–249, 1984.
- [2543] J. Sawada. Studies on the acid-protease of *Paecilomyces varioti* Bainier TPR-220. Part I. Crystallization of the acid-protease of *Paecilomyces varioti* Bainier TPR-220. *Agric. Biol. Chem.*, 27:677–683, 1963.
- [2544] J. Sawada. The acid-protease of *Paecilomyces varioti*. III. The specificity of the crystalline acid-protease on synthetic substrates. *Agric. Biol. Chem.*, 28:869–875, 1964.
- [2545] T. Sawai and Y. Niwa. Transisomaltosylation activity of a bacterial isomaltodextranase. *Agric. Biol. Chem.*, 39:1077–1083, 1975.
- [2546] T. Sawai, K. Toriyama, and K. Yano. A bacterial dextranase releasing only isomaltose from dextrans. *J. Biochem. (Tokyo)*, 75:105–112, 1974.
- [2547] T. Sawai, T. Yamaki, and T. Ohya. Preparation and some properties of *Arthrobacter globiformis* exo-1,6- $\alpha$ -glucosidase. *Agric. Biol. Chem.*, 40:1293–1299, 1976.
- [2548] C.H. Sawyer. Hydrolysis of choline esters by liver. *Science*, 101:385–386, 1945.
- [2549] C.J. Scandella and A. Kornberg. A membrane-bound phospholipase A<sub>1</sub> purified from *Escherichia coli*. *Biochemistry*, 10:4447–4456, 1971.
- [2550] E. Scarano. The enzymatic deamination of 6-aminopyrimidine deoxyribonucleotides. I. The enzymatic deamination of deoxycytidine 5'-phosphate and of 5-methyldeoxycytidine 5-methyldeoxycytidine 5'-phosphate. *J. Biol. Chem.*, 235:706–713, 1960.
- [2551] E. Scarano, L. Bonaduce, and B. de Petrocellis. The enzymatic deamination of 6-aminopyrimidine deoxyribonucleotides. II. Purification and properties of a 6-aminopyrimidine deoxyribonucleoside 5'-phosphate deaminase from unfertilized eggs of sea urchin. *J. Biol. Chem.*, 235:3556–3561, 1960.
- [2552] P. Schack. Fractionation of proteolytic enzymes of dried papaya latex. Isolation and preliminary characterization of a new proteolytic enzyme. *C. R. Trav. Carlesberg*, 36:67–83, 1967.
- [2553] E. Van Schaftingen, D.R. Davies, and H.-G. Hers. Fructose-2,6-bisphosphatase from rat liver. *Eur. J. Biochem.*, 124:143–149, 1982.
- [2554] H.-P. Schär, W. Holzmann, G.M. Ramos Tombo, and O. Ghisalba. Purification and characterization of *N,N*-dimethylformamidase from *Pseudomonas* DMF 3/3. *Eur. J. Biochem.*, 158:469–475, 1986.
- [2555] O. Schatz, J. Mous, and S.F. Le Grice. HIV-1 RT-associated ribonuclease H displays both endonuclease and 3'—5' exonuclease activity. *EMBO J.*, 9:1171–1176, 1990.
- [2556] R. Schauer. Sialic acids. *Adv. Carbohydr. Chem. Biochem.*, 40:131–234, 1982.
- [2557] G. Schiavo, F. Benfenati, B. Poulain, O. Rossetto, P. Polverino de Laureto, B.R. DasGupta, and C. Montecucco. Tetanus and botulinum-B neurotoxins block neurotransmitter release by proteolytic cleavage of synaptobrevin. *Nature*, 359:832–834, 1992.
- [2558] G. Schiavo, C. Malizio, W.S. Trimble, P. Polverino De Laureto, G. Milan, H. Sugiyama, E.A. Johnson, and C. Montecucco. Botulinum G neurotoxin cleaves VAMP/synaptobrevin at a single Ala-Ala peptide bond. *J. Biol. Chem.*, 269:20213–20216, 1994.

- [2559] G. Schiavo and C. Montecucco. Tetanus and botulism neurotoxins. *Methods Enzymol.*, 248:643–652, 1995.
- [2560] G. Schiavo, O. Rossetto, S. Catsicas, P. Polverino De Laureto, B.R. DasGupta, F. Benfenati, and C. Montecucco. Identification of the nerve-terminal targets of botulinum neurotoxins serotypes A, D and E. *J. Biol. Chem.*, 268:23784–23787, 1993.
- [2561] G. Schiavo, O. Rossetto, A. Santucci, B.R. DasGupta, and C. Montecucco. Botulinum neurotoxins are zinc proteins. *J. Biol. Chem.*, 267:23479–23483, 1992.
- [2562] G. Schiavo, A. Santucci, B.R. DasGupta, P.P. Mehta, J. Jontes, F. Benfenati, M.C. Wilson, and C. Montecucco. Botulinum neurotoxins serotypes A and E cleave SNAP-25 at distinct COOH-terminal peptide bonds. *FEBS Lett.*, 335:99–103, 1993.
- [2563] G. Schiavo, C.C. Shone, O. Rossetto, F.C.G. Alexander, and C. Montecucco. Botulinum neurotoxin serotype F is a zinc endopeptidase specific for VAMP/synaptobrevin. *J. Biol. Chem.*, 268:11516–11519, 1993.
- [2564] S. Schiffer, M. Helm, A. Theobald-Dietrich, R. Giege, and A. Marchfelder. The plant tRNA 3' processing enzyme has a broad substrate spectrum. *Biochemistry*, 40:8264–8272, 2001.
- [2565] S. Schiffer, S. Rösch, and A. Marchfelder. Assigning a function to a conserved group of proteins: the tRNA 3'-processing enzymes. *EMBO J.*, 21:2769–2777, 2002.
- [2566] A. Schirmer, D. Jendrossek, and H.G. Schlegel. Degradation of poly(3-hydroxyoctanoic acid) [P(3HO)] by bacteria: purification and properties of a P(3HO) depolymerase from *Pseudomonas fluorescens* GK13. *Appl. Environ. Microbiol.*, 59:1220–1227, 1993.
- [2567] S. Schlegel-Haueter, P. Hore, K.R. Kerry, and G. Semenza. The preparation of lactase and glucoamylase of rat small intestine. *Biochim. Biophys. Acta*, 258:506–519, 1972.
- [2568] J. Schmid and C. Weissmann. Induction of mRNA for a serine protease and a  $\beta$ -thromboglobulin-like protein in mitogen-stimulated human leukocytes. *J. Immunol.*, 139:250–256, 1987.
- [2569] P.C. Schmid, M.L. Zuzarte-Augustin, and H.H.O. Schmid. Properties of rat liver *N*-acylethanolamine amidohydrolase. *J. Biol. Chem.*, 260:14145–14149, 1985.
- [2570] A.J. Schmidt, D.A. Ryjenkov, and M. Gomelsky. The ubiquitous protein domain EAL is a cyclic diguanylate-specific phosphodiesterase: enzymatically active and inactive EAL domains. *J. Bacteriol.*, 187:4774–4781, 2005.
- [2571] D.M. Schmidt, B.K. Hubbard, and J.A. Gerlt. Evolution of enzymatic activities in the enolase superfamily: functional assignment of unknown proteins in *Bacillus subtilis* and *Escherichia coli* as L-Ala-D/L-Glu epimerases. *Biochemistry*, 40:15707–15715, 2001.
- [2572] E. Schmidt and H.-J. Knackmuss. Chemical structure and biodegradability of halogenated aromatic compounds. Conversion of chlorinated muconic acids into maleoylactic acid. *Biochem. J.*, 192:339–347, 1980.
- [2573] F.J. Schmidt and W.H. McClain. An *Escherichia coli* ribonuclease which removes an extra nucleotide from a biosynthetic intermediate of bacteriophage T<sub>4</sub> proline transfer RNA. *Nucleic Acids Res.*, 5:4129–4139, 1978.
- [2574] G. Schmidt. Nonspecific acid phosphomonoesterases. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 5, pages 37–47. Academic Press, New York, 2nd edition, 1961.
- [2575] B.L. Schneider, A.K. Kiupakis, and L.J. Reitzer. Arginine catabolism and the arginine succinyltransferase pathway in *Escherichia coli*. *J. Bacteriol.*, 180:4278–4286, 1998.
- [2576] S. Schneider and G. Fuchs. Phenylacetyl-CoA:acceptor oxidoreductase, a new  $\alpha$ -oxidizing enzyme that produces phenylglyoxylate. Assay, membrane localization, and differential production in *Thaueria aromatica*. *Arch. Microbiol.*, 169:509–516, 1998.
- [2577] B. Schöbel and W. Pollmann. Isolation and characterization of a chlorogenic acid esterase from *Aspergillus niger*. *Z. Naturforsch. C: Biosci.*, 35:209–212, 1980.
- [2578] B. Schöbel and W. Pollmann. Weitere Charakterisierung einer Chlorogensäure - Hydrolase aus *Aspergillus niger*. *Z. Naturforsch. C: Biosci.*, 35:699–701, 1980.

- [2579] M. Schober, T. Knaus, M. Toesch, P. Macheroux, U. Wagner, and K. Faber. The substrate spectrum of the inverting *sec*-alkylsulfatase Pisa1. *Adv. Synth. Catal.*, 354:1737–1742, 2012.
- [2580] L. Schoefer, A. Braune, and M. Blaut. Cloning and expression of a phloretin hydrolase gene from *Eubacterium ramulus* and characterization of the recombinant enzyme. *Appl. Environ. Microbiol.*, 70:6131–6137, 2004.
- [2581] I.C. Schoenhofen, D.J. McNally, J.R. Brisson, and S.M. Logan. Elucidation of the CMP-pseudaminic acid pathway in *Helicobacter pylori*: synthesis from UDP-*N*-acetylglucosamine by a single enzymatic reaction. *Glycobiology*, 16:8C–14C, 2006.
- [2582] I.C. Schoenhofen, E. Vinogradov, D.M. Whitfield, J.R. Brisson, and S.M. Logan. The CMP-legionaminic acid pathway in *Campylobacter*: biosynthesis involving novel GDP-linked precursors. *Glycobiology*, 19:715–725, 2009.
- [2583] R. Scholtz, T. Leisinger, F. Suter, and A.M. Cook. Characterization of 1-chlorohexane halidohydrolase, a dehalogenase of wide substrate range from an *Arthrobacter sp.* *J. Bacteriol.*, 169:5016–5021, 1987.
- [2584] J.E. Schriener, W. Yi, , and S.L. cDNA and genomic cloning of human palmitoyl-protein thioesterase (PPT), the enzyme defective in infantile neuronal ceroid lipofuscinosis. *Genomics*, 34:317–322, 1996.
- [2585] H.C. Schröder, K. Dose, R.K. Zahn, and E.G. Müller. Isolation and characterization of the novel polyadenylate- and polyuridylylate-degrading acid endoribonuclease V from calf thymus. *J. Biol. Chem.*, 255:5108–5112, 1980.
- [2586] H.C. Schröder, R.K. Zahn, K. Dose, and E.G. Müller. Purification and characterization of a poly(A)-specific exoribonuclease from calf thymus. *J. Biol. Chem.*, 255:4535–4538, 1980.
- [2587] H. Schübel, J. Stöckigt, R. Feicht, and H. Simon. Partial-purification and characterization of raucaffricine  $\beta$ -D-glucosidase from plant cell-suspension cultures of *Rauwolfia serpentina* benth. *Helv. Chim. Acta*, 69:538–547, 1986.
- [2588] J. Schultz and K. Mosbach. Studies on lichen enzymes. Purification and properties of an orsellinate depside hydrolase obtained from *Lasallia pustulata*. *Eur. J. Biochem.*, 22:153–157, 1971.
- [2589] S.J. Schultz and J.J. Champoux. RNase H activity: structure, specificity, and function in reverse transcription. *Virus Res.*, 134:86–103, 2008.
- [2590] S.J. Schultz, M. Zhang, and J.J. Champoux. Recognition of internal cleavage sites by retroviral RNases H. *J. Mol. Biol.*, 344:635–652, 2004.
- [2591] S.J. Schultz, M. Zhang, C.D. Kelleher, and J.J. Champoux. Analysis of plus-strand primer selection, removal, and reutilization by retroviral reverse transcriptases. *J. Biol. Chem.*, 275:32299–32309, 2000.
- [2592] O. Schulz, H.F. Sewell, and F. Shakib. A sensitive fluorescent assay for measuring the cysteine protease activity of Der p 1, a major allergen from the dust mite *Dermatophagoides pteronyssinus*. *Mol. Pathol.*, 51:222–224, 1998.
- [2593] O. Schulz, H.F. Sewell, and F. Shakib. Proteolytic cleavage of CD25, the  $\alpha$  subunit of the human T cell interleukin 2 receptor, by Der p 1, a major mite allergen with cysteine protease activity. *J. Exp. Med.*, 187:271–275, 1998.
- [2594] M. Schurmann, R.M. Demming, M. Krewing, J. Rose, J.H. Wubbeler, and A. Steinbuchel. Identification of 3-sulfino-propionyl coenzyme A (CoA) desulfonases within the Acyl-CoA dehydrogenase superfamily. *J. Bacteriol.*, 196:882–893, 2014.
- [2595] M. Schurmann, A. Deters, J.H. Wubbeler, and A. Steinbuchel. A novel 3-sulfino-propionyl coenzyme A (3SP-CoA) desulfonase from *Advenella mimigardefordensis* strain DPN7T acting as a key enzyme during catabolism of 3,3'-dithiodipropionic acid is a member of the acyl-CoA dehydrogenase superfamily. *J. Bacteriol.*, 195:1538–1551, 2013.
- [2596] A.M. Schweingruber, J. Dlugonski, E. Edenharter, and M.E. Schweingruber. Thiamine in *Schizosaccharomyces pombe*: dephosphorylation, intracellular pool, biosynthesis and transport. *Curr. Genet.*, 19:249–254, 1991.
- [2597] K. Schwibbert, A. Marin-Sanguino, I. Bagyan, G. Heidrich, G. Lentzen, H. Seitz, M. Rampp, S.C. Schuster, H.P. Klenk, F. Pfeiffer, D. Oesterhelt, and H.J. Kunte. A blueprint of ectoine metabolism from the genome of the industrial producer *Halomonas elongata* DSM 2581 T. *Environ. Microbiol.*, 13:1973–1994, 2011.
- [2598] S. Schwimmer and A.K. Balls. Isolation and properties of crystalline  $\alpha$ -amylase from germinated barley. *J. Biol. Chem.*, 179:1063–1074, 1949.

- [2599] F.P. Seebeck. *In vitro* reconstitution of mycobacterial ergothioneine biosynthesis. *J. Am. Chem. Soc.*, 132:6632–6633, 2010.
- [2600] E. Seemüller, A. Lupas, D. Stock, J. Löwe, R. Huber, and W. Baumeister. Proteasome from *Thermoplasma acidophilum*: a threonine protease. *Science*, 268:579–582, 1995.
- [2601] J.L. Seffernick, A. Aleem, J.P. Osborne, G. Johnson, M.J. Sadowsky, and L.P. Wackett. Hydroxyatrazine *N*-ethylaminohydrolase (AtzB): an amidohydrolase superfamily enzyme catalyzing deamination and dechlorination. *J. Bacteriol.*, 189:6989–6997, 2007.
- [2602] J.L. Seffernick, M.L. Souza, M.J. Sadowsky, and L.P. Wackett. Melamine deaminase and atrazine chlorohydrolase: 98 percent identical but functionally different. *J. Bacteriol.*, 183:2405–2410, 2001.
- [2603] H.L. Segal and B.M. Brenner. 5'-Nucleotidase of rat liver microsomes. *J. Biol. Chem.*, 235:471–474, 1960.
- [2604] N.G. Seidah and M. Chrétien. Proprotein and prohormone convertases of the subtilisin family: recent developments and future perspectives. *Trends Endocrinol. Metab.*, 3:133–140, 1992.
- [2605] N.G. Seidah and M. Chrétien. Pro-protein convertases of the subtilisin/kexin family. *Methods Enzymol.*, 244:175–188, 1994.
- [2606] N.G. Seidah and M. Chrétien. Proprotein convertase SKI-1/S1P. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, volume 2, pages 1845–1847. Elsevier, London, 2nd edition, 2004.
- [2607] N.G. Seidah, L. Gaspar, P. Mion, M. Marcinkiewicz, M. Mbikay, and M. Chrétien. cDNA sequence of two distinct pituitary proteins homologous to Kex2 and furin gene products: tissue-specific mRNAs encoding candidates for pro-hormone processing proteinases. *DNA Cell Biol.*, 9:415–424, 1990.
- [2608] N.G. Seidah, R. Ladenheim, M. Mbikay, J. Hamelin, G. Lutfalla, F. Rougeon, C. Lazure, and M. Chrétien. The cDNA structure of rat plasma kallikrein. *DNA*, 8:563–574, 1989.
- [2609] H. Sekine. Neutral proteinases I and II of *Aspergillus sojae* action on various substrates. *Agric. Biol. Chem.*, 40:703–709, 1976.
- [2610] T. Sekiya, R. Contreras, T. Takeya, and H.G. Khorana. Total synthesis of a tyrosine suppressor transfer RNA gene. XVII. Transcription, *in vitro*, of the synthetic gene and processing of the primary transcript to transfer RNA. *J. Biol. Chem.*, 254:5802–5816, 1979.
- [2611] J.H. Seol, S.K. Woo, E.M. Jung, S.J. Yoo, C.S. Lee, K.J. Kim, K. Tanaka, A. Ichihara, D.B. Ha, and C.H. Chung. Protease Do is essential for survival of *Escherichia coli* at high temperatures: its identity with the *htrA* gene product. *Biochem. Biophys. Res. Commun.*, 176:730–736, 1991.
- [2612] R.C. Sergott, L.J. Debeer, and M.J. Bessman. On the regulation of a bacterial deoxycytidylate deaminase. *J. Biol. Chem.*, 246:7755–7758, 1971.
- [2613] F. Serventi, I. Ramazzina, I. Lamberto, V. Puggioni, R. Gatti, and R. Percudani. Chemical basis of nitrogen recovery through the ureide pathway: formation and hydrolysis of *S*-ureidoglycine in plants and bacteria. *ACS Chem. Biol.*, 5:203–214, 2010.
- [2614] A. Sevanian and L.L. McLeod. Catalytic properties and inhibition of hepatic cholesterol-epoxide hydrolase. *J. Biol. Chem.*, 261:54–59, 1986.
- [2615] N. Shah, D.A. Kuntz, and D.R. Rose. Golgi  $\alpha$ -mannosidase II cleaves two sugars sequentially in the same catalytic site. *Proc. Natl Acad. Sci. USA*, 105:9570–9575, 2008.
- [2616] R. Shah, R. Cosstick, and S.C. West. The RuvC protein dimer resolves Holliday junctions by a dual incision mechanism that involves base-specific contacts. *EMBO J.*, 16:1464–1472, 1997.
- [2617] P.N. Shaklee, J.H. Glaser, and H.E. Conrad. A sulfatase specific for glucuronic acid 2-sulfate residues in glycosaminoglycans. *J. Biol. Chem.*, 260:9146–9149, 1985.
- [2618] J.D. Shannon, E.N. Baramova, J.B. Bjarnason, and F.W. Fox. Amino acid sequence of a *Crotalus atrox* venom metallo-proteinase which cleaves type IV collagen and gelatin. *J. Biol. Chem.*, 264:11575–11583, 1989.



- [2619] N. Shapir, G. Cheng, M.J. Sadowsky, and L.P. Wackett. Purification and characterization of TrzF: biuret hydrolysis by allophanate hydrolase supports growth. *Appl. Environ. Microbiol.*, 72:2491–2495, 2006.
- [2620] N. Shapir, J.P. Osborne, G. Johnson, M.J. Sadowsky, and L.P. Wackett. Purification, substrate range, and metal center of AtzC: the *N*-isopropylammelide aminohydrolase involved in bacterial atrazine metabolism. *J. Bacteriol.*, 184:5376–5384, 2002.
- [2621] N. Shapir, M.J. Sadowsky, and L.P. Wackett. Purification and characterization of allophanate hydrolase (AtzF) from *Pseudomonas* sp. strain ADP. *J. Bacteriol.*, 187:3731–3738, 2005.
- [2622] B. Shapiro. Purification and properties of a lysolecithinase from pancreas. *Biochem. J.*, 53:663–666, 1953.
- [2623] S.D. Shapiro, G.L. Griffin, D.J. Gilbert, N.A. Jenkins, N.G. Copeland, H.G. Welgus, R.M. Senior, and T.J. Ley. Molecular cloning, chromosomal localization, and bacterial expression of a murine macrophage metalloelastase. *J. Biol. Chem.*, 267:4664–4671, 1992.
- [2624] S.D. Shapiro, D.K. Kobayashi, and T.J. Ley. Cloning and characterization of a unique elastolytic metalloproteinase produced by human alveolar macrophages. *J. Biol. Chem.*, 268:23824–23829, 1993.
- [2625] R.G. Sheiknejad and P.N. Srivastava. Isolation and properties of a phosphatidylcholine-specific phospholipase C from bull seminal plasma. *J. Biol. Chem.*, 261:7544–7549, 1986.
- [2626] W. Shen, T.S. Fletcher, and C. Largman. Primary structure of human pancreatic protease E determined by sequence analysis of the cloned mRNA. *Biochemistry*, 26:3447–3452, 1987.
- [2627] R.F. Sherwood, R.G. Melton, and S.A. Alwan. Purification and properties of carboxypeptidase G<sub>2</sub> from *Pseudomonas* sp. strain RS-16. *Eur. J. Biochem.*, 148:447–453, 1985.
- [2628] Z.Z. Shi, B. Han, G.M. Habib, M.M. Matzuk, and M.W. Lieberman. Disruption of  $\gamma$ -glutamyl leukotrienase results in disruption of leukotriene D<sub>4</sub> synthesis *in vivo* and attenuation of the acute inflammatory response. *Mol. Cell Biol.*, 21:5389–5395, 2001.
- [2629] I. Shibuya and A. A. Benson. Hydrolysis of  $\alpha$ -sulphoquinovosides by  $\beta$ -galactosidase. *Nature*, 192:1186–1187, 1961.
- [2630] T. Shida, H. Iwasaki, A. Saito, Y. Kyogoku, and H. Shinagawa. Analysis of substrate specificity of the RuvC holliday junction resolvase with synthetic Holliday junctions. *J. Biol. Chem.*, 271:26105–26109, 1996.
- [2631] Y. Shikama, M. Yamada, and T. Miyashita. Caspase-8 and caspase-10 activate NF- $\kappa$ B through RIP, NIK and IKK $\alpha$  kinases. *Eur. J. Immunol.*, 33:1998–2006, 2003.
- [2632] K. Shimada, M. Takahashi, and K. Tanzawa. Cloning and functional expression of endothelin-converting enzyme from rat endothelial cells. *J. Biol. Chem.*, 269:18275–18278, 1994.
- [2633] C. Shimizu, S. Yoshida, M. Shibata, K. Kato, Y. Momota, K. Matsumoto, T. Shiosaka, R. Midorikawa, T. Kamachi, A. Kawabe, and S. Shiosaka. Characterization of recombinant and brain neuropsin, a plasticity-related serine protease. *J. Biol. Chem.*, 273:11189–11196, 1998.
- [2634] S. Shimizu, J. Ogawa, M.C.-M. Chung, , and H. Purification and characterization of a novel enzyme, arylalkyl acylamidase, from *Pseudomonas putida* Sc2. *Eur. J. Biochem.*, 209:375–382, 1992.
- [2635] Y. Shimura, H. Sakano, and F. Nagawa. Specific ribonucleases involved in processing of tRNA precursors of *Escherichia coli*. Partial purification and some properties. *Eur. J. Biochem.*, 86:267–281, 1978.
- [2636] H.J. Shin and J.L. Mego. A rat liver lysosomal membrane flavin-adenine dinucleotide phosphohydrolase: purification and characterization. *Arch. Biochem. Biophys.*, 267:95–103, 1988.
- [2637] I. Shin, R. Percudani, and S. Rhee. Structural and functional insights into (*S*)-ureidoglycine aminohydrolase, key enzyme of purine catabolism in *Arabidopsis thaliana*. *J. Biol. Chem.*, 287:18796–18805, 2012.
- [2638] S. Shinano and K. Fukushima. Studies on lotus seed protease. Part III. Some physicochemical and enzymic properties. *Agric. Biol. Chem.*, 35:1488–1494, 1971.
- [2639] W.E. Shine, M. Mancha, and P.K. Stumpf. Fat metabolism in higher plants. The function of acyl thioesterases in the metabolism of acyl-coenzymes A and acyl-acyl carrier proteins. *Arch. Biochem. Biophys.*, 172:110–116, 1976.

- [2640] T. Shinohara, C. Sakurada, T. Suzuki, O. Takeuchi, W. Campbell, S. Ikeda, N. Okada, and H. Okada. Procarboxypeptidase R cleaves bradykinin following activation. *Int. Arch. Allergy Immunol.*, 103:400–404, 1994.
- [2641] H. Shiokawa and L. Noda. Isolation and crystallization of acyl phosphatase from rabbit muscle. *J. Biol. Chem.*, 245:669–673, 1970.
- [2642] S. Shipkowski and J.E. Brenchley. Bioinformatic, genetic, and biochemical evidence that some glycoside hydrolase family 42  $\beta$ -galactosidases are arabinogalactan type I oligomer hydrolases. *Appl. Environ. Microbiol.*, 72:7730–7738, 2006.
- [2643] Y. Shirasu, H. Yoshida, S. Matsuki, K. Takemura, N. Ikeda, Y. Shimada, T. Ozawa, T. Mikayama, H. Iijima, A. Ishida, Y. Sato, Y. Tamai, J. Tanaka, and H. Ikenaga. Molecular cloning and expression in *Escherichia coli* of a cDNA encoding human pancreatic elastase 2. *J. Biochem. (Tokyo)*, 102:1555–1563, 1987.
- [2644] K. Shishido, M. Kato, and Y. Ikeda. Isolation of thymidylic acid-rich fragments from double-stranded deoxyribonucleic acids. *J. Biochem. (Tokyo)*, 65:479–481, 1968.
- [2645] J. Shlomain and A. Kornberg. Deoxyuridine triphosphatase of *Escherichia coli*. Purification, properties, and use as a reagent to reduce uracil incorporation into DNA. *J. Biol. Chem.*, 253:3305–3312, 1978.
- [2646] D.M. Shotton. Elastase. *Methods Enzymol.*, 19:113–140, 1970.
- [2647] M. Shoyab, T.C. Warren, and G.J. Todaro. Isolation and characterization of an ester hydrolase active on phorbol diesters from murine liver. *J. Biol. Chem.*, 256:12529–12534, 1981.
- [2648] A.K. Shukla and R. Schauer. High performance liquid chromatography of enzymes of sialic acid metabolism. *Hoppe-Seyler's Z. Physiol. Chem.*, 363:1039–1040, 1982.
- [2649] N.P. Shull, S.L. Spinelli, and E.M. Phizicky. A highly specific phosphatase that acts on ADP-ribose 1''-phosphate, a metabolite of tRNA splicing in *Saccharomyces cerevisiae*. *Nucleic Acids Res.*, 33:650–660, 2005.
- [2650] L. Shuster and N.O. Kaplan. A specific b nucleotidase. *J. Biol. Chem.*, 201:535–546, 1953.
- [2651] V.K. Shvyadas, I.Yu. Galaev, and E.V. Kozlova. Preparation and characterization of L- $\alpha$ -aminocaprolactam hydrolase from cells of *Cryptococcus laurentii*. *Biochemistry (Moscow)*, 49:1268–1273, 1984.
- [2652] D.C. Sideris and E.G. Fragoulis. Purification and characterization of a ribonuclease specific for poly(U) and poly(C) from the larvae of *Ceratitidis capitata*. *Eur. J. Biochem.*, 164:309–315, 1987.
- [2653] W. Sidler, E. Niederer, F. Suter, and H. Zuber. The primary structure of *Bacillus cereus* neutral proteinase and comparison with thermolysin and *Bacillus subtilis* neutral proteinase. *Biol. Chem. Hoppe-Seyler*, 367:643–657, 1986.
- [2654] W. Sidorowicz, W.-C. Hsia, M. Maslej-Zownir, and F.J. Behal. Multiple molecular forms of human alanine aminopeptidase: immunochemical properties. *Clin. Chim. Acta*, 107:245–256, 1980.
- [2655] A.R. Sielecki, K. Hayakawa, M. Fujinaga, M.E.P. Murphy, M. Fraser, A.K. Muir, C.T. Carilli, J.A. Lewicki, J.D. Baxter, and M.N.G. James. Structure of recombinant human renin, a target for cardiovascular-active drugs, at 2.5 Å resolution. *Science*, 243:1346–1351, 1989.
- [2656] A.R. Sielecki, W.A. Hendrickson, C.G. Broughton, L.T.J. Delbaere, G.D. Brayer, and M.N.G. James. Protein structure refinement: *Streptomyces griseus* serine protease A at 1.8 Å resolution. *J. Mol. Biol.*, 134:781–804, 1979.
- [2657] H. Sigrist, P. Ronner, and G. Semenza. A hydrophobic form of the small-intestinal sucrase-isomaltase complex. *Biochim. Biophys. Acta*, 406:433–446, 1975.
- [2658] A. Sik, B.J. Passer, E.V. Koonin, and L. Pellegrini. Self-regulated cleavage of the mitochondrial intramembrane-cleaving protease PARL yields P $\beta$ , a nuclear-targeted peptide. *J. Biol. Chem.*, 279:15323–15329, 2004.
- [2659] M.A.G. Sillero, R. Villalba, A. Morena, M. Quintanilla, C.D. Lobatón, and A. Sillero. Dinucleosidetriphosphatase from rat liver. Purification and properties. *Eur. J. Biochem.*, 76:331–337, 1977.
- [2660] S. Silver and G. Ji. Newer systems for bacterial resistance to toxic heavy metals. *Environ. Health Perspect.* 102, Suppl., 3:107–113, 1994.

- [2661] S. Silver, T.K. Misra, and R.A. Laddaga. DNA sequence analysis of bacterial toxic heavy metal resistance. *Biol. Trace Elem. Res.*, 21:145–163, 1989.
- [2662] M. Silverberg and A.P. Kaplan. Human Hageman factor and its fragments. *Methods Enzymol.*, 163:68–80, 1988.
- [2663] M. Silverberg and A.P. Kaplan. Prekallikrein. *Methods Enzymol.*, 163:85–95, 1988.
- [2664] R.B. Sim. The human complement system serine proteases  $C^{1r}$  and  $C^{1s}$  and their proenzymes. *Methods Enzymol.*, 80:26–42, 1981.
- [2665] G. Simmons, M. Bundalian, I. Theodor, J. Martinoli, and H. Pirkle. Action of crotalase, an enzyme with thrombin-like and kallikrein-like specificities, on tripeptide nitroanilide derivatives. *Thromb. Res.*, 40:555–561, 1985.
- [2666] W.H. Simmons and R. Walter. Carboxamidopeptidase: purification and characterization of a neurohypophyseal hormone inactivating peptidase from toad skin. *Biochemistry*, 19:39–48, 1980.
- [2667] M.M. Simon, H. Hoschützky, U. Fruth, H.-G. Simon, and M.D. Kramer. Purification and characterization of a T cell specific serine proteinase (TSP-1) from cloned cytolytic T lymphocytes. *EMBO J.*, 5:3267–3274, 1986.
- [2668] M.F. Singer and J.S. Fruton. Some properties of beef spleen phosphoamidase. *J. Biol. Chem.*, 229:111–119, 1957.
- [2669] T.P. Singer and B.H.J. Hofstee. Studies on wheat germ lipase. I. Methods of estimation, purification and general properties of the enzyme. *Arch. Biochem.*, 18:229–243, 1948.
- [2670] T.P. Singer and B.H.J. Hofstee. Studies on wheat germ lipase. II. Kinetics. *Arch. Biochem.*, 18:245–259, 1948.
- [2671] R.M.M. Singh and E. Adams. Enzymatic deamination of  $\Delta^1$ -pyrroline-4-hydroxy-2-carboxylate to 2,5-dioxovalerate ( $\alpha$ -ketoglutaric semialdehyde). *J. Biol. Chem.*, 240:4344–4351, 1965.
- [2672] R.M.M. Singh and E. Adams. Isolation and identification of 2,5-dioxovalerate, an intermediate in the bacterial oxidation of hydroxyproline. *J. Biol. Chem.*, 240:4352–4356, 1965.
- [2673] A.K. Sinha and G.C. Chatterjee. Metabolism of pyrithiamine by the pyrithiamine-requiring mutant of *Staphylococcus aureus*. *Biochem. J.*, 107:165–169, 1968.
- [2674] C.F. Sio, L.G. Otten, R.H. Cool, S.P. Diggle, P.G. Braun, R. Bos, M. Daykin, M. Cámara, P. Williams, and W.J. Quax. Quorum quenching by an *N*-acyl-homoserine lactone acylase from *Pseudomonas aeruginosa* PAO1. *Infect. Immun.*, 74:1673–1682, 2006.
- [2675] S. Sivikami and A.N. Radhakrishnan. Purification of rabbit intestinal glucoamylase by affinity chromatography on Sephadex G-200. *Indian J. Biochem. Biophys.*, 10:283–284, 1973.
- [2676] I. Sjöholm. Biochemical studies on oxytocin and oxytocinase. *Acta Pharm. Suec.*, 4:81–96, 1967.
- [2677] I. Sjöholm and L. Yman. Degradation of oxytocin, lysine vasopressin, angiotensin II, and angiotensin II amide by oxytocinase (cystine aminopeptidase). *Acta Pharm. Suec.*, 4:65–76, 1967.
- [2678] H. Sjöström, O. Norén, L. Christiansen, H. Wacker, and G. Semenza. A fully active, two-active-site, single-chain sucrase-isomaltase from pig small intestine. Implications for the biosynthesis of a mammalian integral stalked membrane protein. *J. Biol. Chem.*, 255:11332–11338, 1980.
- [2679] H. Sjöström, O. Norén, L. Jeppesen, M. Staun, B. Svensson, and L. Christiansen. Purification of different amphiphilic forms of a microvillus aminopeptidase from pig small intestine using immunoadsorbent chromatography. *Eur. J. Biochem.*, 88:503–511, 1978.
- [2680] H. Sjöström, O. Norén, and L. Josefsson. Purification and specificity of pig intestinal prolidase. *Biochim. Biophys. Acta*, 327:457–470, 1973.
- [2681] R.A. Skidgel. Basic carboxypeptidases: regulators of peptide hormone activity. *Trends Pharmacol. Sci.*, 9:301–303, 1988.
- [2682] R.A. Skidgel. Basic carboxypeptidases: Regulators of peptide hormone activity. *Trends Pharmacol. Sci.*, 9:303–304, 1988.

- [2683] R.A. Skidgel, R.M. Davis, and F. Tan. Human carboxypeptidase M. Purification and characterization of membrane-bound carboxypeptidase that cleaves peptide hormones. *J. Biol. Chem.*, 264:2236–2241, 1989.
- [2684] M.T. Skoog, S. Mehdi, J.S., Bey Wiseman, , and C. 1 esterase and acrosin, for peptide *p*-nitroanilide substrates. *Biochim. Biophys. Acta*, 996:89–94, 1989.
- [2685] H. Skovbjerg, O. Norén, H. Sjöström, E.M. Danielsen, and B.S. Enevoldsen. Further characterization of intestinal lactase/phlorizin hydrolase. *Biochim. Biophys. Acta*, 707:89–97, 1982.
- [2686] H. Skovbjerg, H. Sjöström, and O. Norén. Purification and characterization of amphiphilic lactase-phlorizin hydrolase from human small-intestine. *Eur. J. Biochem.*, 114:653–661, 1981.
- [2687] A. Skytt, M. Strömqvist, and T. Egelrud. Primary substrate specificity of recombinant human stratum corneum chymotryptic enzyme. *Biochem. Biophys. Res. Commun.*, 211:586–589, 1995.
- [2688] J.M. Slack, J. Kuzio, and P. Faulkner. Characterization of V-cath, a cathepsin L-like proteinase expressed by the baculovirus *Autographa californica* multiple nuclear polyhedrosis-virus. *J. Gen. Virol.*, 76:1091–1098, 1995.
- [2689] E.E. Slater. Renin. *Methods Enzymol.*, 80:427–442, 1981.
- [2690] S.N. Slilaty and J.W. Little. Lysine-156 and serine-119 are required for LexA repressor cleavage: a possible mechanism. *Proc. Natl. Acad. Sci. USA*, 84:3987–3991, 1987.
- [2691] W.S. Sly and E.R. Stadtman. Formate metabolism. I. Formyl coenzyme A, an intermediate in the formate-dependent decomposition of acetyl phosphate in *Clostridium kluyveri*. *J. Biol. Chem.*, 238:2632–2638, 1963.
- [2692] G.D. Small and C. Cooper. Purification and properties of nucleoside tetraphosphate hydrolase from rabbit muscle. *Biochemistry*, 5:14–26, 1966.
- [2693] S.P. Smeekens, A.S. Avruch, J. LaMendola, S.J. Chan, and D.F. Steiner. Identification of a cDNA encoding a second putative prohormone convertase related to PC2 in AtT20 cells and islets of Langerhans. *Proc. Natl. Acad. Sci. USA*, 88:340–344, 1991.
- [2694] S.P. Smeekens and D.F. Steiner. Identification of a human insulinoma cDNA encoding a novel mammalian protein structurally related to the yeast dibasic processing protease Kex2. *J. Biol. Chem.*, 265:2997–3000, 1990.
- [2695] M.J. Smeulders, T.R. Barends, A. Pol, A. Scherer, M.H. Zandvoort, A. Udvarhelyi, A.F. Khadem, A. Menzel, J. Hermans, R.L. Shoeman, H.J. Wessels, L.P. van den Heuvel, L. Russ, I. Schlichting, M.S. Jetten, and H.J. Op den Camp. Evolution of a new enzyme for carbon disulphide conversion by an acidothermophilic archaeon. *Nature*, 478:412–416, 2011.
- [2696] C.L. Smith, J. Ghosh, J.S. Elam, J.S. Pinkner, S.J. Hultgren, M.G. Caparon, and T. Ellenberger. Structural basis of *Streptococcus pyogenes* immunity to its NAD<sup>+</sup> glycohydrolase toxin. *Structure*, 19:192–202, 2011.
- [2697] D.W.E. Smith and B.N. Ames. Phosphoribosyladenosine monophosphate, an intermediate in histidine biosynthesis. *J. Biol. Chem.*, 240:3056–3063, 1965.
- [2698] J.M. Smith, K. Harrison, and J. Colby. Purification and characterization of D-2-haloacid dehalogenase from *Pseudomonas putida* strain AJ1/23. *J. Gen. Microbiol.*, 136:881–886, 1990.
- [2699] R.A.G. Smith, J. Green, and P.H. Kopper. The purification and properties of a fibrinolytic neutral metalloendopeptidase from *Streptococcus faecalis*. *Arch. Biochem. Biophys.*, 202:629–638, 1980.
- [2700] S.W. Smith, S.B. Weiss, and E.P. Kennedy. The enzymatic dephosphorylation of phosphatidic acids. *J. Biol. Chem.*, 228:915–922, 1957.
- [2701] T.A. Smith. Agmatine iminohydrolase in maize. *Phytochemistry*, 8:2111–2117, 1969.
- [2702] T.K. Smith, A. Crossman, C.N. Borissow, M.J. Paterson, A. Dix, J.S. Brimacombe, and M.A.J. Ferguson. Specificity of GlcNAc-PI de-*N*-acetylase of GPI biosynthesis and synthesis of parasite-specific suicide substrate inhibitors. *EMBO J.*, 20:3322–3332, 2001.
- [2703] S.V. Smulevitch, A.L. Osterman, O.V. Galperina, M.V. Matz, O.P. Zagnitko, R.M. Kadyrov, I.A. Tsaplina, N.V. Grishin, G.G. Chestukhina, and V.M. Stepanov. Molecular cloning and primary structure of *Thermoactinomyces vulgaris* carboxypeptidase T: a metalloenzyme endowed with dual substrate specificity. *FEBS Lett.*, 291:75–78, 1991.

- [2704] E.J. Snijder, A.L. Wassenaar, L.C. van Dinten, W.J. Spaan, and A.E. Gorbalenya. The arterivirus nsp4 protease is the prototype of a novel group of chymotrypsin-like enzymes, the 3C-like serine proteases. *J. Biol. Chem.*, 271:4864–4871, 1996.
- [2705] S.H. Snyder, O.L. Silva, and M.W. Kies. The mammalian metabolism of L-histidine. IV. Purification and properties of imidazolone propionic acid hydrolase. *J. Biol. Chem.*, 236:2996–2998, 1961.
- [2706] C. Sobhy. Regulation of fatty acid synthetase activity. The 4'-phosphopantetheine hydrolase of rat liver. *J. Biol. Chem.*, 254:8561–8566, 1979.
- [2707] K. Soda, T. Kurihara, J.-Q. Liu, V. Nardi-Dei, C. Park, M. Miyagi, S. Tsunasawa, and N. Esaki. Bacterial 2-haloacid dehalogenases: Structures and catalytic properties. *Pure Appl. Chem.*, 68:2097–2103, 1996.
- [2708] M.L. Sogin, B. Pace, and N.R. Pace. Partial purification and properties of a ribosomal RNA maturation endonuclease from *Bacillus subtilis*. *J. Biol. Chem.*, 252:1350–1357, 1977.
- [2709] A. Sohail, J. Klapacz, M. Samaranyake, A. Ullah, and A.S. Bhagwat. Human activation-induced cytidine deaminase causes transcription-dependent, strand-biased C to U deaminations. *Nucleic Acids Res.*, 31:2990–2994, 2003.
- [2710] K. Solaas, R.J. Sletta, O. Soreide, and B.F. Kase. Presence of choloyl- and chenodeoxycholoyl-coenzyme A thioesterase activity in human liver. *Scand. J. Clin. Lab. Invest.*, 60:91–102, 2000.
- [2711] D.F. Soler and R.B. Harris. Continuous fluorogenic substrates for atrial dipeptidyl carboxylhydrolase. Importance of Ser in the P1 position. *Int. J. Peptide Protein Res.*, 32:35–40, 1988.
- [2712] D.F. Soler and R.B. Harris. Atrial dipeptidyl carboxylhydrolase is a zinc-metallo proteinase which possesses tripeptidyl carboxylhydrolase activity. *Peptides (Fayetteville)*, 10:63–68, 1989.
- [2713] M.R. Sommer and B. Jochimsen. Identification of enzymes involved in indole-3-acetic acid degradation. *Plant Soil*, 186:143–149, 1996.
- [2714] B.H. Song and J. Neuhard. Chromosomal location, cloning and nucleotide sequence of the *Bacillus subtilis* cdd gene encoding cytidine/deoxycytidine deaminase. *Mol. Gen. Genet.*, 216:462–468, 1989.
- [2715] E.K. Song, H. Kim, H.K. Sung, and J. Cha. Cloning and characterization of a levanbiohydrolase from *Microbacterium laevaniformans* ATCC 15953. *Gene*, 291:45–55, 2002.
- [2716] H.N. Song, T.Y. Jung, J.T. Park, B.C. Park, P.K. Myung, W. Boos, E.J. Woo, and K.H. Park. Structural rationale for the short branched substrate specificity of the glycogen debranching enzyme GlgX. *Proteins*, 78:1847–1855, 2010.
- [2717] L. Song. A soluble form of phosphatase in *Saccharomyces cerevisiae* capable of converting farnesyl diphosphate into *E,E*-farnesol. *Appl. Biochem. Biotechnol.*, 128:149–158, 2006.
- [2718] L.X. Song and L.D. Fricker. Purification and characterization of carboxypeptidase D, a novel carboxypeptidase E-like enzyme, from bovine pituitary. *J. Biol. Chem.*, 270:25007–25013, 1995.
- [2719] L.X. Song and L.D. Fricker. Tissue distribution and characterization of soluble and membrane-bound forms of metallo-carboxypeptidase D. *J. Biol. Chem.*, 271:28884–28889, 1996.
- [2720] M.G. Song, Y. Li, and M. Kiledjian. Multiple mRNA decapping enzymes in mammalian cells. *Mol. Cell*, 40:423–432, 2010.
- [2721] S. Song and C. Park. Utilization of D-ribose through D-xylose transporter. *FEMS Microbiol. Lett.*, 163:255–261, 1998.
- [2722] Y.M. Song, Y.S. Kim, D. Kim, D.S. Lee, and H.J. Kwon. Cloning, expression, and biochemical characterization of a new histone deacetylase-like protein from *Thermus caldophilus* GK24. *Biochem. Biophys. Res. Commun.*, 361:55–61, 2007.
- [2723] S. Sonnino, H. Carinatti, and E. Cabib. Guanosine diphosphate D-glucose glucohydrolase. *Arch. Biochem. Biophys.*, 116:26–33, 1966.
- [2724] C.L. Soong, J. Ogawa, E. Sakuradani, and S. Shimizu. Barbiturase, a novel zinc-containing amidohydrolase involved in oxidative pyrimidine metabolism. *J. Biol. Chem.*, 277:7051–7058, 2002.

- [2725] C.L. Soong, J. Ogawa, and S. Shimizu. Novel amidohydrolytic reactions in oxidative pyrimidine metabolism: analysis of the barbiturase reaction and discovery of a novel enzyme, ureidomalonase. *Biochem. Biophys. Res. Commun.*, 286:222–226, 2001.
- [2726] S.H. Sørensen, O. Norén, H. Sjöström, and E.M. Danielsen. Amphiphilic pig intestinal microvillus maltase/glucoamylase. Structure and specificity. *Eur. J. Biochem.*, 126:559–568, 1982.
- [2727] H. Sorimachi, S. Imajoh-Ohmi, Y. Emori, H. Kawasaki, S. Ohno, Y. Minami, and K. Suzuki. Molecular cloning of a novel mammalian calcium-dependent protease distinct from both *m*- and  $\mu$ -types. Specific expression of the mRNA in skeletal muscle. *J. Biol. Chem.*, 264:20106–20111, 1989.
- [2728] H. Sorimachi, K. Kinbara, S. Kimura, M. Takahashi, S. Ishiura, N. Sasagawa, N. Sorimachi, H. Shimada, K. Tagawa, K. Maruyama, and K. Suzuki. Muscle-specific calpain, p94, responsible for limb girdle muscular dystrophy type 2A, associates with connectin through IS2, a p94-specific sequence. *J. Biol. Chem.*, 270:31158–31162, 1995.
- [2729] L. Sottrup-Jensen and H. Birkedal-Hansen. Human fibroblast collagenase- $\alpha$ -macroglobulin interactions. Localization of cleavage sites in the bait regions of five mammalian  $\alpha$ -macroglobulins. *J. Biol. Chem.*, 264:393–401, 1989.
- [2730] F. Soubrier, F. Alhenc-Gelas, C. Hubert, J. Allegrini, M. John, G. Tregear, and P. Corvol. Two putative active centers in human angiotensin I-converting enzyme revealed by molecular cloning. *Proc. Natl. Acad. Sci. USA*, 85:9386–9390, 1988.
- [2731] A. Soucek, C. Michalec, and A. Souckov. Identification and characterization of a new enzyme of the group phospholipase D isolated from *Corynebacterium ovis*. *Biochim. Biophys. Acta*, 227:116–128, 1971.
- [2732] M.F. Souliere, J.P. Perreault, and M. Bisaillon. Characterization of the vaccinia virus D10 decapping enzyme provides evidence for a two-metal-ion mechanism. *Biochem. J.*, 420:27–35, 2009.
- [2733] J. Soutourina, P. Plateau, and S. Blanquet. Metabolism of D-aminoacyl-tRNAs in *Escherichia coli* and *Saccharomyces cerevisiae* cells. *J. Biol. Chem.*, 275:32535–32542, 2000.
- [2734] V.V. Sova, L.A. Elyakova, and V.E. Vaskovsky. Purification and some properties of  $\beta$ -1,3-glucan glucanohydrolase from the crystalline style of bivalvia, *Spisula sachalinensis*. *Biochim. Biophys. Acta*, 212:111–115, 1970.
- [2735] S. Spadari, A.R. Subramanian, and G. Kalnitsky. Highly restricted specificity of the serine proteinase aspergillopeptidase B. *Biochim. Biophys. Acta*, 359:267–272, 1974.
- [2736] P.F. Spahr and R.F. Gesteland. Purification and properties of a new enzyme from *Escherichia coli*: nucleoside phosphoacyl hydrolase. *Eur. J. Biochem.*, 12:270–284, 1970.
- [2737] J.L. Spears, M.A. Rubio, K.W. Gaston, E. Wywiał, A. Strikoudis, J.M. Bujnicki, F.N. Papavasiliou, and J.D. Alfonzo. A single zinc ion is sufficient for an active *Trypanosoma brucei* tRNA editing deaminase. *J. Biol. Chem.*, 286:20366–20374, 2011.
- [2738] G. Speciale, Y. Jin, G.J. Davies, S.J. Williams, and E.D. Goddard-Borger. YihQ is a sulfoquinovosidase that cleaves sulfoquinovosyl diacylglyceride sulfolipids. LID - 10.1038/nchembio.2023 [doi. *Nat. Chem. Biol.*, 2016.
- [2739] H. Speijer, J.W. Govers-Reimslag, R.F. Zwaal, and J. Rosing. Prothrombin activation by an activator from the venom of *Oxyuranus scutellatus* (taipan snake). *J. Biol. Chem.*, 261:13258–13267, 1986.
- [2740] A.M. Spiekerman, K.K. Fredericks, F.W. Wagner, and J.M. Prescott. Leucostoma peptidase A: a metalloprotease from snake venom. *Biochim. Biophys. Acta*, 293:464–475, 1973.
- [2741] M.J. Spiro, V.D. Bhoyroo, and R.G. Spiro. Molecular cloning and expression of rat liver endo- $\alpha$ -mannosidase, an N-linked oligosaccharide processing enzyme. *J. Biol. Chem.*, 272:29356–29363, 1997.
- [2742] M.B. Sporn, H.M. Smith Lazarus, Henderson J.M., and W.R. Studies on nuclear exoribonucleases. 3. Isolation and properties of the enzyme from normal and malignant tissues of the mouse. *Biochemistry*, 8:1698–1706, 1969.
- [2743] S.R. Sprang. G protein mechanisms: insights from structural analysis. *Annu. Rev. Biochem.*, 66:639–678, 1997.
- [2744] A. Spungin and S. Blumberg. *Streptomyces griseus* aminopeptidase is a calcium-activated zinc metalloprotein. Purification and properties of the enzyme. *Eur. J. Biochem.*, 183:471–477, 1989.

- [2745] P.A. Srere, W. Seubert, and F. Lynen. Palmitoyl coenzyme A deacylase. *Biochim. Biophys. Acta*, 33:313–319, 1959.
- [2746] S.M. Srinivasula, S. Gupta, P. Datta, Z. Zhang, R. Hegde, N. Cheong, T. Fernandes-Alnemri, and E.S. Alnemri. Inhibitor of apoptosis proteins are substrates for the mitochondrial serine protease Omi/HtrA2. *J. Biol. Chem.*, 278:31469–31472, 2003.
- [2747] M. Sriram, J. Osipiuk, B. Freeman, R. Morimoto, and A. Joachimiak. Human Hsp70 molecular chaperone binds two calcium ions within the ATPase domain. *Structure*, 5:403–414, 1997.
- [2748] C. Srisomsap, K.L. Richardson, J.C. Jay, and R.B. Marchase. An  $\alpha$ -glucose-1-phosphate phosphodiesterase is present in rat liver cytosol. *J. Biol. Chem.*, 264:20540–20546, 1989.
- [2749] P.K. Srivastava, V.K. Sharma, D.S. Kalonia, and D.F. Grant. Polymorphisms in human soluble epoxide hydrolase: effects on enzyme activity, enzyme stability, and quaternary structure. *Arch. Biochem. Biophys.*, 427:164–169, 2004.
- [2750] R.M. Srivastava, N. Hudson, F.R. Seymour, and B. Weissman. Preparation of (aryl  $\alpha$ -L-idopyranosid)uronic acids. *Carbohydr. Res.*, 60:315–326, 1978.
- [2751] K.S. Srivenugopal and P.R. Adiga. Enzymic conversion of agmatine to putrescine in *Lathyrus sativus* seedlings. Purification and properties of a multifunctional enzyme (putrescine synthase). *J. Biol. Chem.*, 256:9532–9541, 1981.
- [2752] T.C. Stadtman. Alkaline phosphatases. In P.D. Boyer, H. Lardy, and K. Myrback, editors, *The Enzymes*, volume 5, pages 55–71. Academic Press, New York, 2nd edition, 1961.
- [2753] D.K. Staflien and P.P. Cleary. Characterization of the streptococcal C5a peptidase using a C5a-green fluorescent protein fusion protein substrate. *J. Bacteriol.*, 182:3254–3258, 2000.
- [2754] D.M. Stalker, L.D. Malyj, and K.E. McBride. Purification and properties of a nitrilase specific for the herbicide bromoxynil and corresponding nucleotide sequence analysis of the *bxn* gene. *J. Biol. Chem.*, 263:6310–6314, 1988.
- [2755] W.L. Starnes and F.J. Behal. A human liver aminopeptidase. The amino acid and carbohydrate content, and some physical properties of a sialic acid containing glycoprotein. *Biochemistry*, 13:3221–3227, 1974.
- [2756] R. Stasche, S. Hinderlich, C. Weise, K. Effertz, L. Lucka, P. Moormann, and W. Reutter. A bifunctional enzyme catalyzes the first two steps in *N*-acetylneuraminic acid biosynthesis of rat liver. Molecular cloning and functional expression of UDP-*N*-acetyl-glucosamine 2-epimerase/*N*-acetylmannosamine kinase. *J. Biol. Chem.*, 272:24319–24324, 1997.
- [2757] J.G. Stavrianopoulos and E. Chargaff. Purification and properties of ribonuclease H of calf thymus. *Proc. Natl. Acad. Sci. USA*, 70:1959–1963, 1973.
- [2758] D.A. Steege. Emerging features of mRNA decay in bacteria. *RNA*, 6:1079–1090, 2000.
- [2759] M. Steffek, G.L. Newton, Y. Av-Gay, and R.C. Fahey. Characterization of *Mycobacterium tuberculosis* mycothiol *S*-conjugate amidase. *Biochemistry*, 42:12067–12076, 2003.
- [2760] R.L. Stein, A.M. Strimpler, H. Hori, and J.C. Powers. Catalysis by human leukocyte elastase: mechanistic insights into specificity requirements. *Biochemistry*, 26:1301–1305, 1987.
- [2761] D.F. Steiner, S.P. Smeekens, S. Ohagi, and S.J. Chan. The new enzymology of precursor processing endoproteases. *J. Biol. Chem.*, 267:23435–23438, 1992.
- [2762] E.J. Stellwag and P.B. Hylemon. Purification and characterization of bile salt hydrolase from *Bacteroides fragilis* subsp. *fragilis*. *Biochim. Biophys. Acta*, 452:165–176, 1976.
- [2763] S.L. Stephenson and A.J. Kenny. The metabolism of neuropeptides. Hydrolysis of peptides by the phosphoramidon-insensitive rat kidney enzyme ‘endopeptidase-2’ and by rat microvillar membranes. *Biochem. J.*, 255:45–51, 1988.
- [2764] E.E. Sterchi, H.Y. Naim, M.J. Lentze, Hauri, and J.A.M. *N*-Benzoyl-L-tyrosyl-*p*-aminobenzoic acid hydrolase: a metalloendopeptidase of the human intestinal microvillus membrane which degrades biologically active peptides. *Arch. Biochem. Biophys.*, 265:105–118, 1988.
- [2765] M. Sternberg. Bond specificity, active site and milk clotting mechanism of the *Mucor miehei* protease. *Biochim. Biophys. Acta*, 285:383–392, 1972.

- [2766] M. Sternberg and R.G. Shapiro. Studies on the catabolism of the hydroxylysine-linked disaccharide units of basement membranes and collagens. Isolation and characterization of a rat kidney  $\alpha$ -glucosidase of high specificity. *J. Biol. Chem.*, 254:10329–10336, 1979.
- [2767] A. Stevens and R.J. Hilmoe. Studies on a nuclease from *Azotobacter agilis*. I. Isolation and mode of action. *J. Biol. Chem.*, 235:3016–3022, 1960.
- [2768] A. Stevens and R.J. Hilmoe. Studies on a nuclease from *Azotobacter agilis*. II. Hydrolysis of ribonucleic and deoxyribonucleic acids. *J. Biol. Chem.*, 235:3023–3027, 1960.
- [2769] K.J. Stevenson, D. Gibson, and G.H. Dixon. Amino acid analyses of chymotrysin-like proteases from the sea anemone (*Metridium senile*). *Can. J. Biochem.*, 52:93–100, 1974.
- [2770] A.E. Stewart, S.M. Arfin, and R.A. Bradshaw. Protein NH<sub>2</sub>-terminal asparagine deamidase. Isolation and characterization of a new enzyme. *J. Biol. Chem.*, 269:23509–23517, 1994.
- [2771] A.J. Stewart, R. Schmid, C.A. Blindauer, S.J. Paisey, and C. Farquharson. Comparative modelling of human PHOSPHO1 reveals a new group of phosphatases within the haloacid dehalogenase superfamily. *Protein Eng.*, 16:889–895, 2003.
- [2772] S.R. Stitch, I.D.K. Halkerston, and J. Hillman. The enzymic hydrolysis of steroid conjugates. 1. Sulphatase and  $\beta$ -glucuronidase activity of molluscan extracts. *Biochem. J.*, 63:705–710, 1965.
- [2773] J.T. Stivers and A.C. Drohat. Uracil DNA glycosylase: insights from a master catalyst. *Arch. Biochem. Biophys.*, 396:1–9, 2001.
- [2774] K. Stocker and G.H. Barlow. The coagulant enzyme from *Bothrops atrox* venom (batroxobin). *Methods Enzymol.*, 45:214–223, 1976.
- [2775] W. Stöcker, M. Ng, and D.S. Auld. Fluorescent oligopeptide substrates for kinetic characterization of the specificity of *Astacus* protease. *Biochemistry*, 29:10418–10425, 1990.
- [2776] W. Stöcker, R.L. Wolz, R. Zwilling, D.J. Strydom, and D.S. Auld. *Astacus* protease, a zinc metalloenzyme. *Biochemistry*, 27:5026–5032, 1988.
- [2777] S. Stoeva, T. Kleinschmidt, B. Mesrob, and G. Braunitzer. Primary structure of a zinc protease from *Bacillus mesentericus* strain 76. *Biochemistry*, 29:527–534, 1990.
- [2778] J.E. Stok, H. Huang, P.D. Jones, C.E. Wheelock, C. Morisseau, and B.D. Hammock. Identification, expression, and purification of a pyrethroid-hydrolyzing carboxylesterase from mouse liver microsomes. *J. Biol. Chem.*, 279:29863–29869, 2004.
- [2779] E.G. Strauss, R.J. De Groot, R. Levinson, and J.H. Strauss. Identification of the active site residues in the nsP2 proteinase of Sindbis virus. *Virology*, 191:932–940, 1992.
- [2780] F. Strelitz. Studies on cholinesterase. 4. Purification of pseudo-cholinesterase from horse serum. *Biochem. J.*, 38:86–88, 1944.
- [2781] F. Striebel, F. Imkamp, D. Özcelik, and E. Weber-Ban. Pupylation as a signal for proteasomal degradation in bacteria. *Biochim. Biophys. Acta*, 1843:103–113, 2014.
- [2782] F. Striebel, F. Imkamp, M. Sutter, M. Steiner, A. Mamedov, and E. Weber-Ban. Bacterial ubiquitin-like modifier Pup is deamidated and conjugated to substrates by distinct but homologous enzymes. *Nat. Struct. Mol. Biol.*, 16:647–651, 2009.
- [2783] S. Strobl, C. Fernandez-Catalan, M. Braun, R. Huber, H. Masumoto, K. Nakagawa, A. Irie, H. Sorimachi, G. Bourenkow, H. Bartunik, K. Suzuki, and W. Bode. The crystal structure of calcium-free human *m*-calpain suggests an electrostatic switch mechanism for activation by calcium. *Proc. Natl. Acad. Sci. USA*, 97:588–592, 2000.
- [2784] J.-C. Su, C.-C. Li, and C.C. Ting. A new adenylate deaminase from red marine alga *Porphyra crispata*. *Biochemistry*, 5:536–543, 1966.
- [2785] K.P. Subedi, D. Choi, I. Kim, B. Min, and C. Park. Hsp31 of *Escherichia coli* K-12 is glyoxalase III. *Mol. Microbiol.*, 81:926–936, 2011.



- [2786] H. Suda, K. Yamamoto, T. Aoyagi, and H. Umezawa. Purification and properties of *N*-formylmethionine aminopeptidase from rat liver. *Biochim. Biophys. Acta*, 616:60–67, 1980.
- [2787] M. Sue, A. Ishihara, and H. Iwamura. Purification and characterization of a  $\beta$ -glucosidase from rye (*Secale cereale* L.) seedlings. *Plant Sci.*, 155:67–74, 2000.
- [2788] M. Sue, A. Ishihara, and H. Iwamura. Purification and characterization of a hydroxamic acid glucoside  $\beta$ -glucosidase from wheat (*Triticum aestivum* L.) seedlings. *Planta*, 210:432–438, 2000.
- [2789] M. Sue, C. Nakamura, T. Miyamoto, and S. Yajima. Active-site architecture of benzoxazinone-glucoside  $\beta$ -D-glucosidases in Triticeae. *Plant Sci.*, 180:268–275, 2011.
- [2790] M. Sue, K. Yamazaki, S. Yajima, T. Nomura, T. Matsukawa, H. Iwamura, and T. Miyamoto. Molecular and structural characterization of hexameric  $\beta$ -D-glucosidases in wheat and rye. *Plant Physiol.*, 141:1237–1247, 2006.
- [2791] Y. Sugano, H. Kodama, I. Terada, Y. Yamazaki, and M. Noma. Purification and characterization of a novel enzyme,  $\alpha$ -neogaroooligosaccharide hydrolase ( $\alpha$ -NAOS hydrolase), from a marine bacterium, *Vibrio* sp. strain JT0107. *J. Bacteriol.*, 176:6812–6818, 1994.
- [2792] Y. Sugano, I. Terada, M. Arita, M. Noma, and T. Matsumoto. Purification and characterization of a new agarase from a marine bacterium, *Vibrio* sp. strain JT0107. *Appl. Environ. Microbiol.*, 59:1549–1554, 1993.
- [2793] H. Sugihara, M. Moriura, and T. Nikai. Purification and properties of a lethal, hemorrhagic protein, "mucrotoxin A", from the venom of the Chinese habu snake (*Trimeresurus mucrosquamatus*). *Toxicon*, 21:247–255, 1983.
- [2794] K. Sugimura and T. Nishihara. Purification, characterization, and primary structure of *Escherichia coli* protease VII with specificity for paired basic residues: identity of protease VII and *ompT*. *J. Bacteriol.*, 170:5625–5632, 1988.
- [2795] M. Sugiura, A. Ito, and T. Yamaguchi. Studies on dextranase. II. New exo-dextranase from *Brevibacterium fuscum* var. *Dextranlyticum*. *Biochim. Biophys. Acta*, 350:61–70, 1974.
- [2796] K. Suguna, E.A. Padlan, C.W. Smith, W.D. Carlson, and D.R. Davies. Binding of a reduced peptide inhibitor to the aspartic proteinase from *Rhizopus chinensis*: implications for a mechanism of action. *Proc. Natl. Acad. Sci. USA*, 84:7009–7013, 1987.
- [2797] H.M. Suld and P.A. Herbut. Guinea pig serum and liver L-asparaginases. Comparison of serum and papain-digested liver L-asparaginases. *J. Biol. Chem.*, 245:2797–2801, 1970.
- [2798] E. Sulkowski and M. Laskowski. , Sr. Phosphatase-free crystalline micrococcal nuclease. *J. Biol. Chem.*, 241:4386–4388, 1966.
- [2799] J.B. Sumner. Urease. In J.B. Sumner and K. Myrback, editors, *The Enzymes*, volume 1, pages 873–892. Academic Press, New York, 1951.
- [2800] R.A. Sumrada and T.G. Cooper. Urea carboxylase and allophanate hydrolase are components of a multifunctional protein in yeast. *J. Biol. Chem.*, 257:9119–9127, 1982.
- [2801] T.A. Sundarajan and P.S. Sarma. Substrate specificity of phosphoprotein phosphatase from spleen. *Biochem. J.*, 71:537–544, 1959.
- [2802] M. Sundberg, , and K. Purification and properties of two acetylxylan esterases of *Trichoderma reesei*. *Biotechnol. Appl. Biochem.*, 13:1–11, 1991.
- [2803] G. Sundharadas and C. Gilvarg. Biosynthesis of  $\alpha,\epsilon$ -diaminopimelic acid in *Bacillus megaterium*. *J. Biol. Chem.*, 242:3983–3984, 1967.
- [2804] J.S. Sung and D.W. Mosbaugh. *Escherichia coli* double-strand uracil-DNA glycosylase: involvement in uracil-mediated DNA base excision repair and stimulation of activity by endonuclease IV. *Biochemistry*, 39:10224–10235, 2000.
- [2805] S. Supangat, Y.K. Choi, Y.S. Park, D. Son, C.D. Han, and K.H. Lee. Expression, purification, crystallization and preliminary X-ray analysis of sepiapterin reductase from *Chlorobium tepidum*. *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.*, 61:202–204, 2005.

- [2806] I. Sussman and M. Avron. Characterization and partial purification of DL-glycerol-1-phosphatase from *Dunaliella salina*. *Biochim. Biophys. Acta*, 661:199–204, 1981.
- [2807] R. Sütfield and G.H.N. Towers. 5-(4-Acetoxy-1-butynyl)-2,2'-bithiophene:acetate esterase from *Tagetes patula*. *Phytochemistry*, 21:277–279, 1982.
- [2808] D.R. Sutherland, K.M. Abdullah, P. Mellors Cyopick, A. Cleavage of the cell-surface O-sialoglycoproteins CD34, and CD. 44, and CD45 by a novel glycoprotease from *Pasteurella haemolytica*. *J. Immunol.*, 148:1458–1464, 1992.
- [2809] W.D. Sutton. A crude nuclease preparation suitable for use in DNA reassociation experiments. *Biochim. Biophys. Acta*, 240:522–531, 1971.
- [2810] H. Suzuki and H. Kumagai. Autocatalytic processing of  $\gamma$ -glutamyltranspeptidase. *J. Biol. Chem.*, 277:43536–43543, 2002.
- [2811] H. Suzuki, S.-C. Li, and Y.-T. Li.  $\alpha$ -Galactosidase from *Mortierella vinacea*. Crystallization and properties. *J. Biol. Chem.*, 245:781–786, 1970.
- [2812] K. Suzuki, M. Kawada, and N. Shimazono. Soluble lactonase. Identity of lactonase I and aldonolactonase with gluconolactonase. *J. Biochem. (Tokyo)*, 49:448–449, 1961.
- [2813] O. Suzuki, Y. Ishikawa, K. Miyazaki, K. Izu, and T. Matsumoto. Acetylputrescine deacetylase from *Micrococcus luteus* K-11. *Biochim. Biophys. Acta*, 882:140–142, 1986.
- [2814] R. Suzuki, T. Katayama, M. Kitaoka, H. Kumagai, T. Wakagi, H. Shoun, H. Ashida, K. Yamamoto, and S. Fushinobu. Crystallographic and mutational analyses of substrate recognition of endo- $\alpha$ -N-acetylgalactosaminidase from *Bifidobacterium longum*. *J. Biochem.*, 146:389–398, 2009.
- [2815] Y. Suzuki, T. Amano, and Y. Shioi. Characterization and cloning of the chlorophyll-degrading enzyme pheophorbidease from cotyledons of radish. *Plant Physiol.*, 140:716–725, 2006.
- [2816] Y. Suzuki and G.M. Brown. The biosynthesis of folic acid. XII. Purification and properties of dihydroneopterin triphosphate pyrophosphohydrolase. *J. Biol. Chem.*, 249:2405–2410, 1974.
- [2817] Y. Suzuki, M. Doi, and Y. Shioi. Two enzymatic reaction pathways in the formation of pyropheophorbide *a*. *Photosynth. Res.*, 74:225–233, 2002.
- [2818] Y. Suzuki and K. Uchida. Multiple forms of  $\alpha$ -glycosidase from pig duodenum. *Agric. Biol. Chem.*, 49:1573–1581, 1985.
- [2819] I. Svendsen, M.R. Jensen, and K. Breddam. The primary structure of the glutamic acid-specific protease of *Streptomyces griseus*. *FEBS Lett.*, 292:165–167, 1991.
- [2820] K.H. Swamy, C.H. Chung, and A.L. Goldberg. Isolation and characterization of protease Do from *Escherichia coli*, a large serine protease containing multiple subunits. *Arch. Biochem. Biophys.*, 224:543–554, 1983.
- [2821] M.N. Swartz, N.O. Kaplan, and M.F. Lamborg. A "heat-activated" diphosphopyridine nucleotide pyrophosphatase from *Proteus vulgaris*. *J. Biol. Chem.*, 232:1051–1063, 1958.
- [2822] L.S. Symington and R. Kolodner. Partial purification of an enzyme from *Saccharomyces cerevisiae* that cleaves Holliday junctions. *Proc. Natl. Acad. Sci. USA*, 82:7247–7251, 1985.
- [2823] J. Szulmajster. Bacterial degradation of creatinine. II. Creatinine desimidase. *Biochim. Biophys. Acta*, 30:154–163, 1958.
- [2824] T., Dixon Maehama, , and PTEN. /MMAC1, dephosphorylates the lipid second messenger, phosphatidylinositol 3,4,5-trisphosphate. *J. Biol. Chem.*, 273:13375–13378, 1998.
- [2825] I. Tabas and S. Kornfeld. The synthesis of complex-type oligosaccharides. III. Identification of an  $\alpha$ -D-mannosidase activity involved in a late stage of processing of complex-type oligosaccharides. *J. Biol. Chem.*, 253:7779–7786, 1978.
- [2826] I. Tabas and S. Kornfeld. Purification and characterization of a rat liver Golgi  $\alpha$ -mannosidase capable of processing asparagine-linked oligosaccharides. *J. Biol. Chem.*, 254:11655–11663, 1979.

- [2827] H. Tabor and L. Wyngarden. The enzymatic formation of formiminotetrahydrofolic acid, 5,10-methenyltetrahydrofolic acid, and 10-formyltetrahydrofolic acid in the metabolism of formiminoglutamic acid. *J. Biol. Chem.*, 234:1830–1849, 1959.
- [2828] T. Tagami, E. Miyano, J. Sadahiro, M. Okuyama, T. Iwasaki, and A. Kimura. Two novel glycoside hydrolases responsible for the catabolism of cyclobis-(1→6)- $\alpha$ -nigerosyl. *J. Biol. Chem.*, 291:16438–16447, 2016.
- [2829] K. Tagawa and A. Kaji. Preparation of L-arabinose-containing polysaccharides and the action of an  $\alpha$ -L-arabinofuranosidase on these polysaccharides. *Carbohydr. Res.*, 11:293–301, 1969.
- [2830] J. Tai, Y. Kortt, A.A., Liu, T.-Y. and Elliott, S.D. Primary structure of streptococcal proteinase. III. Isolation of cyanogen bromide peptides: complete covalent structure of the polypeptide chain. *J. Biol. Chem.*, 251:1955–1959, 1976.
- [2831] T. Tai, K. Yamashita, M. Ogata-Arakawa, N. Koide, T. Muramatsu, S. Iwashita, Y. Inoue, and A. Kobata. Structural studies of two ovalbumin glycopeptides in relation to the endo- $\beta$ -N-acetylglucosaminidase specificity. *J. Biol. Chem.*, 250:8569–8575, 1975.
- [2832] T. Takada, K. Iida, and J. Moss. Cloning and site-directed mutagenesis of human ADP-ribosylarginine hydrolase. *J. Biol. Chem.*, 268:17837–17843, 1993.
- [2833] M. Takagi, T. Imanaka, and S. Aiba. Nucleotide sequence and promoter region for the neutral protease gene from *Bacillus stearothermophilus*. *J. Bacteriol.*, 163:824–831, 1985.
- [2834] Y. Takagi and B.L. Horecker. Purification and properties of a bacterial riboside hydrolyase. *J. Biol. Chem.*, 225:77–86, 1956.
- [2835] Y. Takagi, D. Setoyama, R. Ito, H. Kamiya, Y. Yamagata, and M. Sekiguchi. Human MTH3 (NUDT18) protein hydrolyzes oxidized forms of guanosine and deoxyguanosine diphosphates: comparison with MTH1 and MTH2. *J. Biol. Chem.*, 287:21541–21549, 2012.
- [2836] A. Takahashi, E.S. Alnemri, Y.A. Lazebnik, T. Fernandes-Alnemri, G. Litwack, R.D. Moir, R.D. Goldman, G.G. Poirier, S.H. Kaufmann, and W.C. Earnshaw. Cleavage of lamin A by Mch2 $\alpha$  but not CPP32: multiple interleukin 1 $\beta$ -converting enzyme-related proteases with distinct substrate recognition properties are active in apoptosis. *Proc. Natl. Acad. Sci. USA*, 93:8395–8400, 1996.
- [2837] K. Takahashi. The structure and function of ribonuclease T<sub>1</sub>. I. Chromatographic purification and properties of ribonuclease T<sub>1</sub>. *J. Biochem. (Tokyo)*, 49:1–8, 1961.
- [2838] K. Takahashi and W.-J. Chang. The structure and function of acid proteases. V. Comparative studies on the specific inhibition of acid proteases by diazoacetyl-DL-norleucine methyl ester, 1,2-epoxy-3-(p-nitrophenoxy)propane and pepstatin. *J. Biochem. (Tokyo)*, 80:497–506, 1976.
- [2839] K. Takahashi, W.-J. Chang, and J.-S. Ko. Specific inhibition of acid proteases from brain, kidney, skeletal muscle, and insectivorous plants by diazoacetyl-DL-norleucine methyl ester and by pepstatin. *J. Biochem. (Tokyo)*, 76:897–899, 1974.
- [2840] M. Takahashi, Y. Matsushita, Y. Iijima, and K. Tanzawa. Purification and characterization of endothelin-converting enzyme from rat lung. *J. Biol. Chem.*, 268:21394–21398, 1993.
- [2841] N. Takahashi. Demonstration of a new amidase acting on glycopeptides. *Biochem. Biophys. Res. Commun.*, 76:1194–1201, 1977.
- [2842] N. Takahashi and F. Egami. Hydrolysis of polysaccharide sulphate esters by a sulphatase preparation from *Charonia lampas*. *Biochem. J.*, 80:384–386, 1961.
- [2843] N. Takahashi and H. Nishibe. Some characteristics of a new glycopeptidase acting on aspartylglycosylamine linkages. *J. Biochem. (Tokyo)*, 84:1467–1473, 1978.
- [2844] T. Takahashi, M. Nakanishi, Y. Yao, I. Uyeda, and N. Serizawa. Direct formation of human interleukin-11 by *cis*-acting system of plant virus protease in *Escherichia coli*. *Biosci. Biotechnol. Biochem.*, 62:953–958, 1998.
- [2845] T. Takahashi and A. Ohsaka. Purification and characterization of a proteinase in the venom of *Trimeresurus flavoviridis*. Complete separation of the enzyme from hemorrhagic activity. *Biochim. Biophys. Acta*, 198:293–307, 1970.

- [2846] T. Takahashi, T. Sugahara, and A. Ohsaka. Purification of *Clostridium perfringens* phospholipase C ( $\alpha$ -toxin) by affinity chromatography on agarose-linked egg-yolk lipoprotein. *Biochim. Biophys. Acta*, 351:155–171, 1974.
- [2847] T. Takahashi and J. Tang. Cathepsin D from porcine and bovine spleen. *Methods Enzymol.*, 80:565–581, 1981.
- [2848] T. Takahashi, Y. Yamazaki, and K. Kato. Substrate specificity of an  $\alpha$ -amino acid ester hydrolase produced by *Acetobacter turbidans* A. T.C.C. 9325. *Biochem. J.*, 137:497–503, 1974.
- [2849] Y. Takahashi, G. Moiseyev, Y. Chen, K. Farjo, O. Nikolaeva, and J.X. Ma. An enzymatic mechanism for generating the precursor of endogenous 13-*cis* retinoic acid in the brain. *FEBS J.*, 278:973–987, 2011.
- [2850] T. Takai, T. Kato, Y. Sakata, H. Yasueda, K. Izuhara, K. Okumura, and H. Ogawa. Recombinant Der p 1 and Der f 1 exhibit cysteine protease activity but no serine protease activity. *Biochem. Biophys. Res. Commun.*, 328:944–952, 2005.
- [2851] H. Takaku, A. Minagawa, M. Takagi, and M. Nashimoto. A candidate prostate cancer susceptibility gene encodes tRNA 3' processing endoribonuclease. *Nucleic Acids Res.*, 31:2272–2278, 2003.
- [2852] S. Takasaki and A. Kobata. Purification and characterization of an endo- $\beta$ -galactosidase produced by *Diplococcus pneumoniae*. *J. Biol. Chem.*, 251:3603–3609, 1976.
- [2853] S. Takasawa, A. Tohgo, N. Noguchi, T. Koguma, K. Nata, T. Sugimoto, H. Yonekura, and H. Okamoto. Synthesis and hydrolysis of cyclic ADP-ribose by human leukocyte antigen CD38 and inhibition of the hydrolysis by ATP. *J. Biol. Chem.*, 268:26052–26054, 1993.
- [2854] I. Takebe. Isolation and characterization of a new enzyme choline sulfatase. *J. Biochem. (Tokyo)*, 50:245–255, 1961.
- [2855] H. Takeda, S. Yamamoto, Y. Kojima, and O. Hayaishi. Studies on monooxygenases. I. General properties of crystalline L-lysine monooxygenase. *J. Biol. Chem.*, 244:2935–2941, 1969.
- [2856] K. Takegawa, S. Miki, T. Jikibara, and S. Iwahara. Purification and characterization of exo- $\alpha$ -D-mannosidase from a *Cellulomonas* sp. *Biochim. Biophys. Acta*, 991:431–437, 1989.
- [2857] S. Takenaka, S. Murakami, Y.J. Kim, and K. Aoki. Complete nucleotide sequence and functional analysis of the genes for 2-aminophenol metabolism from *Pseudomonas* sp. AP-3. *Arch. Microbiol.*, 174:265–272, 2000.
- [2858] M. Takeo, A. Ohara, S. Sakae, Y. Okamoto, C. Kitamura, D. Kato, and S. Negoro. Function of a glutamine synthetase-like protein in bacterial aniline oxidation via  $\gamma$ -glutamylanilide. *J. Bacteriol.*, 195:4406–4414, 2013.
- [2859] H. Takeshima, J. Inokoshi, Y. Takada, H. Tanaka, and S. Omura. A deacylation enzyme for aculeacin A, a neutral lipopeptide antibiotic, from *Actinoplanes utahensis*: purification and characterization. *J. Biochem. (Tokyo)*, 105:606–610, 1989.
- [2860] Y. Takesue. Purification and properties of rabbit intestinal sucrase. *J. Biochem. (Tokyo)*, 65:545–552, 1969.
- [2861] H. Takeya, M. Arakawa, T. Miyata, S. Iwanaga, and T. Omori-Satoh. Primary structure of H<sub>2</sub>-proteinase, a non-hemorrhagic metalloproteinase, isolated from the venom of the habu snake, *Trimeresurus flavoviridis*. *J. Biochem. (Tokyo)*, 106:151–157, 1989.
- [2862] H. Takeya, S. Nishida, T. Miyata, S. Kawada, Y. Saisaka, T. Morita, and S. Iwanaga. Coagulation factor X activating enzyme from Russell's viper venom (RVV-X). A novel metalloproteinase with disintegrin (platelet aggregation inhibitor)-like and C-type lectin-like domains. *J. Biol. Chem.*, 267:14109–14117, 1992.
- [2863] H. Takeya, K. Oda, T. Miyata, T. Omori-Satoh, and S. Iwanaga. The complete amino acid sequence of the high molecular mass hemorrhagic protein HR1B isolated from the venom of *Trimeresurus flavoviridis*. *J. Biol. Chem.*, 265:16068–16073, 1990.
- [2864] H. Takeya, A. Onikura, T. Nikai, H. Iwanaga Sugihara, , and HT. 2, isolated from the venom of *Crotalus ruber ruber*. *J. Biochem. (Tokyo)*, 108:711–719, 1990.
- [2865] A. Tam, W.K. Schmidt, and S. Michaelis. The multispanning membrane protein Ste24p catalyzes CAAX proteolysis and NH<sub>2</sub>-terminal processing of the yeast a-factor precursor. *J. Biol. Chem.*, 276:46798–46806, 2001.
- [2866] R. Tamayo, A.D. Tischler, and A. Camilli. The EAL domain protein VieA is a cyclic diguanylate phosphodiesterase. *J. Biol. Chem.*, 280:33324–33330, 2005.

- [2867] J. Tamura, H. Takagi, and K. Kadowaki. Purification and some properties of the endo  $\alpha$ -1,4-polygalactosaminidase from *Pseudomonas* sp. *Agric. Biol. Chem.*, 52:2475–2484, 1988.
- [2868] A.K. Tan and D.L. Eaton. Activation and characterization of procarboxypeptidase B from human plasma. *Biochemistry*, 34:5811–5816, 1995.
- [2869] A.L. Tan, P. Loke, and T.S. Sim. Molecular cloning and functional characterisation of VanX, a D-alanyl-D-alanine dipeptidase from *Streptomyces coelicolor* A3(2). *Res. Microbiol.*, 153:27–32, 2002.
- [2870] K. Tanaka, T. Nakano, S. Noguchi, and W. Pigman. Purification of  $\alpha$ -L-fucosidase of abalone livers. *Arch. Biochem. Biophys.*, 126:624–633, 1968.
- [2871] N. Tanaka, M. Takeuchi, and E. Ichishima. Purification of an acid proteinase from *Aspergillus saitoi* and determination of peptide bond specificity. *Biochim. Biophys. Acta*, 485:406–416, 1977.
- [2872] S. Tanaka, E.A. Robinson, E. Appella, M. Miller, H.L. Ammon, J. Roberts, I.T. Weber, and A. Wlodawer. Structures of amidohydrolases. Amino acid sequence of a glutaminase-asparaginase from *Acinetobacter glutaminasificans* and preliminary crystallographic data for an asparaginase from *Erwinia chrysanthemi*. *J. Biol. Chem.*, 263:8583–8591, 1988.
- [2873] T. Tanaka, T. Fukui, and T. Imanaka. Different cleavage specificities of the dual catalytic domains in chitinase from the hyperthermophilic archaeon *Thermococcus kodakaraensis* KOD1. *J. Biol. Chem.*, 276:35629–35635, 2001.
- [2874] T. Tanaka and E. Ichishima. Molecular properties of aminopeptidase Ey as a zinc-metalloenzyme. *Int. J. Biochem.*, 25:1681–1688, 1993.
- [2875] T. Tanaka and E. Ichishima. Substrate specificity of aminopeptidase Ey from hen's egg yolk. *Comp. Biochem. Physiol. [B]*, 105:105–110, 1993.
- [2876] T. Tanaka, H. Matsuzawa, S. Kojima, I. Kumagai, K. Miura, and T. Ohta. P1 specificity of aqualysin I (a subtilisin-type serine protease) from *Thermus aquaticus* YT-1, using P1-substituted derivatives of *Streptomyces* subtilisin inhibitor. *Biosci. Biotechnol. Biochem.*, 62:2035–2038, 1998.
- [2877] T. Tanaka, H. Matsuzawa, and T. Ohta. Substrate specificity of aqualysin I, a bacterial thermophilic alkaline serine protease from *Thermus aquaticus* YT-1: Comparison with proteinase K, subtilisin BPN<sup>'</sup> and subtilisin Carlsberg. *Biosci. Biotechnol. Biochem.*, 62:2161–2165, 1998.
- [2878] T. Tanaka, B.J. McRae, K. Cho, R. Cook, J.E. Fraki, D.A. Johnson, and J.C. Powers. Mammalian tissue trypsin-like enzymes. Comparative reactivities of human skin tryptase, human lung tryptase, and bovine trypsin with peptide 4-nitroanilide and thioester substrates. *J. Biol. Chem.*, 258:13552–13557, 1983.
- [2879] T. Tanaka, Y. Minematsu, C.F. Reilly, J. Travis, and J.C. Powers. Human leukocyte cathepsin G. Subsite mapping with 4-nitroanilides, chemical modification, and effect of possible cofactors. *Biochemistry*, 24:2040–2047, 1985.
- [2880] J. Tang. Gastricsin and pepsin. *Methods Enzymol.*, 19:406–421, 1970.
- [2881] J. Tang and R.N.S. Wong. Evolution in the structure and function of aspartic proteases. *J. Cell. Biochem.*, 33:53–63, 1987.
- [2882] B. Tangendjaja, J.B. Lowry, and R.H. Wills. Isolation of a mimosine degrading enzyme from *Leucaena* leaf. *J. Sci. Food Agric.*, 37:523–526, 1986.
- [2883] T. Tani, I. Kawashima, K. Mita, and Y. Takiguchi. Nucleotide sequence of the human pancreatic trypsinogen III cDNA. *Nucleic Acids Res.*, 18:1631–1631, 1990.
- [2884] M.M. Tanizaki, R.B. Zingali, H. Kawazaki, S. Imajoh, S. Yamazaki, and K. Suzuki. Purification and some characteristics of a zinc metalloprotease from the venom of *Bothrops jararaca* (jararaca). *Toxicon*, 27:747–755, 1989.
- [2885] J.A. Tanner, R.M. Watt, Y.B. Chai, L.Y. Lu, M.C. Lin, J.S. Peiris, L.L. Poon, H.F. Kung, and J.D. Huang. The severe acute respiratory syndrome (SARS) coronavirus NTPase/helicase belongs to a distinct class of 5' to 3' viral helicases. *J. Biol. Chem.*, 278:39578–39582, 2003.
- [2886] A.L. Tarentino, C.M. Plummer Gomez, , and Jr. Deglycosylation of asparagine-linked glycans by peptide:N-glycosidase F. *Biochemistry*, 24:4665–4671, 1985.

- [2887] A.L. Tarentino and F. Maley. The purification and properties of a  $\beta$ -aspartyl *N*-acetylglucosylamine amidohydrolase from hen oviduct. *Arch. Biochem. Biophys.*, 130:295–303, 1969.
- [2888] A.L. Tarentino, T.H. Plummer, Maley Jr., and F. The release of intact oligosaccharides from specific glycoproteins by endo- $\beta$ -*N*-acetylglucosaminidase H. *J. Biol. Chem.*, 249:818–824, 1974.
- [2889] A.L. Tarentino, G. Quinones, B.G. Grimwood, C.R., Plummer Hauer, , and Jr. Molecular cloning and sequence analysis of flavastacin: an *O*-glycosylated prokaryotic zinc metalloendopeptidase. *Arch. Biochem. Biophys.*, 319:281–285, 1995.
- [2890] H.L.A. Tarr. Fish muscle riboside hydrolases. *Biochem. J.*, 59:386–391, 1955.
- [2891] H.L.A. Tarr. 1-Methyladenosine hydrolase of starfish (*Pisaster ochraceus*). *J. Fish Res. Board Can.*, 30:1861–1866, 1973.
- [2892] L.D. Tatar, C.L. Marolda, A.N. Polischuk, D. van Leeuwen, and M.A. Valvano. An *Escherichia coli* undecaprenyl-pyrophosphate phosphatase implicated in undecaprenyl phosphate recycling. *Microbiology*, 153:2518–2529, 2007.
- [2893] N. Tautz, K. Elbers, D. Stoll, G. Meyers, and H.J. Thiel. Serine protease of pestiviruses: determination of cleavage sites. *J. Virol.*, 71:5415–5422, 1997.
- [2894] N. Tautz, A. Kaiser, and H.J. Thiel. NS3 serine protease of bovine viral diarrhea virus: characterization of active site residues, NS4A cofactor domain, and protease-cofactor interactions. *Virology*, 273:351–363, 2000.
- [2895] D.L. Taylor, J.C. Ho, A. Oliver, and F.Z. Watts. Cell-cycle-dependent localisation of Ulp1, a *Schizosaccharomyces pombe* Pmt3 (SUMO)-specific protease. *J. Cell Sci.*, 115:1113–1122, 2002.
- [2896] Z.W. Taylor, H.A. Brown, H.M. Holden, and F.M. Raushel. Biosynthesis of nucleoside diphosphoramidates in *Campylobacter jejuni*. *Biochemistry*, 56:6079–6082, 2017.
- [2897] F. Teillet, B. Dublet, J.P. Andrieu, C. Gaboriaud, G.J. Arlaud, and N.M. Thielen. The two major oligomeric forms of human mannan-binding lectin: chemical characterization, carbohydrate-binding properties, and interaction with MBL-associated serine proteases. *J. Immunol.*, 174:2870–2877, 2005.
- [2898] A. Teplyakov, K. Polyakov, G. Obmolova, B. Strokopytov, I. Kuranova, A. Osterman, N. Grishin, S. Smulevitch, O. Zagnitko, O. Galperina, M. Matz, and V. Stepanov. Crystal structure of carboxypeptidase T from *Thermoactinomyces vulgaris*. *Eur. J. Biochem.*, 208:281–288, 1992.
- [2899] A.V. Teplyakov, I.P. Kuranova, E.H. Harutyunyan, B.K. Vainshtein, C. Frömmel, W.E. Höhne, and K.S. Wilson. Crystal structure of thermitase at 1.4 Å resolution. *J. Mol. Biol.*, 214:261–279, 1990.
- [2900] M. Terada, M. Tatibana, and O. Hayaishi. Preparation and properties of nucleoside hydrolase from *Pseudomonas fluorescens*. *J. Biol. Chem.*, 242:5578–5585, 1967.
- [2901] Y. Terao, M. Yamaguchi, S. Hamada, and S. Kawabata. Multifunctional glyceraldehyde-3-phosphate dehydrogenase of *Streptococcus pyogenes* is essential for evasion from neutrophils. *J. Biol. Chem.*, 281:14215–14223, 2006.
- [2902] T. Terashita, K. Oda, M. Kono, and S. Murao. *Streptomyces* pepsin inhibitor-insensitive carboxyl proteinase from *Lentinius edodes*. *Agric. Biol. Chem.*, 45:1937–1943, 1981.
- [2903] T. Terashita, K. Oda, M. Kono, and S. Murao. *Streptomyces* pepsin inhibitor-insensitive carboxyl proteinase from *Ganoderma lucidum*. *Agric. Biol. Chem.*, 48:1029–1035, 1984.
- [2904] N.G. Ternan, J.T. Hamilton, and J.P. Quinn. Initial in vitro characterisation of phosphonopyruvate hydrolase, a novel phosphate starvation-independent, carbon-phosphorus bond cleavage enzyme in *Burkholderia cepacia* Pal6. *Arch. Microbiol.*, 173:35–41, 2000.
- [2905] R. Teufel, V. Mascaraque, W. Ismail, M. Voss, J. Perera, W. Eisenreich, W. Haehnel, and G. Fuchs. Bacterial phenylalanine and phenylacetate catabolic pathway revealed. *Proc. Natl. Acad. Sci. USA*, 107:14390–14395, 2010.
- [2906] N.M. Thanassi and H.I. Nakada. Enzymic degradation of fucoidan by enzymes from the hepatopancreas of abalone, *Halotis* species. *Arch. Biochem. Biophys.*, 118:172–177, 1967.
- [2907] E. Theodoratou, A. Paschos, A. Magalon, E. Fritsche, R. Huber, and A. Bock. Nickel serves as a substrate recognition motif for the endopeptidase involved in hydrogenase maturation. *Eur. J. Biochem.*, 267:1995–1999, 2000.

- [2908] C.J. Thibodeaux and H.W. Liu. Mechanistic studies of 1-aminocyclopropane-1-carboxylate deaminase: characterization of an unusual pyridoxal 5'-phosphate-dependent reaction. *Biochemistry*, 50:1950–1962, 2011.
- [2909] S. Thiel, T. Vorup-Jensen, C.M. Stover, W. Schwaeble, S.B. Laursen, K. Poulsen, A.C. Willis, P. Eggleton, S. Hansen, U. Holmskov, K.B. Reid, and J.C. Jensenius. A second serine protease associated with mannan-binding lectin that activates complement. *Nature*, 386:506–510, 1997.
- [2910] K.V. Thimann and S. Mahadevan. Nitrilase. I. Occurrence, preparation, and general properties of the enzyme. *Arch. Biochem. Biophys.*, 105:133–141, 1964.
- [2911] N.V. Thoai, F. Thome-Beau, and A. Olomucki. [Induction and specificity of enzymes of the new catabolic arginine pathway]. *Biochim. Biophys. Acta*, 115:73–80, 1966.
- [2912] R.W. Thoma and W.H. Peterson. The enzymatic degradation of soluble bound biotin. *J. Biol. Chem.*, 210:569–579, 1954.
- [2913] J. Thomas and J.E. Cronan. The enigmatic acyl carrier protein phosphodiesterase of *Escherichia coli*: genetic and enzymological characterization. *J. Biol. Chem.*, 280:34675–34683, 2005.
- [2914] L. Thomas, C.-H. Yang, and D.A. Goldthwait. Two DNA glycosylases in *Escherichia coli* which release primarily 3-methyladenine. *Biochemistry*, 21:1162–1169, 1982.
- [2915] P.W. Thomas, E.M. Stone, A.L. Costello, D.L. Tierney, and W. Fast. The quorum-quenching lactonase from *Bacillus thuringiensis* is a metalloprotein. *Biochemistry*, 44:7559–7569, 2005.
- [2916] J. Thompson, C.R. Gentry-Weeks, N.Y. Nguyen, J.E. Folk, , and S.A. Purification from *Fusobacterium mortiferum* ATCC 25557 of a 6-phosphoryl-*O*- $\alpha$ -D-glucopyranosyl:6-phosphoglucohydrolase that hydrolyses maltose 6-phosphate and related phospho- $\alpha$ -D-glucosides. *J. Bacteriol.*, 177:2505–2512, 1995.
- [2917] W. Thompson and R.M.C. Dawson. The triphosphoinositide phosphodiesterase of brain tissue. *Biochem. J.*, 91:237–243, 1964.
- [2918] N.A. Thornberry. Interleukin-1 $\beta$  converting enzyme. *Methods Enzymol.*, 244:615–631, 1994.
- [2919] N.A. Thornberry, H.G. Bull, J.R. Calaycay, K.T. Chapman, A.D. Howard, M.J. Kostura, D.K. Miller, S.M. Molineaux, J.R. Weidner, J. Aunins, K.O. Elliston, J.M. Ayala, , and F. J., Chin, J., Ding, G.J.-F., Egger, L.A., Gaffney, E.P., Limjuco, G., Palyha, O.C., Raju, S.M., Rolando, A.M., Salley, J.P., Yamin, T.-T. and Tocci, M.J. A novel heterodimeric cysteine protease is required for interleukin-1 $\beta$  processing in monocytes. *Nature*, 356:768–774, 1992.
- [2920] G. Thumm and F. Götz. Studies on prollystaphin processing and characterization of the lysostaphin immunity factor (Lif) of *Staphylococcus simulans* biovar staphylolyticus. *Mol. Microbiol.*, 23:1251–1265, 1997.
- [2921] G. Tian, A. Bhamidipati, N.J. Cowan, and S.A. Lewis. Tubulin folding cofactors as GTPase-activating proteins. GTP hydrolysis and the assembly of the  $\alpha/\beta$ -tubulin heterodimer. *J. Biol. Chem.*, 274:24054–24058, 1999.
- [2922] U. Tislar and A.J. Barrett. Thiol-dependent metallo-endopeptidase characteristics of Pz-peptidase in rat and rabbit. *Biochem. J.*, 267:531–533, 1990.
- [2923] K. Titani, , and M. A., Ericson, L. H., Walsh, K. A. and Neurath, H. Amino-acid sequence of thermolysin. *Nature New Biol.*, 238:35–37, 1972.
- [2924] K. Titani, H.-J. Torff, S. Hormel, S. Kumar, K.A. Walsh, J. Rödl, H. Neurath, and R. Zwilling. Amino acid sequence of a unique protease from the crayfish *Astacus fluviatilis*. *Biochemistry*, 26:222–226, 1987.
- [2925] R. Tjian. Regulation of viral transcription and DNA replication by the SV40 large T antigen. *Curr. Top. Microbiol. Immunol.*, 93:5–24, 1981.
- [2926] H. Tobe, F. Kojima, T. Aoyagi, and H. Umezawa. Purification by affinity chromatography using amastatin and properties of aminopeptidase A from pig kidney. *Biochim. Biophys. Acta*, 613:459–468, 1980.
- [2927] A. Tocilj, J.D. Schrag, Y. Li, B.L. Schneider, L. Reitzer, A. Matte, and M. Cygler. Crystal structure of *N*-succinylarginine dihydrolase AstB, bound to substrate and product, an enzyme from the arginine catabolic pathway of *Escherichia coli*. *J. Biol. Chem.*, 280:15800–15808, 2005.

- [2928] A. Tohgo, S. Takasawa, N. Noguchi, T. Koguma, K. Nata, T. Sugimoto, Y. Furuya, H. Yonekura, and H. Okamoto. Essential cysteine residues for cyclic ADP-ribose synthesis and hydrolysis by CD38. *J. Biol. Chem.*, 269:28555–28557, 1994.
- [2929] Z.A. Tökés, W.C. Woon, and S.M. Chambers. Digestive enzymes secreted by the carnivorous plant *Nepenthes macfarlanei* L. *Planta*, 119:39–46, 1974.
- [2930] F. Tokunaga, K. Nagasawa, S. Tamura, T. Miyata, S. Iwanaga, and W. Kisiel. The factor V-activating enzyme (RVV-V) from Russell's viper venom. Identification of isoproteins RVV-V $\alpha$ , -V $\beta$  and -V $\gamma$  and their complete amino acid sequences. *J. Biol. Chem.*, 263:17417–17481, 1988.
- [2931] F. Tokunaga, H. Nakajima, and S. Iwanaga. Further studies on lipopolysaccharide-sensitive serine protease zymogen (factor C): its isolation from *Limulus polyphemus* hemocytes and identification as an intracellular zymogen activated by  $\alpha$ -chymotrypsin, not by trypsin. *J. Biochem. (Tokyo)*, 109:150–157, 1991.
- [2932] S. Tombinson and P.W. Taylor. Neuraminidase associated with coliphage E that specifically depolymerizes the *Escherichia coli* K1 capsular polysaccharide. *J. Virol.*, 55:374–378, 1985.
- [2933] N.H. Tomczyk, J.E. Nettleship, R.L. Baxter, H.J. Crichton, S.P. Webster, and D.J. Campopiano. Purification and characterisation of the BIOH protein from the biotin biosynthetic pathway. *FEBS Lett.*, 513:299–304, 2002.
- [2934] F. Tomita and I. Takahashi. A novel enzyme, dCTP deaminase, found in *Bacillus subtilis* infected with phage PBS I. *Biochim. Biophys. Acta*, 179:18–27, 1969.
- [2935] N. Tomita, Y. Izumoto, A. Horii, S. Doi, H. Yokouchi, M. Ogawa, T. Mori, and K. Matsubara. Molecular cloning and nucleotide sequence of human pancreatic prechymotrypsinogen cDNA. *Biochem. Biophys. Res. Commun.*, 158:569–575, 1989.
- [2936] Y. Tomita and Y. Motokawa. Characterization and partial purification of a novel mannosylphosphodolichol phosphodiesterase from chicken liver microsomes. *Eur. J. Biochem.*, 170:363–368, 1987.
- [2937] B. Tomkinson and Ö. Zetterqvist. Immunological cross-reactivity between human tripeptidyl peptidase II and fibronectin. *Biochem. J.*, 267:149–154, 1990.
- [2938] R.V. Tomlinson and C.E. Ballou. Myoinositol polyphosphate intermediates in the dephosphorylation of phytic acid by phytase. *Biochemistry*, 1:166–171, 1962.
- [2939] K. Tomoda and H. Shimazono. Acid protease produced by *Trametes sanguinea* a wood-destroying fungus. Part I. Purification and crystallization of the enzyme. *Agric. Biol. Chem.*, 28:770–773, 1964.
- [2940] A.V. Toms, A.L. Haas, J.H. Park, T.P. Begley, and S.E. Ealick. Structural characterization of the regulatory proteins TenA and TenI from *Bacillus subtilis* and identification of TenA as a thiaminase II. *Biochemistry*, 44:2319–2329, 2005.
- [2941] H. Ton-That, G. Liu, S.K. Mazmanian, K.F. Faull, and O. Schneewind. Purification and characterization of sortase, the transpeptidase that cleaves surface proteins of *Staphylococcus aureus* at the LPXTG motif. *Proc. Natl. Acad. Sci. USA*, 96:12424–12429, 1999.
- [2942] C.G. Tong, M. Dauwalder, G.A. Clawson, C.L. Hatem, and S.J. Roux. The major nucleoside triphosphatase in pea (*Pisum sativum* L.) nuclei and in rat liver nuclei share common epitopes also present in nuclear lamins. *Plant Physiol.*, 101:1005–1011, 1993.
- [2943] L. Tong, G. Wengler, and M.G. Rossmann. Refined structure of Sindbis virus core protein and comparison with other chymotrypsin-like serine proteinase structures. *J. Mol. Biol.*, 230:228–247, 1993.
- [2944] N.T. Tong, A. Tsugita, and V. Keil-Dlouha. Purification and characterization of two high-molecular-mass forms of *Achromobacter collagenase*. *Biochim. Biophys. Acta*, 874:296–304, 1986.
- [2945] N.K. Tonks and P. Cohen. The protein phosphatases involved in cellular regulation. Identification of the inhibitor-2 phosphatases in rabbit skeletal muscle. *Eur. J. Biochem.*, 145:65–70, 1984.
- [2946] H.L. Tookey and A.K. Balls. Plant phospholipase D. I. Studies on cottonseed and cabbage phospholipase D. *J. Biol. Chem.*, 218:213–224, 1956.



- [2947] J. Torabinejad, J.L. Donahue, B.N. Gunesequera, M.J. Allen-Daniels, and G.E. Gillaspay. VTC4 is a bifunctional enzyme that affects myoinositol and ascorbate biosynthesis in plants. *Plant Physiol.*, 150:951–961, 2009.
- [2948] L.L. Torres and G.L. Salerno. A metabolic pathway leading to mannosylfructose biosynthesis in *Agrobacterium tumefaciens* uncovers a family of mannosyltransferases. *Proc. Natl. Acad. Sci. USA*, 104:14318–14323, 2007.
- [2949] A. Torres-Rosado, K.S. O’Shea, A. Tsuji, S.H. Chou, and K. Kurachi. Hepsin, a putative cell-surface serine protease, is required for mammalian cell growth. *Proc. Natl. Acad. Sci. USA*, 90:7181–7185, 1993.
- [2950] F. Toselli, M. Fredenwall, P. Svensson, X.Q. Li, A. Johansson, L. Weidolf, and M.A. Hayes. Oxetane substrates of human microsomal epoxide hydrolase. *Drug Metab. Dispos.*, 45:966–973, 2017.
- [2951] B.B. Touré, J.S. Munzer, A. Basak, S. Benjannet, J. Rochemont, C. Lazure, M. Chrétien, and N.G. Seidah. Biosynthesis and enzymatic characterization of human SKI-1/S1P and the processing of its inhibitory prosegment. *J. Biol. Chem.*, 275:2349–2358, 2000.
- [2952] T. Touze, D. Blanot, and D. Mengin-Lecreulx. Substrate specificity and membrane topology of *Escherichia coli* PgpB, an undecaprenyl pyrophosphate phosphatase. *J. Biol. Chem.*, 283:16573–16583, 2008.
- [2953] E.M. Towler, S.V. Gulnik, T.N. Bhat, D. Xie, E. Gustschina, T.R. Sumpter, N. Robertson, C. Jones, M. Sauter, N. Mueller-Lantzsch, C. Debouck, and J.W. Erickson. Functional characterization of the protease of human endogenous retrovirus, K10: can it complement HIV-1 protease. *Biochemistry*, 37:17137–17144, 1998.
- [2954] P. Towler, B. Staker, S.G. Prasad, S. Menon, J. Tang, T. Parsons, D. Ryan, M. Fisher, D. Williams, N.A. Dales, M.A. Patane, and M.W. Pantoliano. ACE2 X-ray structures reveal a large hinge-bending motion important for inhibitor binding and catalysis. *J. Biol. Chem.*, 279:17996–18007, 2004.
- [2955] C.A. Townsend. New reactions in clavulanic acid biosynthesis. *Curr. Opin. Chem. Biol.*, 6:583–589, 2002.
- [2956] M. Toyoshima and M. Nakajima. Human heparanase. Purification, characterization, cloning, and expression. *J. Biol. Chem.*, 274:24153–24160, 1999.
- [2957] J. Tözsér, I. Bláha, T.D. Copeland, E.M. Wondrak, and S. Oroszlan. Comparison of the HIV-1 and HIV-2 proteinases using oligopeptide substrates representing cleavage sites in Gag and Gag-Pol polyproteins. *FEBS Lett.*, 281:77–80, 1991.
- [2958] M.V. Tracey. Chitinase in some basidiomycetes. *Biochem. J.*, 61:579–586, 1955.
- [2959] K.L. Tran, P.A. Aronov, H. Tanaka, J.W. Newman, B.D. Hammock, and C. Morisseau. Lipid sulfates and sulfonates are allosteric competitive inhibitors of the N-terminal phosphatase activity of the mammalian soluble epoxide hydrolase. *Biochemistry*, 44:12179–12187, 2005.
- [2960] S. Traniello, M. Calcagno, and S. Pontremoli. Fructose 1,6-diphosphatase and sedoheptulose 1,7-diphosphatase from *Candida utilis*: purification and properties. *Arch. Biochem. Biophys.*, 146:603–610, 1971.
- [2961] T.W. Traut and S. Loechel. Pyrimidine catabolism: individual characterization of the three sequential enzymes with a new assay. *Biochemistry*, 23:2533–2539, 1984.
- [2962] D.B. Trentini, J. Fuhrmann, K. Mechtler, and T. Clausen. Chasing phosphoarginine proteins: development of a selective enrichment method using a phosphatase trap. LID - mcp.O113.035790 [pii]. *Mol. Cell. Proteomics*, 2014.
- [2963] F. Trijbels and G.D. Vogels. Allantoicase and ureidoglycolase in *Pseudomonas* and *Penicillium* species. *Biochim. Biophys. Acta*, 118:387–395, 1966.
- [2964] D.M. Trilling and H.V. Aposhian. Sequential cleavage of dinucleotides from DNA by phage Sp3 DNase. *Proc. Natl. Acad. Sci. USA*, 60:214–221, 1968.
- [2965] E.S. Trombetta, J.F. Simons, and A. Helenius. Endoplasmic reticulum glucosidase II is composed of a catalytic subunit, conserved from yeast to mammals, and a tightly bound noncatalytic HDEL-containing subunit. *J. Biol. Chem.*, 271:27509–27516, 1996.
- [2966] P.A. Trudinger. The metabolism of trithionate by *Thiobacillus X*. *Aust. J. Biol. Sci.*, 17:459–468, 1964.

- [2967] A.W. Truman, F. Huang, N.M. Llewellyn, and J.B. Spencer. Characterization of the enzyme BtrD from *Bacillus circulans* and revision of its functional assignment in the biosynthesis of butirosin. *Angew. Chem. Int. Ed. Engl.*, 46:1462–1464, 2007.
- [2968] K.J. Tsai and A.L. Linet. Formation of a phosphorylated enzyme intermediate by the cadA Cd<sup>2+</sup>-ATPase. *Arch. Biochem. Biophys.*, 305:267–270, 1993.
- [2969] S.-C. Tsai and J.L. Gaylor. Testicular sterols. V. Preparation and partial purification of a microsomal prenyl pyrophosphate pyrophosphohydrolase. *J. Biol. Chem.*, 241:4043–4050, 1966.
- [2970] A. Tsalaifouta, E. Psylinakis, E.G. Kapetaniou, D. Kotsifaki, A. Deli, A. Roidis, V. Bouriotis, and M. Kokkinidis. Purification, crystallization and preliminary X-ray analysis of the peptidoglycan *N*-acetylglucosamine deacetylase BC1960 from *Bacillus cereus* in the presence of its substrate (GlcNAc)<sub>6</sub>. *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.*, 64:203–205, 2008.
- [2971] M. Tsang, L.-S. and Schiff, J.A. Properties of enzyme fraction A from *Chlorella* and copurification of 3' (2'), 5'-biphosphonucleoside 3' (2')-phosphohydrolase, adenosine 5'-phosphosulfate sulfohydrolase and adenosine-5'-phosphosulfate cyclase activities. *Eur. J. Biochem.*, 65:113–121, 1976.
- [2972] W.R. Tschantz and R.E. Dalbey. Bacterial leader peptidase I. *Methods Enzymol.*, 244:285–301, 1994.
- [2973] W.R. Tschantz, M. Sung, V.M. Delgado-Partin, and R.E. Dalbey. A serine and a lysine residue implicated in the catalytic mechanism of the *Escherichia coli* leader peptidase. *J. Biol. Chem.*, 268:27349–27354, 1993.
- [2974] K. Tsurulnikov, N. Abuladze, A. Bragin, K. Faull, D. Cascio, R. Damoiseaux, M.J. Schibler, and A. Pushkin. Inhibition of aminoacylase 3 protects rat brain cortex neuronal cells from the toxicity of 4-hydroxy-2-nonenal mercapturate and 4-hydroxy-2-nonenal. *Toxicol. Appl. Pharmacol.*, 263:303–314, 2012.
- [2975] K. Tsurulnikov, N. Abuladze, D. Newman, S. Ryazantsev, T. Wolak, N. Magilnick, M.C. Koag, I. Kurtz, and A. Pushkin. Mouse aminoacylase 3: a metalloenzyme activated by cobalt and nickel. *Biochim. Biophys. Acta*, 1794:1049–1057, 2009.
- [2976] K.K. Tsuboi, G. Wiener, and P.B. Hudson. Acid phosphatase. VII. Yeast phosphomonoesterase; isolation procedure and stability characteristics. *J. Biol. Chem.*, 224:621–635, 1957.
- [2977] T. Tsuchiya, H. Ohta, K. Okawa, A. Iwamatsu, H. Shimada, T. Masuda, and K. Takamiya. Cloning of chlorophyllase, the key enzyme in chlorophyll degradation: finding of a lipase motif and the induction by methyl jasmonate. *Proc. Natl. Acad. Sci. USA*, 96:15362–15367, 1999.
- [2978] Y. Tsuda, N. Teno, Y. Okada, K. Wanaka, M. Bohgaki, A. Hijikata-Okunomiya, U. Okamoto, T. Naito, and S. Okamoto. Synthesis of tripeptide chloromethyl ketones and examination of their inhibitory effects on plasmin and plasma kallikrein. *Chem. Pharm. Bull.*, 37:3108–3111, 1989.
- [2979] M. Tsuji, Y. Nakanishi, H. Habuchi, K. Ishihara, and S. Suzuki. The common identity of UDP-*N*-acetylgalactosamine 4-sulfatase, nitrocatechol sulfatase (arylsulfatase), and chondroitin 4-sulfatase. *Biochim. Biophys. Acta*, 612:373–383, 1980.
- [2980] Y. Tsujisaka, J. Fukimoto, and T. Yamamoto. Specificity of crystalline saccharogenic amylase of moulds. *Nature*, 181:770–771, 1958.
- [2981] T. Tsujita, H. Ninomiya, and H. Okuda. *p*-Nitrophenyl butyrate hydrolyzing activity of hormone-sensitive lipase from bovine adipose tissue. *J. Lipid Res.*, 30:997–1004, 1989.
- [2982] T. Tsukamoto, S. Miura, T. Nakai, S. Yokota, N. Shimozawa, Y. Suzuki, T. Orii, Y. Fujiki, F. Sakai, A. Bogaki, H. Yasuno, and T. Osumi. Peroxisome assembly factor-2, a putative ATPase cloned by functional complementation on a peroxisome-deficient mammalian cell mutant. *Nat. Genet.*, 11:395–401, 1995.
- [2983] Y. Tsumuraya, , and N. , Hashimoto Y. and Kovac, P. Purification of exo-(1→3)-D-galactanase of *Irpex lacteus* (*Polyporus tulipiferae*) and its action on arabinogalactan-proteins. *J. Biol. Chem.*, 265:7207–7215, 1990.
- [2984] S. Tsunasawa, T. Masaki, M. Hirose, M. Soejima, and F. Sakiyama. The primary structure and structural characteristics of *Achromobacter lyticus* protease I, a lysine-specific serine protease. *J. Biol. Chem.*, 264:3832–3839, 1989.

- [2985] S. Tsunasawa, J.W. Stewart, and F. Sherman. Acylamino acid-releasing enzyme from rat liver. *J. Biol. Chem.*, 260:5832–5391, 1985.
- [2986] S. Tsunazawa, K. Narita, and K. Ogata. Acylamino acid-releasing enzyme from rat liver. *J. Biochem. (Tokyo)*, 77:89–102, 1975.
- [2987] D. Tsuru, A. Hattori, H. Tsuji, T. Yamamoto, and J. Fukumoto. Studies on mold proteases. Part II. Substrate specificity of acid protease of *Rhizopus chinensis*. *Agric. Biol. Chem.*, 33:1419–1426, 1969.
- [2988] D. Tsuru, K. Nakamura, T. Yoshimoto, and K. Fujiwara. Pyroglutamyl-peptidase from *Bacillus amyloliquefaciens*. An improved purification method and some properties of the enzyme. *Biochim. Biophys. Acta*, 791:117–122, 1984.
- [2989] D. Tsuru, I. Oka, and T. Yoshimoto. Creatinine decomposing enzymes in *Pseudomonas putida*. *Agric. Biol. Chem.*, 40:1011–1018, 1976.
- [2990] D. Tsuru, S. Shimada, S. Maruta, T. Yoshimoto, K. Oda, S. Murao, T. Miyata, and S. Iwanaga. Isolation and amino acid sequence of a peptide containing an epoxide-reactive residue from the thermolysin-digest of *Scytalidium lignicolum* acid protease B. *J. Biochem. (Tokyo)*, 99:1537–1539, 1986.
- [2991] T. Tsushida and T. Takeo. An enzyme hydrolyzing L-theanine in tea leaves. *Agric. Biol. Chem.*, 49:2913–2917, 1985.
- [2992] M. Tsusué. Studies on sepiapterin deaminase from the silkworm, *Bombyx mori*. Purification and some properties of the enzyme. *J. Biochem. (Tokyo)*, 69:781–788, 1971.
- [2993] D.R.P. Tulsiani, V.P. Coleman, and O. Touster. Asparagine-linked glycoprotein biosynthesis in rat brain: identification of glucosidase I, glucosidase II, and endomannosidase (glucosyl mannosidase). *Arch. Biochem. Biophys.*, 277:114–121, 1990.
- [2994] D.R.P. Tulsiani, S.C. Hubbard, P.W. Robbins, and O. Touster.  $\alpha$ -D-Mannosidases of rat liver Golgi membranes. Mannosidase II is the GlcNAcMAN<sub>5</sub>-cleaving enzyme in glycoprotein biosynthesis and mannosidases IA and IB are the enzymes converting Man<sub>9</sub> precursors to Man<sub>5</sub> intermediates. *J. Biol. Chem.*, 257:3660–3668, 1982.
- [2995] D.R.P. Tulsiani, D.J. Opheim, and O. Touster. Purification and characterization of  $\alpha$ -D-mannosidase from rat liver golgi membranes. *J. Biol. Chem.*, 252:3227–3233, 1977.
- [2996] P. Tumbale, J.S. Williams, M.J. Schellenberg, T.A. Kunkel, and R.S. Williams. Aprataxin resolves adenylated RNA-DNA junctions to maintain genome integrity. *Nature*, 506:111–115, 2014.
- [2997] K. Tung and J.H. Nordin. Structure of the tetrasaccharide produced by the hydrolysis of nigeran by the enzyme mycodextranase. *Biochim. Biophys. Acta*, 158:154–156, 1968.
- [2998] J. Turková, O. Mikes, K. Hayashi, G. Danno, and L. Polgár. Alkaline proteinases of the genus *Aspergillus*. *Biochim. Biophys. Acta*, 257:257–263, 1972.
- [2999] D.H. Turner and J.F. Turner. The hydrolysis of glucose monophosphates by a phosphatase preparation from pea seeds. *Biochem. J.*, 74:486–491, 1960.
- [3000] D.H. Turner and J.F. Turner. Adenylic deaminase of pea seeds. *Biochem. J.*, 79:143–147, 1961.
- [3001] R.T. Turner, J.A. Loy, C. Nguyen, T. Devasamudram, A.K. Ghosh, G. Koelsch, and J. Tang. Specificity of memapsin 1 and its implications on the design of memapsin 2 ( $\beta$ -secretase) inhibitor selectivity. *Biochemistry*, 41:8742–8746, 2002.
- [3002] R.T. Turner, K. oelsch III, Hong G., Castenheira L., Ghosh P., Tang A., and J. Subsite specificity of memapsin 2 ( $\beta$ -secretase): implications for inhibitor design. *Biochemistry*, 40:10001–10006, 2001.
- [3003] T. Turnšek, I. Kregar, and D. Lebez. Acid sulphhydryl protease from calf lymph nodes. *Biochim. Biophys. Acta*, 403:514–520, 1975.
- [3004] A. Turzynski and R. Mentlein. Prolyl aminopeptidase from rat brain and kidney. Action on peptides and identification as leucyl aminopeptidase. *Eur. J. Biochem.*, 190:509–515, 1990.
- [3005] D. Uccelletti, C. O’Callaghan, P. Berninsone, I. Zemtseva, C. Abeijon, and C.B. Hirschberg. ire-1-dependent transcriptional up-regulation of a luminal uridine diphosphatase from *Caenorhabditis elegans*. *J. Biol. Chem.*, 279:27390–27398, 2004.

- [3006] H. Uchida, T. Nanri, Y. Kawabata, I. Kusakabe, , and K. Purification and characterization of intracellular  $\alpha$ -glucuronidase from *Aspergillus niger*. *Biosci. Biotechnol. Biochem.*, 56:1608–1615, 1992.
- [3007] F. Uchino, Y. Kurono, and S. Doi. Purification and some properties of crystalline acid protease from *Acrocyndrium* sp. *Agric. Biol. Chem.*, 31:428–434, 1967.
- [3008] R.L. Ulrich. Quorum quenching: enzymatic disruption of *N*-acylhomoserine lactone-mediated bacterial communication in *Burkholderia thailandensis*. *Appl. Environ. Microbiol.*, 70:6173–6180, 2004.
- [3009] H. Umezawa, Y. Takahashi, A. Fujii, T. Saino, T. Shirai, and T. Takita. Preparation of bleomycinic acid. Hydrolysis of bleomycin B2 by a *Fusarium* acylglutamine amidohydrolase. *J. Antibiot.*, 26:117–119, 1974.
- [3010] A.M. Umnov, S.N. Egorov, S.E. Mansurova, and I.S. Kulaev. A comparative characterisation of the polyphosphatases of *Neurospora crassa* and some other organisms. *Biochemistry (Moscow)*, 39:309–312, 1974.
- [3011] T. Unemoto and M. Hayashi. Chloride ion as a modifier of 2',3'-cyclic phosphodiesterase purified from halophilic *Vibrio alginolyticus*. *Biochim. Biophys. Acta*, 171:89–102, 1969.
- [3012] T. Ungar, M. Nagelschmidt, and H. Struck. *N*-Acetylaminoacyl-*p*-nitranilidase from human placenta. Purification and some properties. *Eur. J. Biochem.*, 97:205–211, 1979.
- [3013] A. Ungewickell, C. Hugge, M. Kisseleva, S.C. Chang, J. Zou, Y. Feng, E.E. Galyov, M. Wilson, and P.W. Majerus. The identification and characterization of two phosphatidylinositol-4,5-bisphosphate 4-phosphatases. *Proc. Natl. Acad. Sci. USA*, 102:18854–18859, 2005.
- [3014] L. Uotila. Preparation and assay of glutathione thiol esters. Survey of human liver glutathione thiol esterases. *Biochemistry*, 12:3938–3943, 1973.
- [3015] L. Uotila. Purification and characterization of *S*-2-hydroxyacylglutathione hydrolase (glyoxalase II) from human liver. *Biochemistry*, 12:3944–3951, 1973.
- [3016] L. Uotila. Purification and properties of *S*-succinylglutathione hydrolase from human liver. *J. Biol. Chem.*, 254:7024–7029, 1979.
- [3017] L. Uotila and M. Koivusalo. Purification and properties of *S*-formylglutathione hydrolase from human liver. *J. Biol. Chem.*, 249:7664–7672, 1974.
- [3018] S. Urban and M. Freeman. Intramembrane proteolysis controls diverse signalling pathways throughout evolution. *Curr. Opin. Genet. Dev.*, 12:512–518, 2002.
- [3019] S. Urban and M. Freeman. Substrate specificity of Rhomboid intramembrane proteases is governed by helix-breaking residues in the substrate transmembrane domain. *Mol. Cell*, 11:1425–1434, 2003.
- [3020] S. Urban, J.R. Lee, and M. Freeman. *Drosophila* Rhomboid-1 defines a family of putative intramembrane serine proteases. *Cell*, 107:173–182, 2001.
- [3021] S. Urban, J.R. Lee, and M. Freeman. A family of Rhomboid intramembrane proteases activates all *Drosophila* membrane-tethered EGF-like ligands. *EMBO J.*, 21:4277–4286, 2002.
- [3022] S. Urban, D. Schlieper, and M. Freeman. Conservation of intramembrane proteolytic activity and substrate specificity in prokaryotic and eukaryotic Rhomboids. *Curr. Biol.*, 12:1507–1512, 2002.
- [3023] S. Urban and M.S. Wolfe. Reconstitution of intramembrane proteolysis in vitro reveals that pure rhomboid is sufficient for catalysis and specificity. *Proc. Natl. Acad. Sci. USA*, 102:1883–1888, 2005.
- [3024] V. Uttamsingh, D.A. Keller, and M.W. Anders. Acylase I-catalyzed deacetylation of *N*-acetyl-L-cysteine and *S*-alkyl-*N*-acetyl-L-cysteines. *Chem. Res. Toxicol.*, 11:800–809, 1998.
- [3025] A.M. Vaccaro, M. Muscillo, and K. Suzuki. Characterization of human glucosylsphingosine glucosyl hydrolase and comparison with glucosylceramidase. *Eur. J. Biochem.*, 146:315–321, 1985.
- [3026] T.I. Vaganova, N.M. Ivanova, and V.M. Stepanov. Isolation and properties of the acid metalloproteinase from *Aspergillus oryzae*. *Biochemistry (Mosc)*, 53:1171–1178, 1988.

- [3027] P.R. Vagelos and A.R. Larrabee. Acyl carrier protein. IX. Acyl carrier protein hydrolase. *J. Biol. Chem.*, 242:1776–1781, 1967.
- [3028] G.V. Vahouny and C.R. Tradwell. Enzymatic synthesis and hydrolysis of cholesterol esters. *Methods Biochem. Anal.*, 16:219–272, 1968.
- [3029] C.M. Vallejo, C.D. Lobatón, M. Quintanilla, A. Sillero, and M.A.G. Sillero. Dinucleosidasetetraphosphatase in rat liver and *Artemia salina*. *Biochim. Biophys. Acta*, 438:304–309, 1976.
- [3030] L.A. Valls, C.P. Hunter, J.H. Rothman, and T.H. Stevens. Protein sorting in yeast: the localization determinant of yeast vacuolar carboxypeptidase Y resides in the propeptide. *Cell*, 48:887–897, 1987.
- [3031] M.A. Valvano, P. Messner, and P. Kosma. Novel pathways for biosynthesis of nucleotide-activated *glycero-manno-heptose* precursors of bacterial glycoproteins and cell surface polysaccharides. *Microbiology*, 148:1979–1989, 2002.
- [3032] H. van den Bosch. Intracellular phospholipases A. *Biochim. Biophys. Acta*, 604:191–246, 1980.
- [3033] H. van den Bosch, A.J. Aarsman, J.G.N. De Jong, and L.L.M. van Deenen. Studies on lysophospholipases. I. Purification and some properties of a lysophospholipase from beef pancreas. *Biochim. Biophys. Acta*, 296:94–104, 1973.
- [3034] H. van den Bosch, A.J. Aarsman, and L.L.M. van Deenen. Isolation and properties of a phospholipase A<sub>1</sub> activity from beef pancreas. *Biochim. Biophys. Acta*, 348:197–209, 1974.
- [3035] H. van den Bosch, G.M. Vianen, and G.P.H. van Heusden. Lysophospholipase-transacylase from rat lung. *Methods Enzymol.*, 71:513–521, 1981.
- [3036] J.M. van den Elsen, D.A. Kuntz, and D.R. Rose. Structure of Golgi  $\alpha$ -mannosidase II: a target for inhibition of growth and metastasis of cancer cells. *EMBO J.*, 20:3008–3017, 2001.
- [3037] C. van der Drift, P.E. van Helvoort, and G.D. Vogels. S-Ureidoglycolate dehydrogenase: purification and properties. *Arch. Biochem. Biophys.*, 145:465–469, 1971.
- [3038] C. van der Drift and G.D. Vogels. Effect of metal and hydrogen ions on the activity and stability of allantoicase. *Biochim. Biophys. Acta*, 198:339–352, 1971.
- [3039] L. van der Drift, G.D. Vogels, and C. van der Drift. Allantoin racemase: a new enzyme from *Pseudomonas* species. *Biochim. Biophys. Acta*, 391:240–248, 1975.
- [3040] C.J. van der Vlugt-Bergmans and M.J. van der Werf. Genetic and biochemical characterization of a novel monoterpene  $\epsilon$ -lactone hydrolase from *Rhodococcus erythropolis* DCL14. *Appl. Environ. Microbiol.*, 67:733–741, 2001.
- [3041] M.J. van der Werf, K.M. Overkamp, and J.A.M. de Bont. Limonene-1,2-epoxide hydrolase from *Rhodococcus erythropolis* DCL14 belongs to a novel class of epoxide hydrolases. *J. Bacteriol.*, 180:5052–5057, 1998.
- [3042] M.J. van der Werf, H.J. Swarts, and J.A. de Bont. *Rhodococcus erythropolis* DCL14 contains a novel degradation pathway for limonene. *Appl. Environ. Microbiol.*, 65:2092–2102, 1999.
- [3043] P. van der Werf, M. Orłowski, and A. Meister. Enzymatic conversion of 5-oxo-L-proline (L-pyrrolidone carboxylate) to L-glutamate coupled with cleavage of adenosine triphosphate to adenosine diphosphate, a reaction in the  $\gamma$ -glutamyl cycle. *Proc. Natl. Acad. Sci. USA*, 68:2982–2985, 1971.
- [3044] E. van Dijk, N. Cougot, S. Meyer, S. Babajko, E. Wahle, and B. Seraphin. Human Dcp2: a catalytically active mRNA decapping enzyme located in specific cytoplasmic structures. *EMBO J.*, 21:6915–6924, 2002.
- [3045] E. van Dijk, H. Le Hir, and B. Seraphin. DcpS can act in the 5'-3' mRNA decay pathway in addition to the 3'-5' pathway. *Proc. Natl. Acad. Sci. USA*, 100:12081–12086, 2003.
- [3046] L.C. van Dinten, S. Rensen, A.E. Gorbalenya, and E.J. Snijder. Proteolytic processing of the open reading frame 1b-encoded part of arterivirus replicase is mediated by nsp4 serine protease and is essential for virus replication. *J. Virol.*, 73:2027–2037, 1999.
- [3047] L.A.H. van Loon-Klaasen, H.T. Cuypers, H. van Westreenen, W.W. de Jong, and H. Bloemendal. The primary structure of bovine lens leucine aminopeptidase. Complete amino acid sequence of the N-terminal cyanogen bromide fragment and site limited tryptic digestion. *Biochem. Biophys. Res. Commun.*, 95:334–341, 1980.

- [3048] M. van Tienhoven, J. Atkins, Y. Li, and P. Glynn. Human neuropathy target esterase catalyzes hydrolysis of membrane lipids. *J. Biol. Chem.*, 277:20942–20948, 2002.
- [3049] L. Vandeputte-Rutten, R.A. Kramer, J. Kroon, N. Dekker, M.R. Egmond, and P. Gros. Crystal structure of the outer membrane protease OmpT from *Escherichia coli* suggests a novel catalytic site. *EMBO J.*, 20:5033–5039, 2001.
- [3050] P. Vanderslice, S.M. Ballinger, E.K. Tam, S.M. Goldstein, C.S. Craik, and G. Caughey. Human mast cell tryptase: multiple cDNAs and genes reveal a multigene serine protease family. *Proc. Natl. Acad. Sci. USA*, 87:3811–3815, 1990.
- [3051] N.F. Vanzo, Y.S. Li, B. Py, E. Blum, C.F. Higgins, L.C. Raynal, H.M. Krisch, and A.J. Carpousis. Ribonuclease E organizes the protein interactions in the *Escherichia coli* RNA degradosome. *Genes Dev.*, 12:2770–2781, 1998.
- [3052] B. Varga, O. Barabas, J. Kovari, J. Toth, E. Hunyadi-Gulyas, E. Klement, K.F. Medzihradsky, F. Tolgyesi, J. Fidy, and B.G. Vertessy. Active site closure facilitates juxtaposition of reactant atoms for initiation of catalysis by human dUTPase. *FEBS Lett.*, 581:4783–4788, 2007.
- [3053] J.E. Varner. Urease. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 4, pages 247–256. Academic Press, New York, 2nd edition, 1960.
- [3054] N. Vasantha, L.D. Thompson, C. Rhodes, C. Banner, J. Nagle, and D. Filpula. Genes for alkaline protease and neutral protease from *Bacillus amyloliquefaciens* contain a large open reading frame between the regions coding for signal sequence and mature protein. *J. Bacteriol.*, 159:811–819, 1984.
- [3055] Y. Vasilopoulos, M.J. Cork, R. Murphy, H.C. Williams, D.A. Robinson, G.W. Duff, S.J. Ward, and R. Tazi-Ahni. Genetic association between an AACC insertion in the 3'UTR of the stratum corneum chymotryptic enzyme gene and atopic dermatitis. *J. Invest. Dermatol.*, 123:62–66, 2004.
- [3056] M. Vaughan, J.E. Berger, and D. Steinberg. Hormone-sensitive lipase and monoglyceride lipase activities in adipose tissue. *J. Biol. Chem.*, 239:401–409, 1964.
- [3057] G. Velasco, A.A. Ferrando, X.S. Puente, L.M. Sanchez, and C. López-Otín. Human cathepsin O. Molecular cloning from a breast carcinoma, production of the active enzyme in *Escherichia coli*, and expression analysis in human tissues. *J. Biol. Chem.*, 269:27136–27142, 1994.
- [3058] J. Verchot. Potyvirus helper component proteinase. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 677–679. Handbook of Proteolytic Enzymes, London, 1998.
- [3059] J.H. Verheijen. Tissue-type plasminogen activator and fast-acting plasminogen activator inhibitor in plasma. *Methods Enzymol.*, 163:302–309, 1988.
- [3060] B. Verjans, F. De Smedt, R. Lecocq, V. Vanweyenberg, C. Moreau, and C. Erneux. Cloning and expression in *Escherichia coli* of a dog thyroid cDNA encoding a novel inositol 1,4,5-trisphosphate 5-phosphatase. *Biochem. J.*, 300:85–90, 1994.
- [3061] L.A. Verkruyse, , and S.L. Lysosomal targeting of palmitoyl-protein thioesterase. *J. Biol. Chem.*, 271:15831–15836, 1996.
- [3062] S. Verma, T. Goldammer, and R. Aitken. Cloning and expression of activation induced cytidine deaminase from *Bos taurus*. *Vet. Immunol. Immunopathol.*, 134:151–159, 2010.
- [3063] W. Versées, K. Decanniere, E. Van Holsbeke, N. Devroede, and J. Steyaert. Enzyme-substrate interactions in the purine-specific nucleoside hydrolase from *Trypanosoma vivax*. *J. Biol. Chem.*, 277:15938–15946, 2002.
- [3064] N.D. Vetter, D.M. Langill, S. Anjum, J. Boisvert-Martel, R.C. Jagdhane, E. Omene, H. Zheng, K.E. van Straaten, I. Asiamah, E.S. Krol, D.A. Sanders, and D.R. Palmer. A previously unrecognized kanosamine biosynthesis pathway in *Bacillus subtilis*. *J. Am. Chem. Soc.*, 135:5970–5973, 2013.
- [3065] C. Vicente and M.E. Legaz. Preparation and properties of agmatine amidinohydrolase of *Evernia prunastri*. *Physiol. Plant.*, 55:335–339, 1982.
- [3066] C. Vickers, P. Hales, V. Kaushik, L. Dick, J. Gavin, J. Tang, K. Godbout, T. Parsons, E. Baronas, F. Hsieh, S. Acton, M. Patane, A. Nichols, and P. Tummino. Hydrolysis of biological peptides by human angiotensin-converting enzyme-related carboxypeptidase. *J. Biol. Chem.*, 277:14838–14843, 2002.

- [3067] M.J. Vincent, A.J. Sanchez, B.R. Erickson, A. Basak, M. Chretien, N.G. Seidah, and S.T. Nichol. Crimean-Congo hemorrhagic fever virus glycoprotein proteolytic processing by subtilase SKI-1. *J. Virol.*, 77:8640–8649, 2003.
- [3068] S. Vincenzetti, A. Cambi, J. Neuhard, K. Schnorr, M. Grelloni, and A. Vita. Cloning, expression, and purification of cytidine deaminase from *Arabidopsis thaliana*. *Protein Expr. Purif.*, 15:8–15, 1999.
- [3069] S. Visser, A.J.P.M. Robben, and C.J. Slangen. Specificity of a cell-envelope-located proteinase (PIII-type) from *Lactococcus lactis* subsp. *cremoris* AM1 in its action on bovine  $\beta$ -casein. *Appl. Microbiol. Biotechnol.*, 35:477–483, 1991.
- [3070] S. Visser, C.J. Slangen, and P.J. van Rooijen. Peptide substrates for chymosin (rennin). Interaction sites in  $\kappa$ -casein-related sequences located outside the (103-108)-hexapeptide region that fits into the enzyme's active-site cleft. *Biochem. J.*, 244:553–558, 1987.
- [3071] N. Vitale, J. Moss, and M. Vaughan. Molecular characterization of the GTPase-activating domain of ADP-ribosylation factor domain protein 1 (ARD1). *J. Biol. Chem.*, 273:2553–2560, 1998.
- [3072] H.J. Vogel. Path of ornithine synthesis in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA*, 39:578–583, 1953.
- [3073] H.J. Vogel and D.M. Bonner. Acetylornithine deacetylase of *Escherichia coli*: partial purification and some properties. *J. Biol. Chem.*, 218:97–106, 1956.
- [3074] G.D. Vogels. Reversible activation of allantoate amidohydrolase by acid-pretreatment and other properties of the enzyme. *Biochim. Biophys. Acta*, 113:277–291, 1966.
- [3075] G.D. Vogels and C. Van der Drift. Degradation of purines and pyrimidines by microorganisms. *Bacteriol. Rev.*, 40:403–468, 1976.
- [3076] V.M. Vogt. Purification and further properties of single-strand-specific nuclease from *Aspergillus oryzae*. *Eur. J. Biochem.*, 33:192–200, 1973.
- [3077] F.N. Vogtle, S. Wortelkamp, R.P. Zahedi, D. Becker, C. Leidhold, K. Gevaert, J. Kellermann, W. Voos, A. Sickmann, N. Pfanner, and C. Meisinger. Global analysis of the mitochondrial N-proteome identifies a processing peptidase critical for protein stability. *Cell*, 139:428–439, 2009.
- [3078] K. von Figura. Human  $\alpha$ -N-acetylglucosaminidase. 1. Purification and properties. *Eur. J. Biochem.*, 80:523–533, 1977.
- [3079] K. von Figura. Human  $\alpha$ -N-acetylglucosaminidase. 2. Activity towards natural substrates and multiple recognition forms. *Eur. J. Biochem.*, 80:535–542, 1977.
- [3080] V. Vreys and G. David. Mammalian heparanase: what is the message. *J. Cell. Mol. Med.*, 11:427–452, 2007.
- [3081] D. Wachsmuth, I. Fritze, and G. Pfeleiderer. An aminopeptidase occurring in pig kidney. I. An improved method of preparation. Physical and enzymatic properties. *Biochemistry*, 5:169–174, 1966.
- [3082] F.W. Wagner, A.M. Spiekerman, and J.M. Prescott. Leucostoma peptidase A. Isolation and physical properties. *J. Biol. Chem.*, 243:4486–4493, 1968.
- [3083] A. Waheed, A. Hasilik, and K. von Figura. Processing of the phosphorylated recognition marker in lysosomal enzymes. Characterization and partial purification of a microsomal  $\alpha$ -N-acetylglucosaminyl phosphodiesterase. *J. Biol. Chem.*, 256:5717–5721, 1981.
- [3084] I. Waizenegger, J. Gimenez-Abian, D. Wernic, and J. Peters. Regulation of human separase by securin binding and autocleavage. *Curr. Biol.*, 12:1368–1378, 2002.
- [3085] M. Wakabayashi and W.H. Fishman. The comparative ability of  $\beta$ -glucuronidase preparations (liver, *Escherichia coli*, *Helix pomatia*, and *Patella vulgata*) to hydrolyze certain steroid glucosiduronic acids. *J. Biol. Chem.*, 236:996–1001, 1961.
- [3086] M. Wakayama, T. Ashika, Y. Miyamoto, T. Yoshikawa, Y. Sonoda, K. Sakai, and M. Moriguchi. Primary structure of N-acyl-D-glutamate amidohydrolase from *Alcaligenes xylosoxydans* subsp. *xylosoxydans* A-6. *J. Biochem. (Tokyo)*, 118:204–209, 1995.

- [3087] M. Wakayama, S. Hayashi, Y. Yatsuda, Y. Katsuno, K. Sakai, and M. Moriguchi. Overproduction of D-aminoacylase from *Alcaligenes xylosoxydans* subsp. *xylosoxydans* A-6 in *Escherichia coli* and its purification. *Protein Expr. Purif.*, 7:395–399, 1996.
- [3088] M. Wakayama, Y. Katsuno, S. Hayashi, Y. Miyamoto, K. Sakai, , and M. Cloning and sequencing of a gene encoding D-aminoacylase from *Alcaligenes xylosoxydans* subsp. *xylosoxydans* A-6 and expression of the gene in *Escherichia coli*. *Biosci. Biotechnol. Biochem.*, 59:2115–2119, 1995.
- [3089] M. Wakayama, Y. Miura, K. Oshima, K. Sakai, , and M. Metal-characterization of *N*-acyl-D-glutamate amidohydrolase from *Pseudomonas* sp. strain 5f-1. *Biosci. Biotechnol. Biochem.*, 59:1489–1492, 1995.
- [3090] M. Wakayama, T. Tsutsumi, H. Yada, K. Sakai, , and M. Chemical modification of histidine residue of *N*-acyl-D-glutamate amidohydrolase from *Pseudomonas* sp. 5f-1. *Biosci. Biotechnol. Biochem.*, 60:650–653, 1996.
- [3091] M. Wakayama, E. Watanabe, Y. Takenaka, Y. Miyamoto, Y. Tau, K. Sakai, , and M. Cloning, expression and nucleotide sequence of the *N*-acyl-D-aspartate amidohydrolase gene from *Alcaligenes xylosoxydans* subsp. *xylosoxydans* A-6. *J. Ferment. Bioeng.*, 80:311–317, 1995.
- [3092] P.R. Wakeley, R. Dorazi, N.T. Hoa, J.R. Bowyer, and S.M. Cutting. Proteolysis of SpoIVB is a critical determinant in signalling of pro- $\sigma^K$  processing in *Bacillus subtilis*. *Mol. Microbiol.*, 36:1336–1348, 2000.
- [3093] D.M. Walker, S. Urbe, S.K. Dove, D. Tenza, G. Raposo, and M.J. Clague. Characterization of MTMR3. an inositol lipid 3-phosphatase with novel substrate specificity. *Curr. Biol.*, 11:1600–1605, 2001.
- [3094] F.J. Walker, W.G. Owen, and C.T. Esmon. Characterization of the prothrombin activator from the venom of *Oxyuranus scutellatus scutellatus* (taipan venom). *Biochemistry*, 19:1020–1023, 1980.
- [3095] G.J. Walker and A. Pulkownik. Degradation of dextrans by an  $\alpha$ -1,6-glucan glucohydrolase from *Streptococcus mitis*. *Carbohydr. Res.*, 29:1–14, 1973.
- [3096] J.B. Walker and M. Skorvaga. Streptomycin biosynthesis and metabolism. Phosphate transfer from dihydrostreptomycin 6-phosphate to inosamines, streptomine, and 2-deoxystreptomine. *J. Biol. Chem.*, 248:2441–2446, 1973.
- [3097] M.S. Walker and J.B. Walker. Streptomycin biosynthesis. Separation and substrate specificities of phosphatases acting on guanidinodeoxy-scyllo-inositol phosphate and streptomycin-(streptidino)phosphate. *J. Biol. Chem.*, 246:7034–7040, 1971.
- [3098] E.F. Wallace, C.J. Evans, S.M. Jurik, I.N. Mefford, and J.D. Barchas. Carboxypeptidase B activity from adrenal medulla. Is it involved in the processing of proenkephalin? *Life Sci.*, 31:1793–1796, 1982.
- [3099] K. Wallenfels and O.P. Malhotra. In P.D. Boyer, H. Lardy, and K. Myrbäck, editors, *The Enzymes*, volume 4, pages 409–430. Academic Press, New York, 2nd edition, 1960.
- [3100] S.R. Wallner, B.M. Nestl, and K. Faber. Highly enantioselective sec-alkyl sulfatase activity of *Sulfolobus acidocaldarius* DSM 639. *Org. Lett.*, 6:5009–5010, 2004.
- [3101] K.A. Walsh. Trypsinogens and trypsins of various species. *Methods Enzymol.*, 19:41–63, 1970.
- [3102] R. Walter, W.H. Simmons, and T. Yoshimoto. Proline specific endo- and exopeptidases. *Mol. Cell. Biochem.*, 30:111–127, 1980.
- [3103] R. Walter and T. Yoshimoto. Postproline cleaving enzyme: kinetic studies of size and stereospecificity of its active site. *Biochemistry*, 17:4139–4144, 1978.
- [3104] B. Wang, G.-P. Shi, P.M. Yao, Z. Li, H.A. Chapman, and D. Brömme. Human cathepsin F. Molecular cloning, functional expression, tissue localization, and enzymatic characterization. *J. Biol. Chem.*, 273:32000–32008, 1998.
- [3105] B.Z. Wang, P. Guo, B.J. Hang, L. Li, J. He, and S.P. Li. Cloning of a novel pyrethroid-hydrolyzing carboxylesterase gene from *Sphingobium* sp. strain JZ-1 and characterization of the gene product. *Appl. Environ. Microbiol.*, 75:5496–5500, 2009.
- [3106] D Wang, H. Yu, J. Song, Y. Xu, , and F. Enzyme kinetics of ginsenosidase type IV hydrolyzing 6-*O*-multi-glycosides of protopanaxatriol type ginsenosides. *Process Biochem.*, 47:133–138, 2012.



- [3107] D.M. Wang, H.S. Yu, J.G. Song, Y.F. Xu, C.Y. Liu, and F.X. Jin. A novel ginsenosidase from an *Aspergillus* strain hydrolyzing 6-*O*-multi-glycosides of protopanaxatriol-type ginsenosides, named ginsenosidase type IV. *J Microbiol Biotechnol*, 21:1057–1063, 2011.
- [3108] G. Wang, A. Olczak, L.S. Forsberg, and R.J. Maier. Oxidative stress-induced peptidoglycan deacetylase in *Helicobacter pylori*. *J. Biol. Chem.*, 284:6790–6800, 2009.
- [3109] H. Wang, H. Pang, M. Bartlam, and Z. Rao. Crystal structure of human E1 enzyme and its complex with a substrate analog reveals the mechanism of its phosphatase/enolase activity. *J. Mol. Biol.*, 348:917–926, 2005.
- [3110] H. Wang, K.I. Piatkov, C.S. Brower, and A. Varshavsky. Glutamine-specific N-terminal amidase, a component of the N-end rule pathway. *Mol. Cell*, 34:686–695, 2009.
- [3111] J. Wang, Y. Okamoto, J. Morishita, K. Tsuboi, A. Miyatake, and N. Ueda. Functional analysis of the purified anandamide-generating phospholipase D as a member of the metallo- $\beta$ -lactamase family. *J. Biol. Chem.*, 281:12325–12335, 2006.
- [3112] J. Wang, S.H. Rho, H.H. Park, and S.H. Eom. Correction of X-ray intensities from an HsIV-HsIU co-crystal containing lattice-translocation defects. *Acta Crystallogr. D Biol. Crystallogr.*, 61:932–941, 2005.
- [3113] L. Wang, H. Huang, H.H. Nguyen, K.N. Allen, P.S. Mariano, and D. Dunaway-Mariano. Divergence of biochemical function in the HAD superfamily: D-glycero-D-manno-heptose-1,7-bisphosphate phosphatase (GmhB). *Biochemistry*, 49:1072–1081, 2010.
- [3114] L. Wang, Z. Lu, K.N. Allen, P.S. Mariano, and D. Dunaway-Mariano. Human symbiont *Bacteroides thetaiotaomicron* synthesizes 2-keto-3-deoxy-D-glycero-D-galacto-nononic acid (KDN). *Chem. Biol.*, 15:893–897, 2008.
- [3115] L. Wang, M. Miura, L. Bergeron, H. Zhu, and J. Yuan. *Ich-1*, an *Ice/ced-3*-related gene, encodes both positive and negative regulators of programmed cell death. *Cell*, 78:739–750, 1994.
- [3116] L.H. Wang, L.X. Weng, Y.H. Dong, and L.H. Zhang. Specificity and enzyme kinetics of the quorum-quenching *N*-acyl homoserine lactone lactonase (AHL-lactonase). *J. Biol. Chem.*, 279:13645–13651, 2004.
- [3117] S. Wang, A.M. Haapalainen, F. Yan, Q. Du, P.C. Tyler, G.B. Evans, A. Rinaldo-Matthis, R.L. Brown, G.E. Norris, S.C. Almo, and V.L. Schramm. A picomolar transition state analogue inhibitor of MTAN as a specific antibiotic for *Helicobacter pylori*. *Biochemistry*, 51:6892–6894, 2012.
- [3118] S. Wang, M. Miura, Y.K. Jung, H. Zhu, E. Li, and J. Yuan. Murine caspase-11, an ICE-interacting protease, is essential for the activation of ICE. *Cell*, 92:501–509, 1998.
- [3119] T.F. Wang and G. Guidotti. CD39 is an ecto-(Ca<sup>2+</sup>, Mg<sup>2+</sup>)-apyrase. *J. Biol. Chem.*, 271:9898–9901, 1996.
- [3120] T.P. Wang, H.Z. Sable, and J.O. Lampen. Enzymatic deamination of cytosine nucleosides. *J. Biol. Chem.*, 184:17–28, 1950.
- [3121] W. Wang, D.F. Hendriks, and S. Scharpé. Carboxypeptidase U, a plasma carboxypeptidase with high affinity for plasminogen. *J. Biol. Chem.*, 269:15937–15944, 1994.
- [3122] W. Wang, M. Maniar, R. Jain, J. Jacobs, J. Trias, and Z. Yuan. A fluorescence-based homogeneous assay for measuring activity of UDP-3-*O*-(*R*-3-hydroxymyristoyl)-*N*-acetylglucosamine deacetylase. *Anal. Biochem.*, 290:338–346, 2001.
- [3123] Y. Wang, J. Niemi, K. Airas, K. Ylihonko, J. Hakala, and P. Mantsala. Modifications of aclacinomycin T by aclacinomycin methyl esterase (RdmC) and aclacinomycin-10-hydroxylase (RdmB) from *Streptomyces purpurascens*. *Biochim. Biophys. Acta*, 1480:191–200, 2000.
- [3124] Y. Wang, Z. Nimec, T.J. Ryan, J.A. Dias, and J. Galivan. The properties of the secreted  $\gamma$ -glutamyl hydrolases from H35 hepatoma cells. *Biochim. Biophys. Acta*, 1164:227–235, 1993.
- [3125] Y. Wang, Y. Zhang, and Y. Ha. Crystal structure of a rhomboid family intramembrane protease. *Nature*, 444:179–180, 2006.
- [3126] J.B. Ward, C.A.M. Curtis, C. Taylor, and R.S. Buxton. Purification and characterization of two phage PBSX-induced lytic enzymes of *Bacillus subtilis* 168: an *N*-acetylmuramoyl-L-alanine amidase and an *N*-acetylmuramidase. *J. Gen. Microbiol.*, 128:1171–1178, 1982.

- [3127] F. Warnaar. Triterpene ester synthesis in latex of *Euphorbia* species. *Phytochemistry*, 26:2715–2721, 1987.
- [3128] A.H. Warner and F.J. Finamore. Isolation, purification, and characterization of  $P^1, P^4$ -diguanosine 5'-tetraphosphate asymmetrical-pyrophosphohydrolase from brine shrimp eggs. *Biochemistry*, 4:1568–1575, 1965.
- [3129] H.R. Warner and W.E.M. Lands. The metabolism of plasmalogen: enzymatic hydrolysis of the vinyl ether. *J. Biol. Chem.*, 236:2404–2409, 1961.
- [3130] D.E. Warnock and S.L. Schmid. Dynamin GTPase, a force-generating molecular switch. *Bioessays*, 18:885–893, 1996.
- [3131] M.J. Warren, E. Raux, H.L. Schubert, and J.C. Escalante-Semerena. The biosynthesis of adenosylcobalamin (vitamin B<sub>12</sub>). *Nat. Prod. Rep.*, 19:390–412, 2002.
- [3132] H.D. Van Wart and D.R. Steinbrink. Complementary substrate specificities of class I and class II collagenases from *Clostridium histolyticum*. *Biochemistry*, 24:6520–6526, 1985.
- [3133] R. Watanabe, K. Ohishi, Y. Maeda, N. Nakamura, and T. Kinoshita. Mammalian PIG-L and its yeast homologue Gpi12p are *N*-acetylglucosaminylphosphatidylinositol de-*N*-acetylases essential in glycosylphosphatidylinositol biosynthesis. *Biochem. J.*, 339:185–192, 1999.
- [3134] T.R. Waters and P.F. Swann. Kinetics of the action of thymine DNA glycosylase. *J. Biol. Chem.*, 273:20007–20014, 1998.
- [3135] T.R. Waters and P.F. Swann. Thymine-DNA glycosylase and G to A transition mutations at CpG sites. *Mutat. Res.*, 462:137–147, 2000.
- [3136] L.M. Watkins, R. Rodriguez, D. Schneider, R. Broderick, M. Cruz, R. Chambers, E. Ruckman, M. Cody, and G.T. Mrachko. Purification and characterization of the aromatic desulfonase, 2-(2'-hydroxyphenyl)benzenesulfinate desulfonase. *Arch. Biochem. Biophys.*, 415:14–23, 2003.
- [3137] G.H. Watson, C. Houghton, and R.B. Cain. Microbial metabolism of the pyridine ring. The metabolism of pyridine-3,4-diol (3,4-dihydroxypyridine) by *Agrobacterium* sp. *Biochem. J.*, 140:277–292, 1974.
- [3138] N. Watson and D. Apirion. Ribonuclease F, a putative processing endoribonuclease from *Escherichia coli*. *Biochem. Biophys. Res. Commun.*, 103:543–551, 1981.
- [3139] C. Vander Wauven and V. Stalon. Occurrence of succinyl derivatives in the catabolism of arginine in *Pseudomonas cepacia*. *J. Bacteriol.*, 164:882–886, 1985.
- [3140] A. Wawrzynow, D. Wojtkowiak, J. Marszalek, B. Banecki, M. Jonsen, B. Graves, C. Georgopoulos, and M. Zylicz. The ClpX heat-shock protein of *Escherichia coli*, the ATP-dependent substrate specificity component of the ClpP-ClpX protease, is a novel molecular chaperone. *EMBO J.*, 14:1867–1877, 1995.
- [3141] E.C. Webb and P.F.W. Morrow. The activation of an arylsulphatase from ox liver by chloride and other anions. *Biochem. J.*, 73:7–15, 1959.
- [3142] J.M. Weber. Adenovirus protease. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 741–743. Academic Press, London, 1998.
- [3143] A. Webster, R.T. Hay, and G. Kemp. The adenovirus protease is activated by a virus-coded disulphide-linked peptide. *Cell*, 72:274–275, 1993.
- [3144] G.R. Webster, E.A. Marples, and R.H.S. Thompson. Glycerylphosphorylcholine diesterase activity in nervous tissue. *Biochem. J.*, 65:374–377, 1957.
- [3145] W.J. Wechter, A.J. Mikulski, and M. Laskowski. , Sr. Gradation of specificity with regard to sugar among nucleases. *Biochem. Biophys. Res. Commun.*, 30:318–322, 1968.
- [3146] J.F. Wedgwood and J.L. Strominger. Enzymatic activities in cultured human lymphocytes that dephosphorylate dolichyl pyrophosphate and dolichyl phosphate. *J. Biol. Chem.*, 255:1120–1123, 1980.
- [3147] L. Wei, E. Clauser, F. Alhenc-Gelas, and P. Corvol. The two homologous domains of human angiotensin I-converting enzyme interact differently with competitive inhibitors. *J. Biol. Chem.*, 267:13398–13405, 1992.

- [3148] S. Wei, K. Lai, S. Patel, R. Piantedosi, H. Shen, V. Colantuoni, F.B. Kraemer, and W.S. Blaner. Retinyl ester hydrolysis and retinol efflux from BFC-1 $\beta$  adipocytes. *J. Biol. Chem.*, 272:14159–14165, 1977.
- [3149] A.J. Weightman, A.L. Weightman, and J.H. Slater. Stereospecificity of 2-monochloropropionate dehalogenation by the two dehalogenases of *Pseudomonas* P3: evidence for two different dehalogenation mechanisms. *J. Gen. Microbiol.*, 128:1755–1762, 1982.
- [3150] J. Weigl and W. Yashe. The enzymic hydrolysis of carrageenan by *Pseudomonas carrageenovora*: purification of a  $\kappa$ -carrageenase. *Can. J. Microbiol.*, 12:939–947, 1966.
- [3151] L. Weignerova, T. Filipi, D. Manglova, and V. Kren. Induction, purification and characterization of  $\alpha$ -*N*-acetylgalactosaminidase from *Aspergillus niger*. *Appl. Microbiol. Biotechnol.*, 79:769–774, 2008.
- [3152] W. Weil-Malherbe and R.H. Green. Ammonia formation in brain. 2. Brain adenylic deaminase. *Biochem. J.*, 61:218–224, 1955.
- [3153] R. Weimberg and M. Doudoroff. The oxidation of L-arabinose by *Pseudomonas saccharophila*. *J. Biol. Chem.*, 217:607–624, 1955.
- [3154] L. Weinstein and P. Albersheim. Structure of plant cell walls. IX. Purification and partial characterization of a wall-degrading endo-arabinase and an arabinosidase from *Bacillus subtilis*. *Plant Physiol.*, 63:425–432, 1979.
- [3155] B. Weissmann. The transglycosylative action of testicular hyaluronidase. *J. Biol. Chem.*, 216:783–794, 1955.
- [3156] B. Weissmann, H. Chao, and P. Chow. A glucosamine O,*N*-disulfate O-sulfohydrolase with a probable role in mammalian catabolism of heparan sulfate. *Biochem. Biophys. Res. Commun.*, 97:827–833, 1980.
- [3157] B. Weissmann, G. Rowen, J. Marshall, and D. Friederici. Mammalian  $\alpha$ -acetylglucosaminidase. Enzymic properties, tissue distribution, and intracellular localization. *Biochemistry*, 6:207–214, 1967.
- [3158] H.G. Welgus and G.A. Grant. Degradation of collagen substrates by a trypsin-like serine protease from the fiddler crab *Uca pugnator*. *Biochemistry*, 22:2228–2233, 1983.
- [3159] H.G. Welgus, G.A. Grant, J.J. Jeffrey, and A.Z. Eisen. Substrate specificity of the collagenolytic serine protease from *Uca pugnator*: studies with collagenous substrates. *Biochemistry*, 21:5183–5189, 1982.
- [3160] L. Wells, Y. Gao, J.A. Mahoney, K. Vosseller, C. Chen, A. Rosen, and G.W. Hart. Dynamic O-glycosylation of nuclear and cytosolic proteins: further characterization of the nucleocytoplasmic  $\beta$ -*N*-acetylglucosaminidase, O-GlcNAcase. *J. Biol. Chem.*, 277:1755–1761, 2002.
- [3161] X.E. Wells and E.M. Lees. Ureidoglycolate amidohydrolase from developing French bean fruits (*Phaseolus vulgaris* [L.]). *Arch. Biochem. Biophys.*, 287:151–159, 1991.
- [3162] A.K. Werner, T. Romeis, and C.P. Witte. Ureide catabolism in *Arabidopsis thaliana* and *Escherichia coli*. *Nat. Chem. Biol.*, 6:19–21, 2010.
- [3163] E. Werries, E. Wollek, A. Gottschalk, and E. Buddecke. Separation of *N*-acetyl- $\alpha$ -glucosaminidase and *N*-acetyl- $\alpha$ -galactosaminidase from ox spleen. Cleavage of the O-glycosidic linkage between carbohydrate and polypeptide in ovine and bovine submaxillary glycoprotein by *N*-acetyl- $\alpha$ -galactosaminidase. *Eur. J. Biochem.*, 10:445–449, 1969.
- [3164] D.W.S. Westlake. Microbiological degradation of quercitrin. *Can. J. Microbiol.*, 9:211–220, 1963.
- [3165] J. Westling, A.J. Fosang, K. Last, V.P. Thompson, K.N. Tomkinson, T. Hebert, T. McDonagh, L.A. Collins-Racie, E.R. LaVallie, E.A. Morris, and J.D. Sandy. ADAMTS4 cleaves at the aggrecanase site (Glu<sup>373</sup>-Ala<sup>374</sup>) and secondarily at the matrix metalloproteinase site (Asn<sup>341</sup>-Phe<sup>342</sup>) in the aggrecan interglobular domain. *J. Biol. Chem.*, 277:16059–16066, 2002.
- [3166] T. Wex, B. Levy, H. Wex, and D. Brömme. Human cathepsins F and W: A new subgroup of cathepsins. *Biochem. Biophys. Res. Commun.*, 259:401–407, 1999.
- [3167] D.E. Wexler, D.E. Chenoweth, and P.P. Cleary. Mechanism of action of the group A streptococcal C5a inactivator. *Proc. Natl. Acad. Sci. USA*, 82:8144–8148, 1985.

- [3168] R.L. Whistler and E. Masek. Enzymatic hydrolysis of xylan. *J. Am. Chem. Soc.*, 77:1241–1243, 1955.
- [3169] D.R. Whitaker, K.R. Hanson, and P.K. Datta. Improved procedures for preparation and characterization of myrothecium cellulase. 2. Purification procedures. *Can. J. Biochem. Physiol.*, 41:671–696, 1963.
- [3170] D.R. Whitaker and C. Roy. Concerning the nature of the  $\alpha$ - and  $\beta$ -lytic proteases of *Sorangium* sp. *Can. J. Biochem.*, 45:911–911, 1967.
- [3171] D.R. Whitaker, C. Roy, C.S. Tsai, and L. Juraöek. Lytic enzymes of *Sorangium* sp. A comparison of the proteolytic properties of the  $\alpha$ - and  $\beta$ -lytic proteases. *Can. J. Biochem.*, 43:1961–1970, 1965.
- [3172] J.R. Whitaker. Protease of *Endothia parasitica*. *Methods Enzymol.*, 19:436–445, 1970.
- [3173] K.N. White and D.B. Hope. Partial purification and characterization of a microsomal carboxylesterase specific for salicylate esters from guinea-pig liver. *Biochim. Biophys. Acta*, 785:138–147, 1984.
- [3174] R.J. White and C.A. Pasternak. The purification and properties of *N*-acetylglucosamine 6-phosphate deacetylase from *Escherichia coli*. *Biochem. J.*, 105:121–125, 1967.
- [3175] J.L. Whittingham, J.P. Turkenburg, C.S. Verma, M.A. Walsh, and G. Grogan. The 2-Å crystal structure of 6-oxo camphor hydrolase. New structural diversity in the crotonase superfamily. *J. Biol. Chem.*, 278:1744–1750, 2003.
- [3176] D.A. Whittington, K.M. Rusche, H. Shin, C.A. Fierke, and D.W. Christianson. Crystal structure of LpxC, a zinc-dependent deacetylase essential for endotoxin biosynthesis. *Proc. Natl. Acad. Sci. USA*, 100:8146–8150, 2003.
- [3177] S. Wickham, M.B. West, P.F. Cook, and M.H. Hanigan. Gamma-glutamyl compounds: substrate specificity of  $\gamma$ -glutamyl transpeptidase enzymes. *Anal. Biochem.*, 414:208–214, 2011.
- [3178] R.B. Wickner and H. Tabor. *N*-Formimino-L-glutamate iminohydrolase (*Pseudomonas* sp.). *Methods Enzymol.*, 17B:80–84, 1971.
- [3179] E. Widemann, L. Miesch, R. Lugan, E. Holder, C. Heinrich, Y. Aubert, M. Miesch, F. Pinot, and T. Heitz. The amidohydrolases IAR3 and ILL6 contribute to jasmonoyl-isoleucine hormone turnover and generate 12-hydroxyjasmonic acid upon wounding in *Arabidopsis* leaves. *J. Biol. Chem.*, 288:31701–31714, 2013.
- [3180] J.R. Widhalm, C. van Oostende, F. Furt, and G.J. Basset. A dedicated thioesterase of the Hotdog-fold family is required for the biosynthesis of the naphthoquinone ring of vitamin K<sub>1</sub>. *Proc. Natl. Acad. Sci. USA*, 106:5599–5603, 2009.
- [3181] G. Wiederschain and E. Beyer. [Interrelation of  $\alpha$ -D-fucosidase and  $\alpha$ -D-galactosidase activities in man and animals]. *Dokl. Akad. Nauk S.S.S.R.*, 231:486–488, 1976.
- [3182] G. Wiederschain and A. Prokopenkov.  $\beta$ -D-Galactosidase and  $\beta$ -D-fucosidase of pig kidney. *Arch. Biochem. Biophys.*, 158:539–543, 1973.
- [3183] G.Y. Wiederschain, E.M. Beyer, B.A. Klyaschitsy, and A.S. Shashkov. Specificity patterns of different types of human fucosidase. Recognition of a certain region of the pyranose ring in sugars by the enzymes. *Biochim. Biophys. Acta*, 659:434–444, 1981.
- [3184] P.E. Wilcox. Chymotrypsinogens - chymotrypsins. *Methods Enzymol.*, 19:64–108, 1970.
- [3185] S.M. Wilhelm, I.E. Collier, B.L. Marmer, A.Z. Eisen, G.A. Grant, and G.I. Goldberg. SV40-transformed human lung fibroblasts secrete a 92-kDa type IV collagenase which is identical to that secreted by normal human macrophages. *J. Biol. Chem.*, 264:17213–17221, 1989.
- [3186] S. Wilk, E. Wilk, and R.P. Magnusson. Purification, characterization and cloning of a cytosolic aspartyl aminopeptidase. *J. Biol. Chem.*, 273:15961–15970, 1998.
- [3187] S. Wilk and E.K. Wilk. Pyroglutamyl peptidase II, a thyrotropin releasing hormone degrading enzyme: purification and specificity studies of the rabbit brain enzyme. *Neurochem. Int.*, 15:81–89, 1989.
- [3188] S.H. Wilkes, M.E. Bayliss, and J.M. Prescott. Critical ionizing groups in *Aeromonas* neutral protease. *J. Biol. Chem.*, 263:1821–1825, 1988.

- [3189] S.H. Wilkes and J.M. Prescott. Aeromonas neutral protease. *Methods Enzymol.*, 45:404–415, 1976.
- [3190] B.M. Wilkinson, J. Purswani, and C.J. Stirling. Yeast GTB1 encodes a subunit of glucosidase II required for glycoprotein processing in the endoplasmic reticulum. *J. Biol. Chem.*, 281:6325–6333, 2006.
- [3191] K.D. Wilkinson. Ubiquitin C-terminal hydrolase. In A.J. Barrett, N.D. Rawlings, and J.F. Woessner, editors, *Handbook of Proteolytic Enzymes*, pages 470–472. Academic Press, London, 1998.
- [3192] D.C. Williams, J.R. Whitaker, and P.V. Caldwell. Hydrolysis of peptide bonds of the oxidized B-chain of insulin by *Endothia parasitica* protease. *Arch. Biochem. Biophys.*, 149:52–61, 1972.
- [3193] T.W. Willis and A.T. Tu. Purification and biochemical characterization of atroxase, a nonhemorrhagic fibrinolytic protease from western diamondback rattlesnake venom. *Biochemistry*, 27:4769–4777, 1988.
- [3194] B. Winchester. Role of  $\alpha$ -D-mannosidases in the biosynthesis and catabolism of glycoproteins. *Biochem. Soc. Trans.*, 12:522–524, 1984.
- [3195] D.R. Wines, J.M. Brady, D.B. Pritchett, J.L. Roberts, and R.J. MacDonald. Organization and expression of the rat kallikrein gene family. *J. Biol. Chem.*, 264:7653–7662, 1989.
- [3196] J. Winkelman and A.L. Lehninger. Aldono- and uronolactonase of animal tissues. *J. Biol. Chem.*, 233:794–799, 1958.
- [3197] R.G. Winkler, D.G. Blevins, and D.D. Randall. Ureide catabolism in soybeans. III. Ureidoglycolate amidohydrolase and allantoate amidohydrolase are activities of an allantoate degrading enzyme complex. *Plant Physiol.*, 86:1084–1088, 1988.
- [3198] C.J. Winrow, M.L. Hemming, D.M. Allen, G.B. Quistad, J.E. Casida, and C. Barlow. Loss of neuropathy target esterase in mice links organophosphate exposure to hyperactivity. *Nat. Genet.*, 33:477–485, 2003.
- [3199] C. Wirblich, M. Sibilica, M.B. Boniotti, C. Rossi, H.J. Thiel, and G. Meyers. 3C-like protease of rabbit hemorrhagic disease virus: identification of cleavage sites in the ORF1 polyprotein and analysis of cleavage specificity. *J. Virol.*, 69:7159–7168, 1995.
- [3200] M. Wiskerchen and M.S. Collett. Pestivirus gene expression: protein p80 of bovine viral diarrhea virus is a proteinase involved in polyprotein processing. *Virology*, 184:341–350, 1991.
- [3201] O. Wiss and F. Weber. Die Reindarstellung der Kynureninase. *Hoppe-Seyler's Z. Physiol. Chem.*, 304:232–240, 1956.
- [3202] C. Witte, R.E. Jensen, M.P. Yaffe, and G. Schatz. MAS1, a gene essential for yeast mitochondrial assembly, encodes a subunit of the mitochondrial processing protease. *EMBO J.*, 7:1439–1447, 1988.
- [3203] A. Wlodawer, M. Li, Z. Dauter, A. Gustchina, K. Uchida, H. Oyama, B.M. Dunn, and K. Oda. Carboxyl proteinase from *Pseudomonas* defines a novel family of subtilisin-like enzymes. *Nat. Struct. Biol.*, 8:442–446, 2001.
- [3204] A. Wlodawer, M. Li, A. Gustchina, H. Oyama, B.M. Dunn, and K. Oda. Structural and enzymatic properties of the sedolisin family of serine-carboxyl peptidases. *Acta Biochim. Pol.*, 50:81–102, 2003.
- [3205] J.F. Woessner, Taplin Jr., and C.J. Purification and properties of a small latent matrix metalloproteinase of the rat uterus. *J. Biol. Chem.*, 263:16918–16925, 1988.
- [3206] D.H. Wolf and C. Ehmann. Carboxypeptidase S from yeast: regulation of its activity during vegetative growth and differentiation. *FEBS Lett.*, 91:59–62, 1978.
- [3207] J. Wolf, A.P. Gerber, and W. Keller. *tadA*, an essential tRNA-specific adenosine deaminase from *Escherichia coli*. *EMBO J.*, 21:3841–3851, 2002.
- [3208] W.A. Wolf and G.M. Brown. The biosynthesis of folic acid. X. Evidence for an Amadori rearrangement in the enzymatic formation of dihydroneopterin triphosphate from GTP. *Biochim. Biophys. Acta*, 192:468–478, 1969.
- [3209] J.B. Wolfe, B.B. Britton, , and H.I. Glucosamine degradation by *Escherichia coli*. III. Isolation and studies of "phosphoglucosaminisomerase". *Arch. Biochem. Biophys.*, 66:333–339, 1957.
- [3210] R.L. Wolz and J.S. Bond. Meprins A and B. *Methods Enzymol.*, 248:325–345, 1995.

- [3211] S.K. Wong, S. Sato, and D.W. Lazinski. Substrate recognition by ADAR1 and ADAR2. *RNA*, 7:846–858, 2001.
- [3212] R.G. Woodbury, M. Everitt, and H. Neurath. Mast cell proteases. *Methods Enzymol.*, 80:588–609, 1981.
- [3213] J.D. Woodson and J.C. Escalante-Semerena. CbiZ, an amidohydrolase enzyme required for salvaging the coenzyme B<sub>12</sub> precursor cobinamide in archaea. *Proc. Natl. Acad. Sci. USA*, 101:3591–3596, 2004.
- [3214] R. Woscholski and P.J. Parker. Inositol phosphatases: constructive destruction of phosphoinositides and inositol phosphates. In S. Cockcroft, editor, *Biology of Phosphoinositides*, pages 320–338. Biology of Phosphoinositides, Oxford, 2000.
- [3215] J.W. Wray and R.H. Abeles. The methionine salvage pathway in *Klebsiella pneumoniae* and rat liver. Identification and characterization of two novel dioxynases. *J. Biol. Chem.*, 270:3147–3153, 1995.
- [3216] M. Wright, G. Buttin, and J. Hurwitz. The isolation and characterization from *Escherichia coli* of an adenosine triphosphate-dependent deoxyribonuclease directed by *rec B, C* genes. *J. Biol. Chem.*, 246:6543–6555, 1971.
- [3217] B. Wu, Y. Liu, Q. Zhao, S. Liao, J. Zhang, M. Bartlam, W. Chen, and Z. Rao. Crystal structure of RS21-C<sub>6</sub>, involved in nucleoside triphosphate pyrophosphohydrolysis. *J. Mol. Biol.*, 367:1405–1412, 2007.
- [3218] J. Wu, A.K. Bera, R.J. Kuhn, and J.L. Smith. Structure of the Flavivirus helicase: implications for catalytic activity, protein interactions, and proteolytic processing. *J. Virol.*, 79:10268–10277, 2005.
- [3219] L.C. Wu, D.R. Pfeiffer, E.A. Calhoon, F. Madiari, G. Marcucci, S. Liu, and M.S. Jurkowitz. Purification, identification, and cloning of lysoplasmalogenase, the enzyme that catalyzes hydrolysis of the vinyl ether bond of lysoplasmalogen. *J. Biol. Chem.*, 286:24916–24930, 2011.
- [3220] P.C. Wu, Y.H. Liu, Z.Y. Wang, X.Y. Zhang, H. Li, W.Q. Liang, N. Luo, J.M. Hu, J.Q. Lu, T.G. Luan, and L.X. Cao. Molecular cloning, purification, and biochemical characterization of a novel pyrethroid-hydrolyzing esterase from *Klebsiella* sp. strain ZD112. *J. Agric. Food Chem.*, 54:836–842, 2006.
- [3221] Q. Wu, J.M. Lahti, G.M. Air, P.D. Burrows, and M.D. Cooper. Molecular cloning of the murine BP-1/6C3 antigen: a member of the zinc-dependent metallopeptidase family. *Proc. Natl. Acad. Sci. USA*, 87:993–997, 1990.
- [3222] T.K. Wu, R.W. Busby, T.A. Houston, D.B. McIlwaine, L.A. Egan, and C.A. Townsend. Identification, cloning, sequencing, and overexpression of the gene encoding proclavaminic amidino hydrolase and characterization of protein function in clavulanic acid biosynthesis. *J. Bacteriol.*, 177:3714–3720, 1995.
- [3223] W.I. Wu, Y. Liu, B. Riedel, J.B. Wissing, A.S. Fischl, and G.M. Carman. Purification and characterization of diacylglycerol pyrophosphate phosphatase from *Saccharomyces cerevisiae*. *J. Biol. Chem.*, 271:1868–1876, 1996.
- [3224] Z. Wu, G.D. Wright, and C.T. Walsh. Overexpression, purification, and characterization of VanX, a D-, D-dipeptidase which is essential for vancomycin resistance in *Enterococcus faecium* BM4147. *Biochemistry*, 34:2455–2463, 1995.
- [3225] S. Wydau, M.L. Ferri-Fioni, S. Blanquet, and P. Plateau. GEK1, a gene product of *Arabidopsis thaliana* involved in ethanol tolerance, is a D-aminoacyl-tRNA deacylase. *Nucleic Acids Res.*, 35:930–938, 2007.
- [3226] R.L. Wykle and J.M. Schremmer. A lysophospholipase D pathway in the metabolism of ether-linked lipids in brain microsomes. *J. Biol. Chem.*, 249:1742–1746, 1974.
- [3227] A. Wypijewska, E. Bojarska, M. Lukaszewicz, J. Stepinski, J. Jemielity, R.E. Davis, and E. Darzynkiewicz. 7-Methylguanosine diphosphate (m<sup>7</sup>GDP) is not hydrolyzed but strongly bound by decapping scavenger (DcpS) enzymes and potently inhibits their activity. *Biochemistry*, 51:8003–8013, 2012.
- [3228] Y.N. Xia, R. Morgan, I. Schildkraut, , and J.L. A site-specific single-strand endonuclease activity induced by NYs-1 virus-infection of a *Chlorella*-like green-alga. *Nucleic Acids Res.*, 16:9477–9487, 1988.
- [3229] Y. Xing, Z. Li, Y. Chen, J.B. Stock, P.D. Jeffrey, and Y. Shi. Structural mechanism of demethylation and inactivation of protein phosphatase 2A. *Cell*, 133:154–163, 2008.
- [3230] W. Xiong, L.-M. Chen, and J. Chao. Purification and characterization of a kallikrein-like T-kininogenase. *J. Biol. Chem.*, 265:2822–2827, 1990.

- [3231] D. Xu, N. Emoto, A. Giaid, C. Slaughter, S. Kaw, D. DeWit, and M. Yanagisawa. ECE-1: A membrane-bound metalloprotease that catalyzes the proteolytic activation of big endothelin-1. *Cell*, 78:473–485, 1994.
- [3232] J. Xu, E. Mendez, P.R. Caron, C. Lin, M.A. Murcko, M.S. Collett, and C.M. Rice. Bovine viral diarrhea virus NS3 serine proteinase: polyprotein cleavage sites, cofactor requirements, and molecular model of an enzyme essential for pestivirus replication. *J. Virol.*, 71:5312–5322, 1997.
- [3233] J. Xu, J.Y. Yang, Q.W. Niu, and N.H. Chua. *Arabidopsis* DCP2, DCP1, and VARICOSE form a decapping complex required for postembryonic development. *Plant Cell*, 18:3386–3398, 2006.
- [3234] Q. Xu, P. Abdubek, T. Astakhova, H.L. Axelrod, C. Bakolitsa, X. Cai, D. Carlton, C. Chen, H.J. Chiu, M. Chiu, T. Clayton, D. Das, M.C. Deller, L. Duan, K. Ellrott, C.L. Farr, J. Feuerhelm, J.C. Grant, A. Grzechnik, G.W. Han, L. Jaroszewski, K.K. Jin, H.E. Klock, M.W. Knuth, P. Kozbial, S.S. Krishna, A. Kumar, W.W. Lam, D. Marciano, M.D. Miller, A.T. Morse, E. Nigoghossian, A. Nopakun, L. Okach, C. Puckett, R. Reyes, H.J. Tien, C.B. Trame, H. van den Bedem, D. Weekes, T. Wooten, A. Yeh, K.O. Hodgson, J. Wooley, M.A. Elsliger, A.M. Deacon, A. Godzik, S.A. Lesley, and I.A. Wilson. Structure of the  $\gamma$ -D-glutamyl-L-diamino acid endopeptidase YkfC from *Bacillus cereus* in complex with L-Ala- $\gamma$ -D-Glu: insights into substrate recognition by NlpC/P60 cysteine peptidases. *Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun.*, 66:1354–1364, 2010.
- [3235] W. Xu, Y.H. Chooi, J.W. Choi, S. Li, J.C. Vederas, N.A. Da Silva, and Y. Tang. LovG: the thioesterase required for dihydromonacolin L release and lovastatin nonaketide synthase turnover in lovastatin biosynthesis. *Angew. Chem. Int. Ed. Engl.*, 52:6472–6475, 2013.
- [3236] W. Xu, C.A. Dunn, and M.J. Bessman. Cloning and characterization of the NADH pyrophosphatases from *Caenorhabditis elegans* and *Saccharomyces cerevisiae*, members of a Nudix hydrolase subfamily. *Biochem. Biophys. Res. Commun.*, 273:753–758, 2000.
- [3237] W. Xu, C.R. Jones, C.A. Dunn, and M.J. Bessman. Gene *ytkD* of *Bacillus subtilis* encodes an atypical nucleoside triphosphatase member of the Nudix hydrolase superfamily. *J. Bacteriol.*, 186:8380–8384, 2004.
- [3238] K.K. Yabusaki and C.E. Ballou. Long-chain fatty acyl-CoA thioesterases from *Mycobacterium smegmatis*. *Methods Enzymol.*, 71:242–246, 1981.
- [3239] F. Yagi, J. Fan, K. Tadera, and A. Kobayashi. Purification and characterization of carboxyl proteinase from *Aspergillus kawachii*. *Agric. Biol. Chem.*, 50:1029–1033, 1986.
- [3240] T. Yahraus, N. Braverman, G. Dodt, J.E. Kalish, J.C. Morrell, H.W. Moser, D., Gould Valle, , and PXAAA. 1, encodes a cytoplasmic ATPase required for stability of the PTS1 receptor. *EMBO J.*, 15:2914–2923, 1996.
- [3241] F. Yamada, N. Takahashi, and T. Murachi. Purification and characterization of a proteinase from pineapple fruit, fruit bromelain FA2. *J. Biochem. (Tokyo)*, 79:1223–1234, 1976.
- [3242] H. Yamada, K. Okamoto, K. Kodama, F. Noguchi, and S. Tanaka. Enzymatic studies on mannitol formation by *Piricularia oryzae*. *J. Biochem. (Tokyo)*, 49:404–410, 1961.
- [3243] K. Yamagami. Isolation of a choriolytic enzyme (hatching enzyme) of the teleost, *Oryzias latipes*. *Dev. Biol.*, 29:343–348, 1972.
- [3244] T. Yamagata, H. Saito, O. Habuchi, and S. Suzuki. Purification and properties of bacterial chondroitinases and chondrosulfatases. *J. Biol. Chem.*, 243:1523–1535, 1968.
- [3245] I. Yamaguchi, H. Shibata, H. Seto, and T. Misato. Isolation and purification of blasticidin S deaminase from *Aspergillus terreus*. *J. Antibiot.*, 28:7–14, 1975.
- [3246] Y. Yamakawa and T. Omori-Satoh. A protease in the venom of king cobra (*Ophiophagus hannah*): purification, characterization and substrate specificity on oxidized insulin B-chain. *Toxicon*, 26:1145–1155, 1988.
- [3247] Y. Yamakawa and T. Omori-Satoh. The sites of cleavage in oxidized insulin-B chain by a hemorrhagic protease derived from the venom of the habu (*Trimeresurus flavoviridis*). *Toxicon*, 26:227–231, 1988.
- [3248] K. Yamamoto, A. Nakayama, Y. Yamamoto, and S. Tabata. Val<sup>216</sup> decides the substrate specificity of  $\alpha$ -glucosidase in *Saccharomyces cerevisiae*. *Eur. J. Biochem.*, 271:3414–3420, 2004.

- [3249] K. Yamamoto, K. Oishi, I. Fujimatsu, and K. Komatsu. Production of R-(-)-mandelic acid from mandelonitrile by *Alcaligenes faecalis* ATCC 8750. *Appl. Environ. Microbiol.*, 57:3028–3032, 1991.
- [3250] S. Yamamoto-Katayama, M. Ariyoshi, K. Ishihara, T. Hirano, H. Jingami, and K. Morikawa. Crystallographic studies on human BST-1/CD157 with ADP-ribosyl cyclase and NAD glycohydrolase activities. *J. Mol. Biol.*, 316:711–723, 2002.
- [3251] N. Yamano, Y. Matsushita, Y. Kamada, S. Fujishima, , and M. Purification and characterization of *N*-acetylglucosamine 6-phosphate deacetylase with activity against *N*-acetylglucosamine from *Vibrio cholerae* non-O1. *Biosci. Biotechnol. Biochem.*, 60:1320–1323, 1996.
- [3252] Q. Yan, W. Zhou, X. Li, M. Feng, and P. Zhou. Purification method improvement and characterization of a novel ginsenoside-hydrolyzing  $\beta$ -glucosidase from *Paecilomyces Bainier* sp. 229. *Biosci. Biotechnol. Biochem.*, 72:352–359, 2008.
- [3253] H. Yanagisawa and Y. Suzuki. Preparation and properties of *N*-carbamylputrescine amidohydrolase from maize shoots. *Phytochemistry*, 21:2201–2203, 1982.
- [3254] T. Yanagita and J.W. Foster. A bacterial riboflavin hydrolase. *J. Biol. Chem.*, 221:593–607, 1956.
- [3255] T. Yanai and M. Sato. Purification and characterization of an  $\alpha$ -L-rhamnosidase from *Pichia angusta* X349. *Biosci. Biotechnol. Biochem.*, 64:2179–2185, 2000.
- [3256] H. Yanase, T. Sakai, and K. Tonomuro. Purification, crystallization and some properties of  $\beta$ -cyano-L-alanine-degrading enzyme in *Pseudomonas* sp. 13. *Agric. Biol. Chem.*, 47:473–482, 1983.
- [3257] F. Yang, W. Hu, H. Xu, C. Li, B. Xia, and C. Jin. Solution structure and backbone dynamics of an endopeptidase HycI from *Escherichia coli*: implications for mechanism of the [NiFe] hydrogenase maturation. *J. Biol. Chem.*, 282:3856–3863, 2007.
- [3258] F. Yang, L.H. Wang, J. Wang, Y.H. Dong, J.Y. Hu, and L.H. Zhang. Quorum quenching enzyme activity is widely conserved in the sera of mammalian species. *FEBS Lett.*, 579:3713–3717, 2005.
- [3259] M.Y. Yang, E. Ferrari, and D.J. Henner. Cloning of the neutral protease gene of *Bacillus subtilis* and the use of the cloned gene to create an in vitro-derived deletion mutation. *J. Bacteriol.*, 160:15–21, 1984.
- [3260] T. Yang, J. Li, H.X. Wang, and Y. Zeng. A geraniol-synthase gene from *Cinnamomum tenuipilum*. *Phytochemistry*, 66:285–293, 2005.
- [3261] C. Yao, M.R. Karabasil, N. Purwanti, X. Li, T. Akamatsu, N. Kanamori, and K. Hosoi. Tissue kallikrein mK13 is a candidate processing enzyme for the precursor of interleukin-1 $\beta$  in the submandibular gland of mice. *J. Biol. Chem.*, 281:7968–7976, 2006.
- [3262] M. Yao, T. Ose, H. Sugimoto, A. Horiuchi, A. Nakagawa, S. Wakatsuki, D. Yokoi, T. Murakami, M. Honma, and I. Tanaka. Crystal structure of 1-aminocyclopropane-1-carboxylate deaminase from *Hansenula saturnus*. *J. Biol. Chem.*, 275:34557–34565, 2000.
- [3263] R. Yao, Z. Nimec, T.J. Ryan, and J. Galivan. Identification, cloning, and sequencing of a cDNA coding for rat  $\gamma$ -glutamyl hydrolase. *J. Biol. Chem.*, 271:8525–8528, 1996.
- [3264] R. Yao, M.S. Rhee, and J. Galivan. Effects of  $\gamma$ -glutamyl hydrolase on folyl and antifolylpolyglutamates in cultured H35 hepatoma cells. *Mol. Pharmacol.*, 48:505–511, 1995.
- [3265] R. Yao, E. Schneider, T.J. Ryan, and J. Galivan. Human  $\gamma$ -glutamyl hydrolase: cloning and characterization of the enzyme expressed in vitro. *Proc. Natl. Acad. Sci. USA*, 93:10134–10138, 1996.
- [3266] K. Yaoi and Y. Mitsuishi. Purification, characterization, cloning, and expression of a novel xyloglucan-specific glycosidase, oligoxyloglucan reducing end-specific cellobiohydrolase. *J. Biol. Chem.*, 277:48276–48281, 2002.
- [3267] A. Yaron. Dipeptidyl carboxypeptidase from *Escherichia coli*. *Methods Enzymol.*, 45:599–610, 1976.
- [3268] A. Yaron and A. Berger. Aminopeptidase-P. *Methods Enzymol.*, 19:522–534, 1970.
- [3269] A. Yaron and D. Mlynar. Aminopeptidase-P. *Biochem. Biophys. Res. Commun.*, 32:658–663, 1968.



- [3270] T. Yasuhara, T. Nakai, and A. Ohashi. Aminopeptidase Y, a new aminopeptidase from *Saccharomyces cerevisiae*. Purification, properties, localization, and processing by protease B. *J. Biol. Chem.*, 269:13644–13650, 1994.
- [3271] K. Yasuhira, Y. Tanaka, H. Shibata, Y. Kawashima, A. Ohara, D. Kato, M. Takeo, and S. Negoro. 6-Aminohexanoate oligomer hydrolases from the alkalophilic bacteria *Agromyces* sp. strain KY5R and *Kocuria* sp. strain KY2. *Appl. Environ. Microbiol.*, 73:7099–7102, 2007.
- [3272] S. Yasumasu, I. Iuchi, and K. Yamagami. Medaka hatching enzyme consists of two kinds of proteases which act cooperatively. *Zool. Sci.*, 5:191–195, 1988.
- [3273] S. Yasumasu, I. Iuchi, and K. Yamagami. Isolation and some properties of low choriolytic enzyme (LCE), a component of the hatching enzyme of the teleost, *Oryzias latipes*. *J. Biochem. (Tokyo)*, 105:212–218, 1989.
- [3274] S. Yasumasu, I. Iuchi, and K. Yamagami. Purification and partial characterization of high choriolytic enzyme (HCE), a component of the hatching enzyme of the teleost, *Oryzias latipes*. *J. Biochem. (Tokyo)*, 105:204–211, 1989.
- [3275] S. Yasumasu, S. Katow, T.S. Hamazaki, I. Iuchi, and K. Yamagami. Two constituent proteases of a teleostean hatching enzyme: concurrent syntheses and packaging in the same secretory granules in discrete arrangement. *Dev. Biol.*, 149:349–356, 1992.
- [3276] S. Yasumasu, S. Katow, Y. Umino, I. Iuchi, and K. Yamagami. A unique proteolytic action of HCE, a constituent protease of a fish hatching enzyme: tight binding to its natural substrate, egg envelope. *Biochem. Biophys. Res. Commun.*, 162:58–63, 1989.
- [3277] S. Yasumasu, K. Yamada, K. Akasaka, K. Mitsunaga, I. Iuchi, H. Shimada, and K. Yamagami. Isolation of cDNAs for LCE and HCE, two constituent proteases of the hatching enzyme of *Oryzias latipes*, and concurrent expression of their mRNAs during development. *Dev. Biol.*, 153:250–258, 1992.
- [3278] M.G. Yates. A non-specific adenine nucleotide deaminase from *Desulfovibrio desulfuricans*. *Biochim. Biophys. Acta*, 171:299–310, 1969.
- [3279] E. Yavin and S. Gatt. Enzymatic hydrolysis of sphingolipids. 8. Further purification and properties of rat brain ceramidase. *Biochemistry*, 8:1692–1698, 1969.
- [3280] S. Yazawa, R. Madiyalakan, R.P. Chawda, and K.L. Matta.  $\alpha$ -L-Fucosidase from *Aspergillus niger*: demonstration of a novel  $\alpha$ -L-(1→6)-fucosidase acting on glycopeptides. *Biochem. Biophys. Res. Commun.*, 136:563–569, 1986.
- [3281] J. Ye, R.B. Rawson, R. Komuro, X. Chen, U.P. Dave, R. Prywes, M.S. Brown, and J.L. Goldstein. ER stress induces cleavage of membrane-bound ATF6 by the same proteases that process SREBPs. *Mol. Cell*, 6:1355–1364, 2000.
- [3282] S.J. Yeaman. Hormone-sensitive lipase - new roles for an old enzyme. *Biochem. J.*, 379:11–22, 2004.
- [3283] G. Yeung, J.J. Mulero, D.W. McGowan, S.S. Bajwa, and J.E. Ford. CD39L2, a gene encoding a human nucleoside diphosphatase, predominantly expressed in the heart. *Biochemistry*, 39:12916–12923, 2000.
- [3284] Q. Yin, H.H. Park, J.Y. Chung, S.C. Lin, Y.C. Lo, L.S. da Graca, X. Jiang, and H. Wu. Caspase-9 holoenzyme is a specific and optimal procaspase-3 processing machine. *Mol. Cell*, 22:259–268, 2006.
- [3285] L. Yman. Studies on human serum aminopeptidase. Some properties of oxytocinase, human serum aminopeptidase A and leucine aminopeptidase and their purification from retroplacental serum. *Acta Pharm. Suec.*, 7:75–86, 1970.
- [3286] K. Yokobayashi, A. Misaki, and T. Harada. Purification and properties of *Pseudomonas* isoamylase. *Biochim. Biophys. Acta*, 212:458–469, 1970.
- [3287] T. Yokota, T. Omori, and T. Kodama. Purification and properties of haloalkane dehalogenase from *Corynebacterium* sp. strain m15-3. *J. Bacteriol.*, 169:4049–4054, 1987.
- [3288] K. Yokoyama, Y. Yamamoto, F. Kudo, and T. Eguchi. Involvement of two distinct *N*-acetylglucosaminyltransferases and a dual-function deacetylase in neomycin biosynthesis. *ChemBioChem*, 9:865–869, 2008.
- [3289] S. Yonekura, U. Sanada, and Q.M. Zhang-Akiyama. CiMutT, an asidian MutT homologue, has a 7, 8-dihydro-8-oxo-dGTP pyrophosphohydrolase activity responsible for sanitization of oxidized nucleotides in *Ciona intestinalis*. *Genes Genet Syst*, 85:287–295, 2010.

- [3290] S. Yonezawa, K. Fujii, Y. Maejima, K. Tamoto, Y. Mori, and N. Muto. Further studies on rat cathepsin E: subcellular localization and existence of the active subunit form. *Arch. Biochem. Biophys.*, 267:176–183, 1988.
- [3291] S.J. Yoo, J.H. Seol, I.S. Seong, M.S. Kang, and C.H. Chung. ATP binding, but not its hydrolysis, is required for assembly and proteolytic activity of the HslVU protease in *Escherichia coli*. *Biochem. Biophys. Res. Commun.*, 238:581–585, 1997.
- [3292] S.J. Yoo, J.H. Seol, D.H. Shin, M. Rohrwild, M.S. Kang, K. Tanaka, A.L. Goldberg, and C.H. Chung. Purification and characterization of the heat shock proteins HslV and HslU that form a new ATP-dependent protease in *Escherichia coli*. *J. Biol. Chem.*, 271:14035–14040, 1996.
- [3293] T. Yorifuji, M. Kaneoke, E. Shimizu, K. Shiota, and R. Matsuo. Degradation of  $\alpha,\omega$ -diguanoalkanes and a novel enzyme, diguanidinobutane amidohydrolase, in *Pseudomonas putida*. *Agric. Biol. Chem.*, 53:3003–3009, 1989.
- [3294] T. Yorifuji, M. Kato, T. Kobayashi, S. Ozaki, and S. Ueno. 4-Guanidinobutyrate amidohydrolase from *Pseudomonas* sp ATCC 14676: purification to homogeneity and properties. *Agric. Biol. Chem.*, 44:1127–1134, 1980.
- [3295] T. Yorifuji, T. Kobayashi, A. Tabuchi, Y. Shiritani, and K. Yonaha. Distribution of amidohydrolases among *Pseudomonas* and comparative studies of some purified enzymes by one-dimensional peptide mapping. *Agric. Biol. Chem.*, 47:2825–2830, 1983.
- [3296] T. Yorifuji, I. Sugai, H. Matsumoto, and A. Tabuchi. Characterization of 3-guanidinopropionate amidohydrolase from *Pseudomonas aeruginosa* and a comparative study with 4-guanidinobutyrate amidohydrolase from another *Pseudomonas*. *Agric. Biol. Chem.*, 46:1361–1363, 1982.
- [3297] T. Yorifuji, H. Tamai, and H. Usami. Purification, crystallization and properties of  $Mn^{2+}$  dependent guanidoacetate amidohydrolase from a *Pseudomonas*. *Agric. Biol. Chem.*, 41:959–966, 1977.
- [3298] A. Yoshida and M. Lin.  $NH_2$ -terminal formylmethionine- and  $NH_2$ -terminal methionine-cleaving enzymes in rabbits. *J. Biol. Chem.*, 247:952–957, 1972.
- [3299] K. Yoshida, M. Yamaguchi, T. Morinaga, M. Kinehara, M. Ikeuchi, H. Ashida, and Y. Fujita. *myo*-Inositol catabolism in *Bacillus subtilis*. *J. Biol. Chem.*, 283:10415–10424, 2008.
- [3300] N. Yoshida, S. Tsuruyama, K. Nagata, K. Hirayama, K. Noda, and S. Makisumi. Purification and characterization of an acidic amino acid specific endopeptidase of *Streptomyces griseus* obtained from a commercial preparation (Pronase). *J. Biochem. (Tokyo)*, 104:451–456, 1988.
- [3301] S. Yoshida, K. Hiraga, T. Takehana, I. Taniguchi, H. Yamaji, Y. Maeda, K. Toyohara, K. Miyamoto, Y. Kimura, and K. Oda. A bacterium that degrades and assimilates poly(ethylene terephthalate). *Science*, 351:1196–1199, 2016.
- [3302] T. Yoshikawa, G. Turner, L.E. Esterling, A.R. Sanders, and S.D. Detera-Wadleigh. A novel human *myo*-inositol monophosphatase gene, IMP.18p, maps to a susceptibility region for bipolar disorder. *Mol. Psychiatry*, 2:393–397, 1997.
- [3303] H. Yoshima, S. Takasaki, S. Ito-Mega, and A. Kobata. Purification of almond emulsin  $\alpha$ -L-fucosidase I by affinity chromatography. *Arch. Biochem. Biophys.*, 194:394–398, 1979.
- [3304] T. Yoshimoto, I. Oka, and D. Tsuru. Purification, crystallization, and some properties of creatine amidohydrolase from *Pseudomonas putida*. *J. Biochem. (Tokyo)*, 79:1381–1383, 1976.
- [3305] K. Yoshimura, T. Ogawa, Y. Ueda, and S. Shigeoka. AtNUDX1, an 8-oxo-7,8-dihydro-2'-deoxyguanosine 5'-triphosphate pyrophosphohydrolase, is responsible for eliminating oxidized nucleotides in *Arabidopsis*. *Plant Cell Physiol.*, 48:1438–1449, 2007.
- [3306] I.G. Young and F. Gibson. Regulation of the enzymes involved in the biosynthesis of 2,3-dihydroxybenzoic acid in *Aerobacter aerogenes* and *Escherichia coli*. *Biochim. Biophys. Acta*, 177:401–411, 1969.
- [3307] G.M. Yousef, A. Scorilas, A. Magklara, A. Soosaipillai, and E.P. Diamandis. The *KLK7* (*PRSS6*) gene, encoding for the stratum corneum chymotryptic enzyme is a new member of the human kallikrein gene family - genomic characterization, mapping, tissue expression and hormonal regulation. *Gene*, 254:119–128, 2000.
- [3308] H. Yu, Q. Liu, C. Zhang, M. Lu, Y. Fu, W.-T. Im, S.-T. Lee, and F. Jin. A new ginsenosidase from *Aspergillus* strain hydrolyzing 20-O-multi-glycoside of PPD ginsenoside. *Process Biochem.*, 44:772–775, 2009.

- [3309] H. Yu, C. Zhang, M. Lu, F. Sun, Y. Fu, and F. Jin. Purification and characterization of new special ginsenosidase hydrolyzing multi-glycosides of protopanaxadiol ginsenosides, ginsenosidase type I. *Chem Pharm Bull (Tokyo)*, 55:231–235, 2007.
- [3310] S. Yu, V. Jensen, J. Seeliger, I. Feldmann, S. Weber, E. Schleicher, S. Haussler, and W. Blankenfeldt. Structure elucidation and preliminary assessment of hydrolase activity of PqsE, the *Pseudomonas* quinolone signal (PQS) response protein. *Biochemistry*, 48:10298–10307, 2009.
- [3311] X.C. Yu and W. Margolin. Ca<sup>2+</sup>-mediated GTP-dependent dynamic assembly of bacterial cell division protein FtsZ into asters and polymer networks in vitro. *EMBO J.*, 16:5455–5463, 1997.
- [3312] Z. Yu, F. Xu, L.M. Huse, C. Morisseau, A.J. Draper, J.W. Newman, C. Parker, L. Graham, M.M. Engler, B.D. Hammock, D.C. Zeldin, and D.L. Kroetz. Soluble epoxide hydrolase regulates hydrolysis of vasoactive epoxyeicosatrienoic acids. *Circ. Res.*, 87:992–998, 2000.
- [3313] E.C. Yurewicz, M.A. Ghalambor, D.H. Duckworth, and E.C. Heath. Catalytic and molecular properties of a phage-induced capsular polysaccharide depolymerase. *J. Biol. Chem.*, 246:5607–5716, 1971.
- [3314] E.C. Yurewicz, M.A. Ghalambor, and E.C. Heath. The structure of *Aerobacter aerogenes* capsular polysaccharide. *J. Biol. Chem.*, 246:5596–5606, 1971.
- [3315] T. Yutaka, A.L. Fluharty, R.L. Stevens, and H. Kihara. Purification and some properties of human liver iduronate sulfatase. *J. Biochem. (Tokyo)*, 91:433–441, 1982.
- [3316] T. Yutaka, S. Okada, T. Kato, K. Inui, and H. Yabuchi. Galactose 6-sulfate sulfatase activity in Morquio syndrome. *Clin. Chim. Acta*, 122:169–180, 1982.
- [3317] G.T. Zancan, E.F. Recondo, and L.F. Leloir. Enzymic dephosphorylation of adenosine diphosphate phosphoglyceric acid. *Biochim. Biophys. Acta*, 92:125–131, 1964.
- [3318] C.L. Zayas and J.C. Escalante-Semerena. Reassessment of the late steps of coenzyme B<sub>12</sub> synthesis in *Salmonella enterica*: evidence that dephosphorylation of adenosylcobalamin-5'-phosphate by the CobC phosphatase is the last step of the pathway. *J. Bacteriol.*, 189:2210–2218, 2007.
- [3319] L. Zechmeister and G. Tóth. Chromatographic adsorption of the enzymes of emulsin which act on chitins. *Enzymologia*, 7:165–169, 1939.
- [3320] D.C. Zeldin, S. Wei, J.R. Falck, B.D. Hammock, J.R. Snapper, and J.H. Capdevila. Metabolism of epoxyeicosatrienoic acids by cytosolic epoxide hydrolase: substrate structural determinants of asymmetric catalysis. *Arch. Biochem. Biophys.*, 316:443–451, 1995.
- [3321] C. Zevaco, J. Hermier, and J.-C. Gripon. Le système protéolytique de *Penicillium roqueforti*. II - Purification et propriétés de la protéase acide. *Biochimie*, 55:1353–1360, 1973.
- [3322] C. Zevaco, V. Monnet, and J.-C. Gripon. Intracellular X-prolyl dipeptidyl peptidase from *Lactococcus lactis* spp. lactis: purification and properties. *J. Appl. Bacteriol.*, 68:357–366, 1990.
- [3323] M. Zha, C. Zhong, Y. Peng, H. Hu, and J. Ding. Crystal structures of human NUDT5 reveal insights into the structural basis of the substrate specificity. *J. Mol. Biol.*, 364:1021–1033, 2006.
- [3324] C. Zhang, Y. Zhang, J. Chen, and X. Liang. Purification and characterization of baicalin-β-D-glucuronidase hydrolyzing baicalin to baicalein from fresh roots of *Scutellaria viscidula* Bge. *Proc. Biochem.*, 40:1911–1915, 2005.
- [3325] J. Zhang and M. Inouye. MazG, a nucleoside triphosphate pyrophosphohydrolase, interacts with Era, an essential GTPase in *Escherichia coli*. *J. Bacteriol.*, 184:5323–5329, 2002.
- [3326] J. Zhang, T. Kasciukovic, and M.F. White. The CRISPR associated protein Cas4 Is a 5' to 3' DNA exonuclease with an iron-sulfur cluster. *PLoS One*, 7:e47232–e47232, 2012.
- [3327] J.R. Zhang and M.P. Deutscher. Cloning, characterization, and effects of overexpression of the *Escherichia coli* rnd gene encoding RNase D. *J. Bacteriol.*, 170:522–527, 1988.

- [3328] K. Zhang and A.W. Nicholson. Regulation of ribonuclease III processing by double-helical sequence antideterminants. *Proc. Natl. Acad. Sci. USA*, 94:13437–13441, 1997.
- [3329] L. Zhang, X. Xu, Z. Luo, D. Shen, and H. Wu. Identification of an unusual AT(D)Pase-like activity in multifunctional NAD glycohydrolase from the venom of *Agkistrodon acutus*. *Biochimie*, 91:240–251, 2009.
- [3330] R. Zhang, T. Minh, L. Lezondra, S. Korolev, S.F. Moy, F. Collart, and A. Joachimiak. 1.6 Å crystal structure of YteR protein from *Bacillus subtilis*, a predicted lyase. *Proteins*, 60:561–565, 2005.
- [3331] B.G. Zhao, C.A. Janson, B.Y. Amegadzie, K. D'Alessio, C. Griffin, C.R. Hanning, C. Jones, J. Kurdyla, M. McQueney, X.Y. Qiu, W.W. Smith, and S.S. Abdel-Meguid. Crystal structure of human osteoclast cathepsin K complex with E-64. *Nature Struct. Biol.*, 4:109–111, 1997.
- [3332] X.-J. Zhao and H.C. Wu. Nucleotide sequence of the *Staphylococcus aureus* signal peptidase II (lsp) gene. *FEBS Lett.*, 299:80–84, 1992.
- [3333] W. Zheng, S.A. Johnston, and L. Joshua-Tor. The unusual active site of Gal<sub>6</sub>/bleomycin hydrolase can act as a carboxypeptidase, aminopeptidase, and peptide ligase. *Cell*, 93:103–109, 1998.
- [3334] B. Zhivotovsky and S. Orrenius. Caspase-2 function in response to DNA damage. *Biochem. Biophys. Res. Commun.*, 331:859–867, 2005.
- [3335] J. Zhou, W.L. Kelly, B.O. Bachmann, M. Gunsior, C.A. Townsend, and E.I. Solomon. Spectroscopic studies of substrate interactions with clavaminase synthase 2, a multifunctional  $\alpha$ -KG-dependent non-heme iron enzyme: Correlation with mechanisms and reactivities. *J. Am. Chem. Soc.*, 123:7388–7398, 2001.
- [3336] J.Q. Zhou, H. Qi, V.P. Schulz, M.K. Mateyak, E.K. Monson, and V.A. Zakian. *Schizosaccharomyces pombe* pfh1+ encodes an essential 5' to 3' DNA helicase that is a member of the PIF1 subfamily of DNA helicases. *Mol. Biol. Cell*, 13:2180–2191, 2002.
- [3337] N.Y. Zhou, S.L. Fuenmayor, and P.A. Williams. nag genes of *Ralstonia* (formerly *Pseudomonas*) sp. strain U2 encoding enzymes for gentisate catabolism. *J. Bacteriol.*, 183:700–708, 2001.
- [3338] T. Zhou, B.P. Rosen, and D.L. Gatti. Crystallization and preliminary X-ray analysis of the catalytic subunit of the ATP-dependent arsenite pump encoded by the *Escherichia coli* plasmid R773. *Acta Crystallogr. D Biol. Crystallogr.*, 55:921–924, 1999.
- [3339] A. Zhu, C. Monahan, Z.K. Wang, and J. Goldstein. Expression, purification, and characterization of recombinant  $\alpha$ -N-acetylgalactosaminidase produced in the yeast *Pichia pastoris*. *Protein Expr. Purif.*, 8:456–462, 1996.
- [3340] A. Zhukov, U. Hellman, and M. Ingelman-Sundberg. Purification and characterization of hepsin from rat liver microsomes. *Biochim. Biophys. Acta*, 1337:85–95, 1997.
- [3341] M. Ziak, M. Meier, K.S. Etter, and J. Roth. Two isoforms of trimming glucosidase II exist in mammalian tissues and cell lines but not in yeast and insect cells. *Biochem. Biophys. Res. Commun.*, 280:363–367, 2001.
- [3342] M.D. Zimmerman, M. Proudfoot, A. Yakunin, and W. Minor. Structural insight into the mechanism of substrate specificity and catalytic activity of an HD-domain phosphohydrolase: the 5'-deoxyribonucleotidase YfbR from *Escherichia coli*. *J. Mol. Biol.*, 378:215–226, 2008.
- [3343] S.B. Zimmerman and A. Kornberg. Deoxycytidine di- and triphosphate cleavage by an enzyme formed in bacteriophage-infected *Escherichia coli*. *J. Biol. Chem.*, 236:1480–1486, 1961.
- [3344] C.A. Zittle, E.S. DellaMonica, J.H. Custer, and R. Krikorian. Purification of human red cell acetylcholinesterase by electrophoresis, ultracentrifugation and gradient extraction. *Arch. Biochem. Biophys.*, 56:469–475, 1955.
- [3345] A. Zlotnick, V.S. Reddy, R. Dasgupta, A. Schneemann, W.J. Ray, Rueckert Jr., Johnson R.R., and J.E. Capsid assembly in a family of animal viruses primes an autoproteolytic maturation that depends on a single aspartic acid residue. *J. Biol. Chem.*, 269:13680–13684, 1994.
- [3346] Y. Zong, T.W. Bice, H. Ton-That, O. Schneewind, and S.V. Narayana. Crystal structures of *Staphylococcus aureus* sortase A and its substrate complex. *J. Biol. Chem.*, 279:31383–31389, 2004.

- [3347] Y. Zong, S.K. Mazmanian, O. Schneewind, and S.V. Narayana. The structure of sortase B, a cysteine transpeptidase that tethers surface protein to the *Staphylococcus aureus* cell wall. *Structure*, 12:105–112, 2004.
- [3348] B.J.M. Zonneveld. A new type of enzyme, and exo-splitting  $\alpha$ -1,3 glucanase from non-induced cultures of *Aspergillus nidulans*. *Biochim. Biophys. Acta*, 258:541–547, 1972.
- [3349] J. Zou, J. Marjanovic, M.V. Kisseleva, M. Wilson, and P.W. Majerus. Type I phosphatidylinositol-4,5-bisphosphate 4-phosphatase regulates stress-induced apoptosis. *Proc. Natl. Acad. Sci. USA*, 104:16834–16839, 2007.
- [3350] S. Zucker, D.J. Buttle, M.J.H. Nicklin, and A.J. Barrett. Proteolytic activities of papain, chymopapain and papaya proteinase III. *Biochim. Biophys. Acta*, 828:196–204, 1985.
- [3351] J.J. Zulty and M.K. Speedie. Purification and characterization of *S*-adenosylhomocysteine deaminase from streptonigrin-producing *Streptomyces flocculus*. *J. Bacteriol.*, 171:6840–6844, 1989.
- [3352] V.V. Zverlov, K.P. Fuchs, and W.H. Schwarz. Chi18A, the endochitinase in the cellulosome of the thermophilic, cellulolytic bacterium *Clostridium thermocellum*. *Appl. Environ. Microbiol.*, 68:3176–3179, 2002.
- [3353] V.V. Zverlov, C. Hertel, K. Bronnenmeier, A. Hroch, J. Kellermann, and W.H. Schwarz. The thermostable  $\alpha$ -L-rhamnosidase RamA of *Clostridium stercorarium*: biochemical characterization and primary structure of a bacterial  $\alpha$ -L-rhamnoside hydrolase, a new type of inverting glycoside hydrolase. *Mol. Microbiol.*, 35:173–179, 2000.

## Index

- $\alpha$ -*N*-acetylgalactosaminidase, 100  
 $\alpha$ -*N*-acetylglucosaminidase, 101  
 $\alpha$ -D-ribose 1-methylphosphonate 5-triphosphate diphosphatase, 314  
 $\alpha$ -D-xyloside xylohydrolase, 129  
 $\alpha$ -L-fucosidase, 101  
 $\alpha$ -L-rhamnosidase, 98  
 $\beta$ -1,2-mannosidase, 135  
 $\beta$ -Ala-His dipeptidase, 162  
 $\beta$ -*N*-acetylgalactosaminidase, 101  
 $\beta$ -*N*-acetylhexosaminidase, 101  
 $\beta$ -D-fucosidase, 98  
 $\beta$ -D-glucopyranosyl abscisate  $\beta$ -glucosidase, 128  
 $\beta$ -L-rhamnosidase, 99  
 $\gamma$ -D-glutamyl-*meso*-diaminopimelate peptidase, 174  
 $\gamma$ -D-glutamyl-L-lysine dipeptidyl-peptidase, 165  
(*R*)-2-haloacid dehalogenase, 330  
(*S*)-2-haloacid dehalogenase, 328  
(*S*)-*N*-acetyl-1-phenylethylamine hydrolase, 268  
(+)-kolavelool synthase, 70  
*Aspergillus* deoxyribonuclease K<sub>1</sub>, 78  
*Aspergillus* nuclease S<sub>1</sub>, 90  
*N*<sup>4</sup>-( $\beta$ -*N*-acetylglucosaminyl)-L-asparaginase, 256  
*Physarum polycephalum* ribonuclease, 86  
*Serratia marcescens* nuclease, 90  
*all-trans*-retinyl ester 13-*cis* isomerohydrolase, 21  
*endo*-1,3- $\beta$ -xylanase, 97  
*trans*-2,3-dihydro-3-hydroxyanthranilic acid synthase, 149  
D-Ala-D-Ala dipeptidase, 162  
D-*glycero*- $\alpha$ -D-*manno*-heptose 1,7-bisphosphate 7-phosphatase, 49  
D-*glycero*- $\beta$ -D-*manno*-heptose 1,7-bisphosphate 7-phosphatase, 49  
1,2- $\alpha$ -L-fucosidase, 103  
1,3- $\alpha$ -L-fucosidase, 113  
1,3- $\alpha$ -isomaltosidase, 136  
1,6- $\alpha$ -D-mannosidase, 125  
1,6- $\alpha$ -L-fucosidase, 117  
11-*cis*-retinyl-palmitate hydrolase, 15  
2'-*N*-acetylparomamine deacetylase, 275  
2,6- $\beta$ -fructan 6-levanbiohydrolase, 104  
20-*O*-multi-glycoside ginsenosidase, 134  
3D-(3,5/4)-trihydroxycyclohexane-1,2-dione acylhydrolase (ring-opening), 328  
4- $\alpha$ -D-(1 $\rightarrow$ 4)- $\alpha$ -D-glucanotrehalose trehalohydrolase, 120  
5' to 3' exodeoxyribonuclease (nucleoside 3'-phosphate-forming), 73  
5'-(*N*<sup>7</sup>-methyl 5'-triphosphoguanosine)-[mRNA] diphosphatase, 313  
5'-(*N*<sup>7</sup>-methylguanosine 5'-triphospho)-[mRNA] hydrolase, 314  
5-(3,4-diacetoxybut-1-ynyl)-2,2'-bithiophene deacetylase, 15  
4-acetamidobutyrate deacetylase, 263  
4-acetamidobutyryl-CoA deacetylase, 261  
2-(acetamidomethylene)succinate hydrolase, 257  
acetoacetyl-CoA hydrolase, 26  
acetoxybutynylbithiophene deacetylase, 13  
*N*-acetyl- $\beta$ -alanine deacetylase, 255  
*N*-acetyl-1-D-*myo*-inositol-2-amino-2-deoxy- $\alpha$ -D-glucopyranoside deacetylase, 272  
2'''-acetyl-6'''-hydroxyneomycin C deacetylase, 275  
*N*-acetyl-D-muramate 6-phosphate phosphatase, 54  
*N*<sup>2</sup>-acetyl-L-2,4-diaminobutanoate deacetylase, 278  
[acetyl-CoA carboxylase]-phosphatase, 40  
acetyl-CoA hydrolase, 25  
acetylajmaline esterase, 19  
acetylalkylglycerol acetylhydrolase, 16  
acetylcholinesterase, 4  
*N*-acetyldiaminopimelate deacetylase, 260  
acetylestase, 4  
*N*-acetylgalactosamine-4-sulfatase, 66  
*N*-acetylgalactosamine-6-sulfatase, 64  
*N*-acetylgalactosaminoglycan deacetylase, 14  
*N*-acetylglucosamine deacetylase, 257  
*N*-acetylglucosamine-1-phosphodiester  $\alpha$ -*N*-acetylglucosaminidase, 60  
*N*-acetylglucosamine-6-phosphate deacetylase, 256  
*N*-acetylglucosamine-6-sulfatase, 66  
*N*-acetylglucosaminylphosphatidylinositol deacetylase, 269  
6-acetylglucose deacetylase, 9  
*N*-acetylmuramoyl-L-alanine amidase, 256  
acetylmithine deacetylase, 254  
*N*-acetylphosphatidylethanolamine-hydrolysing phospholipase D, 62  
acetylputrescine deacetylase, 263  
acetylpyruvate hydrolase, 324  
acetylsalicylate deacetylase, 13  
acetylspemidine deacetylase, 260  
acetylxyylan esterase, 16  
acid phosphatase, 31  
acireductone synthase, 47  
aclacinomycin methylesterase, 23  
acrocylindropepsin, 223  
acrosin, 178  
actinidain, 204  
actinomycin lactonase, 10  
aculeacin-A deacylase, 265  
*N*-acyl-D-amino-acid deacylase, 267  
*N*-acyl-D-aspartate deacylase, 267  
*N*-acyl-D-glutamate deacylase, 267  
*N*-acyl-aliphatic-L-amino acid amidohydrolase, 254  
*N*-acyl-aromatic-L-amino acid amidohydrolase, 275  
[acyl-carrier-protein] phosphodiesterase, 56  
acyl-CoA hydrolase, 28  
acyl-homoserine-lactone acylase, 271  
acyl-lysine deacylase, 254

acylagmatine amidase, 258  
 acylaminoacyl-peptidase, 172  
 acylcarnitine hydrolase, 8  
 acylglycerol lipase, 7  
*N*-acylneuramate-9-phosphatase, 37  
 acyloxyacyl hydrolase, 18  
 acylphosphatase, 304  
 5'-acylphosphoadenosine hydrolase, 306  
 acylpyruvate hydrolase, 324  
 ADAM 17 endopeptidase, 247  
 ADAM10 endopeptidase, 246  
 adamalysin, 238  
 ADAMTS-4 endopeptidase, 246  
 ADAMTS13 endopeptidase, 247  
 adenain, 208  
 adenine deaminase, 289  
 adenine glycosylase, 144  
 adenosine deaminase, 289  
 adenosine nucleosidase, 139  
 adenosine-5'-diphospho-5'-[DNA] diphosphatase, 72  
 adenosine-phosphate deaminase, 292  
 adenosine-tetraphosphatase, 305  
 adenosinetriphosphatase, 303  
*S*-adenosyl-L-methionine hydrolase (*L*-homoserine-forming), 145  
*S*-adenosyl-L-methionine hydrolase (adenosine-forming), 335  
 adenosylcobalamin/ $\alpha$ -ribazole phosphatase, 46  
 adenosylcobinamide hydrolase, 269  
 adenosylhomocysteinase, 145  
*S*-adenosylhomocysteine deaminase, 294  
 adenosylhomocysteine nucleosidase, 139  
 adenylylsulfatase, 315  
 ADP deaminase, 290  
 ADP-dependent medium-chain-acyl-CoA hydrolase, 28  
 ADP-dependent short-chain-acyl-CoA hydrolase, 28  
 ADP-phosphoglycerate phosphatase, 36  
 ADP-ribose 1''-phosphate phosphatase, 49  
 ADP-ribose diphosphatase, 305  
 ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase, 139  
 ADP-ribosyl-[dinitrogen reductase] hydrolase, 143  
 ADP-sugar diphosphatase, 306  
 $\alpha$ -agarase, 124  
 $\beta$ -agarase, 107  
 agmatinase, 285  
 agmatine deiminase, 285  
 alanine carboxypeptidase, 168  
 aliphatic nitrilase, 299  
 alkaline phosphatase, 31  
 1-alkyl-2-acetylgllycerophosphocholine esterase, 11  
 alkylacetylgllycerophosphatase, 43  
 alkylamidase, 258  
 alkylgllycerophosphoethanolamine phosphodiesterase, 59  
 alkylhalidase, 328  
 allantoate deiminase, 285  
 allantoicase, 284  
 allantoinase, 280  
 allophanate hydrolase, 261  
 alternative-complement-pathway C3/C5 convertase, 184  
 $\omega$ -amidase, 252  
 (*R*)-amidase, 272  
 amidase, 252  
 amidinoaspartase, 286  
 2-amino-5-formylamino-6-ribosylaminopyrimidin-4(3*H*)-one 5'-monophosphate deformylase, 272  
 5-amino-6-(5-phospho-D-ribitylamino)uracil phosphatase, 54  
 $\alpha$ -amino-acid esterase, 11  
 D-aminoacyl-tRNA deacylase, 23  
 aminoacyl-tRNA hydrolase, 8  
 2-aminobenzoylacetyl-CoA thioesterase, 31  
 1-aminocyclopropane-1-carboxylate deaminase, 301  
 aminodeoxyfucalose deaminase, 297  
 aminodeoxyfucalose nucleosidase, 144  
 6-aminohexanoate-cyclic-dimer hydrolase, 281  
 6-aminohexanoate-oligomer endohydrolase, 276  
 6-aminohexanoate-oligomer exohydrolase, 259  
 aminoimidazolase, 290  
 2-aminomuconate deaminase, 300  
 2-aminomuconate deaminase (2-hydroxyomuconate-forming), 302  
 5-aminopentanamidase, 257  
 aminopeptidase B, 154  
 aminopeptidase Ey, 157  
 aminopeptidase I, 157  
 aminopeptidase S, 158  
 aminopeptidase Y, 156  
*N*<sup>1</sup>-aminopropylagmatine ureohydrolase, 288  
 aminopyrimidine aminohydrolase, 300  
 AMP deaminase, 289  
 AMP nucleosidase, 138  
 amygdalin  $\beta$ -glucosidase, 115  
 $\alpha$ -amylase, 91  
 $\beta$ -amylase, 91  
 amylo- $\alpha$ -1,6-glucosidase, 97  
 ananain, 206  
 angiotensin-converting enzyme 2, 172  
 anthrax lethal factor endopeptidase, 246  
 $\beta$ -apiosyl- $\beta$ -glucosidase, 125  
 apyrase, 303  
 aqualysin 1, 199  
 (Ara-f)<sub>3</sub>-Hyp  $\beta$ -L-arabinobiosidase, 132  
 arabinan endo-1,5- $\alpha$ -L-arabinanase, 111  
 arabinogalactan endo- $\beta$ -1,4-galactanase, 109  
 D-arabinonolactonase, 8  
 L-arabinonolactonase, 6  
 D-arginase, 285  
 arginase, 283  
 arginine deiminase, 284  
 arsenite-transporting ATPase, 317  
 aryl-acylamidase, 253  
 arylacetone nitrilase, 299  
 arylalkyl acylamidase, 266  
 aryldialkylphosphatase, 70  
 arylesterase, 3  
 arylformamidase, 253

arylsulfatase, 64  
 asclepain, 203  
 asparaginase, 251  
 aspartoacylase, 254  
 aspartyl aminopeptidase, 157  
 $\beta$ -aspartyl-*N*-acetylglucosaminidase, 140  
 $\beta$ -aspartyl-peptidase, 173  
 aspergillopepsin I, 220  
 aspergillopepsin II, 220  
 assemblin, 195  
 astacin, 232  
 ATP deaminase, 292  
 ATP diphosphatase, 304  
 atrazine chlorohydrolase, 330  
 atrolysin A, 228  
 atrolysin B, 237  
 atrolysin C, 237  
 atrolysin E, 237  
 atrolysin F, 238  
 atroxase, 237  
 aureolysin, 234  
 avenacosidase, 132  
  
 bacillolysin, 234  
 bacterial leucyl aminopeptidase, 155  
 baicalin- $\beta$ -D-glucuronidase, 126  
 barbiturase, 279  
 barrierpepsin, 224  
 D-benzoylarginine-4-nitroanilide amidase, 265  
*N*-benzyloxycarbonylglycine hydrolase, 262  
*N* <sup>$\alpha$</sup> -benzyloxycarbonylleucine hydrolase, 263  
 biotinidase, 253  
 bis(2-ethylhexyl)phthalate esterase, 14  
 bis(5'-adenosyl)-triphosphatase, 308  
 bis(5'-nucleosyl)-tetraphosphatase (asymmetrical), 306  
 bis(5'-nucleosyl)-tetraphosphatase (symmetrical), 309  
 3'(2'),5'-bisphosphate nucleotidase, 32  
 2,3-bisphosphoglycerate 3-phosphatase, 48  
 biuret amidohydrolase, 267  
 blasticidin-S deaminase, 293  
 bleomycin hydrolase, 208  
 blood-group-substance endo-1,4- $\beta$ -galactosidase, 112  
 bontoxilysin, 243  
 bothrolysin, 239  
 bothropasin, 239  
 brachyurin, 181  
 branched-dextran exo-1,2- $\alpha$ -glucosidase, 114  
 bromoxynil nitrilase, 299  
  
 C-terminal processing peptidase, 196  
 C5a peptidase, 199  
 caldesmon-phosphatase, 42  
 calicivirin, 216  
 calpain-1, 211  
 calpain-2, 212  
 calpain-3, 212  
  
 cAMP deaminase, 298  
 cancer procoagulant, 205  
 candidapepsin, 222  
 capsular-polysaccharide endo-1,3- $\alpha$ -galactosidase, 108  
*N*-carbamoyl-D-amino-acid hydrolase, 266  
*N*-carbamoyl-L-amino-acid hydrolase, 268  
*N*-carbamoylputrescine amidase, 261  
*N*-carbamoylsarcosine amidase, 262  
 carbon disulfide hydrolase, 334  
 carbonyl sulfide hydrolase, 335  
 2-carboxy-D-arabinitol-1-phosphatase, 44  
 carboxylesterase, 3  
 carboxymethylcellulase, 136  
 carboxymethylenebutenolidase, 11  
 carboxymethylhydantoinase, 280  
 carboxypeptidase A, 167  
 carboxypeptidase A<sub>2</sub>, 170  
 carboxypeptidase B, 167  
 carboxypeptidase C, 167  
 carboxypeptidase D, 167  
 carboxypeptidase E, 169  
 carboxypeptidase M, 169  
 carboxypeptidase T, 171  
 carboxypeptidase Taq, 171  
 carboxypeptidase U, 171  
 caricain, 206  
 carnitinamidase, 265  
 $\iota$ -carrageenase, 124  
 $\kappa$ -carrageenase, 107  
 $\lambda$ -carrageenase, 125  
 caspase-1, 207  
 caspase-10, 215  
 caspase-11, 215  
 caspase-2, 212  
 caspase-3, 212  
 caspase-4, 213  
 caspase-5, 213  
 caspase-6, 213  
 caspase-7, 214  
 caspase-8, 214  
 caspase-9, 215  
 cathepsin B, 202  
 cathepsin D, 219  
 cathepsin E, 224  
 cathepsin F, 209  
 cathepsin G, 179  
 cathepsin H, 204  
 cathepsin K, 208  
 cathepsin L, 204  
 cathepsin O, 209  
 cathepsin S, 205  
 cathepsin T, 205  
 cathepsin V, 209  
 cathepsin X, 172  
 CC-preferring endodeoxyribonuclease, 77  
 Cd<sup>2+</sup>-exporting ATPase, 316



CDP-diacylglycerol diphosphatase, 308  
 CDP-glycerol diphosphatase, 306  
 cellulase, 92  
 cellulose 1,4- $\beta$ -cellobiosidase (non-reducing end), 109  
 cellulose 1,4- $\beta$ -cellobiosidase (reducing end), 128  
 cellulose-polysulfatase, 65  
 cephalosporin-C deacetylase, 10  
 ceramidase, 255  
 cerebroside-sulfatase, 65  
 cerevisin, 184  
 cetraxate benzylesterase, 16  
 chenodeoxycholytaurine hydrolase, 265  
 chitin deacetylase, 259  
 chitin disaccharide deacetylase, 273  
 chitinase, 93  
 chitosanase, 118  
 4-chlorobenzoate dehalogenase, 329  
 4-chlorobenzoyl-CoA dehalogenase, 329  
 chlorogenate hydrolase, 11  
 chlorophyllase, 6  
 chlorophyllide *a* hydrolase, 24  
 cholesterol-5,6-oxide hydrolase, 148  
 choline-sulfatase, 65  
 cholinesterase, 5  
 choloyl-CoA hydrolase, 29  
 choloylglycine hydrolase, 256  
 chondro-4-sulfatase, 65  
 chondro-6-sulfatase, 65  
 choriolysin H, 243  
 choriolysin L, 242  
 chorismatase, 149  
 chymase, 182  
 chymopapain, 203  
 chymosin, 218  
 chymotrypsin, 176  
 chymotrypsin C, 176  
 citrate-lyase deacetylase, 27  
 citrullinase, 255  
 Cl<sup>-</sup>-transporting ATPase, 317  
 classical-complement-pathway C3/C5 convertase, 183  
 clostridial aminopeptidase, 155  
 clostripain, 203  
 CMP-*N*-acylneuraminate phosphodiesterase, 59  
 coagulation factor IXa, 179  
 coagulation factor VIIa, 179  
 coagulation factor Xa, 177  
 coagulation factor XIa, 180  
 coagulation factor XIIa, 182  
 cobalt-precorrin 5A hydrolase, 326  
 cocaine esterase, 20  
 coccolysin, 234  
 complement factor D, 183  
 complement factor I, 183  
 complement subcomponent C<sup>1s</sup>, 183  
 complement subcomponent C<sup>1r</sup>, 182  
 coniferin  $\beta$ -glucosidase, 117  
 creatinase, 284  
 creatininase, 281  
 creatinine deaminase, 292  
 crossover junction endodeoxyribonuclease, 78  
 cruzipain, 211  
 cucumisin, 180  
 cutinase, 17  
 cyanoalanine nitrilase, 299  
 cyanophycinase, 166  
 cyanuric acid amidohydrolase, 282  
 cyclamate sulfohydrolase, 332  
 3',5'-cyclic-AMP phosphodiesterase, 62  
 3',5'-cyclic-GMP phosphodiesterase, 58  
 cyclic-guanylate-specific phosphodiesterase, 62  
 2',3'-cyclic-nucleotide 2'-phosphodiesterase, 57  
 2',3'-cyclic-nucleotide 3'-phosphodiesterase, 59  
 3',5'-cyclic-nucleotide phosphodiesterase, 57  
 cyclohexane-1,2-dione hydrolase, 325  
 cyclohexane-1,3-dione hydrolase, 325  
 cyclomaltodextrinase, 101  
 [CysO sulfur-carrier protein]-*S*-L-cysteine hydrolase, 334  
 cystinyl aminopeptidase, 154  
*N*<sup>5</sup>-(cytidine 5'-diphosphoramidyl)-L-glutamine hydrolase, 279  
 cytidine deaminase, 289  
 cytosine deaminase, 288  
 cytosol alanyl aminopeptidase, 156  
 cytosol nonspecific dipeptidase, 161  
  
 dactylysin, 241  
 dCMP deaminase, 291  
 dCTP deaminase, 291  
 dCTP deaminase (dUMP-forming), 294  
 (d)CTP diphosphatase, 314  
 dCTP diphosphatase, 305  
 deaminated glutathione amidase, 279  
 3-deoxy-2-octulosonidase, 116  
 6-deoxy-6-sulfogluconolactonase, 23  
 3-deoxy-*manno*-octulosonate-8-phosphatase, 40  
 3-deoxy-*D-glycero-D-galacto*-nonulopyranosonate 9-phosphatase, 54  
 5'-deoxyadenosine deaminase, 297  
 2-deoxyglucose-6-phosphatase, 45  
 2-deoxyglucosidase, 114  
 deoxylimonate A-ring-lactonase, 11  
 5'-deoxynucleotidase, 51  
 deoxynucleotide 3'-phosphatase, 38  
 3-deoxyoctulosonase, 120  
 deoxyribodipyrimidine endonucleosidase, 141  
 deoxyribonuclease I, 75  
 deoxyribonuclease II, 78  
 deoxyribonuclease IV, 75  
 deoxyribonuclease (pyrimidine dimer), 86  
 deoxyribonuclease V, 77  
 deoxyribonuclease X, 78  
 desampylase, 175  
 deuterolysin, 236

dextran 1,6- $\alpha$ -isomaltotriosidase, 110  
dextranase, 93  
dGTPase, 64  
diacylglycerol diphosphate phosphatase, 48  
diadenosine hexaphosphate hydrolase (AMP-forming), 313  
diadenosine hexaphosphate hydrolase (ATP-forming), 313  
diaminohydroxyphosphoribosylaminopyrimidine deaminase, 293  
difructose-anhydride synthase, 118  
diguanidinobutanase, 287  
dihydrocoumarin hydrolase, 9  
dihydromonacolin L-[lovastatin nonaketide synthase] thioesterase, 30  
7,8-dihydroneopterin 2',3'-cyclic phosphate phosphodiesterase, 63  
dihydroneopterin triphosphate diphosphatase, 315  
dihydroorotase, 280  
dihydropyrimidinase, 279  
1,4-dihydroxy-2-naphthoyl-CoA hydrolase, 30  
2,6-dihydroxypseudoxynicotine hydrolase, 327  
diisopropyl-fluorophosphatase, 71  
 $\beta$ -diketone hydrolase, 324  
dimethylargininase, 286  
*N,N*-dimethylformamidase, 262  
2,4-dinitroanisole *O*-demethylase, 149  
dioscin glycosidase (3-*O*- $\beta$ -D-Glc-diosgenin-forming), 132  
dioscin glycosidase (diosgenin-forming), 132  
2,6-dioxo-6-phenylhexa-3-enoate hydrolase, 325  
2,5-dioxopiperazine hydrolase, 281  
dipeptidase E, 162  
dipeptidyl-dipeptidase, 163  
dipeptidyl-peptidase I, 162  
dipeptidyl-peptidase II, 163  
dipeptidyl-peptidase III, 163  
dipeptidyl-peptidase IV, 163  
diphosphoinositol-polyphosphate diphosphatase, 311  
4,5:9,10-diseco-3-hydroxy-5,9,17-trioxoandrosta-1(10),2-diene-4-oate hydrolase, 326  
disulfoglucosamine-6-sulfatase, 66  
DNA helicase, 321  
DNA-3'-diphospho-5'-guanosine diphosphatase, 73  
DNA-3-methyladenine glycosylase I, 142  
DNA-3-methyladenine glycosylase II, 142  
DNA-deoxyinosine glycosylase, 141  
DNA-formamidopyrimidine glycosylase, 142  
dodecanoyl-[acyl-carrier-protein] hydrolase, 28  
dolichyl-phosphatase, 41  
dolichyldiphosphatase, 310  
dolichylphosphate-glucose phosphodiesterase, 61  
dolichylphosphate-mannose phosphodiesterase, 61  
double-stranded RNA adenine deaminase, 296  
double-stranded uracil-DNA glycosylase, 144  
dUTP diphosphatase, 307  
dynamamin GTPase, 323  
ectoine hydrolase, 298  
enamidase, 282  
endo- $\alpha$ -*N*-acetylgalactosaminidase, 110  
endo- $\alpha$ -sialidase, 117  
endo-1,3(4)- $\beta$ -glucanase, 92  
endo-1,4- $\beta$ -xylanase, 92  
endo-chitodextrinase, 136  
endogalactosaminidase, 113  
endoglycosylceramidase, 116  
endopeptidase Clp, 194  
endopeptidase La, 185  
endopeptidase So, 188  
endopolyphosphatase, 304  
endothelin-converting enzyme 1, 244  
endothiapsin, 221  
enteropeptidase, 178  
envelysin, 230  
*trans*-epoxysuccinate hydrolase, 146  
equine arterivirus serine peptidase, 200  
exo- $\alpha$ -sialidase, 94  
exo-1,4- $\beta$ -D-glucosaminidase, 126  
exo-chitinase (non-reducing end), 135  
exo-chitinase (reducing end), 136  
exo-poly- $\alpha$ -galacturonosidase, 107  
exodeoxyribonuclease I, 71  
exodeoxyribonuclease III, 71  
exodeoxyribonuclease ( $\lambda$ -induced), 71  
exodeoxyribonuclease (phage SP<sub>3</sub>-induced), 72  
exodeoxyribonuclease V, 72  
exodeoxyribonuclease VII, 72  
exopolyphosphatase, 304  
exoribonuclease H, 74  
exoribonuclease II, 73  
FAD diphosphatase, 306  
farnesyl diphosphatase, 69  
fatty acid amide hydrolase, 271  
fatty-acyl-ethyl-ester synthase, 15  
feruloyl esterase, 17  
*N*-feruloylglycine deacylase, 265  
fibrolase, 244  
ficain, 203  
flavastacin, 245  
flavivirin, 193  
fluoroacetyl-CoA thioesterase, 30  
FMN hydrolase, 53  
folate  $\gamma$ -glutamyl hydrolase, 174  
formamidase, 260  
formimidoylaspartate deiminase, 284  
formimidoylglutamase, 284  
formimidoylglutamate deiminase, 285  
formyl-CoA hydrolase, 26  
formylaspartate deformylase, 253  
*N*-formylglutamate deformylase, 264  
*S*-formylglutathione hydrolase, 27  
formylmethionine deformylase, 257  
*N*-formylmethionyl-peptidase, 174  
formyltetrahydrofolate deformylase, 253

fragilysin, 244  
 fructan  $\beta$ -(2,1)-fructosidase, 122  
 fructan  $\beta$ -(2,6)-fructosidase, 123  
 fructan  $\beta$ -fructosidase, 107  
 $\beta$ -fructofuranosidase, 96  
 fructose-2,6-bisphosphate 2-phosphatase, 40  
 fructose-2,6-bisphosphate 6-phosphatase, 42  
 fructose-bisphosphatase, 33  
 fruit bromelain, 207  
 fucoidanase, 99  
 fumarylacetoacetase, 323  
 3-fumarylpyruvate hydrolase, 327  
 fumonisin B1 esterase, 21  
 furin, 190  
 fusarinine-C ornithinesterase, 12  
 futasoline hydrolase, 143  
  
 galactan 1,3- $\beta$ -galactosidase, 121  
 galactan endo- $\beta$ -1,3-galactanase, 130  
 galactan endo-1,6- $\beta$ -galactosidase, 125  
 $\beta$ -galactofuranosidase, 121  
 galactolipase, 8  
 D-galactose 1-phosphate phosphatase, 52  
 L-galactose 1-phosphate phosphatase, 51  
 $\alpha$ -galactosidase, 95  
 $\beta$ -galactosidase, 95  
 galactosylceramidase, 100  
 galactosylgalactosylglucosylceramidase, 100  
 galacturan 1,4- $\alpha$ -galacturonidase, 104  
 gametolysin, 236  
 gastricsin, 218  
 GDP-glucosidase, 99  
 gelatinase A, 233  
 gelatinase B, 235  
 gellan tetrasaccharide unsaturated glucuronosyl hydrolase, 129  
 geranyl diphosphate diphosphatase, 70  
 geranyl diphosphate phosphohydrolase, 315  
 geranylgeranyl diphosphate diphosphatase, 69  
 gingipain K, 210  
 gingipain R, 208  
 ginsenosidase type I, 133  
 ginsenosidase type III, 133  
 ginsenosidase type IV, 134  
 ginsenoside Rb1  $\beta$ -glucosidase, 133  
 Glu-Glu dipeptidase, 160  
 glucan 1,3- $\alpha$ -glucosidase, 108  
 glucan 1,3- $\beta$ -glucosidase, 102  
 glucan 1,4- $\alpha$ -glucosidase, 91  
 glucan 1,4- $\alpha$ -maltohexaosidase, 111  
 glucan 1,4- $\alpha$ -maltohydrolase, 118  
 glucan 1,4- $\alpha$ -maltotetrahydrolase, 103  
 glucan 1,4- $\alpha$ -maltotrihydrolase, 115  
 glucan 1,4- $\beta$ -glucosidase, 106  
 glucan 1,6- $\alpha$ -glucosidase, 105  
 glucan 1,6- $\alpha$ -isomaltosidase, 109  
 glucan endo-1,2- $\beta$ -glucosidase, 105  
  
 glucan endo-1,3- $\alpha$ -glucosidase, 103  
 glucan endo-1,3- $\beta$ -D-glucosidase, 98  
 glucan endo-1,6- $\beta$ -glucosidase, 106  
 gluconolactonase, 6  
 glucosamine-6-phosphate deaminase, 301  
 glucose-1-phosphatase, 33  
 glucose-1-phospho-D-mannosylglycoprotein phosphodiesterase, 62  
 glucose-6-phosphatase, 33  
 $\alpha$ -glucosidase, 95  
 $\beta$ -glucosidase, 95  
 glucosinolate  $\gamma$ -glutamyl hydrolase, 176  
 glucosyl-3-phosphoglycerate phosphatase, 50  
 glucosylceramidase, 99  
 glucosylglycerate hydrolase, 137  
 glucosylglycerol 3-phosphatase, 45  
 glucuronate-2-sulfatase, 67  
 $\alpha$ -glucuronidase, 119  
 $\beta$ -glucuronidase, 96  
 glucuronoarabinoxylan endo-1,4- $\beta$ -xylanase, 119  
 glucuronosyl-disulfoglucosamine glucuronidase, 102  
 glutamate carboxypeptidase, 169  
 glutamate carboxypeptidase II, 171  
 glutamin-(asparagin-)ase, 258  
 D-glutaminase, 258  
 glutaminase, 252  
 glutamyl aminopeptidase, 155  
 glutamyl endopeptidase, 179  
 glutamyl endopeptidase II, 191  
 $\gamma$ -glutamyl hercynylcysteine S-oxide hydrolase, 276  
 $\gamma$ -glutamyl- $\gamma$ -aminobutyrate hydrolase, 270  
 $\gamma$ -glutamylanilide hydrolase, 277  
 glutaryl-7-aminocephalosporanic-acid acylase, 270  
 glutathione hydrolase, 175  
 glutathione thiolesterase, 26  
 glutathionylspermidine amidase, 266  
 Gly-Xaa carboxypeptidase, 168  
 glycerol-1,2-cyclic-phosphate 2-phosphodiesterase, 60  
 glycerol-1-phosphatase, 35  
 glycerol-2-phosphatase, 35  
 glycerophosphocholine cholinephosphodiesterase, 59  
 glycerophosphocholine phosphodiesterase, 55  
 glycerophosphodiester phosphodiesterase, 61  
 glycerophosphoinositol glycerophosphodiesterase, 60  
 glycerophosphoinositol inositolphosphodiesterase, 60  
 [glycogen-synthase-D] phosphatase, 39  
 glycoprotein endo- $\alpha$ -1,2-mannosidase, 118  
 glycosphingolipid deacylase, 264  
 glycosulfatase, 64  
 glycosylceramidase, 103  
 glycosylphosphatidylinositol phospholipase D, 61  
 glycy endopeptidase, 205  
 glycyrrhizin hydrolase, 117  
 gpr endopeptidase, 245  
 granzyme A, 191  
 granzyme B, 191

GTP cyclohydrolase I, 291  
 GTP cyclohydrolase II, 293  
 GTP cyclohydrolase IIa, 294  
 GTP cyclohydrolase IV, 296  
 guanidinoacetase, 283  
 guanidinobutyrase, 284  
 guanidinodeoxy-*scyllo*-inositol-4-phosphatase, 39  
 guanidinopropionase, 286  
 guanine deaminase, 289  
 guanosine deaminase, 291  
 guanosine-3',5'-bis(diphosphate) 3'-diphosphatase, 68  
 guanosine-5'-diphospho-5'-[DNA] diphosphatase, 73  
 guanosine-5'-triphosphate,3'-diphosphate phosphatase, 309  
 guanosine-diphosphatase, 310  
  
 haloacetate dehalogenase, 329  
 2-haloacid dehalogenase (configuration-inverting), 330  
 2-haloacid dehalogenase (configuration-retaining), 330  
 haloalkane dehalogenase, 329  
 helper-component proteinase, 210  
 hepacivirin, 195  
 heparanase, 126  
 hepoxilin-epoxide hydrolase, 147  
 hepsin, 197  
 hesperidin 6-*O*- $\alpha$ -L-rhamnosyl- $\beta$ -D-glucosidase, 127  
 heterotrimeric G-protein GTPase, 322  
 hippurate hydrolase, 257  
 histidinol-phosphatase, 34  
 histolysain, 207  
 histone deacetylase, 271  
 HIV-1 retropepsin, 220  
 HIV-2 retropepsin, 227  
 hormone-sensitive lipase, 18  
 horrilysin, 238  
 HslU—HslV peptidase, 248  
 HtrA2 peptidase, 198  
 human endogenous retrovirus K endopeptidase, 228  
 hyaluronoglucosaminidase, 97  
 hyaluronoglucuronidase, 98  
 HycI peptidase, 228  
 2-hydroxy-3-keto-5-methylthiopentenyl-1-phosphate phosphatase, 50  
 2-hydroxy-6-oxo-6-(2-aminophenyl)hexa-2,4-dienoate hydrolase, 326  
 2-hydroxy-6-oxonona-2,4-dienedioate hydrolase, 326  
 4-hydroxy-7-methoxy-3-oxo-3,4-dihydro-2*H*-1,4-benzoxazin-2-yl glucoside  $\beta$ -D-glucosidase, 130  
*N*<sup>ω</sup>-hydroxy-L-arginine amidinohydrolase, 288  
 2-hydroxy-dATP diphosphatase, 312  
 hydroxyacylglutathione hydrolase, 26  
 4-hydroxybenzoyl-CoA thioesterase, 29  
 2'-hydroxybiphenyl-2-sulfinate desulfinate, 334  
 hydroxybutyrate-dimer hydrolase, 7  
 hydroxydechloroatrazine ethylaminohydrolase, 297  
 3-hydroxyisobutyryl-CoA hydrolase, 25  
 hydroxyisourate hydrolase, 282  
 2-(hydroxymethyl)-3-(acetamidomethylene)succinate hydrolase, 264  
 hydroxymethylglutaryl-CoA hydrolase, 26  
 [hydroxymethylglutaryl-CoA reductase (NADPH)]-phosphatase, 40  
 2-hydroxymuconate-6-semialdehyde hydrolase, 325  
 hypodermin C, 184  
  
 iduronate-2-sulfatase, 66  
 L-iduronidase, 106  
 IgA-specific metalloendopeptidase, 230  
 IgA-specific serine endopeptidase, 189  
 imidazolonepropionase, 280  
 2-iminobutanoate/2-iminopropanoate deaminase, 302  
 IMP cyclohydrolase, 290  
 IMP-specific 5'-nucleotidase, 53  
 infectious pancreatic necrosis birnavirus Vp4 peptidase, 200  
 inorganic diphosphatase, 302  
 inosinate nucleosidase, 140  
 inosine diphosphate phosphatase, 314  
 inosine nucleosidase, 138  
 inositol-1,4-bisphosphate 1-phosphatase, 43  
 inositol-phosphate phosphatase, 36  
 inositol-polyphosphate 5-phosphatase, 43  
 insulysin, 240  
 intermediate cleaving peptidase 55, 158  
 interstitial collagenase, 229  
 inulinase, 92  
 isatin hydrolase, 283  
 isoamylase, 104  
 isochorismatase, 146  
 isomaltose glucohydrolase, 137  
*N*-isopropylammelide isopropylaminohydrolase, 297  
 isopullulanase, 102  
 isotuberculosinol synthase, 69  
  
 jararhagin, 244  
 jasmonoyl-L-amino acid hydrolase, 278  
 juvenile-hormone esterase, 14  
  
 kallikrein 13, 201  
 kallikrein 8, 201  
 kanosamine-6-phosphate phosphatase, 51  
 keratan-sulfate endo-1,4- $\beta$ -galactosidase, 112  
 kexin, 186  
 kynureninase, 324  
  
 (13*E*)-labda-7,13-dien-15-ol synthase, 70  
 $\beta$ -lactamase, 280  
 lactase, 113  
 D-lactate-2-sulfatase, 67  
 lacto-*N*-biosidase, 120  
 lactocepin, 195  
 1,4-lactonase, 7  
 legumain, 207  
 leishmanolysin, 235  
 leucolysin, 229

leucyl aminopeptidase, 153  
 leucyl endopeptidase, 185  
 leukocyte elastase, 182  
 leukotriene-A<sub>4</sub> hydrolase, 147  
 leukotriene-C<sub>4</sub> hydrolase, 175  
 levanase, 104  
 licheninase, 105  
 limit dextrin  $\alpha$ -1,6-maltotetraose-hydrolase, 134  
 limit dextrinase, 120  
 limonene-1,2-epoxide hydrolase, 147  
 limonin-D-ring-lactonase, 9  
 limulus clotting enzyme, 192  
 limulus clotting factor  $\bar{B}$ , 192  
 limulus clotting factor  $\bar{C}$ , 192  
 lipid-phosphate phosphatase, 47  
 lipoprotein lipase, 9  
*N*-(long-chain-acyl)ethanolamine deacylase, 263  
 long-chain-fatty-acyl-glutamate deacylase, 262  
 Lys-Lys/Arg-Xaa endopeptidase, 202  
 lysine carboxypeptidase, 168  
 L-lysine-lactamase, 281  
 lysophospholipase, 4  
 lysoplasmalogenase, 146  
 lysosomal Pro-Xaa carboxypeptidase, 166  
 lysostaphin, 245  
 lysozyme, 94  
 lysyl endopeptidase, 184  
 $\alpha$ -lytic endopeptidase, 178  
 $\beta$ -lytic metalloendopeptidase, 235  
  
 macrophage elastase, 242  
 magnolysin, 241  
 maleamate amidohydrolase, 273  
 maleimide hydrolase, 282  
 maleylpyruvate hydrolase, 328  
*N*-malonylurea hydrolase, 270  
 maltose 6'-phosphate phosphatase, 51  
 maltose-6'-phosphate glucosidase, 116  
 (3*S*)-malyl-CoA thioesterase, 30  
 mandelamide amidase, 268  
 mannan 1,2-(1,3)- $\alpha$ -mannosidase, 106  
 mannan 1,4-mannobiosidase, 111  
 $\alpha$ -mannan endo-1,2- $\alpha$ -mannanase, 135  
 mannan endo-1,4- $\beta$ -mannosidase, 106  
 mannan endo-1,6- $\alpha$ -mannosidase, 111  
 mannan exo-1,2-1,6- $\alpha$ -mannosidase, 119  
 mannan-binding lectin-associated serine protease-2, 197  
 mannitol-1-phosphatase, 35  
 $\alpha$ -mannosidase, 95  
 $\beta$ -mannosidase, 96  
 mannosyl-3-phosphoglycerate phosphatase, 45  
 mannosyl-glycoprotein endo- $\beta$ -*N*-acetylglucosaminidase, 110  
 mannosyl-oligosaccharide  $\alpha$ -1,3-glucosidase, 137  
 mannosyl-oligosaccharide 1,2- $\alpha$ -mannosidase, 114  
 mannosyl-oligosaccharide 1,3-1,6- $\alpha$ -mannosidase, 114  
 mannosyl-oligosaccharide glucosidase, 112  
  
 mannosylfructose-phosphate phosphatase, 48  
 mannosylglycerate hydrolase, 127  
 mannosylglycoprotein endo- $\beta$ -mannosidase, 122  
 matrilysin, 232  
 matriptase, 198  
 melamine deaminase, 298  
 memapsin 1, 226  
 memapsin 2, 227  
 membrane alanyl aminopeptidase, 153  
 membrane dipeptidase, 161  
 membrane Pro-Xaa carboxypeptidase, 170  
 membrane-type matrix metalloproteinase-1, 246  
 meprin A, 231  
 meprin B, 242  
 Met-Xaa dipeptidase, 160  
 metallocarboxypeptidase D, 172  
 methenyltetrahydrofolate cyclohydrolase, 290  
 methenyltetrahydromethanopterin cyclohydrolase, 294  
 methionyl aminopeptidase, 156  
*N*-methyl nucleosidase, 143  
 [3-methyl-2-oxobutanoate dehydrogenase (2-methylpropanoyl-transferring)]-phosphatase, 42  
*N*-methyl-2-oxoglutaramate hydrolase, 258  
*S*-methyl-5'-thioadenosine deaminase, 295  
 1-methyladenosine nucleosidase, 140  
 methylated diphthine methylhydrolase, 23  
 methylenediurea deaminase, 287  
 4-methyleneglutaminase, 264  
 methylguanidinase, 286  
 7-methylguanosine nucleotidase, 51  
 (*S*)-methylmalonyl-CoA hydrolase, 28  
 4-methyloxaloacetate esterase, 11  
 methylphosphothioglycerate phosphatase, 33  
 methylthioadenosine nucleosidase, 141  
 methylumbelliferyl-acetate deacetylase, 13  
 metridin, 177  
 microbial collagenase, 229  
 micrococcal nuclease, 90  
 microsomal epoxide hydrolase, 147  
 mimosinase, 263  
 mitochondrial intermediate peptidase, 241  
 mitochondrial processing peptidase, 242  
 Mn<sup>2+</sup>-dependent ADP-ribose/CDP-alcohol diphosphatase, 311  
 mono(ethylene terephthalate) hydrolase, 24  
 monomethyl-sulfatase, 67  
 monosaccharide-transporting ATPase, 317  
 monoterpene  $\epsilon$ -lactone hydrolase, 20  
 monoterpényl-diphosphatase, 68  
 mRNA(cytosine<sup>666</sup>) deaminase, 296  
 mucorpepsin, 221  
 mucrolysin, 240  
 multiple inositol-polyphosphate phosphatase, 44  
 muramoylpentapeptide carboxypeptidase, 169  
 muramoyltetrapeptide carboxypeptidase, 170  
 mycodextranase, 103  
 mycolysin, 234



mycophenolic acid acyl-glucuronide esterase, 22  
 mycothiol *S*-conjugate amidase, 275  
 myeloblastin, 190  
 [myosin-light-chain] phosphatase, 42  
  
 N-formylmaleamate deformylase, 273  
 N-methylhydantoinase (ATP-hydrolysing), 281  
 NAD<sup>+</sup> diphosphatase, 307  
 NAD<sup>+</sup> glycohydrolase, 138  
 nardilysin, 241  
 $\alpha$ -neoagaro-oligosaccharide hydrolase, 124  
 neopullulanase, 119  
 nepenthesin, 219  
 neprilysin, 230  
 neurolysin, 231  
 neutrophil collagenase, 235  
 nicotinamidase, 255  
 nicotinamide-nucleotide amidase, 259  
 nitrilase, 298  
 5-nitroanthranilic acid aminohydrolase, 301  
 2-nitroimidazole nitrohydrolase, 301  
 4-nitrophenylphosphatase, 39  
 NMN nucleosidase, 140  
 nodavirus endopeptidase, 226  
 non-chaperonin molecular chaperone ATPase, 321  
 non-reducing end  $\alpha$ -L-arabinofuranosidase, 102  
 non-reducing end  $\beta$ -L-arabinofuranosidase, 131  
 non-reducing end  $\beta$ -L-arabinopyranosidase, 108  
 non-stereospecific dipeptidase, 161  
 nuclear-inclusion-a endopeptidase, 209  
 3',5'-nucleoside bisphosphate phosphatase, 52  
 nucleoside diphosphate phosphatase, 303  
 nucleoside phosphoacylhydrolase, 307  
 nucleoside-triphosphate phosphatase, 305  
 3'-nucleotidase, 32  
 5'-nucleotidase, 32  
 nucleotidase, 37  
 nucleotide diphosphatase, 304  
  
 oleoyl-[acyl-carrier-protein] hydrolase, 27  
 oleuropein  $\beta$ -glucosidase, 137  
 oligo-1,6-glucosidase, 93  
 oligonucleotidase, 74  
 oligopeptidase A, 243  
 oligopeptidase B, 192  
 oligosaccharide reducing-end xylanase, 123  
 oligosaccharide-diphosphodolichol diphosphatase, 310  
 oligoxyloglucan  $\beta$ -glycosidase, 115  
 oligoxyloglucan reducing-end-specific cellobiohydrolase, 122  
 omptin, 227  
 ophiolysin, 239  
*pro*-opiomelanocortin converting enzyme, 220  
 orsellinate-depside hydrolase, 10  
 oryzin, 187  
 oviductin, 202  
 oxaloacetase, 323  
  
 oxamate amidohydrolase, 278  
 oxepin-CoA hydrolase, 149  
 2-oxo-3-(5-oxofuran-2-ylidene)propanoate lactonase, 22  
 8-oxo-dGDP phosphatase, 312  
 8-oxo-dGTP diphosphatase, 312  
 3-oxoadipate enol-lactonase, 7  
 6-oxocamphor hydrolase, 327  
 6-oxocyclohex-1-ene-1-carbonyl-CoA hydratase, 327  
 2-oxoglutaramate amidase, 274  
 8-oxoguanine deaminase, 295  
 5-oxoprolinase (ATP-hydrolysing), 281  
  
 palmitoyl-CoA hydrolase, 25  
 palmitoyl[protein] hydrolase, 29  
 pancreatic elastase, 181  
 pancreatic elastase II, 189  
 pancreatic endopeptidase E, 189  
 pantetheine hydrolase, 270  
 pantothenase, 255  
 papain, 202  
 pappalysin-1, 245  
 pectinesterase, 5  
 penicillin amidase, 253  
 penicillopepsin, 221  
 pentanamidase, 260  
 PepB aminopeptidase, 157  
 pepsin A, 218  
 pepsin B, 218  
 L-peptidase, 210  
 peptidase 1 (mite), 216  
 peptidase Do, 198  
 peptidase K, 187  
 peptide deformylase, 268  
 peptide-*N*<sup>4</sup>-(*N*-acetyl- $\beta$ -glucosaminyl)asparagine amidase, 261  
 peptidoglycan  $\beta$ -*N*-acetylmuramidase, 109  
 peptidoglycan-*N*-acetylglucosamine deacetylase, 273  
 $\beta$ -peptidyl aminopeptidase, 158  
 peptidyl-Asp metalloendopeptidase, 235  
 peptidyl-dipeptidase A, 165  
 peptidyl-dipeptidase B, 165  
 peptidyl-dipeptidase Dcp, 166  
 peptidyl-glutaminase, 259  
 peptidyl-glycinamidase, 173  
 peptidyl-Lys metalloendopeptidase, 232  
 peroxisome-assembly ATPase, 320  
 peroxyureidoacrylate/ureidoacrylate amidohydrolase, 274  
 pestivirus NS3 polyprotein peptidase, 200  
 phenylacetyl-CoA hydrolase, 29  
 pheophorbide, 19  
 phloretin hydrolase, 324  
 phorbol-diester hydrolase, 12  
 phosphatidate phosphatase, 31  
 phosphatidylglycerophosphatase, 36  
 phosphatidylinositol deacylase, 12  
 phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase, 45  
 phosphatidylinositol-3,4,5-trisphosphate 5-phosphatase, 50

phosphatidylinositol-3,4-bisphosphate 4-phosphatase, 45  
 phosphatidylinositol-3,5-bisphosphate 3-phosphatase, 52  
 phosphatidylinositol-3-phosphatase, 44  
 phosphatidylinositol-4,5-bisphosphate 4-phosphatase, 48  
 6-phospho- $\beta$ -galactosidase, 108  
 6-phospho- $\beta$ -glucosidase, 108  
 5-phospho-D-xylono-1,4-lactonase, 25  
 phospho*enol*pyruvate phosphatase, 44  
 phosphoadenylylsulfatase, 315  
 phosphoamidase, 331  
 phosphodiesterase I, 54  
 phosphoethanolamine/phosphocholine phosphatase, 47  
 6-phosphogluconolactonase, 9  
 3-phosphoglycerate phosphatase, 39  
 phosphoglycerate phosphatase, 35  
 phosphoglycolate phosphatase, 34  
 phosphohistidine phosphatase, 331  
 phosphoinositide 5-phosphatase, 38  
 phosphoinositide phospholipase C, 56  
 phospholipase A<sub>1</sub>, 9  
 phospholipase A<sub>2</sub>, 4  
 phospholipase C, 55  
 phospholipase D, 55  
 phosphonoacetaldehyde hydrolase, 332  
 phosphonoacetate hydrolase, 332  
 phosphonopyruvate hydrolase, 333  
 5''-phosphoribostamycin phosphatase, 50  
 phosphoribosyl 1,2-cyclic phosphate 1,2-diphosphodiesterase, 63  
 phosphoribosyl 1,2-cyclic phosphate phosphodiesterase, 63  
 phosphoribosyl-AMP cyclohydrolase, 292  
 phosphoribosyl-ATP diphosphatase, 308  
 [phosphorylase] phosphatase, 34  
 phosphoserine phosphatase, 31  
 2-phosphosulfolactate phosphatase, 46  
 $\alpha,\alpha$ -phosphotrehalase, 109  
 phthalyl amidase, 266  
 physarolisin, 196  
 3-phytase, 32  
 4-phytase, 36  
 5-phytase, 46  
 phytepsin, 225  
 picornain 2A, 206  
 picornain 3C, 206  
 pimelyl-[acyl-carrier protein] methyl ester esterase, 20  
 pitrilysin, 240  
 plasma kallikrein, 181  
 plasmepsin I, 225  
 plasmepsin II, 225  
 plasmin, 178  
*t*-plasminogen activator, 188  
 plasminogen activator Pla, 227  
 poly(3-hydroxybutyrate) depolymerase, 17  
 poly(3-hydroxyoctanoate) depolymerase, 17  
 poly(A)-specific ribonuclease, 74  
 poly(ADP-ribose) glycohydrolase, 120  
 poly(ethylene terephthalate) hydrolase, 24  
 polygalacturonase, 94  
 polymannuronate hydrolase, 116  
 polynucleotide-aldehyde esterase, 18  
 polynucleotide 3'-phosphatase, 37  
 polynucleotide 5'-phosphatase, 37  
 polyporopepsin, 223  
 $\beta$ -porphyranase, 129  
 preflagellin peptidase, 228  
 prenyl-diphosphatase, 68  
 prepilin peptidase, 226  
 $\beta$ -primeverosidase, 121  
 Pro-Pro endopeptidase, 247  
 proclavamate amidohydrolase, 287  
 procollagen *N*-endopeptidase, 231  
 procollagen C-endopeptidase, 232  
 L-proline amide hydrolase, 272  
 prolyl aminopeptidase, 154  
 prolyl oligopeptidase, 180  
 proprotein convertase 1, 194  
 proprotein convertase 2, 194  
 proteasome endopeptidase complex, 248  
 protein *O*-GlcNAcase, 127  
 [protein ADP-ribosylarginine] hydrolase, 141  
 protein arginine phosphatase, 331  
 protein C (activated), 188  
 protein deglycase, 278  
 protein N-terminal asparagine amidohydrolase, 277  
 protein N-terminal glutamine amidohydrolase, 277  
 protein phosphatase methylesterase-1, 21  
 protein-arginine deiminase, 286  
 protein-glucosylgalactosylhydroxylysine glucosidase, 113  
 protein-glutamate methylesterase, 14  
 protein-glutamine glutaminase, 259  
 protein-serine/threonine phosphatase, 34  
 protein-synthesizing GTPase, 322  
 protein-tyrosine-phosphatase, 41  
 protodioscin 26-*O*- $\beta$ -D-glucosidase, 131  
 prunasin  $\beta$ -glucosidase, 115  
 pseudolysin, 233  
 pseudouridine 5'-phosphatase, 52  
 pterin deaminase, 291  
 pullulanase, 99  
 Pup amidohydrolase, 276  
 purine nucleosidase, 138  
 pycnoporopepsin, 223  
 pyrethroid hydrolase, 21  
 pyridoxal phosphatase, 46  
 4-pyridoxolactonase, 8  
 pyrimidine-5'-nucleotide nucleosidase, 140  
 pyriothiamine deaminase, 292  
 pyroglutamyl-peptidase I, 173  
 pyroglutamyl-peptidase II, 173  
 2-pyrone-4,6-dicarboxylate lactonase, 13  
 1-pyrroline-4-hydroxy-2-carboxylate deaminase, 293  
 [pyruvate dehydrogenase (acetyl-transferring)]-phosphatase, 40

[pyruvate kinase]-phosphatase, 41

quercitrinase, 104

quorum-quenching *N*-acyl-homoserine lactonase, 19

raucaffricine  $\beta$ -glucosidase, 116

$\gamma$ -renin, 185

renin, 219

repressor LexA, 193

retinoid isomerohydrolase, 15

retroviral ribonuclease H, 88

rhamnogalacturonan acetyltransferase, 21

rhamnogalacturonan galacturonohydrolase, 128

rhamnogalacturonan hydrolase, 127

rhamnogalacturonan rhamnohydrolase, 128

L-rhamnono-1,4-lactonase, 15

rhizopuspepsin, 221

rhodotorulapepsin, 222

rhomboid protease, 197

riboflavinase, 300

ribonuclease  $\alpha$ , 86

ribonuclease D, 74

ribonuclease E, 88

ribonuclease F, 89

ribonuclease H, 86

ribonuclease III, 86

ribonuclease IV, 87

ribonuclease IX, 88

ribonuclease M5, 87

ribonuclease P, 87

ribonuclease P4, 87

ribonuclease [poly-(U)-specific], 87

ribonuclease T<sub>1</sub>, 89

ribonuclease V, 89

ribosylpyrimidine nucleosidase, 139

ricinine nitrilase, 298

RNA 2',3'-cyclic 3'-phosphodiesterase, 63

RNA helicase, 321

rRNA *N*-glycosylase, 142

rRNA endonuclease, 90

ruberlysin, 238

russellysin, 240

S2P endopeptidase, 247

saccharolysin, 236

saccharopepsin, 222

SARS coronavirus main proteinase, 217

scutellarin, 186

scytalidopepsin A, 223

scytalidopepsin B, 224

sedoheptulose-bisphosphatase, 38

sedolisin, 196

semenogelase, 190

separase, 211

sepiapterin deaminase, 293

serine-ethanolaminephosphate phosphodiesterase, 56

serine-type D-Ala-D-Ala carboxypeptidase, 166

serralysin, 236

sialate *O*-acetyltransferase, 13

*O*-sialoglycoprotein endopeptidase, 240

signal peptidase I, 193

signal peptidase II, 224

signal-recognition-particle GTPase, 323

sinapine esterase, 12

single-stranded DNA cytosine deaminase, 296

site-1 protease, 199

small monomeric GTPase, 322

snake venom factor V activator, 194

snopalysin, 245

soluble epoxide hydrolase, 148

sorbitol-6-phosphatase, 41

sortase A, 217

sortase B, 217

(*R*)-specific secondary-alkylsulfatase, 67

spermosin, 195

sphingomyelin deacylase, 274

sphingomyelin phosphodiesterase, 56

sphingomyelin phosphodiesterase D, 59

spleen exonuclease, 75

SpoIVB peptidase, 200

staphopain, 210

Ste24 endopeptidase, 246

stem bromelain, 206

D-stereospecific aminopeptidase, 157

steroid-lactonase, 10

sterol esterase, 5

steryl- $\beta$ -glucosidase, 112

steryl-sulfatase, 64

stratum corneum chymotryptic enzyme, 201

streptogrisin A, 191

streptogrisin B, 191

streptomycin-6-phosphatase, 39

streptopain, 203

streptothricin hydrolase, 283

3 $\alpha$ (*S*)-strictosidine  $\beta$ -glucosidase, 112

stromelysin 1, 231

stromelysin 2, 232

*N*-substituted formamide deformylase, 269

subtilisin, 187

succinyl-CoA hydrolase, 25

succinyl-diaminopimelate desuccinylase, 255

*N*-succinylarginine dihydrolase, 287

succinylglutamate desuccinylase, 270

*S*-succinylglutathione hydrolase, 27

sucrose  $\alpha$ -glucosidase, 100

sucrose-phosphate phosphatase, 35

sugar-phosphatase, 35

sugar-terminal-phosphatase, 43

3-sulfinoopropanoyl-CoA desulfinase, 334

*N*-sulfoglucosamine sulfohydrolase, 332

*N*-sulfoglucosamine-3-sulfatase, 67

4-sulfomuconolactone hydrolase, 22

sulfoquinovosidase, 135



*S*-sulfosulfanyl-L-cysteine sulfohydrolase, 68  
 T<sub>4</sub> deoxyribonuclease II, 77  
 T<sub>4</sub> deoxyribonuclease IV, 77  
 tannase, 7  
 teichoic acid D-alanine hydrolase, 24  
 tentoxilysin, 243  
 theanine hydrolase, 264  
 thermitase, 188  
 thermolysin, 233  
 thermomycolin, 188  
 thermopsin, 226  
 thiamine phosphate phosphatase, 53  
 thiamine-triphosphatase, 308  
 thimet oligopeptidase, 231  
 thiocyanate hydrolase, 300  
 thioglucosidase, 121  
 thrombin, 177  
 thymidine-triphosphatase, 309  
 thymidylate 5'-phosphatase, 38  
 thymine-DNA glycosylase, 144  
 tissue kallikrein, 181  
 togavirin, 193  
 α,α-trehalase, 96  
 trehalose-phosphatase, 33  
 triacetate-lactonase, 10  
 triacylglycerol lipase, 3  
 trimerelysin I, 239  
 trimerelysin II, 239  
 trimetaphosphatase, 303  
 tripeptide aminopeptidase, 154  
 tripeptidyl-peptidase I, 164  
 tripeptidyl-peptidase II, 164  
 triphosphatase, 307  
 trithionate hydrolase, 333  
 tRNA<sup>Ala</sup>(adenine<sup>37</sup>) deaminase, 295  
 tRNA(adenine<sup>34</sup>) deaminase, 295  
 tRNA(cytosine<sup>8</sup>) deaminase, 295  
 tRNase Z, 88  
 tropinesterase, 5  
 trypsin, 177  
 tryptase, 186  
 tryptophanamidase, 262  
 tryptophanyl aminopeptidase, 156  
 tuberculosinol synthase, 69  
 tubulin GTPase, 323  
 tubuliny-Tyr carboxypeptidase, 170  
 type I site-specific deoxyribonuclease, 76  
 type II site-specific deoxyribonuclease, 76  
 type III site-specific deoxyribonuclease, 76  
 u-plasminogen activator, 189  
 ubiquitinyl hydrolase 1, 175  
 UDP-2,3-diacetylglucosamine diphosphatase, 311  
 UDP-2,4-diacetamido-2,4,6-trideoxy-β-L-altropyranose hydro-  
 lase, 312  
 UDP-3-*O*-acyl-*N*-acetylglucosamine deacetylase, 274  
 UDP-*N,N'*-diacetylbacillosamine 2-epimerase (hydrolysing), 131  
 UDP-*N*-acetylglucosamine 2-epimerase (hydrolysing), 131  
 UDP-sugar diphosphatase, 310  
 UDP-sulfoquinovose synthase, 333  
 Ulp1 peptidase, 217  
 undecaprenyl-diphosphate phosphatase, 308  
 unsaturated chondroitin disaccharide hydrolase, 130  
 unsaturated rhamnogalacturonyl hydrolase, 128  
 uracil-DNA glycosylase, 143  
 urease, 252  
 (*S*)-ureidoglycine aminohydrolase, 288  
 ureidoglycolate amidohydrolase, 276  
 β-ureidopropionase, 252  
 ureidosuccinase, 252  
 urethanase, 266  
 uridine nucleosidase, 138  
 uronolactonase, 6  
 V-cath endopeptidase, 211  
 validoxylamine A 7'-phosphate phosphatase, 53  
 venom exonuclease, 75  
 venombin A, 190  
 venombin AB, 185  
 versiconal hemiacetal acetate esterase, 22  
 vesicle-fusing ATPase, 320  
 vibriolysin, 233  
 vicianin β-glucosidase, 115  
 wax-ester hydrolase, 12  
 [Wnt protein] *O*-palmitoleoyl-L-serine hydrolase, 23  
 Xaa-Arg dipeptidase, 159  
 Xaa-methyl-His dipeptidase, 160  
 Xaa-Pro aminopeptidase, 155  
 Xaa-Pro dipeptidase, 160  
 Xaa-Pro dipeptidyl-peptidase, 164  
 Xaa-Trp aminopeptidase, 156  
 Xaa-Xaa-Pro tripeptidyl-peptidase, 164  
 xanthomonalisin, 196  
 XTP/dITP diphosphatase, 315  
 xylan α-1,2-glucuronosidase, 118  
 xylan 1,3-β-xylosidase, 105  
 xylan 1,4-β-xylosidase, 98  
 xyloglucan-specific endo-β-1,4-glucanase, 122  
 xyloglucan-specific exo-β-1,4-glucanase, 123  
 xylono-1,4-lactonase, 16  
 yapsin 1, 225  
 yeast ribonuclease, 75  
 zinc D-Ala-D-Ala carboxypeptidase, 170  
 zingipain, 216